

< 根拠文献 >

- 1 ) Kotoh K, Nakamuta M, Morizono S, Kohjima M, Arimura E, Fukushima M, Enjoji M, Sakai H and Nawata H: A multi-step, incremental expansion method for radio frequency ablation: optimization of the procedure to prevent increases in intra-tumor pressure and to reduce the ablation time, *Liver International*; 2005; 25; 542-547
- 2 ) Tito Livraghi, Luigi Solbiati, M.Franca Meloni, G. Scott Gazelle, Elkan F.Halpern and S. Nahum Goldberg: Treatment of Focal Liver Tumors with Percutaneous Radio-frequency Ablation: Complications Encountered in a Multicenter Study; *Radiology*; 2003; 226; 2; 441-451
- 3 ) 今村 也寸志, 小原 一憲, 柴藤 俊彦, 馬場 芳郎, 田原 憲治, 窪園 修: ラジオ波焼灼療法後に急速に悪化した肝細胞癌の 2 症例, *日本消化器病学会雑誌*, 2002; 99; 40-44
- 4 ) Josep M. Llovet, Ramon Vilana, Concepció Brú, Lluís Bianchi, Joan Manuel Salmeron, Loreto Boix, Sergi Ganau, Margarita Sala, Mario Pagès, Carmen Ayuso, Manel Solè, Joan Rodés and Jordi Bruix: Increased Risk of Tumor Seeding After Percutaneous Radiofrequency Ablation for Single Hepatocellular Carcinoma: *Hepatology*; 2001; 33; 1124-1129
- 5 ) 中井 資貴, 白木 達也, 東 克彦, 前田 雅子, 佐原 伸也, 竹内 希, 木村 誠志, 寺田 正樹, 佐藤 守男: 肝細胞癌に対する TACE 併用低出力ラジオ波凝固療法, *日本医学放射線学会雑誌*, 2005; 65; 124-125
- 6 ) Toshihiko Kawasaki, Masatoshi Kubo, Hobyung Chung and Yasunori Minami: Hepatocellular carcinoma that ruptured during radiofrequency ablation therapy: *Journal of Gastroenterology*; 2004; 39; 1015-1016