



This English version is intended to be a reference material to provide convenience for users. In the event of inconsistency between the Japanese original and this English translation, the former shall prevail.

Revision of Precautions Candesartan Cilexetil/Amlodipine Besilate

January 12, 2016

Non-proprietary name

Candesartan Cilexetil/Amlodipine Besilate

Safety measure

Precautions should be revised in the package insert.

In the Clinically significant adverse reaction subsection regarding hepatic function disorder and jaundice of the Adverse Reaction section, the following text should be revised (underlined parts are revised):

Fulminant hepatitis, hepatic function disorder, and jaundice:

Fulminant hepatitis, hepatic function disorder associated with increased levels of AST (GOT), ALT (GPT), γ -GTP, etc. or jaundice may occur. Patients should be carefully monitored. If any abnormalities are observed, administration of this drug should be discontinued and appropriate measures should be adopted.

In the Clinically significant adverse reaction subsection regarding rhabdomyolysis of the Adverse Reaction section, the following text should be revised (underlined parts are revised):

Rhabdomyolysis:

Rhabdomyolysis characterized by myalgia, feelings of weakness, increased creatinine kinase (creatine phosphokinase), or increased blood and urine myoglobin may occur. Patients should be carefully monitored. If these symptoms are observed, administration of this drug should be discontinued immediately and appropriate measures should be adopted.

In addition, caution should be exercised for development of acute kidney injury due to rhabdomyolysis.

Pharmaceuticals and Medical Devices Agency

Office of Safety I

3-3-2 Kasumigaseki, Chiyoda-ku, Tokyo 100-0013 Japan

E-mail: safety.info@pmda.go.jp