

Published by
Ministry of Health, Labour and Welfare



Translated by
Pharmaceuticals and Medical Devices Agency



This English version is intended to be a reference material to provide convenience for users. In the event of inconsistency between the Japanese original and this English translation, the former shall prevail.

Revision of Precautions

Talipexole hydrochloride

August 22, 2019

Therapeutic category

Antiparkinsonian agents

Non-proprietary name

Talipexole hydrochloride

Safety measure

Precautions should be revised in the package insert.

Pharmaceuticals and Medical Devices Agency

3-3-2 Kasumigaseki, Chiyoda-ku, Tokyo 100-0013 Japan
E-mail: safety.info@pmda.go.jp

Revision in line with the Instructions for Package Inserts of Prescription Drugs, PAB Notification No. 606 by the Director General of Pharmaceutical Affairs Bureau, MHW, dated April 25, 1997 (Old instructions): Revised language is underlined.

Current	Revision
<p>Important Precautions</p> <p>When dose reduction or discontinuation of this drug is necessary, the dose should be gradually reduced.</p>	<p>Important Precautions</p> <p>When dose reduction or discontinuation of this drug is necessary, the dose should be gradually reduced. <u>Rapid dose reduction or discontinuation may cause syndrome malin with symptoms such as pyrexia, disturbed consciousness, akinetic mutism, severe muscle stiffness, involuntary movement, dysphagia, tachycardia, blood pressure fluctuation, sweating, increased serum CK (CPK). In addition, rapid dose reduction or discontinuation of dopamine receptor agonists may cause drug withdrawal syndrome (characterized by apathy, anxiety, depression, fatigue, sweating, pain, etc.).</u></p>
<p>Adverse Reactions</p> <p>Other Adverse Reactions</p> <p>(N/A)</p>	<p>Adverse Reactions</p> <p>Other Adverse Reactions</p> <p><u>Drug withdrawal syndrome* (apathy, anxiety, depression, fatigue, sweating, pain, etc.)</u></p> <p><u>*When any abnormalities are observed, appropriate measures should be taken such as resuming administration or returning the dose to the level prior to reduction.</u></p>

N/A: Not Applicable, because the section is not included in the current package insert.

Pharmaceuticals and Medical Devices Agency

3-3-2 Kasumigaseki, Chiyoda-ku, Tokyo 100-0013 Japan
E-mail: safety.info@pmda.go.jp