Published by Ministry of Health, Labour and Welfare Translated by Pharmaceuticals and Medical Devices Agency





This English version is intended to be a reference material to provide convenience for users. In the event of inconsistency between the Japanese original and this English translation, the former shall prevail.

Revision of Precautions Ofloxacin (oral dosage form)

September 24, 2019

Therapeutic category

Synthetic antibiotics

Non-proprietary name

Ofloxacin

Safety measure

Precautions should be revised in the package insert.

Pharmaceuticals and Medical Devices Agency

Revision in line with the Instructions for Package Inserts of Prescription Drugs, PAB Notification No. 606 by the Director General of Pharmaceutical Affairs Bureau, MHW, dated April 25, 1997 (Old instructions):

Revised language is underlined.

Current	Revision
Adverse Reactions	Adverse Reactions
Clinically Significant Adverse Reactions	Clinically Significant Adverse Reactions
Tendon disorders such as Achilles tendonitis and tendon rupture:	Tendon disorders such as Achilles tendonitis and tendon rupture
Symptoms: Pain, edema around the tendon	Tendon disorders such as Achilles tendonitis and tendon rupture
These tendon disorders are more likely to occur in patients aged 60	may occur. If symptoms such as pain, edema, and redness around
years or older, patients who concomitantly use corticosteroids, and	the tendon are observed, administration of this drug should be
patients with a history of organ transplant.	discontinued and appropriate measures should be taken. These
	tendon disorders are more likely to occur in patients with a history
	of organ transplant.
(N/A)	Peripheral neuropathy
	Peripheral neuropathy may occur. If symptoms such as numbness,
	muscle weakness, and pain are observed, administration of this
	drug should be discontinued and appropriate measures should be
	taken.

N/A: Not Applicable, because the section is not included in the current package insert.