Published by Ministry of Health, Labour and Welfare Translated by Pharmaceuticals and Medical Devices Agency





This English version is intended to be a reference material to provide convenience for users. In the event of inconsistency between the Japanese original and this English translation, the former shall prevail.

Revision of Precautions Moxifloxacin hydrochloride (oral dosage form)

September 24, 2019

Therapeutic category

Synthetic antibiotics

Non-proprietary name

Moxifloxacin hydrochloride

Safety measure

Precautions should be revised in the package insert.

Revision in line with the Instructions for Package Inserts of Prescription Drugs, PAB Notification No. 606 by the Director General of Pharmaceutical Affairs Bureau, MHW, dated April 25, 1997 (Old instructions):

Revised language is underlined.

Current	Revision
Adverse Reactions	Adverse Reactions
Clinically Significant Adverse Reactions	Clinically Significant Adverse Reactions
Tendon disorders such as tendonitis and tendon rupture:	Tendon disorders such as Achilles tendonitis and tendon rupture:
Tendon disorders such as tendonitis and tendon rupture may occur.	Tendon disorders such as Achilles tendonitis and tendon rupture
If pain or inflammation in the tendon is observed, administration of	may occur. If symptoms such as pain, edema, and redness around
this drug should be discontinued and appropriate measures should	the tendon are observed, administration of this drug should be
be taken. Cases of these symptoms that developed several months	discontinued and appropriate measures should be taken. Cases of
after the termination of this drug have been reported overseas.	these symptoms that developed several months after the
	termination of this drug have been reported overseas.