Published by Ministry of Health, Labour and Welfare Translated by Pharmaceuticals and Medical Devices Agency





This English version is intended to be a reference material to provide convenience for users. In the event of inconsistency between the Japanese original and this English translation, the former shall prevail.

Revision of Precautions

Elbasvir

Grazoprevir hydrate

February 25, 2020

Therapeutic category

Aitivirals

Non-proprietary name

Elbasvir

Grazoprevir hydrate

Safety measure

Precautions should be revised in the package insert.

Pharmaceuticals and Medical Devices Agency

3-3-2 Kasumigaseki, Chiyoda-ku, Tokyo 100-0013 Japan E-mail: safety.info@pmda.go.jp

Revision in line with the Instructions for Package Inserts of Prescription Drugs, etc. PSEHB Notification No. 0608-1 by the Director of Pharmaceutical Safety and Environmental Health Bureau, MHLW, dated June 8, 2017 (New instructions):

Revised language is underlined.

Current	Revision
8. IMPORTANT PRECAUTIONS	8. IMPORTANT PRECAUTIONS
(N/A)	Cases where a dose increase of warfarin or tacrolimus or a
	reduction of insulin or other antidiabetic agents due to
	hypoglycemia were required following initiation of a direct-acting
	antiviral(s) for hepatitis C have been reported. Dose adjustment for
	concomitant drugs may be required in association with anti-viral
	treatment with this drug. In particular, if patients on warfarin,
	tacrolimus or other drugs with a narrow therapeutic window that are
	metabolized by the liver, or on antidiabetic agents are initiated on
	this drug, their prescribing physicians of such drugs in principle
	should be informed of the initiation and the patients should be
	carefully monitored for their conditions through methods such as
	frequent monitoring of PT-INR, blood drug concentration, or blood
	sugar levels.

N/A: Not Applicable, because the section is not included in the current package insert.