Published by Ministry of Health, Labour and Welfare

62

Translated by Pharmaceuticals and Medical Devices Agency



This English version is intended to be a reference material to provide convenience for users. In the event of inconsistency between the Japanese original and this

English translation, the former shall prevail.

Revision of Precautions Baloxavir marboxil

March 31, 2020

Therapeutic category

Antivirals

Non-proprietary name

Baloxavir marboxil

Safety measure Precautions should be revised in the package insert.

Pharmaceuticals and Medical Devices Agency

3-3-2 Kasumigaseki, Chiyoda-ku, Tokyo 100-0013 Japan E-mail: <u>safety.info@pmda.go.jp</u> Revision in line with the Instructions for Package Inserts of Prescription Drugs, PAB Notification No. 606 by the Director General ofPharmaceutical Affairs Bureau, MHW, dated April 25, 1997 (Old instructions):Revised language is underlined.

Current	Revision
Adverse Reactions	Adverse Reactions
Clinically Significant Adverse Reactions	Clinically Significant Adverse Reactions
(N/A)	Ischaemic colitis:
	Ischaemic colitis may occur. If abnormalities such as abdominal
	pain, diarrhoea, and bloody stool are observed, appropriate
	measures should be taken.

Revision in line with the Instructions for Package Inserts of Prescription Drugs, etc. PSEHB Notification No. 0608-1 by the Director of

Pharmaceutical Safety and Environmental Health Bureau, MHLW, dated June 8, 2017 (New instructions):

: Revised language is underlined.

Current	Revision
11. ADVERSE REACTIONS	11. ADVERSE REACTIONS
11.1. Clinically Significant Adverse Reactions	11.1. Clinically Significant Adverse Reactions
(N/A)	Ischaemic colitis
	If abnormalities such as abdominal pain, diarrhoea, and bloody
	stool are observed, appropriate measures should be taken.

N/A: Not Applicable, because the section is not included in the current package insert.

Pharmaceuticals and Medical Devices Agency

3-3-2 Kasumigaseki, Chiyoda-ku, Tokyo 100-0013 Japan E-mail: <u>safety.info@pmda.go.jp</u>