

Q-05B

Revision 1 Correction 2

July 2023

PHARMACOPOEIAL DISCUSSION GROUP**REVISION 1 CORRECTION 2****CODE: Q-05B****NAME: MICROBIAL ENUMERATION TESTS****(Correction of the sign-off document Revision 1 Correction 1 signed on 10 June 2010)****Items to be corrected:**

- Section 4-6 'Results and interpretation', clarification of the reading procedure to be performed when verifying the suitability of the membrane filtration method.
- Section 5-3 'Interpretation of the results', correction by adding the 's' to reflect the plural of the word 'moulds'.


It is understood that sign-off covers the technical content of the draft and each party will adapt it as necessary to conform to the usual presentation of the pharmacopoeia in question; such adaptation includes stipulation of the particular pharmacopoeia's reference materials and general chapters.

European Pharmacopoeia

Signature

Name

Date

DocuSigned by:

E6AA71F148A7464...

C. Vielle


17/08/2023

Japanese Pharmacopoeia

Signature

Name

Date


for K. Nakai

Yoshiro Saito


Aug. 22, 2023

United States Pharmacopoeia

Signature

Name

Date

DocuSigned by:

A7467E52FCC94E9...

Kevin Moore

8/4/2023

1 **MICROBIOLOGICAL EXAMINATION OF**
2 **NON-STERILE PRODUCTS: MICROBIAL ENUMERATION**
3 **TESTS**

4
5 1 INTRODUCTION

6
7 The tests described hereafter will allow quantitative enumeration of mesophilic bacteria and fungi
8 which may grow under aerobic conditions.

9
10 The tests are designed primarily to determine whether a substance or preparation complies with
11 an established specification for microbiological quality. When used for such purposes follow
12 the instructions given below, including the number of samples to be taken and interpret
13 the results as stated below.

14
15 The methods are not applicable to products containing viable micro-organisms as active
16 ingredients.

17
18 Alternative microbiological procedures, including automated methods, may be used, provided
19 that their equivalence to the Pharmacopoeial method has been demonstrated.

20
21 2 GENERAL PROCEDURES

22
23 Carry out the determination under conditions designed to avoid extrinsic microbial contamination
24 of the product to be examined. The precautions taken to avoid contamination must be such that
25 they do not affect any micro-organisms which are to be revealed in the test.

26
27 If the product to be examined has antimicrobial activity, this is insofar as possible removed or
28 neutralised. If inactivators are used for this purpose their efficacy and their absence of toxicity for
29 micro-organisms must be demonstrated.

30
31 If surface-active substances are used for sample preparation, their absence of toxicity for micro-
32 organisms and their compatibility with inactivators used must be demonstrated.

33
34 3 ENUMERATION METHODS

35
36 Use the membrane filtration method, or the plate-count methods, as prescribed. The most
37 probable number (MPN) method is generally the least accurate method for microbial counts,
38 however, for certain product groups with very low bioburden, it may be the most appropriate
39 method.

40
41 The choice of a method is based on factors such as the nature of the product and the required
42 limit of micro-organisms. The method chosen must allow testing of a sufficient sample size to
43 judge compliance with the specification. The suitability of the chosen method must be
44 established.

45

1 4 GROWTH PROMOTION TEST, SUITABILITY OF THE COUNTING METHOD AND
2 NEGATIVE CONTROLS

3
4 4-1 GENERAL CONSIDERATIONS

5
6 The ability of the test to detect micro-organisms in the presence of product to be tested must be
7 established.

8
9 Suitability must be confirmed if a change in testing performance, or the product, which may
10 affect the outcome of the test is introduced.

11
12 4-2 PREPARATION OF TEST STRAINS

13
14 Use standardised stable suspensions of test strains or prepare as stated below. Seed lot culture
15 maintenance techniques (seed-lot systems) are used so that the viable micro-organisms used for
16 inoculation are not more than 5 passages removed from the original master seed-lot. Grow each
17 of the bacterial and fungal test strains separately as described in Table 1.
18

19 *Table 1. – Preparation and Use of Test Micro-organisms*

Micro-organism	Preparation of test strain	Growth promotion		Suitability of counting method in the presence of the product	
		Total aerobic microbial count	Total yeasts and moulds count	Total aerobic microbial count	Total yeasts and moulds count
<i>Staphylococcus aureus</i> such as ATCC 6538, NCIMB 9518, CIP 4.83 or NBRC 13276	<i>Casein soya bean digest agar</i> or <i>casein soya bean digest broth</i> 30 - 35 °C 18-24 h	<i>Casein soya bean digest agar</i> and <i>casein soya bean digest broth</i> ≤ 100 CFU/ 30 - 35 °C ≤ 3 days		<i>Casein soya bean digest agar</i> /MPN <i>casein soya bean digest broth</i> ≤ 100 CFU/ 30 - 35 °C ≤ 3 days	
<i>Pseudomonas aeruginosa</i> such as ATCC 9027, NCIMB 8626, CIP 82.118 or NBRC 13275	<i>Casein soya bean digest agar</i> or <i>casein soya bean digest broth</i> 30 - 35 °C 18-24 h	<i>Casein soya bean digest agar</i> and <i>casein soya bean digest broth</i> ≤ 100 CFU/ 30 - 35 °C ≤ 3 days		<i>Casein soya bean digest agar</i> /MPN <i>casein soya bean digest broth</i> ≤ 100 CFU/ 30 - 35 °C ≤ 3 days	

Revision 1 Correction 2

Q-05b (CP:EP)
July 2023

<i>Bacillus subtilis</i> such as ATCC 6633, NCIMB 8054, CIP 52.62 or NBRC 3134	<i>Casein soya bean digest agar</i> or <i>casein soya bean digest broth</i> 30 - 35 °C 18-24 h	<i>Casein soya bean digest agar</i> and <i>casein soya bean digest broth</i> ≤ 100 CFU 30 - 35 °C ≤ 3 days		<i>Casein soya bean digest agar</i> /MPN <i>casein soya bean digest broth</i> ≤ 100 CFU 30 - 35 °C ≤ 3 days	
<i>Candida albicans</i> such as ATCC 10231, NCPF 3179, IP 48.72 or NBRC 1594	<i>Sabouraud-dextrose agar</i> or <i>Sabouraud-dextrose broth</i> 20 -25 °C 2 – 3 days	<i>Casein soya bean digest agar</i> ≤ 100 CFU 30 - 35 °C ≤ 5 days	<i>Sabouraud-dextrose agar</i> ≤ 100 CFU 20 – 25 °C ≤ 5 days	<i>Casein soya bean digest agar</i> ≤ 100 CFU 30 - 35 °C ≤ 5 days MPN: not applicable	<i>Sabouraud-dextrose agar</i> ≤ 100 CFU/ 20 – 25 °C ≤ 5 days,
<i>Aspergillus brasiliensis</i> such as ATCC 16404, IMI 149007, IP 1431.83 or NBRC 9455	<i>Sabouraud-dextrose agar</i> or <i>potato-dextrose agar</i> 20 -25 °C 5 - 7 days, or until good sporulation is achieved	<i>Casein soya bean digest agar</i> ≤ 100 CFU 30 - 35 °C ≤ 5 days	<i>Sabouraud-dextrose agar</i> ≤ 100 CFU 20 – 25 °C ≤ 5 days	<i>Casein soya bean digest agar</i> ≤ 100 CFU 30-35 °C ≤ 5 days MPN: not applicable	<i>Sabouraud-dextrose agar</i> ≤ 100 CFU 20 – 25 °C ≤ 5 days,

1
2 Use *buffered sodium chloride-peptone solution pH 7.0* or *phosphate buffer solution pH 7.2* to
3 make test suspensions; to suspend *A. brasiliensis* spores, 0.05 per cent of polysorbate 80 may be
4 added to the buffer. Use the suspensions within 2 h or within 24 h if stored at 2-8 °C. As an
5 alternative to preparing and then diluting a fresh suspension of vegetative cells of *A. brasiliensis*
6 or *B. subtilis*, a stable spore suspension is prepared and then an appropriate volume of the spore
7 suspension is used for test inoculation. The stable spore suspension may be maintained at 2-8 °C
8 for a validated period of time.

9
10 **4-3 NEGATIVE CONTROL**

11
12 To verify testing conditions a negative control is performed using the chosen diluent in place of
13 the test preparation. There must be no growth of micro-organisms. A negative control is also
14 performed when testing the products as described under 5. A failed negative control requires an
15 investigation.

16
17
18 **4-4 GROWTH PROMOTION OF THE MEDIA**

19
20 Test each batch of ready-prepared medium and each batch of medium, prepared either from
21 dehydrated medium or from the ingredients described

22
23 Inoculate portions/plates of *casein soya bean digest broth* and *casein soya bean digest agar* with
24 a small number (not more than 100 CFU) of the micro-organisms indicated in Table 1, using a
25 separate portion/plate of medium for each. Inoculate plates of *Sabouraud-dextrose agar* with a

1 small number (not more than 100 CFU) of the micro-organisms indicated in Table 1, using a
2 separate plate of medium for each. Incubate in the conditions described in Table 1.

3
4 For solid media, growth obtained must not differ by a factor greater than 2 from the calculated
5 value for a standardised inoculum. For a freshly prepared inoculum, growth of the micro-
6 organisms comparable to that previously obtained with a previously tested and approved batch of
7 medium occurs. Liquid media are suitable if clearly visible growth of the micro-organisms
8 comparable to that previously obtained with a previously tested and approved batch of medium
9 occurs.

10 11 **4-5 SUITABILITY OF THE COUNTING METHOD IN THE PRESENCE OF PRODUCT**

12
13 **4-5-1 Preparation of the sample.** The method for sample preparation depends on the physical
14 characteristics of the product to be tested. If none of the procedures described below can be
15 demonstrated to be satisfactory, an alternative procedure must be developed.

16
17 *Water-soluble products.* Dissolve or dilute (usually a 1 in 10 dilution is prepared) the product to
18 be examined in *buffered sodium chloride-peptone solution pH 7.0, phosphate buffer solution pH*
19 *7.2 or casein soya bean digest broth.* If necessary adjust to pH 6-8. Further dilutions, where
20 necessary, are prepared with the same diluent.

21
22 *Non-fatty products insoluble in water.* Suspend the product to be examined (usually a 1 in 10
23 dilution is prepared) in *buffered sodium chloride-peptone solution pH 7.0, phosphate buffer*
24 *solution pH 7.2 or casein soya bean digest broth.* A surface-active agent such as 1 g/l of
25 polysorbate 80 may be added to assist the suspension of poorly wettable substances. If necessary
26 adjust to pH 6-8. Further dilutions, where necessary, are prepared with the same diluent.

27
28 *Fatty products.* Dissolve in isopropyl myristate, sterilised by filtration or mix the product to be
29 examined with the minimum necessary quantity of sterile polysorbate 80 or another non-
30 inhibitory sterile surface-active reagent, heated if necessary to not more than 40 °C, or in
31 exceptional cases to not more than 45 °C. Mix carefully and if necessary maintain
32 the temperature in a water-bath. Add sufficient of the pre-warmed chosen diluent to make a 1 in
33 10 dilution of the original product. Mix carefully whilst maintaining the temperature for the
34 shortest time necessary for the formation of an emulsion. Further serial tenfold dilutions may be
35 prepared using the chosen diluent containing a suitable concentration of sterile polysorbate 80 or
36 another non-inhibitory sterile surface-active reagent.

37
38 *Fluids or solids in aerosol form.* Aseptically transfer the product into a membrane filter apparatus
39 or a sterile container for further sampling. Use either the total contents or a defined number of
40 metered doses from each of the containers tested.

41
42 *Transdermal patches.* Remove the protective cover sheets ("release liner") of the transdermal
43 patches and place them, adhesive side upwards, on sterile glass or plastic trays. Cover
44 the adhesive surface with sterile porous material, for example sterile gauze, to prevent the patches
45 from sticking together, and transfer the patches to a suitable volume of the chosen diluent
46 containing inactivators such as polysorbate 80 and/or lecithin. Shake the preparation vigorously
47 for at least 30 min.

1
2 **4-5-2 Inoculation and dilution.** Add to the sample prepared as described above (4-5-1) and to a
3 control (with no test material included) a sufficient volume of the microbial suspension to obtain
4 an inoculum of not more than 100 CFU. The volume of the suspension of the inoculum should
5 not exceed 1 per cent of the volume of diluted product.
6

7 To demonstrate acceptable microbial recovery from the product, the lowest possible dilution
8 factor of the prepared sample must be used for the test. Where this is not possible due to
9 antimicrobial activity or poor solubility, further appropriate protocols must be developed. If
10 inhibition of growth by the sample cannot otherwise be avoided, the aliquot of the microbial
11 suspension may be added after neutralisation, dilution or filtration.
12

13 **4-5-3 Neutralisation/removal of antimicrobial activity.** The number of micro-organisms
14 recovered from the prepared sample diluted as described in 4-5-2 and incubated following
15 the procedure described in 4-5-4, is compared to the number of micro-organisms recovered from
16 the control preparation.

17 If growth is inhibited (reduction by a factor greater than 2), then modify the procedure for
18 the particular enumeration test to ensure the validity of the results. Modification of the procedure
19 may include, for example, (1) an increase in the volume of the diluent or culture medium, (2)
20 incorporation of a specific or general neutralising agents into the diluent, (3) membrane filtration
21 or (4) a combination of the above measures.
22

23 *Neutralising agents.* Neutralising agents may be used to neutralise the activity of antimicrobial
24 agents (Table 2). They may be added to the chosen diluent or the medium preferably before
25 sterilisation. If used, their efficacy and their absence of toxicity for micro-organisms must be
26 demonstrated by carrying out a blank with neutraliser and without product.
27

28 Table 2. – *Common neutralising agents for interfering substances*

29 Interfering substance	Potential neutralising method
30 Glutaraldehyde, Mercurials	Sodium hydrogensulphite (Sodium bisulphite)
31 Phenolics, Alcohol, Aldehydes, Sorbate	Dilution
32 Aldehydes	Glycine
33 Quaternary Ammonium Compounds (QACs), Lecithin	
34 Parahydroxybenzoates (Parabens), Bis-biguanides	
35 QAC, Iodine, Parabens	Polysorbate
36 Mercurials	Thioglycollate
37 Mercurials, Halogens, Aldehydes	Thiosulphate
38 EDTA (edetate)	Mg or Ca ions

39
40 If no suitable neutralising method can be found, it can be assumed that the failure to isolate
41 the inoculated organism is attributable to the microbicidal activity of the product. This
42 information serves to indicate that the article is not likely to be contaminated with the given
43 species of the micro-organism. However, it is possible that the product only inhibits some of
44 the micro-organisms specified herein, but does not inhibit others not included amongst the test
45 strains or for which the latter are not representative. Then, perform the test with the highest
46 dilution factor compatible with microbial growth and the specific acceptance criterion.
47

48 **4-5-4 Recovery of micro-organism in the presence of product.** For each of the micro-
49 organisms listed, separate tests are performed. Only micro-organisms of the added test strain are

1 counted.

2

3 4-5-4-1 *Membrane filtration*

4

5 Use membrane filters having a nominal pore size not greater than 0.45 µm. The type of filter
6 material is chosen in such a way that the bacteria-retaining efficiency is not affected by
7 the components of the sample to be investigated. For each of the micro-organisms listed, one
8 membrane filter is used.

9

10 Transfer a suitable amount of the sample prepared as described under 4-5-1 to 4-5-3 (preferably
11 representing 1 g of the product, or less if large numbers of CFU are expected) to the membrane
12 filter, filter immediately and rinse the membrane filter with an appropriate volume of diluent.

13

14 For the determination of total aerobic microbial count (TAMC), transfer the membrane filter to
15 the surface of *casein soya bean digest agar*. For the determination of total combined
16 yeasts/moulds count (TYMC) transfer the membrane to the surface of *Sabouraud-dextrose agar*.
17 Incubate the plates as indicated in Table 1. Perform the counting.

18

19 4-5-4-2 *Plate-count methods*

20

21 Perform plate-count methods at least in duplicate for each medium and use the mean count of
22 the result.

23

24 4-5-4-2-1 *Pour-plate method*

25

26 For Petri dishes 9 cm in diameter, add to the dish 1 ml of the sample prepared as described under
27 4-5-1 to 4-5-3 and 15-20 ml of *casein soya bean digest agar* or *Sabouraud-dextrose agar*, both
28 media being at not more than 45 °C. If larger Petri dishes are used, the amount of agar medium is
29 increased accordingly. For each of the micro-organisms listed in Table 1, at least 2 Petri dishes
30 are used.

31

32 Incubate the plates as indicated in Table 1. Take the arithmetic mean of the counts per medium
33 and calculate the number of CFU in the original inoculum.

34

35 4-5-4-2-2 *Surface-spread method*

36

37 For Petri dishes 9 cm in diameter, add 15-20 ml of *casein soya bean digest agar* or *Sabouraud-*
38 *dextrose agar* at about 45 °C to each Petri dish and allow to solidify. If larger Petri dishes are
39 used, the volume of the agar is increased accordingly. Dry the plates, for example in a laminar-
40 air-flow cabinet or in an incubator. For each of the micro-organisms listed in Table 1, at least 2
41 Petri dishes are used. Spread a measured volume of not less than 0.1 ml of the sample prepared as
42 described under 4-5-1 to 4-5-3 over the surface of the medium. Incubate and count as prescribed
43 under 4-5-4-2-1.

44

45 4-5-4-3 *Most-probable-number (MPN) method*

46

47 The precision and accuracy of the MPN method is less than that of the membrane filtration

1 method or the plate-count method. Unreliable results are obtained particularly for
2 the enumeration of moulds. For these reasons the MPN method is reserved for the enumeration of
3 TAMC in situations where no other method is available. If the use of the method is justified,
4 proceed as follows.

5
6 Prepare a series of at least 3 serial tenfold dilutions of the product as described under 4-5-1 to 4-
7 5-3. From each level of dilution, 3 aliquots of 1 g or 1 ml are used to inoculate 3 tubes with 9-10
8 ml of *casein soya bean digest broth*. If necessary a surface-active agent such as polysorbate 80,
9 or an inactivator of antimicrobial agents may be added to the medium. Thus, if 3 levels of
10 dilution are prepared 9 tubes are inoculated.

11
12 Incubate all tubes at 30-35 °C for not more than 3 days. If reading of the results is difficult or
13 uncertain owing to the nature of the product to be examined, subculture in the same broth, or
14 *casein soya bean digest agar*, for 1-2 days at the same temperature and use these results.
15 Determine the most probable number of micro-organisms per gram or millilitre of the product to
16 be examined from Table 3.

17 18 4-6 RESULTS AND INTERPRETATION

19 When verifying the suitability of the membrane filtration method, the count of any of the test
20 organisms must not differ by a factor greater than 2 from the value of the control defined in 4-5-2
21 in the absence of the product. When verifying the suitability of the plate-count method, a mean
22 count of any of the test organisms not differing by a factor greater than 2 from the value of
23 the control defined in 4-5-2 in the absence of the product must be obtained. When verifying
24 the suitability of the MPN method the calculated value from the inoculum must be within 95 per
25 cent confidence limits of the results obtained with the control.

26
27 If the above criteria cannot be met for one or more of the organisms tested with any of the
28 described methods, the method and test conditions that come closest to the criteria are used to test
29 the product.

30 31 5 TESTING OF PRODUCTS

32 33 5-1 AMOUNT USED FOR THE TEST

34 Unless otherwise prescribed, use 10 g or 10 ml of the product to be examined taken with
35 the precautions referred to above. For fluids or solids in aerosol form, sample 10 containers. For
36 transdermal patches, sample 10 patches.

37
38 The amount to be tested may be reduced for active substances that will be formulated in
39 the following conditions: the amount per dosage unit (e.g. tablet, capsule, injection) is less than
40 or equal to 1 mg or the amount per gram or millilitre (for preparations not presented in dose
41 units) is less than 1 mg. In these cases, the amount of sample to be tested is not less than
42 the amount present in 10 dosage units or 10 g or 10 ml of the product.

43
44 For materials used as active substances where sample quantity is limited or batch size is
45 extremely small (i.e. less than 1000 ml or 1000 g), the amount tested shall be 1 per cent of
46 the batch unless a lesser amount is prescribed or justified and authorised.

1 For products where the total number of entities in a batch is less than 200 (e.g. samples used in
2 clinical trials), the sample size may be reduced to 2 units, or 1 unit if the size is less than 100.

3
4 Select the sample(s) at random from the bulk material or from the available containers of
5 the preparation. To obtain the required quantity, mix the contents of a sufficient number of
6 containers to provide the sample.

7 -2 EXAMINATION OF THE PRODUCT

9
10 **5-2-1 Membrane filtration.** Use a filtration apparatus designed to allow the transfer of the filter
11 to the medium. Prepare the sample using a method that has been shown suitable as described in
12 section 4 and transfer the appropriate amount to each of 2 membrane filters and filter
13 immediately. Wash each filter following the procedure shown to be suitable.

14
15 For the determination of TAMC, transfer one of the membrane filters to the surface of *casein*
16 *soya bean digest agar*. For the determination of TYMC, transfer the other membrane to
17 the surface of *Sabouraud-dextrose agar*. Incubate the plate of *casein soya bean digest agar* at 30-
18 35 °C for 3-5 days and the plate of *Sabouraud-dextrose agar* at 20-25 °C for 5-7 day Calculate
19 the number of CFU per gram or per millilitre of product.

20
21 When examining transdermal patches, filter 10 per cent of the volume of the preparation
22 described under 4-5-1 separately through each of 2 sterile filter membranes. Transfer one
23 membrane to *casein soya bean digest agar* for TAMC and the other membrane to *Sabouraud-*
24 *dextrose agar* for TYMC.

25 26 **5-2-2 Plate-count methods**

27 28 *5-2-2-1 Pour-plate method*

29
30 Prepare the sample using a method that has been shown to be suitable as described in section 4.
31 Prepare for each medium at least 2 Petri dishes for each level of dilution. Incubate the plates of
32 *casein soya bean digest agar* at 30-35 °C for 3-5 days and the plates of *Sabouraud-dextrose agar*
33 at 20-25 °C for 5-7 days. Select the plates corresponding to a given dilution and showing
34 the highest number of colonies less than 250 for TAMC and 50 for TYMC. Take the arithmetic
35 mean per culture medium of the counts and calculate the number of CFU per gram or per
36 millilitre of product.

37 38 *5-2-2-2 Surface-spread method*

39
40 Prepare the sample using a method that has been shown to be suitable as described in section 4.
41 Prepare at least 2 Petri dishes for each medium and each level of dilution. For incubation and
42 calculation of the number of CFU proceed as described for the pour-plate method.

43 44 *5-2-2-3 Most-probable-number method*

45
46 Prepare and dilute the sample using a method that has been shown to be suitable as described in
47 section 4. Incubate all tubes for 3-5 days at 30-35 °C. Subculture if necessary, using

1 the procedure shown to be suitable. Record for each level of dilution the number of tubes
2 showing microbial growth. Determine the most probable number of micro-organisms per gram or
3 millilitre of the product to be examined from Table 3.

4

5 Table 3. – *Most-probable-number values of micro-organisms*

6

Observed combinations of numbers of tubes showing growth in each set			MPN per g or per ml of product	95 per cent confidence limits
Number of g or ml of product per tube				
0.1	0.01	0.001		
0	0	0	Less than 3	0-9.4
0	0	1	3	0.1-9.5
0	1	0	3	0.1-10
0	1	1	6.1	1.2-17
0	2	0	6.2	1.2-17
0	3	0	9.4	3.5-35
1	0	0	3.6	0.2-17
1	0	1	7.2	1.2-17
1	0	2	11	4-35
1	1	0	7.4	1.3-20
1	1	1	11	4-35
1	2	0	11	4-35
1	2	1	15	5-38
1	3	0	16	5-38
2	0	0	9.2	1.5-35
2	0	1	14	4-35
2	0	2	20	5-38
2	1	0	15	4-38
2	1	1	20	5-38
2	1	2	27	9-94
2	2	0	21	5-40
2	2	1	28	9-94
2	2	2	35	9-94
2	3	0	29	9-94
2	3	1	36	9-94
3	0	0	23	5-94
3	0	1	38	9-104
3	0	2	64	16-181
3	1	0	43	9-181
3	1	1	75	17-199
3	1	2	120	30-360
3	1	3	160	30-380
3	2	0	93	18-360
3	2	1	150	30-380
3	2	2	210	30-400
3	2	3	290	90-990
3	3	0	240	40-990
3	3	1	460	90-1980
3	3	2	1100	200-4000
3	3	3	More than 1100	

7

1 **5-3 INTERPRETATION OF THE RESULTS**

2

3 The total aerobic microbial count (TAMC) is considered to be equal to the number of CFU found
4 using *casein soya bean digest agar*; if colonies of fungi are detected on this medium, they are
5 counted as part of TAMC. The total combined yeasts/moulds count (TYMC) is considered to be
6 equal to the number of CFU found using *Sabouraud-dextrose agar*; if colonies of bacteria are
7 detected on this medium, they are counted as part of TYMC. When the TYMC is expected to
8 exceed the acceptance criterion due to the bacterial growth, *Sabouraud-dextrose agar* containing
9 antibiotics may be used. If the count is carried out by the MPN method the calculated value is
10 the TAMC.

11

12 When an acceptance criterion for microbiological quality is prescribed it is interpreted as follows:

13 — 10^1 micro-organisms: maximum acceptable count = 20,

14 — 10^2 micro-organisms: maximum acceptable count = 200,

15 — 10^3 micro-organisms: maximum acceptable count = 2000, and so forth.

16

17 The recommended solutions and media are described in *Tests for specified micro-organisms*.

