参考文献

Dursun R, Alpaslan M, Caliskan M, Ciftci O, Kulaksizoglu S, Seckin D, et al. Isotretinoin does not prolong QT intervals and QT dispersion in patients with severe acne: a surprising finding for a drug with numerous side effects. J Drugs Dermatol. 2011 Jul;10(7):710-4.

Gollnick H. Current concepts of the pathogenesis of acne: implications for drug treatment. Drugs. 2003;63(15):1579-96.

早川律子、赤岡家雄、菅原幸子、他、経皮吸収型製剤の評価方法 – 第二部 臨床試験、Therapeutic Research. 1990;11(3):251-7.

林伸和、赤松浩彦、岩月啓氏、他、尋常性痤瘡治療ガイドライン、日皮会誌 2008; 118(10): 1893-1923

Karadag AS, Gumrukcuoglu HA, Gunes Bilgili SG, Ozkol HU, Ertugrul DT, Simsek H, et al. Does isotretinoin therapy have any effects on electrocardiography, heart rate and blood pressure? J Dermatolog Treat. 2012 Jun;23(3):168-71.

川村太郎、笹川正二、増田勉、他、貼布試験標準化の基礎的研究、日皮会誌 1970; 80(5): 301-14

Kawashima M, Harada S, Loesche C, Miyachi Y. Adapalene gel 0.1% is effective and safe for Japanese patients with acne vulgaris: a randomized, multicenter, investigator-blinded, controlled study. J Dermatol Sci. 2008 Mar;49(3):241-8.

川島眞、佐藤伸一、古川福実、他、過酸化ベンゾイルゲルの尋常性痤瘡を対象とした第Ⅱ/Ⅲ臨床試験 ― プラセボ対照、ランダム化、二重盲検、並行群間比較、多施設共同試験 ―、 臨床医薬 2014; 30(8): 651-68

川島眞、流利孝、桂巻常夫、尋常性痤瘡患者での過酸化ベンゾイルゲル長期投与時（52 週間）の安全性および有効性評価 ― 非盲検、ランダム化、多施設共同第Ⅲ相臨床試験 ―、 臨床医薬 2014; 30(8): 669-89

Kubota K, Ishizaki T. Dose-dependent pharmacokinetics of benzoic acid following oral administration of sodium benzoate to humans. Eur J Clin Pharmacol. 1991;41(4):363-8.

Morsches B, Holzmann H. [Studies on the percutaneous absorption of benzoyl peroxide (author's transl)]. Arzneimittelforschung. 1982;32(3):298-300. German.

Muizzuddin N, Hellemans L, Van Overloop L, Corstjens H, Declercq L, Maes D. Structural and functional differences in barrier properties of African American, Caucasian and East Asian skin. J Dermatol Sci. 2010 Aug;59(2):123-8.

Nacht S, Yeung D, Beasley JN Jr, Anjo MD, Maibach HI. Benzoyl peroxide: percutaneous penetration and metabolic disposition. J Am Acad Dermatol. 1981 Jan;4(1):31-7.

薬事・食品衛生審議会食品衛生分科会添加物部会（平成 25 年 10 月 30 日）資料、平成 24 年度マーケットバスケット方式による保存料及び着色料の摂取量調査の結果について、http://www.mhlw.go.jp/file/06-Seisakujouhou-11130500-Shokuhinanzenbu/0000041954.pdf

Selcoki Y, Gorpelioglu C, Turgut F, Sarifakioglu E, Ozkara A, Tekin O, et al. Isotretinoin: is there any arrhythmic effect? Int J Dermatol. 2008 Feb;47(2):195-7.

Sieber R, Bütikofer U, Bosset JO, Rüegg M. [Benzoic acid as a natural integral part of foods - an overview]. Mitt Gebiete der Lebensm Hyg. 1989 Nov;80:345–62. German.

Sieber R, Bütikofer U, Baumann E, Bosset JO. [Occurrence of benzoic acid in fermented milk products and cheese]. 1990;81:484–93. German.

Tan J, Gollnick HP, Loesche C, Ma YM, Gold LS. Synergistic efficacy of adapalene 0.1%-benzoyl peroxide 2.5% in the treatment of 3855 acne vulgaris patients. J Dermatolog Treat. 2011 Aug;22(4):197-205.

Thiboutot D, Gollnick H, Bettoli V, Dréno B, Kang S, Leyden JJ, et al. New insights into the management of acne: an update from the Global Alliance to Improve Outcomes in Acne group. J Am Acad Dermatol. 2009 May;60(5 Suppl):S1-50.

Wepierre J, Corroller M, Didry JR. Distribution and dissociation of benzoyl peroxide in cutaneous tissue after application on skin in the hairless rat. Int J Cosmet Sci. 1986 Jun;8(3):97-104.

Yeung D, Nacht S, Bucks D, Maibach HI. Benzoyl peroxide: percutaneous penetration and metabolic disposition. II. Effect of concentration. J Am Acad Dermatol. 1983 Dec;9(6):920-4.