乾燥濃縮人プロトロンビン複合体

ケイセントラ静注用 500

ケイセントラ静注用 1000

2.7　臨床概要

2.7.5　参考文献

CSL ベーリング株式会社

2.7.5　参考文献

Almus FE, Rao LVM, Rapaport SI. Functional properties of factor VIIa / tissue factor formed with purified tissue factor and with tissue factor expressed on cultured endothelial cells. Thromb Haemost 1989;62:1067-73.

Ansell J, Hirsh J, Hylek E, Jacobson A, Crowther M, Palareti G. Pharmacology and Management of the Vitamin K Antagonists, American College of Chest Physicians (ACCP) Evidence-Based Clinical Practice Guidelines (8th edition), Chest 2008; 133:160S-198S.

Bertina RM. Hereditary protein S deficiency. Haemostasis 1985;15:241-6.

Bom VJJ, van Hinsbergh VWM, Reinalda-Poot HH, Mohanlal RW, Bertina RM. Extrinsic activation of human coagulation factors IX and X on the endothelial surface. Thromb Haemost 1991;66:283-91.

Clouse LH, Comp PC. The regulation of hemostasis: the protein C system. N Engl J Med 1986;314:1298-1304.

Colman RW, Clowes AW, George JN, Hirsh J, Marder VJ. Overview of Hemostasis. In: Colman RW, Hirsh J, Marder VJ, Clowes AW, George JN, editors. Hemostasis and thrombosis - basic principles and clinical practice. 4th ed. Philadelphia: Lippincott Williams and Wilkins; 2001; pp 3-16.

Committee for Proprietary Medicinal Products (CPMP). Note for guidance on the investigation of human plasma derived factor VIII and IX products: CPMP/BPWG/198/95 rev. 1. October 2000.

Crochemore T, de Toledo Piza FM, Silva E, Corrêa TD. Thromboelastometry-guided hemostatic therapy: an efficacious approach to manage bleeding risk in acute fatty liver of pregnancy: a case report. J Med Case Rep. 2015; 9:202.

D'Angelo A, Vigano-D'Angelo S, Esmon CT, Comp PC. Acquired deficiencies of protein S: Protein S activity during oral anticoagulation, in liver disease, and in disseminated intravascular coagulation. J Clin Invest 1988;81:1445-54.

Eaton D, Rodriguez H, Vehar GA. Proteolytic processing of human factor VIII. Correlation of specific cleavages by thrombin, factor Xa, and activated protein C with activation and inactivation of factor VIII coagulant activity. Biochemistry 1986;25:505-12.

Esmon CT. Protein C, Protein S, and Thrombomodulin.　In: Colman RW, Hirsh J, Marder VJ, Clowes AW, George JN, editors. Hemostasis and thrombosis - basic principles and clinical practice. 4th ed. Philadelphia: Lippincott Williams and Wilkins; 2001 p. 335-53.

Evans G, Luddington R. Beriplex P/N reverses severe warfarin-induced overanticoagulation immediately and completely in patients presenting with major bleeding. Br J Haematol 2001;115:998-1001.

Executive Board and Scientific Advisory Committee of the German Federal Medical Association. “Leitlinien zur Therapie mit Blutkomponenten und Plasmaderivaten” [Guidelines for therapy with blood components and plasma derivatives]. 3rd ed. Cologne: Deutscher Ärzte-Verlag; 2003.

Fickenscher K. Analysis of individual coagulation factors. In: Thomas L, editor. Clinical laboratory diagnostics - use and assessment of clinical laboratory results. 1st ed. Frankfurt/Main, Germany: TH-Books Verlagsgesellschaft;1998. p.607-9.

Friedman KD, Menitove JE. Preparation and clinical use of plasma and plasma fractions. In: Beutler E, Lichtman MA, Coller BS, Kipps TJ, Seligsohn U, editors. Williams hematology. 6th ed. New York: McGraw Hill; 2001. p. 1917-34.

Furie B, Furie BC. Molecular and cellular biology of blood coagulation (Review). N Engl J Med 1992;326:800-6.

Greenberg DL, Davie EW. Blood coagulation factors: their complementary DNAs, genes, and expression. In: Colman RW, Hirsh J, Marder VJ, Clowes AW, George JN, editors. Hemostasis and thrombosis - basic principles and clinical practice. 4th ed. Philadelphia: Lippincott Williams and Wilkins; 2001 p. 21-57.

Griffin JH. Control of coagulation reactions. In: Beutler E, Lichtman MA, Coller BS, Kipps TJ, Seligsohn U, editors. Williams hematology. 6th ed. New York: McGraw Hill; 2001. p. 1435-49.

Gulati G, Hevelow M, George M, Behling E, Siegel J. International Normalized Ratio versus plasma levels of coagulation factors in patients on Vitamin K antagonist therapy. Arch Pathol Lab med 2011; 135:490-494.

Handin RI, Lux SE, Stossel TP, eds. Blood: Principles and Practice of Hematology. 2nd ed. Lippincott Williams & Wilkins: Philadelphia; 2003.

Jerkeman A, Astermark J, Hedner U, Lethagen S, Olsson CG, Berntorp E. Correlation between different intensities of anti-vitamin K treatment and coagulation parameters. Thromb Res 2000; 98:467-471.

Kisiel W, Canfield WM, Ericsson LH, Davie EW. Anticoagulant properties of bovine plasma protein C following activation by thrombin. Biochemistry 1977; 16:5824-31.

Köhler M. Thrombogenicity of prothrombin complex concentrates. Thromb Res. 1999; 95(4 Suppl 1):S13-7.

Lane DA, Philippou H, Huntington JA. Directing thrombin. Blood 2005; 106(8):2605-2612.

Lippi G, Favaloro E, Franchini M, Guidi GC. Milestones and perspectives in coagulation and hemostasis. Semin Thromb Hemost 2009; 35(1):9-22.

Lockley C. Managing the unpredictable: Challenges in severe factor X deficiency. Haemophilia 2014; 20(Suppl. 3):1-186.

Nadler SB, Hidalgo JU, Bloch T. Prediction of blood volume in normal human adults. Surgery 1962; 51:224-232.

Owen WG. Protein C. In: Colman RW, Hirsh J, Marder VJ, Salzman EW, editors. Hemostasis and Thrombosis. 2nd ed., Philadelphia: JB Lippincott Co.; 1987; pp 235-41.

Paul B, Oxley A, Brigham K, Cox T, Hamilton PJ. Factor II, VII, IX and X concentrations in patients receiving long term warfarin. J clin pathol 1987; 40:94-98.

Preston FE, Laidlaw ST, Sampson B, Kitchen S. Rapid reversal of oral anticoagulation with warfarin by a prothrombin complex concentrate (Beriplex): efficacy and safety in 42 patients. Br J Haematol 2002;116:619-24.

Rao LVM, Rapaport SI. Activation of factor VII bound to tissue factor: a key early step in the tissue factor pathway of blood coagulation. Proc Natl Acad Sci USA 1988;85:6687-91.

Rao LVM, Robinson T, Hoang AD. Factor VIIa / tissue factor-catalyzed activation of factors IX and X on a cell surface and in suspension: a kinetic study. Thromb Haemost 1992;67:654-9.

Steinberg M, Nemerson Y. Activation of factor X. In: Colman RW, Hirsh J, Marder VJ, Salzman EW, editors. Hemostasis and Thrombosis. 2nd ed., Philadelphia: JB Lippincott Co.; 1987 p. 112-19.

Suzuki K, Stenflo J, Dahlbäck B, Teodorsson B. Inactivation of human coagulation factor V by activated protein C. J Biol Chem 1983;258:1914-20.

Watala C, Glanski J, Kardas P. Multivariante relationships between international normalized ratio and vitamin K-dependent coagulation-derived parameters in normal healthy donors and oral anticoagulant therapy patients. Thrombosis J 2003; 1:7.

White GC 2nd, Shapiro AD, Kurczynski EM, Kim HC, Bergman GE, the Mononine Study Group. Variability of in vivo recovery of factor IX after infusion of monoclonal antibody purified factor IX concentrates in patients with hemophilia B. Thromb Haemost 1995;73:779-84.