Bingman AE, Secrest P, Diaz SL, Varki NM, Varki A. A red meat-derived glycan promotes inflammation and cancer progression. Proc Natl Acad Sci U S A. 2015 Jan 13;112(2):542-7.

75) Hogan-Ryan A, Fennelly JJ, Jones M, Cantwell B, Duffy MJ. Serum sialic acid and CEA concentrations in human breast cancer. Br J Cancer. 1980 Apr;41(4):587-92.

76) Romppanen J, Eskelinen M, Tikanoja S, Mononen I. Total and lipid-bound serum sialic acid in benign and malignant breast disease. Anticancer Res. 1997 Mar-Apr;17(2B):1249-53.

77) Gatchev O, Råstam L, Lindberg G, Gullberg B, Eklund GA, Törnberg S. Tumours of the central nervous system and serum sialic acid concentration in men and women. Br J Cancer. 1993 Aug;68(2):425-7.

78) Tomaszewska R, Sonta-Jakimczyk D, Dyduch A, Olejnik I, Mazur B. Sialic acid concentration in different stages of malignant lymphoma and leukemia in children. Acta Paediatr Jpn. 1997 Aug;39(4):448-50.

79) Berbeć H, Paszkowska A, Siwek B, Gradziel K, Cybulski M. Total serum sialic acid concentration as a supporting marker of malignancy in ovarian neoplasia. Eur J Gynaecol Oncol. 1999;20(5-6):389-92.

80) Kongtawelert P, Tangkijvanich P, Ong-Chai S, Poovorawan Y. Role of serumtotal sialic acid in differentiating cholangiocarcinoma from hepatocellularcarcinoma. World J Gastroenterol. 2003 Oct;9(10):2178-81.

81) Chrostek L, Cylwik B, Panasiuk A, Brodowska-Adamusiak D, Gruszewska E. Lipid-bound sialic acid (LSA) in liver diseases of different etiologies. Ann Hepatol. 2011 Apr-Jun;10(2):150-4.

82) Upadhya S, Upadhya S, Prabhu KS. Serum glycoconjugates and ceruloplasmin incancer of uterine cervix. Indian J Clin Biochem. 2002 Jan;17(1):20-4.

83) Sandhu R, Lal H, Kundu ZS, Kharb S. Serum fluoride and sialic acid levels in osteosarcoma. Biol Trace Elem Res. 2011 Dec;144(1-3):1-5.

84) Ohkawa M, Okasho A, Fuse H, Orito M, Sugata T, Hisazumi H. [Serum sialic acid levels in patients with acute prostatitis or acute epididymitis]. Hinyokika Kiyo. 1985 Dec;31(12):2161-8.

85) Kalela A, Pönniö M, Koivu TA, Höyhtyä M, Huhtala H, Sillanaukee P, Nikkari ST.

Association of serum sialic acid and MMP-9 with lipids and inflammatory markers. Eur J Clin Invest. 2000 Feb;30(2):99-104.

86) Chrostek L, Cylwik B, Gindzienska-Sieskiewicz E, Gruszewska E, Szmitkowski M,Sierakowski S. Sialic acid level reflects the disturbances of glycosylation andacute-phase reaction in rheumatic diseases. Rheumatol Int. 2014 Mar;34(3):393-9.

87) Lindberg G, Eklund GA, Gullberg B, Råstam L. Serum sialic acid concentration and cardiovascular mortality. BMJ. 1991 Jan 19;302(6769):143-6.

88) Lindberg G, Råstam L, Gullberg B, Eklund GA. Serum sialic acid concentration predicts both coronary heart disease and stroke mortality: multivariate analysis including 54,385 men and women during 20.5 years follow-up. Int J Epidemiol. 1992 Apr;21(2):253-7.

89) Crook MA, Earle K, Morocutti A, Yip J, Viberti G, Pickup JC. Serum sialic acid, a risk factor for cardiovascular disease, is increased in IDDM patients with microalbuminuria and clinical proteinuria. Diabetes Care. 1994 Apr;17(4):305-10.

90) Wakabayashi I, Sakamoto K, Yoshimoto S, Kakishita E. Serum sialic acid concentration and atherosclerotic risk factors. J Atheroscler Thromb. 1994;1(2):113-7.

91) Pickup JC, Mattock MB, Crook MA, Chusney GD, Burt D, Fitzgerald AP. Serum sialic acid concentration and coronary heart disease in NIDDM. Diabetes Care. 1995 Aug;18(8):1100-3.

92) Råstam L, Lindberg G, Folsom AR, Burke GL, Nilsson-Ehle P, Lundblad A. Association between serum sialic acid concentration and carotid atherosclerosis measured by B-mode ultrasound. The ARIC Investigators. Atherosclerosis Risk in Communities Study. Int J Epidemiol. 1996 Oct;25(5):953-8.

93) Crook M, Lumb P, Andrews V, Swaminathan R. Serum total sialic acid, a reputed cardiovascular risk factor, and its relationship to lipids, plasma fasting insulin, blood pressure and body mass index in normal individuals. Clin Sci (Lond). 1998 Jul;95(1):53-7.

94) Wakabayashi I. Age-related change in relationship between body-mass index, serum sialic acid, and atherogenic risk factors. J Atheroscler Thromb. 1998;5(2):60-5.

95) Kalela A, Pönniö M, Koivu TA, Höyhtyä M, Huhtala H, Sillanaukee P, Nikkari ST. Association of serum sialic acid and MMP-9 with lipids and inflammatory markers. Eur J Clin Invest. 2000 Feb;30(2):99-104.

96) Gokmen SS, Kilicli G, Ozcelik F, Ture M, Gulen S. Association between serum total and lipid-bound sialic acid concentration and the severity of coronary atherosclerosis. J Lab Clin Med. 2002 Aug;140(2):110-8.

97) Crook MA, Miell J, Ameerally P, Lumb P, Singh N, Russell-Jones D, Goldsmith L. Serum sialic acid, a reputed cardiovascular risk factor, is related to serum leptin concentrations in Fijians. Clin Chim Acta. 2003 May;331(1-2):1-5.

98) Serdar Z, Yeşilbursa D, Dirican M, Sarandöl E, Serdar A. Sialic acid and oxidizability of lipid and proteins and antioxidant status in patients with coronary artery disease. Cell Biochem Funct. 2007 Nov-Dec;25(6):655-64.

99) Tseke P, Grapsa E, Stamatelopoulos K, Samouilidou E, Rammos G, Papamichael C,Zakopoulos N. Correlations of sialic acid with markers of inflammation,atherosclerosis and cardiovascular events in hemodialysis patients. Blood Purif. 2008;26(3):261-6.

100) Pönniö M, Alho H, Nikkari ST, Olsson U, Rydberg U, Sillanaukee P. Serum sialic acid in a random sample of the general population. Clin Chem. 1999 Oct;45(10):1842-9.

101) Jinghua L, Tie Z, Ping W, Yongtong C. The relationship between serum sialicacid and high-sensitivity C-reactive protein with prehypertension. Med Sci Monit.2014 Apr 3;20:551-5.

102) Chen J, Gall MA, Yokoyama H, Jensen JS, Deckert M, Parving HH. Raised serum sialic acid concentration in NIDDM patients with and without diabetic nephropathy. Diabetes Care. 1996 Feb;19(2):130-4.

103) Yokoyama H, Jensen JS, Myrup B, Mathiesen ER, Rønn B, Deckert T. Raised serum sialic acid concentration precedes onset of microalbuminuria in IDDM. A 10-year follow-up study. Diabetes Care. 1996 May;19(5):435-40.

104) Wakabayashi I, Masuda H. Age-dependent relation of serum sialic acid concentration to aortic pulse wave velocity in type 2 diabetes. Diabetes Metab. 2004 Nov;30(5):441-9.

105) Nayak SB, Bhaktha G. Relationship between sialic acid and metabolic variablesin Indian type 2 diabetic patients. Lipids Health Dis. 2005 Aug 10;4:15.

106) Cylwik B, Chrostek L, Zalewski B, Dabrowski A, Szmitkowski M. Serum total sialic

acid in differential diagnostics of jaundice caused by malignant and nonmalignant diseases: a ROC curve analysis. Dig Dis Sci. 2007 Sep;52(9):2317-22.

107) Gruszewska E, Cylwik B, Panasiuk A, Szmitkowski M, Flisiak R, Chrostek L.Total and free serum sialic acid concentration in liver diseases. Biomed Res Int.2014; 2014:876096.

108) Alvi MH, Amer NA, Sumerin I. Serum 5-nucleotidase and serum sialic acid in pregnancy. Obstet Gynecol. 1988 Aug;72(2):171-4.

109) Crook M, Constable S, Lumb P, Rymer J. Elevated serum sialic acid in pregnancy. J Clin Pathol. 1997 Jun;50(6):494-5.

110) Cylwik B, Chrostek L, Brodowska-Adamusiak D, Gruszewska E, Daniluk M, Szmitkowski M. The changes of sialic acid concentration and content in apolipoprotein B-containing lipoproteins in the sera of alcoholics. Alcohol Alcohol. 2010 Sep-Oct;45(5):422-6.

111) Chrostek L, Cylwik B, Korcz W, Krawiec A, Koput A, Supronowicz Z, Szmitkowski M. Serum free sialic acid as a marker of alcohol abuse. Alcohol Clin Exp Res. 2007 Jun;31(6):996-1001.

112) Chrostek L, Cylwik B, Krawiec A, Korcz W, Szmitkowski M. Relationship between serum sialic acid and sialylated glycoproteins in alcoholics. Alcohol Alcohol. 2007 Nov-Dec;42(6):588-92.

113) Lindberg G, Råstam L, Gullberg B, Eklund GA, Törnberg S. Serum sialic acid concentration and smoking: a population based study. BMJ. 1991 Nov 23;303(6813):1306-7.