














Acebutolol Hydrochloride**Acetohexamide 1****Acetohexamide 2**


Aclarubicin Hydrochloride**Acrinol Hydrate****Actinomycin D**

Afloqualone**Alimemazine Tartrate****Allopurinol**


Alprazolam**Alprenolol Hydrochloride****Alprostadil**

Alprostadil Alfadex


A solution prepared as follows: To 10 mL of a solution in dilute ethanol (3 in 10,000) add 1 mL of potassium hydroxide-ethanol TS, and allow to stand for 15 minutes.

Amibenonium Chloride


A solution in methanol (1 in 5000)

Amitriptyline Hydrochloride


An aqueous solution (1 in 100,000)

Amoxapine




A solution in 0.1 mol/L hydrochloric acid TS (1 in 50,000)




Amphotericin B




A solution prepared as follows: Dissolve 25 mg in 5 mL of dimethylsulfoxide, and add methanol to make 50 mL. To 1 mL of this solution add methanol to make 50 mL.




Arotinolol Hydrochloride




A solution in methanol (1 in 75,000)


Aspoxicillin Hydrate**Atenolol****Azathioprine**

Aztreonam**Bacampicillin Hydrochloride****Baclofen**


Bamethan Sulfate**Benidipine Hydrochloride****Benserazide Hydrochloride**

Benzalkonium Chloride**Benzbromarone****Benzethonium Chloride**


Benzylpenicillin Benzathine Hydrate**Benzylpenicillin Potassium****Berberine Chloride Hydrate**

Berberine Tannate

A solution prepared as follows: Dissolve 0.01 g in 10 mL of methanol and 0.4 mL of 1 mol/L hydrochloric acid TS, and add water to make 200 mL. To 8 mL of this solution add water to make 25 mL.




Betahistine Mesilate




A solution in 0.1 mol/L hydrochloric acid TS (1 in 50,000)


Betamethasone

A solution prepared as follows: Dissolve 1.0 mg in 10 mL of ethanol (95). To 2 mL of this solution, add 10 mL of phenylhydrazine hydrochloride TS, shake, heat in a water bath at 60°C for 20 minutes, and allow it to cool.


Blank: A solution obtained by proceeding in the same manner with 2.0 mL of ethanol (95).

Betamethasone Dipropionate**Bezafibrate****Bifonazole**


Biperiden Hydrochloride**Bisacodyl****Bleomycin Hydrochloride**

Bleomycin Sulfate




A solution prepared as follows: To 4 mg add 5 μ L of copper (II) sulfate TS, and dissolve in water to make 100 mL.




Bromazepam



A solution in ethanol (99.5) (1 in 200,000)

Bromhexine Hydrochloride

A solution in 0.01 mol/L hydrochloric acid TS (3 in 100,000)

Bromocriptine Mesilate**Bucumolol Hydrochloride****Bufetolol Hydrochloride**

Bufexamac**Bumetanide****Bupranolol Hydrochloride**

Butropium Bromide 1**Butropium Bromide 2****Calcium Folate**