


## Dantrolene Sodium Hydrate

ダントロレンナトリウム水和物



$C_{14}H_9N_4NaO_5 \cdot 3\frac{1}{2}H_2O$ : 399.29

Monosodium 3-[5-(4-nitrophenyl)furan-2-ylmethylene]amino-2,5-dioxo-1,3-imidazolidinate hemiheptahydrate [14663-23-1, anhydride]

Dantrolene Sodium Hydrate contains not less than 98.0% of dantrolene sodium ( $C_{14}H_9N_4NaO_5$ : 336.23), calculated on the anhydrous basis.

**Description** Dantrolene Sodium Hydrate occurs as a yellowish orange to deep orange, crystalline powder.

It is soluble in propylene glycol, sparingly soluble in methanol, slightly soluble in ethanol (95), very slightly soluble in water and in acetic acid (100), and practically insoluble in acetone, in tetrahydrofuran and in diethyl ether.

**Identification (1)** Determine the absorption spectrum of a solution of Dantrolene Sodium Hydrate in methanol (1 in 100,000) as directed under Ultraviolet-visible Spectrophotometry <2.24>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wavelengths.

**(2)** Determine the infrared absorption spectrum of Dantrolene Sodium Hydrate as directed in the potassium bromide disk method under Infrared Spectrophotometry <2.25>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wave numbers.

**(3)** To 0.1 g of Dantrolene Sodium Hydrate add 20 mL of water and 2 drops of acetic acid (100), shake well, and filter: the filtrate responds to the Qualitative Tests <1.09> (1) for sodium salt.

**Purity (1)** Alkalinity—To 0.7 g of Dantrolene Sodium Hydrate add 10 mL of water, shake well, and centrifuge or filter through a membrane filter. To 5 mL of the supernatant or the filtrate add 45 mL of water, 3 drops of phenolphthalein TS and 0.10 mL of 0.1 mol/L hydrochloric acid VS: a red color is not produced.

**(2)** Heavy metals <1.07>—Proceed with 1.0 g of Dantrolene Sodium Hydrate according to Method 2, and perform the test. Prepare the control solution with 2.0 mL of Standard Lead Solution (not more than 20 ppm).

**(3)** Related Substances—Dissolve 50 mg of Dantrolene Sodium Hydrate in 20 mL of tetrahydrofuran and 2 mL of acetic acid (100), add ethanol (99.5) to make 100 mL, and use this solution as the sample solution. Pipet 1 mL of the sample solution, add ethanol (99.5) to make exactly 50 mL, and use this solution as the standard solution. Perform the test with exactly 10  $\mu$ L of the sample solution and standard solution as directed under Liquid Chromatography <2.01> according to the following conditions. Determine each peak area from these solutions by the automatic integration method: the total

area of all peaks other than the peak of dantrolene from the sample solution is not larger than the peak area of dantrolene from the standard solution.

**Operating conditions—**

**Detector:** An ultraviolet absorption photometer (wavelength: 300 nm).

**Column:** A stainless steel column about 4 mm in inside diameter and about 15 cm in length, packed with silica gel for liquid chromatography (5  $\mu$ m in particle diameter).

**Column temperature:** A constant temperature of about 30°C.

**Mobile phase:** A mixture of hexane, acetic acid (100) and ethanol (99.5) (90:10:9).

**Flow rate:** Adjust the flow rate so that the retention time of dantrolene is about 8 minutes.

**Selection of column:** Dissolve 5 mg of Dantrolene Sodium Hydrate and 0.1 g of theophylline in 20 mL of tetrahydrofuran and 2 mL of acetic acid (100), and add ethanol (99.5) to make 100 mL. To 10 mL of this solution add ethanol (99.5) to make 100 mL. Proceed with 10  $\mu$ L of this solution under the above operating conditions, and calculate the resolution. Use a column giving elution of theophylline and dantrolene in this order with the resolution between these peaks being not less than 6.

**Detection sensitivity:** Adjust the detection sensitivity so that the peak height of dantrolene from 10  $\mu$ L of the standard solution is 10% to 40% of the full scale.

**Time span of measurement:** About twice as long as the retention time of dantrolene beginning after the solvent peak.

**Water <2.48>** 14.5 – 17.0% (0.2 g, direct titration).


**Assay** Weigh accurately about 0.7 g of Dantrolene Sodium Hydrate, dissolve in 180 mL of a mixture of propylene glycol and acetone (1:1), and titrate <2.50> with 0.1 mol/L perchloric acid VS (potentiometric titration). Perform a blank determination, and make any necessary correction.

Each mL of 0.1 mol/L perchloric acid VS  
= 33.624 mg of  $C_{14}H_9N_4NaO_5$

**Containers and storage** Containers—Tight containers.

## Daunorubicin Hydrochloride

ダウノルビシン塩酸塩



$C_{27}H_{29}NO_{10} \cdot HCl$ : 563.98

(2S,4S)-2-Acetyl-4-(3-amino-2,3,6-trideoxy- $\alpha$ -L-lyxohexopyranosyloxy)-2,5,12-trihydroxy-7-methoxy-1,2,3,4-tetrahydrotetracycline-6,11-dione monohydrochloride [23541-50-6]

Daunorubicin Hydrochloride is the hydrochloride of

an anthracycline substance having antitumor activity produced by the growth of *Streptomyces peucetius*.

It contains not less than 940  $\mu\text{g}$  (potency) and not more than 1050  $\mu\text{g}$  (potency) per mg, calculated on the dried basis. The potency of Daunorubicin Hydrochloride is expressed as mass (potency) of daunorubicin hydrochloride ( $\text{C}_{27}\text{H}_{29}\text{NO}_{10}\cdot\text{HCl}$ ).

**Description** Daunorubicin Hydrochloride occurs as a red powder.

It is soluble in water and in methanol, and slightly soluble in ethanol (99.5).

**Identification (1)** Determine the absorption spectrum of a solution of Daunorubicin Hydrochloride in methanol (1 in 100,000) as directed under Ultraviolet-visible Spectrophotometry <2.24>, and compare the spectrum with the Reference Spectrum or the spectrum of a solution of Daunorubicin Hydrochloride Reference Standard prepared in the same manner as the sample solution: both spectra exhibit similar intensities of absorption at the same wavelengths.

(2) A solution of Daunorubicin Hydrochloride (1 in 50) responds to the Qualitative Tests <1.09> (2) for chloride.

**Optical rotation** <2.49>  $[\alpha]_{\text{D}}^{20}$ : +250 – +275° (15 mg calculated on the dried basis, methanol, 10 mL, 100 mm).

**pH** <2.54> Dissolve 0.15 g of Daunorubicin Hydrochloride in 30 mL of water: the pH of the solution is between 4.5 and 6.0.

**Purity (1)** Clarity and color of solution—Dissolve 20 mg of Daunorubicin Hydrochloride in 10 mL of water: the solution is clear and red.

(2) Heavy metals <1.07>—Proceed with 1.0 g of Daunorubicin Hydrochloride according to Method 2, and perform the test. Prepare the control solution with 2.0 mL of Standard Lead Solution (not more than 20 ppm).

(3) Related substances—Dissolve 10 mg of Daunorubicin Hydrochloride in 5 mL of methanol, and use this solution as the sample solution. Pipet 3 mL of the sample solution, add methanol to make exactly 100 mL, and use this solution as the standard solution. Perform the test with these solutions as directed under Thin-layer Chromatography <2.03>. Spot 10  $\mu\text{L}$  each of the sample solution and standard solution on a plate of silica gel for thin-layer chromatography. Develop the plate with a mixture of chloroform, methanol, water and acetic acid (100) (15:5:1:1) to a distance of about 10 cm, and air-dry the plate. Examine the spots with the naked eye: the spot other than the principal spot obtained from the sample solution is not more intense than the spot from the standard solution.

**Loss on drying** <2.41> Not more than 7.5% (0.1 g, reduced pressure not exceeding 0.67 kPa, 60°C, 3 hours).

**Assay** Weigh accurately an amount of Daunorubicin Hydrochloride and Daunorubicin Hydrochloride Reference Standard, equivalent to about 20 mg (potency), dissolve each in a suitable amount of the mobile phase, add exactly 4 mL of the internal standard solution and the mobile phase to make 20 mL, and use these solutions as the sample solution and the standard solution, respectively. Perform the test with 5  $\mu\text{L}$  each of the sample solution and standard solution as directed under Liquid Chromatography <2.01> according to the following conditions, and determine the ratios,  $Q_T$  and  $Q_S$ , of

the peak area of daunorubicin to that of the internal standard.

$$\begin{aligned} &\text{Amount } [\mu\text{g (potency)}] \text{ of } \text{C}_{27}\text{H}_{29}\text{NO}_{10}\cdot\text{HCl} \\ &= W_S \times (Q_T/Q_S) \times 1000 \end{aligned}$$

$W_S$ : Amount [mg (potency)] of Daunorubicin Hydrochloride Reference Standard

**Internal standard solution**—A solution of 2-naphthalenesulfonic acid in the mobile phase (1 in 100).

**Operating conditions**—

Detector: An ultraviolet absorption photometer (wavelength: 254 nm).

Column: A stainless steel column 4.6 mm in inside diameter and 30 cm in length, packed with octadecylsilanized silica gel for liquid chromatography (10  $\mu\text{m}$  in particle diameter).

Column temperature: A constant temperature of about 25°C.

Mobile phase: Adjust the pH of a mixture of water and acetonitrile (31:19) to 2.2 with phosphoric acid.

Flow rate: Adjust the flow rate so that the retention time of daunorubicin is about 9 minutes.

**System suitability**—


System performance: When the procedure is run with 5  $\mu\text{L}$  of the standard solution under the above operating conditions, the internal standard and daunorubicin are eluted in this order with the resolution between these peaks being not less than 2.0.

System repeatability: When the test is repeated 6 times with 5  $\mu\text{L}$  of the standard solution under the above operating conditions, the relative standard deviation of the ratios of the peak area of daunorubicin to that of the internal standard is not more than 2.0%.

**Containers and storage** Containers—Tight containers.

## Deferoxamine Mesilate

デフェロキサミンメシル酸塩



$\text{C}_{25}\text{H}_{48}\text{N}_6\text{O}_8\cdot\text{CH}_4\text{O}_3\text{S}$ : 656.79

*N*-[5-(Acetylhydroxyamino)pentyl]-*N'*-(5-{3-[(5-aminopentyl)hydroxycarbonyl]propanoylamino}pentyl)-*N'*-hydroxysuccinamide monomethanesulfonate [138-14-7]

Deferoxamine Mesilate contains not less than 98.0% and not more than 102.0% of  $\text{C}_{25}\text{H}_{48}\text{N}_6\text{O}_8\cdot\text{CH}_4\text{O}_3\text{S}$ , calculated on the anhydrous basis.

**Description** Deferoxamine Mesilate occurs as a white to pale yellowish white, crystalline powder.

It is freely soluble in water, and practically insoluble in ethanol (99.5), in 2-propanol and in diethyl ether.

Melting point: about 147°C (with decomposition).

**Identification (1)** To 5 mL of a solution of Deferoxamine Mesilate (1 in 500) add 1 drop of iron (III) chloride TS: a deep red color develops.

**(2)** To 0.05 g of Deferoxamine Mesilate add 0.2 g of sodium hydroxide, melt by heating over a small flame, and heat further for 2 to 3 seconds. To the residue add 0.5 mL of water, acidify with dilute hydrochloric acid, and warm: the gas evolved changes moistened potassium iodate-starch paper to blue.

**(3)** Determine the infrared absorption spectrum of Deferoxamine Mesilate as directed in the potassium bromide disk method under Infrared Spectrophotometry <2.25>, and compare the spectrum with the Reference Spectrum or the spectrum of Deferoxamine Mesilate Reference Standard: both spectra exhibit similar intensities of absorption at the same wave numbers.

**pH** <2.54> Dissolve 1.0 g of Deferoxamine Mesilate in 10 mL of water: the pH of this solution is between 3.5 and 5.5.

**Purity (1)** Clarity and color of solution—Dissolve 1.0 g of Deferoxamine Mesilate in 10 mL of water: the solution is clear and colorless to pale yellow.

**(2)** Chloride <1.03>—Perform the test with 1.0 g of Deferoxamine Mesilate. Prepare the control solution with 0.90 mL of 0.01 mol/L hydrochloric acid VS (not more than 0.032%).

**(3)** Sulfate <1.14>—Perform the test with 0.6 g of Deferoxamine Mesilate. Prepare the control solution with 0.50 mL of 0.005 mol/L sulfuric acid VS (not more than 0.040%).

**(4)** Heavy metals <1.07>—Proceed with 2.0 g of Deferoxamine Mesilate according to Method 4, and perform the test. Prepare the control solution with 2.0 mL of Standard Lead Solution (not more than 10 ppm).

**(5)** Arsenic <1.11>—Prepare the test solution with 1.0 g of Deferoxamine Mesilate according to Method 3, and perform the test. Use a solution of magnesium nitrate hexahydrate in ethanol (95) (1 in 10) (not more than 2 ppm).

**(6)** Related substances—Dissolve 50 mg of Deferoxamine Mesilate in 50 mL of the mobile phase, and use this solution as the sample solution. Pipet 3 mL of the sample solution, add the mobile phase to make exactly 50 mL, and use this solution as the standard solution. Perform the test with exactly 20  $\mu$ L each of the sample solution and standard solution as directed under Liquid Chromatography <2.01> according to the following conditions. Determine each peak area of both solutions by the automatic integration method: the total area of all peaks other than the peak of deferoxamine is not larger than the peak area of deferoxamine from the standard solution.

**Operating conditions—**

**Detector:** An ultraviolet absorption photometer (wavelength: 230 nm).

**Column:** A stainless steel column 4 mm in inside diameter and 20 cm in length, packed with octadecylsilanized silica gel for liquid chromatography (10  $\mu$ m in particle diameter).

**Column temperature:** A constant temperature of about 40°C.

**Mobile phase:** Dissolve 1.32 g of diammonium hydrogen phosphate, 0.37 g of disodium dihydrogen ethylenediamine tetraacetate dihydrate and 1.08 g of sodium 1-heptanesulfonate in 950 mL of water, and adjust the pH of this solution

to 2.8 with phosphoric acid. To 800 mL of this solution add 100 mL of 2-propanol.

**Flow rate:** Adjust the flow rate so that the retention time of deferoxamine is about 15 minutes.

**Time span of measurement:** About two times as long as the retention time of deferoxamine beginning after the solvent peak.

**System suitability—**

**Test for required detection:** To exactly 2 mL of the standard solution add the mobile phase to make exactly 100 mL. Confirm that the peak area of deferoxamine obtained from 20  $\mu$ L of this solution is equivalent to 1.5 to 2.5% of that of deferoxamine obtained from 20  $\mu$ L of the standard solution.

**System performance:** Dissolve 16 mg of Deferoxamine Mesilate and 4 mg of methyl parahydroxybenzoate in 50 mL of the mobile phase. When the procedure is run with 20  $\mu$ L of this solution under the above operating conditions, deferoxamine and methyl parahydroxybenzoate are eluted in this order with the resolution between these peaks being not less than 4.

**System repeatability:** When the test is repeated 6 times with 20  $\mu$ L of the standard solution under the above operating conditions, the relative standard deviation of the peak areas of deferoxamine is not more than 3.0%.

**Water** <2.48> Not more than 2.0% (0.2 g, direct titration).

**Residue on ignition** <2.44> Not more than 0.1% (1 g).

**Assay** Weigh accurately about 60 mg of Deferoxamine Mesilate and Deferoxamine Mesilate Reference Standard (previously determine the water <2.48> in the same manner as Deferoxamine Mesilate), dissolve each in 20 mL of water, add 10 mL of 0.05 mol/L sulfuric acid TS, and add water to make exactly 50 mL. Pipet 5 mL each of these solutions, add exactly 5 mL of 0.05 mol/L sulfuric acid TS and exactly 0.2 mL of iron (III) chloride TS, then add water to make exactly 50 mL, and use these solutions as the sample solution and the standard solution, respectively. Perform the test with these solutions as directed under Ultraviolet-visible Spectrophotometry <2.24>, using a solution prepared by adding 0.05 mol/L sulfuric acid TS to 0.2 mL of iron (III) chloride TS to make exactly 50 mL as the blank, and determine the absorbances,  $A_T$  and  $A_S$ , of each solution from the sample solution and the standard solution at 430 nm.


$$\begin{aligned} \text{Amount (mg) of } C_{25}H_{48}N_6O_8 \cdot CH_4O_3S \\ = W_S \times (A_T/A_S) \end{aligned}$$

$W_S$ : Amount (mg) of Deferoxamine Mesilate Reference Standard, calculated on the anhydrous basis

**Containers and storage** Containers—Tight containers.

## Dehydrocholic Acid

デヒドロコール酸



$C_{24}H_{34}O_5$ : 402.52

3,7,12-Trioxo-5β-cholan-24-oic acid [81-23-2]

Dehydrocholic Acid, when dried, contains not less than 98.5% of  $C_{24}H_{34}O_5$ .

**Description** Dehydrocholic Acid occurs as a white, crystalline powder. It is odorless, and has a bitter taste.

It is sparingly soluble in 1,4-dioxane, slightly soluble in ethanol (95), and practically insoluble in water and in diethyl ether.

It dissolves in sodium hydroxide TS.

**Identification (1)** Dissolve 5 mg of Dehydrocholic Acid in 1 mL of sulfuric acid and 1 drop of formaldehyde solution, and allow to stand for 5 minutes. Add 5 mL of water to the solution: the solution shows a yellow color and a blue-green fluorescence.

(2) To 0.02 g of Dehydrocholic Acid add 1 mL of ethanol (95), shake, add 5 drops of 1,3-dinitrobenzene TS and 0.5 mL of a solution of sodium hydroxide (1 in 8), and allow to stand: a purple to red-purple color develops, and gradually changes to brown.

**Optical rotation** <2.49>  $[\alpha]_D^{20}$ : +29 – +32° (after drying, 0.2 g, 1,4-dioxane, 10 mL, 100 mm).

**Melting point** <2.60> 233 – 242°C

**Purity (1)** Odor—To 2.0 g of Dehydrocholic Acid add 100 mL of water, and boil for 2 minutes: the solution is odorless.

(2) Clarity and color of solution—To 0.10 g of Dehydrocholic Acid, previously powdered in a mortar, add 30 mL of ethanol (95), and dissolve by shaking for 10 minutes: the solution is clear and colorless.

(3) Chloride <1.03>—To 2.0 g of Dehydrocholic Acid add 100 mL of water, shake for 5 minutes and filter, and use this filtrate as the sample solution. To 25 mL of the sample solution add 6 mL of dilute nitric acid, heat in a water bath for 6 minutes, filter after cooling, and collect the clear filtrate. Wash the residue with 10 mL of water, combine the washings and the filtrate, dilute with water to 50 mL, and perform the test using this solution as the test solution. Prepare the control solution with 0.30 mL of 0.01 mol/L hydrochloric acid VS (not more than 0.021%).

(4) Sulfate <1.14>—Add 1 mL of dilute hydrochloric acid to 25 mL of the sample solution obtained in (3), heat in a water bath for 6 minutes, filter after cooling, and collect the clear filtrate. Wash the residue with 10 mL of water, combine the washings and the filtrate, dilute with water to 50 mL, and perform the test using this solution as the test solution. Pre-

pare the control solution with 0.50 mL of 0.005 mol/L sulfuric acid VS (not more than 0.048%).

(5) Heavy metals <1.07>—Proceed with 1.0 g of Dehydrocholic Acid according to Method 2, and perform the test. Prepare the control solution with 2.0 mL of Standard Lead Solution (not more than 20 ppm).

(6) Barium—To the solution obtained in (1) add 2 mL of hydrochloric acid, and boil for 2 minutes. Cool, filter, and wash with water until 100 mL of the filtrate is obtained. To 10 mL of the filtrate add 1 mL of dilute sulfuric acid: no turbidity is produced.

**Loss on drying** <2.41> Not more than 1.0% (1 g, 105°C, 2 hours).

**Residue on ignition** <2.44> Not more than 0.2% (1 g).


**Assay** Weigh accurately about 0.5 g of Dehydrocholic Acid, previously dried, add 40 mL of neutralized ethanol and 20 mL of water, and dissolve by warming. Add 2 drops of phenolphthalein TS, titrate <2.50> with 0.1 mol/L sodium hydroxide VS, adding 100 mL of freshly boiled and cooled water as the end point is approached, and continue the titration.

Each mL of 0.1 mol/L sodium hydroxide VS  
= 40.25 mg of  $C_{24}H_{34}O_5$

**Containers and storage** Containers—Well-closed containers.

## Purified Dehydrocholic Acid

精製デヒドロコール酸



$C_{24}H_{34}O_5$ : 402.52

3,7,12-Trioxo-5β-cholan-24-oic acid [81-23-2]

Purified Dehydrocholic Acid, when dried, contains not less than 99.0% of  $C_{24}H_{34}O_5$ .

**Description** Purified Dehydrocholic Acid occurs as a white, crystalline powder. It is odorless, and has a bitter taste.

It is sparingly soluble in 1,4-dioxane, slightly soluble in ethanol (95), and practically insoluble in water and in diethyl ether.

It dissolves in sodium hydroxide TS.

**Identification (1)** Dissolve 5 mg of Purified Dehydrocholic Acid in 1 mL of sulfuric acid and 1 drop of formaldehyde solution, and allow to stand for 5 minutes. Add 5 mL of water to the solution: the solution shows a yellow color and blue-green fluorescence.

(2) To 0.02 g of Purified Dehydrocholic Acid add 1 mL of ethanol (95), shake, add 5 drops of 1,3-dinitrobenzene TS and 0.5 mL of a solution of sodium hydroxide (1 in 8), and allow to stand: a purple to red-purple color develops, and gradually changes to brown.

**Optical rotation** <2.49>  $[\alpha]_{\text{D}}^{20}$ : +29 – +32° (after drying, 0.2 g, 1,4-dioxane, 10 mL, 100 mm).

**Melting point** <2.60> 237 – 242°C

**Purity (1) Odor**—To 2.0 g of Purified Dehydrocholic Acid add 100 mL of water, and boil for 2 minutes: the solution is odorless.

(2) Clarity and color of solution—Dissolve 0.10 g of Purified Dehydrocholic Acid, previously powdered in a mortar, in 30 mL of ethanol (95) by shaking for 10 minutes: the solution is clear and colorless.

(3) Chloride <1.03>—To 2.0 g of Purified Dehydrocholic Acid add 100 mL of water, shake for 5 minutes and filter, and use this filtrate as the sample solution. To 25 mL of the sample solution add 6 mL of dilute nitric acid, heat in a water bath for 6 minutes, filter after cooling, and collect the clear filtrate. Wash the residue with 10 mL of water, combine the washings and the filtrate, dilute with water to 50 mL, and perform the test using this solution as the test solution. Prepare the control solution with 0.30 mL of 0.01 mol/L hydrochloric acid VS (not more than 0.021%).

(4) Sulfate <1.14>—Add 1 mL of dilute hydrochloric acid to 25 mL of the sample solution obtained in (3), heat in a water bath for 6 minutes, filter after cooling, and collect the clear filtrate. Wash the residue with 10 mL of water, combine the washings and the filtrate, dilute with water to 50 mL, and perform the test using this solution as the test solution. Prepare the control solution with 0.50 mL of 0.005 mol/L sulfuric acid VS (not more than 0.048%).

(5) Heavy metals <1.07>—Proceed with 1.0 g of Purified Dehydrocholic Acid according to Method 2, and perform the test. Prepare the control solution with 2.0 mL of Standard Lead Solution (not more than 20 ppm).

(6) Barium—To the solution obtained in (1) add 2 mL of hydrochloric acid, and boil for 2 minutes, cool, filter, and wash the filter with water until 100 mL of the filtrate is obtained. To 10 mL of the filtrate add 1 mL of dilute sulfuric acid: no turbidity is produced.

**Loss on drying** <2.41> Not more than 1.0% (1 g, 105°C, 2 hours).

**Residue on ignition** <2.44> Not more than 0.2% (1 g).

**Assay** Weigh accurately about 0.5 g of Purified Dehydrocholic Acid, previously dried, add 40 mL of neutralized ethanol and 20 mL of water, and dissolve by warming. Add 2 drops of phenolphthalein TS, then titrate <2.50> with 0.1 mol/L sodium hydroxide VS, adding 100 mL of freshly boiled and cooled water as the end point is approached, and continue the titration.

Each mL of 0.1 mol/L sodium hydroxide VS  
= 40.25 mg of  $\text{C}_{24}\text{H}_{34}\text{O}_5$

**Containers and storage** Containers—Well-closed containers.

## Dehydrocholic Acid Injection

### Dehydrocholate Sodium Injection

デヒドロコール酸注射液

Dehydrocholic Acid Injection is an aqueous solution for injection.

It contains not less than 95% and not more than 105% of the labeled amount of dehydrocholic acid ( $\text{C}_{24}\text{H}_{34}\text{O}_5$ ; 402.52).

**Method of preparation** Dissolve Purified Dehydrocholic Acid in a solution of Sodium Hydroxide, and prepare as directed under Injections.

**Description** Dehydrocholic Acid Injection is a clear, colorless to light yellow liquid, and has a bitter taste.

pH: 9 – 11

**Identification** Transfer a volume of Dehydrocholic Acid Injection, equivalent to 0.1 g of Purified Dehydrocholic Acid according to the labeled amount, to a separator, and add 10 mL of water and 1 mL of dilute hydrochloric acid: a white precipitate is produced. Extract the mixture with three 15-mL portions of chloroform, combine all the chloroform extracts, evaporate the chloroform on a water bath, and dry the residue at 105°C for 1 hour: the residue so obtained melts <2.60> between 235°C and 242°C.

**Purity** Heavy metals <1.07>—Evaporate a volume of Dehydrocholic Acid Injection, equivalent to 1.0 g of Purified Dehydrocholic Acid according to the labeled amount, on a water bath to dryness. Proceed with the residue according to Method 2, and perform the test. Prepare the control solution with 2.0 mL of Standard Lead Solution (not more than 20 ppm).

**Bacterial endotoxins** <4.01> Less than 0.30 EU/mg.

**Extractable volume** <6.05> It meets the requirement.

**Assay** Transfer an exactly measured volume of Dehydrocholic Acid Injection, equivalent to about 0.5 g of dehydrocholic acid ( $\text{C}_{24}\text{H}_{34}\text{O}_5$ ), to a 100-mL separator, and add, if necessary, water to make 25 mL. Add 2 mL of hydrochloric acid, and extract with 25-mL, 20-mL and 15-mL portions of chloroform successively. Combine the chloroform extracts, wash with cold water until the washings become negative to acid, and evaporate the chloroform on a water bath. Dissolve the residue in 40 mL of neutralized ethanol and 20 mL of water by warming. Add 2 drops of phenolphthalein TS to this solution, titrate <2.50> with 0.1 mol/L sodium hydroxide VS, adding 100 mL of freshly boiled and cooled water as the end point is approached, and continue the titration.


Each mL of 0.1 mol/L sodium hydroxide VS  
= 40.25 mg of  $\text{C}_{24}\text{H}_{34}\text{O}_5$

**Containers and storage** Containers—Hermetic containers, and colored containers may be used.

Storage—Light-resistant.

## Demethylchlortetracycline Hydrochloride

デメチルクロルテトラサイクリン塩酸塩



$C_{21}H_{21}ClN_2O_8 \cdot HCl$ : 501.31  
(4S,4aS,5aS,6S,12aS)-7-Chloro-4-dimethylamino-3,6,10,12,12a-pentahydroxy-1,11-dioxo-1,4,4a,5,5a,6,11,12a-octahydrotetracycline-2-carboxamide monohydrochloride [64-73-3]

Demethylchlortetracycline Hydrochloride is the hydrochloride of a tetracycline substance having antibacterial activity produced by the growth of the mutant of *Streptomyces aureofaciens*.

It contains not less than 900  $\mu$ g (potency) and not more than 1010  $\mu$ g (potency) per mg, calculated on the dried basis. The potency of Demethylchlortetracycline Hydrochloride is expressed as mass (potency) of demethylchlortetracycline hydrochloride ( $C_{21}H_{21}ClN_2O_8 \cdot HCl$ ).

**Description** Demethylchlortetracycline Hydrochloride occurs as a yellow crystalline powder.

It is soluble in water, and slightly soluble in ethanol (99.5).

**Identification (1)** Dissolve 40 mg of Demethylchlortetracycline Hydrochloride in 250 mL of water. To 10 mL of this solution add 85 mL of water and 5 mL of a solution of sodium hydroxide (1 in 5). Determine the absorption spectrum of this solution as directed under Ultraviolet-visible Spectrophotometry <2.24>, and compare the spectrum with the Reference Spectrum or the spectrum of a solution of Demethylchlortetracycline Hydrochloride Reference Standard prepared in the same manner as the sample solution: both spectra exhibit similar intensities of absorption at the same wavelengths.

(2) Determine the infrared absorption spectrum of Demethylchlortetracycline Hydrochloride as directed in the potassium chloride disk method under Infrared Spectrophotometry <2.25>, and compare the spectrum with the Reference Spectrum or the spectrum of Demethylchlortetracycline Hydrochloride Reference Standard: both spectra exhibit similar intensities of absorption at the same wave numbers.

(3) A solution of Demethylchlortetracycline Hydrochloride (1 in 100) responds to the Qualitative Tests <1.09> (2) for chloride.

**Optical rotation** <2.49>  $[\alpha]_D^{20}$ :  $-248 - -263^\circ$  (0.25 g calculated on the dried basis, 0.1 mol/L hydrochloric acid TS, 25 mL, 100 mm).

**pH** <2.54> Dissolve 1.0 g of Demethylchlortetracycline Hydrochloride in 100 mL of water: the pH of the solution is between 2.0 and 3.0.

**Purity (1)** Heavy metals <1.07>—Proceed with 1.0 g of

Demethylchlortetracycline Hydrochloride according to Method 2, and perform the test. Prepare the control solution with 2.0 mL of Standard Lead Solution (not more than 20 ppm).

(2) Arsenic <1.11>—Prepare the test solution with 1.0 g of Demethylchlortetracycline Hydrochloride according to Method 4, and perform the test (not more than 2 ppm).

(3) Related substances—Dissolve 25 mg of Demethylchlortetracycline Hydrochloride in 50 mL of 0.01 mol/L hydrochloric acid TS, and use this solution as the sample solution. Pipet 5 mL of the sample solution, add 0.01 mol/L hydrochloric acid TS to make exactly 100 mL, and use this solution as the standard solution. Perform the test with exactly 20  $\mu$ L each of the sample solution and standard solution as directed under Liquid Chromatography <2.01> according to the following conditions. Determine each peak area obtained from the chromatograms of these solutions by the automatic integration method: the peak area other than demethylchlortetracycline obtained from the sample solution is not more than 6/5 times that of demethylchlortetracycline from the standard solution, and the sum of the areas of the peaks other than demethylchlortetracycline from the sample solution is not more than 2 times the peak area of demethylchlortetracycline from the standard solution.

**Operating conditions**—

Detector, column, column temperature, mobile phase, and flow rate: Proceed as directed in the operating conditions in the Assay.

Time span of measurement: About 2 times as long as the retention time of demethylchlortetracycline beginning after the solvent peak.

**System suitability**—

Test for required detectability: Measure exactly 10 mL of the standard solution, add 0.01 mol/L hydrochloric acid TS to make exactly 50 mL, and use this solution as the solution for system suitability test. Pipet 5 mL of the solution for system suitability test, and add 0.01 mol/L hydrochloric acid TS to make exactly 50 mL. Confirm that the peak area of demethylchlortetracycline obtained from 20  $\mu$ L of this solution is equivalent to 7 to 13% of that from 20  $\mu$ L of the solution for system suitability test.

System performance: Proceed as directed in the system suitability in the Assay.

System repeatability: When the test is repeated 6 times with 20  $\mu$ L of the standard solution under the above operating conditions, the relative standard deviation of the peak area of demethylchlortetracycline is not more than 1.0%.

**Loss on drying** <2.41> Not more than 2.0% (1 g, in vacuum, 60°C, 3 hours).

**Residue on ignition** <2.44> Not more than 0.2% (1 g).

**Assay** Weigh accurately an amount of Demethylchlortetracycline Hydrochloride and Demethylchlortetracycline Hydrochloride Reference Standard, equivalent to about 25 mg (potency), dissolve each in 0.01 mol/L hydrochloric acid TS to make exactly 50 mL, and use these solutions as the sample solution and standard solution, respectively. Perform the test with exactly 20  $\mu$ L each of the sample solution and standard solution as directed under Liquid Chromatography <2.01> according to the following conditions, and determine the peak areas,  $A_T$  and  $A_S$ , of demethylchlortetracycline.

Amount [ $\mu$ g (potency)] of  $C_{21}H_{21}ClN_2O_8 \cdot HCl$

$$= W_S \times (A_T/A_S) \times 1000$$

$W_S$ : Amount [mg (potency)] of Demethylchlortetracycline Hydrochloride Reference Standard

#### Operating conditions—

**Detector:** An ultraviolet absorption photometer (wavelength: 254 nm).

**Column:** A stainless steel column 4.1 mm in inside diameter and 25 cm in length, packed with styrene-divinylbenzene copolymer for liquid chromatography (10  $\mu$ m in particle diameter).

**Column temperature:** A constant temperature of about 60°C.

**Mobile phase:** Dissolve 3.5 g of dipotassium hydrogen phosphate, 1.5 g of tetrabutylammonium hydrogensulfate and 0.4 g of disodium dihydrogen ethylenediamine tetraacetate dihydrate in 300 mL of water, and adjust the pH to 8.5 with sodium hydroxide TS. To this solution add 75.0 g of *t*-butanol and water to make 1000 mL.

**Flow rate:** Adjust the flow rate so that the retention time of demethylchlortetracycline is about 8 minutes.

#### System suitability—

**System performance:** Heat 10 mL of the standard solution on a water bath for 60 minutes. When the procedure is run with 20  $\mu$ L of this solution so obtained under the above operating conditions, 4-epidemethylchlortetracycline and demethylchlortetracycline are eluted in this order with the resolution between these peaks being not less than 3. The relative retention time of 4-epidemethylchlortetracycline with respect to demethylchlortetracycline is about 0.7.


**System repeatability:** When the test is repeated 6 times with 20  $\mu$ L of the standard solution under the above operating conditions, the relative standard deviation of the peak area of demethylchlortetracycline is not more than 1.0%.

**Containers and storage** Containers—Tight containers.

Storage—Light-resistant.

## Deslanoside

デスラノシド



$C_{47}H_{74}O_{19}$ : 943.08

3 $\beta$ -[ $\beta$ -D-Glucopyranosyl-(1 $\rightarrow$ 4)-2,6-dideoxy-

$\beta$ -D-ribo-hexopyranosyl-(1 $\rightarrow$ 4)-2,6-dideoxy-  
 $\beta$ -D-ribo-hexopyranosyl-(1 $\rightarrow$ 4)-2,6-dideoxy- $\beta$ -D-ribo-  
hexopyranosyloxy]-12 $\beta$ ,14-dihydroxy-5 $\beta$ ,14 $\beta$ -card-  
20(22)-enolide [17598-65-1]

Deslanoside, when dried, contains not less than 90.0% and not more than 102.0% of  $C_{47}H_{74}O_{19}$ .

**Description** Deslanoside occurs as colorless or white crystals or a white, crystalline powder. It is odorless.

It is freely soluble in anhydrous pyridine, sparingly soluble in methanol, slightly soluble in ethanol (95), and practically insoluble in water and in diethyl ether.

It is hygroscopic.

**Identification** Transfer 1 mg of Deslanoside to a small test tube about 10 mm in inside diameter, dissolve in 1 mL of a solution of iron (III) chloride hexahydrate in acetic acid (100) (1 in 1000), and underlay gently with 1 mL of sulfuric acid: at the zone of contact of two liquids a brown ring is produced, and the color of the upper layer near to the contact zone changes gradually to blue through purple, and the entire acetic acid layer shows a blue-green color through a deep blue color.

**Purity (1)** Clarity and color of solution—Dissolve 20 mg of Deslanoside in 10 mL of ethanol (95) and 3 mL of water by warming, cool, and dilute to 100 mL with water: the solution is clear and colorless.

**(2)** Related substances—Dissolve 10 mg of Deslanoside in exactly 5 mL of methanol, and use this solution as the sample solution. Dissolve 1.0 mg of Deslanoside Reference Standard in exactly 5 mL of methanol, and use this solution as the standard solution. Perform the test with these solutions as directed under Thin-layer Chromatography <2.03>. Spot 20  $\mu$ L each of the sample solution and standard solution on a plate of silica gel for thin-layer chromatography. Develop the plate with a mixture of dichloromethane, methanol and water (84:15:1) to a distance of about 13 cm, and air-dry the plate. Spray evenly dilute sulfuric acid on the plate, and heat the plate at 110°C for 10 minutes: the spots other than the principal spot from the sample solution are not larger than and not more intense than the spot from the standard solution.

**Optical rotation** <2.49>  $[\alpha]_D^{20}$ : +6.5 – +8.5° (after drying, 0.5 g, anhydrous pyridine, 25 mL, 100 mm).

**Loss on drying** <2.41> Not more than 8.0% (0.5 g, in vacuum, phosphorus (V) oxide, 60°C, 4 hours).

**Residue on ignition** <2.44> Not more than 0.5% (0.1 g).

**Assay** Dissolve about 12 mg each of Deslanoside and Deslanoside Reference Standard, previously dried and accurately weighed, in 20 mL each of methanol, add water to make exactly 100 mL, and use these solutions as the sample solution and the standard solution, respectively. Pipet 5 mL each of these solutions, transfer to light-resistant, 25-mL volumetric flasks, shake well with 5 mL each of 2,4,6-trinitrophenol TS and 0.5 mL each of a solution of sodium hydroxide (1 in 10), add diluted methanol (1 in 4) to make 25 mL, and allow to stand at a temperature between 18°C and 22°C for 25 minutes. Determine the absorbances,  $A_T$  and  $A_S$ , of the subsequent solutions of the sample solution and standard solution, respectively, at 485 nm as directed under Ultraviolet-visible Spectrophotometry <2.24>, using a solution prepared

with 5 mL of diluted methanol (1 in 5) in the same manner as the blank.

$$\begin{aligned} & \text{Amount (mg) of } C_{47}H_{74}O_{19} \\ & = W_S \times (A_T/A_S) \end{aligned}$$

$W_S$ : Amount (mg) of Deslanoside Reference Standard

**Containers and storage** Containers—Tight containers.

## Deslanoside Injection

デスラノシド注射液

Deslanoside Injection is an aqueous solution for injection.

It contains not less than 90% and not more than 110% of the labeled amount of deslanoside ( $C_{47}H_{74}O_{19}$ ; 943.08).

**Method of preparation** Dissolve Deslanoside in 10 vol% ethanol and prepare as directed under Injections. It may contain Glycerin. It may be prepared with a suitable amount of Ethanol and Water for Injection.

**Description** Deslanoside Injection is a clear and colorless liquid.

pH: 5.0 – 7.0

**Identification (1)** Place a volume of Deslanoside Injection, equivalent to 2 mg of Deslanoside according to the labeled amount, in a separator, add sodium chloride in the ratio of 0.2 g to each mL of this solution, and extract with three 10-mL portions of chloroform. Combine the chloroform extracts, mix uniformly, pipet 15 mL of this solution, and evaporate the chloroform under reduced pressure. Proceed with the residue as directed in the Identification under Deslanoside.

(2) Place a volume of Deslanoside Injection, equivalent to 1 mg of Deslanoside according to the labeled amount, in a separator, add sodium chloride in the ratio of 0.2 g to each mL of this solution, and extract with three 5-mL portions of chloroform. Combine the chloroform extracts, evaporate the chloroform under reduced pressure, dissolve the residue in 5 mL of methanol, and use this solution as the sample solution. Separately, dissolve 1 mg of Deslanoside Reference Standard in 5 mL of methanol, and use this solution as the standard solution. Perform the test with these solutions as directed under Thin-layer Chromatography <2.03>. Spot 20  $\mu$ L each of these solutions on a plate of silica gel for thin-layer chromatography. Develop the plate with a mixture of dichloromethane, methanol and water (84:15:1) to a distance of about 13 cm, and air-dry the plate. Spray evenly dilute sulfuric acid upon the plate, and heat the plate at 110°C for 10 minutes: the spots from the sample solution and the standard solution show a black color and have the same  $R_f$  value.

**Extractable volume** <6.05> It meets the requirement.

**Assay** Measure exactly a volume of Deslanoside Injection, equivalent to about 3 mg of deslanoside ( $C_{47}H_{74}O_{19}$ ). Add 5 mL of methanol and water to make 25 mL. Use this solution as the sample solution, and proceed as directed in the Assay under Deslanoside.

$$\text{Amount (mg) of deslanoside } (C_{47}H_{74}O_{19})$$


$$= W_S \times (A_T/A_S) \times (1/4)$$

$W_S$ : Amount (mg) of Deslanoside Reference Standard

**Containers and storage** Containers—Hermetic containers.  
Storage—Light-resistant.

## Dexamethasone

デキサメタゾン



$C_{22}H_{29}FO_5$ ; 392.46

9-Fluoro-11 $\beta$ ,17,21-trihydroxy-16 $\alpha$ -methylpregna-1,4-diene-3,20-dione [50-02-2]

Dexamethasone, when dried, contains not less than 97.0% and not more than 102.0% of  $C_{22}H_{29}FO_5$ .

**Description** Dexamethasone occurs as white to pale yellow crystals or crystalline powder.

It is sparingly soluble in methanol, in ethanol (95) and in acetone, slightly soluble in acetonitrile, and practically insoluble in water.

Melting point: about 245°C (with decomposition).

**Identification (1)** Proceed with 10 mg of Dexamethasone as directed under Oxygen Flask Combustion Method, using a mixture of 0.5 mL of 0.01 mol/L sodium hydroxide TS and 20 mL of water as the absorbing liquid: the solution obtained responds to the Qualitative Tests <1.09> for fluoride

(2) Dissolve 1 mg of Dexamethasone in 10 mL of ethanol (95). Mix 2 mL of the solution with 10 mL of phenylhydrazine hydrochloride TS, heat in a water bath at 60°C for 20 minutes, and cool the solution. Determine the absorption spectrum of the solution as directed under Ultraviolet-visible Spectrophotometry <2.24>, using as the blank the solution prepared with 2 mL of ethanol (95) in the same manner as the former solution, and compare the spectrum with the Reference Spectrum or the spectrum of a solution of Dexamethasone Reference Standard prepared in the same manner as the sample solution: both spectra exhibit similar intensities of absorption at the same wavelengths.

(3) Determine the infrared absorption spectrum of Dexamethasone, previously dried, as directed in the potassium bromide disk method under Infrared Spectrophotometry <2.25>, and compare the spectrum with the Reference Spectrum or the spectrum of previously dried Dexamethasone Reference Standard: both spectra exhibit similar intensities of absorption at the same wave numbers. If any difference appears between the spectra, dissolve Dexamethasone and Dexamethasone Reference Standard in acetone, respectively, then evaporate the acetone to dryness, and repeat the test on the residues.

**Optical rotation** <2.49>  $[\alpha]_D^{20}$ : +86 – +94° (after drying, 0.1 g, methanol, 10 mL, 100 mm).

**Purity (1)** Heavy metals <1.07>—Proceed with 1.0 g of



Dexamethasone according to Method 2, and perform the test. Prepare the control solution with 3.0 mL of Standard Lead Solution (not more than 30 ppm).

(2) Related substances—Dissolve 0.18 g of Dexamethasone in 100 mL of acetonitrile. To 33 mL of this solution add a solution, prepared by dissolving 1.32 g of ammonium formate in water to make 1000 mL and adjusted to pH 3.6 with formic acid, to make 100 mL, and use this solution as the sample solution. To exactly 1 mL of the sample solution add the mobile phase to make exactly 100 mL, and use this solution as the standard solution. Perform the test with exactly 10  $\mu$ L each of the sample solution and standard solution as directed under Liquid Chromatography <2.01> according to the following conditions, and determine each peak area by the automatic integration method: the peak area other than dexamethasone is not larger than the peak area of dexamethasone obtained from the standard solution, and the total area of the peaks other than the peak of dexamethasone from the sample solution is not larger than 2 times the peak area of dexamethasone from the standard solution

**Operating conditions—**

Detector: An ultraviolet absorption photometer (wavelength: 254 nm).

Column: A stainless steel column 4.6 mm in inside diameter and 25 cm in length, packed with phenylsilylanized silica gel for liquid chromatography (5  $\mu$ m in particle diameter).

Column temperature: A constant temperature of about 25°C.

Mobile phase: Dissolve 1.32 g of ammonium formate in 1000 mL of water, and adjust the pH to 3.6 with formic acid. To 670 mL of this solution add 330 mL of acetonitrile.

Flow rate: Adjust the flow rate so that the retention time of dexamethasone is about 13 minutes.

Time span of measurement: About 4 times as long as the retention time of dexamethasone beginning after the solvent peak.

**System suitability—**

Test for required detectability: To exactly 1 mL of the standard solution add the mobile phase to make exactly 10 mL. Confirm that the peak area of dexamethasone obtained with 10  $\mu$ L of this solution is equivalent to 8 to 12% of that with 10  $\mu$ L of the standard solution.

System performance: When the procedure is run with 10  $\mu$ L of the standard solution under the above operating conditions, the number of theoretical plates and the symmetry factor of the peak of dexamethasone are not less than 5000 and not more than 1.5, respectively.

System repeatability: When the test is repeated 6 times with 10  $\mu$ L of the standard solution under the above operating conditions, the relative standard deviation of the peak area of dexamethasone is not more than 1.0%.

**Loss on drying** <2.41> Not more than 0.5% (0.2 g, 105°C, 3 hours).

**Residue on ignition** <2.44> Not more than 0.1% (0.2 g, platinum crucible).

**Assay** Dissolve about 10 mg each of Dexamethasone and Dexamethasone Reference Standard, previously dried and accurately weighed, in 70 mL each of diluted methanol (1 in 2), add exactly 5 mL each of the internal standard solution, then add diluted methanol (1 in 2) to make 100 mL, and use these solutions as the sample solution and standard solution.

Perform the test with 10  $\mu$ L each of these solutions as directed under Liquid Chromatography <2.01> according to the following conditions, and calculate the ratios,  $Q_T$  and  $Q_S$ , of the peak area of dexamethasone to that of the internal standard, respectively.

$$\text{Amount (mg) of } C_{22}H_{29}FO_5 = W_S \times Q_T / Q_S$$

$W_S$ : Amount (mg) of Dexamethasone Reference Standard

**Internal standard solution**—A solution of propyl parahydroxybenzoate in diluted methanol (1 in 2) (1 in 1000).

**Operating conditions—**

Detector: An ultraviolet absorption photometer (wavelength: 254 nm).

Column: A stainless steel column 4 mm in inside diameter and 30 cm in length, packed with octadecylsilylanized silica gel for liquid chromatography (10  $\mu$ m in particle diameter).

Column temperature: A constant temperature of about 25°C.

Mobile phase: A mixture of water and acetonitrile (2:1).

Flow rate: Adjust the flow rate so that the retention time of dexamethasone is about 6 minutes.

**System suitability—**

System performance: When the procedure is run with 10  $\mu$ L of the standard solution under the above operating conditions, dexamethasone and the internal standard are eluted in this order with the resolution between these peaks being not less than 6.

System repeatability: When the test is repeated 6 times with 10  $\mu$ L of the standard solution under the above operating conditions, the relative standard deviation of the ratio of the peak area of dexamethasone to that of the internal standard is not more than 1.0%.

**Containers and storage** Containers—Tight containers.

Storage—Light-resistant.

## Dextran 40

### デキストラン 40

Dextran 40 is a product obtained by partial decomposition of polysaccharide, which is produced by fermentation of sucrose with *Leuconostoc mesenteroides* van Tieghem (*Lactobacillaceae*), and the average molecular mass is about 40,000.

When dried, it contains not less than 98.0% and not more than 102.0% of dextran 40.

**Description** Dextran 40 occurs as a white, amorphous powder. It is odorless and tasteless.

It is practically insoluble in ethanol (95) and in diethyl ether.

It dissolves gradually in water.

It is hygroscopic.

**Identification** To 1 mL of a solution of Dextran 40 (1 in 3000) add 2 mL of anthrone TS: a blue-green color develops and turns gradually dark blue-green. Then to this solution add 1 mL of diluted sulfuric acid (1 in 2) or 1 mL of acetic acid (100): the solution does not change in color.

**pH** <2.54> Dissolve 1.0 g of Dextran 40 in 10 mL of water:

the pH of this solution is between 5.0 and 7.0.

**Purity** (1) Clarity and color of solution—Dissolve 1.0 g of Dextran 40 in 10 mL of water by warming; the solution is clear and colorless.

(2) Chloride  $\langle 1.03 \rangle$ —Perform the test with 2.0 g of Dextran 40. Prepare the control solution with 1.0 mL of 0.01 mol/L hydrochloric acid (not more than 0.018%).

(3) Heavy metals  $\langle 1.07 \rangle$ —Proceed with 1.0 g of Dextran 40 according to Method 1, and perform the test. Prepare the control solution with 2.0 mL of Standard Lead Solution (not more than 20 ppm).

(4) Arsenic  $\langle 1.11 \rangle$ —Prepare the test solution with 1.5 g of Dextran 40 according to Method 1, and perform the test (not more than 1.3 ppm).

(5) Nitrogen—Weigh accurately about 2 g of Dextran 40, previously dried, and perform the test as directed under Nitrogen Determination  $\langle 1.08 \rangle$ , where 10 mL of sulfuric acid is used for decomposition, and 45 mL of a solution of sodium hydroxide (2 in 5) is added; the amount of nitrogen (N: 14.01) is not more than 0.010%.

(6) Reducing substances—Weigh exactly 3.00 g of Dextran 40, previously dried, dissolve in water to make exactly 50 mL, and use this solution as the sample solution. Separately, weigh exactly 0.450 g of glucose, previously dried, dissolve in water to make exactly 500 mL, and use this solution as the control solution. Pipet 5 mL each of the sample solution and the control solution, and add water to make exactly 50 mL, respectively. Pipet 5 mL each of these solutions, add 5 mL of alkaline copper TS, exactly measured, and heat for 15 minutes in a water bath. After cooling, add 1 mL of a solution of potassium iodine (1 in 40) and 1.5 mL of dilute sulfuric acid, and titrate  $\langle 2.50 \rangle$  with 0.005 mol/L sodium thiosulfate VS (indicator: 2 mL of starch TS).

The titrant consumed for the sample solution is not less than that for the control solution.

**Loss on drying**  $\langle 2.41 \rangle$  Not more than 5.0% (1 g, 105°C, 6 hours).

**Residue on ignition**  $\langle 2.44 \rangle$  Not more than 0.1% (1 g).

**Viscosity**  $\langle 2.53 \rangle$  (1) Dextran 40—Weigh accurately 0.2 to 0.5 g of Dextran 40, previously dried, dissolve in water to make exactly 100 mL, and use this solution as the sample solution. Perform the test with the sample solution and with water as directed in Method 1 at 25°C: the intrinsic viscosity is between 0.16 and 0.19.

(2) High-molecular fraction—Weigh accurately about 6 g of Dextran 40, previously dried, dissolve in water to make exactly 100 mL, and transfer to a flask. Add slowly enough methanol to get 7% to 10% of the precipitate (usually 80 to 90 mL) at  $25 \pm 1^\circ\text{C}$  with stirring. Dissolve the precipitate at 35°C in a water bath with occasional shaking, and allow to stand for more than 15 hours at  $25 \pm 1^\circ\text{C}$ . Remove the supernatant liquid by decantation, and heat the precipitate of the lower layer to dryness on a water bath. Dry the residue, and determine the intrinsic viscosity of the dried substance as directed in (1): the value is not more than 0.27.

(3) Low-molecular fraction—Weigh accurately about 6 g of Dextran 40, previously dried, dissolve in water to make exactly 100 mL, and transfer to a flask. Add slowly enough methanol to get 90% to 93% of the precipitate (usually 115 to 135 mL) at  $25 \pm 1^\circ\text{C}$  with stirring, centrifuge at 25°C, and evaporate the supernatant liquid to dryness on a water bath.

Dry the residue, and determined the intrinsic viscosity of the dried substance as directed in (1): the value is not less than 0.09.

**Antigenicity** Dissolve 10.0 g of Dextran 40 in isotonic sodium chloride solution to make 100 mL, sterilize, and use this solution as the sample solution. Inject 1.0 mL of the sample solution on 3 occasions at intervals of 2 days into the peritoneal cavity of each of 4 well-nourished, healthy guinea pigs weighing 250 to 300 g. Inject 0.10 mL of horse serum into the peritoneal cavity of each of 4 guinea pigs of another group as a control. Inject 0.20 mL of the sample solution intravenously to each of 2 guinea pigs of the first group 14 days after the first intraperitoneal injection and into each of the remaining 2 guinea pigs 21 days after the injection, and inject 0.20 mL of horse serum intravenously in the same manner into each guinea pig of the second group. Observe the signs of respiratory distress, collapse or death of the animals for 30 minutes after each intravenous injection and 24 hours later: the animals of the first group exhibit no signs mentioned above.

All the animals of the second group exhibit symptoms of respiratory distress or collapse and not less than 3 animals are killed.

**Pyrogen**  $\langle 4.04 \rangle$  Dissolve 10.0 g of Dextran 40 in isotonic sodium chloride solution to make 100 mL, and perform the test: this solution meets the requirement.

**Assay** Weigh accurately about 3 g of Dextran 40, previously dried, dissolve in water to make exactly 50 mL, and use this solution as the sample solution. Determine the optical rotation  $\alpha_D$  with the sample solution as directed under Optical Rotation Determination  $\langle 2.49 \rangle$  in a 100-mL cell at  $20 \pm 1^\circ\text{C}$ .

$$\text{Amount (mg) of dextran 40} = \alpha_D \times 253.8$$

**Containers and storage** Containers—Tight containers.

## Dextran 40 Injection

### デキストラン 40 注射液

Dextran 40 Injection is an aqueous solution for injection.

It contains not less than 9.5 w/v% and not more than 10.5 w/v% of dextran 40.

#### Method of preparation

|                                   |                       |
|-----------------------------------|-----------------------|
| Dextran 40                        | 10 g                  |
| Isotonic Sodium Chloride Solution | a sufficient quantity |
| To make 100 mL                    |                       |

Prepare as directed under Injections, with the above ingredients.

No preservative is added.

**Description** Dextran 40 Injection is a clear and colorless liquid. It is slightly viscous.

**Identification** (1) Dilute 1 mL of Dextran 40 Injection with water to 200 mL, and to 1 mL of the diluted solution add 2 mL of anthrone TS: a blue-green color develops and

turns gradually dark blue-green. Add 1 mL of diluted sulfuric acid (1 in 2) or 1 mL of acetic acid (100) to this solution: the solution does not change in color.

(2) Dextran 40 Injection responds to the Qualitative Tests <1.09> for sodium salt and for chloride.

**pH** <2.54> 4.5 – 7.0

**Bacterial endotoxins** <4.01> Less than 0.50 EU/mL.

**Extractable volume** <6.05> It meets the requirement.

**Viscosity** <2.53> Measure exactly 2 to 5 mL of Dextran 40 Injection, add isotonic sodium chloride solution to make exactly 100 mL, and use this solution as the sample solution. Perform the test with the sample solution and with isotonic sodium chloride solution as directed in Method 1 at 25°C: the intrinsic viscosity is between 0.16 and 0.19. Calculate the concentration of the sample solution (g/100 mL) as directed in the Assay.

**Assay** To exactly 30 mL of Dextran 40 Injection add water to make exactly 50 mL, and use this solution as the sample solution. Determine the optical rotation  $\alpha_D$  with the sample solution as directed under Optical Rotation Determination <2.49> in a 100-mL cell at  $20 \pm 1^\circ\text{C}$ .

$$\begin{aligned} &\text{Amount (mg) of dextran 40 in 100 mL of} \\ &\text{Dextran 40 Injection} \\ &= \alpha_D \times 846.0 \end{aligned}$$

**Containers and storage** Containers—Hermetic containers. Plastic containers for aqueous injections may be used.

Storage—Avoid exposure to undue fluctuations in temperature.

## Dextran 70

デキストラン 70

Dextran 70 is a product obtained by partial decomposition of polysaccharide, which is produced by fermentation of sucrose with *Leuconostoc mesenteroides* van Tieghem (*Lactobacillaceae*), and the average molecular mass is about 70,000.

When dried, it contains not less than 98.0% and not more than 102.0% of dextran 70.

**Description** Dextran 70 occurs as a white, amorphous powder. It is odorless and tasteless.

It is practically insoluble in ethanol (95) and in diethyl ether.

It dissolves gradually in water.

It is hygroscopic.

**Identification** To 1 mL of a solution of Dextran 70 (1 in 3000) add 2 mL of anthrone TS: a blue-green color develops and turns gradually dark blue-green. Then to this solution add 1 mL of diluted sulfuric acid (1 in 2) or 1 mL of acetic acid (100): the solution does not change in color.

**pH** <2.54> Dissolve 3.0 g of Dextran 70 in 50 mL of water: the pH of this solution is between 5.0 and 7.0.

**Purity** (1) Clarity and color of solution—Dissolve 1.0 g of Dextran 70 in 10 mL of water with warming: the solution is

clear and colorless.

(2) Chloride <1.03>—With 2.0 g of Dextran 70, perform the test. Prepare the control solution with 1.0 mL of 0.01 mol/L hydrochloric acid VS (not more than 0.018%).

(3) Heavy metals <1.07>—Proceed with 1.0 g of Dextran 70 according to Method 1, and perform the test. Prepare the control solution with 2.0 mL of Standard Lead Solution (not more than 20 ppm).

(4) Arsenic <1.11>—Prepare the test solution with 1.5 g of Dextran 70 according to Method 1, and perform the test (not more than 1.3 ppm).

(5) Nitrogen—Weigh accurately about 2 g of Dextran 70, previously dried, perform the test as directed under Nitrogen Determination <1.08>, where 10 mL of sulfuric acid is used for decomposition, and 45 mL of a solution of sodium hydroxide (2 in 5) is added: the amount of nitrogen (N: 14.007) is not more than 0.010%.

(6) Reducing substances—Weigh exactly 3.00 g of Dextran 70, previously dried, dissolve in water to make exactly 50 mL, and use this solution as the sample solution. Separately, weigh exactly 0.300 g of glucose, previously dried, dissolve in water to make exactly 500 mL, and use this solution as the control solution. Pipet 5 mL each of the sample solution and the control solution, and add water to make exactly 50 mL, respectively. Pipet 5 mL of these diluted solutions, add exactly 5 mL of alkaline copper TS, and heat for 15 minutes in a water bath. After cooling, add 1 mL of a solution of potassium iodide (1 in 40) and 1.5 mL of dilute sulfuric acid, and titrate <2.50> with 0.005 mol/L sodium thiosulfate VS (indicator: 2 mL of starch TS).

The titrant consumed for the sample solution is not less than that for the control solution.

**Loss on drying** <2.41> Not more than 5.0% (1 g, 105°C, 6 hours).

**Residue on ignition** <2.44> Not more than 0.10% (1 g).

**Viscosity** <2.53> (1) Dextran 70—Weigh accurately 0.2 to 0.5 g of Dextran 70, previously dried, dissolve in water to make exactly 100 mL, and use this solution as the sample solution. Perform the test with the sample solution and with water as directed in Method 1 at 25°C: the intrinsic viscosity is between 0.21 and 0.26.

(2) High-molecular fraction—Weigh accurately about 6 g of Dextran 70, previously dried, dissolve in water to make exactly 100 mL, and transfer to a flask. Add slowly enough methanol to get 7% to 10% of the precipitate (usually, 75 to 85 mL) at  $25 \pm 1^\circ\text{C}$  with stirring. Dissolve the precipitate in a water bath at 35°C with occasional shaking, and allow to stand for more than 15 hours at  $25 \pm 1^\circ\text{C}$ . Remove the supernatant liquid by decantation, and heat the precipitate of the lower layer on a water bath to dryness. Dry the residue, and determine the intrinsic viscosity of the dried residue as directed in (1): the value is not more than 0.35.

(3) Low-molecular fraction—Weigh accurately about 6 g of Dextran 70, previously dried, dissolve in water to make exactly 100 mL, and transfer to a flask. Add slowly enough methanol to get 90% to 93% of the precipitate (usually 110 to 130 mL) at  $25 \pm 1^\circ\text{C}$  with stirring, centrifuge at 25°C, and evaporate the supernatant liquid to dryness on a water bath. Dry the residue, and determine the intrinsic viscosity of the dried residue as directed in (1): the value is not less than 0.10.

**Antigenicity** Dissolve 6.0 g of Dextran 70 in isotonic sodi-

um chloride solution to make 100 mL, sterilize, and use this solution as the sample solution. Inject 1.0 mL of the sample solution on 3 occasions at intervals of 2 days into the peritoneal cavity of each of 4 well-nourished, healthy guinea pigs weighing 250 to 300 g. Separately, inject 0.10 mL of horse serum into the peritoneal cavity of each of 4 guinea pigs of another group as a control. Inject 0.20 mL of the sample solution intravenously to each of 2 guinea pigs of the first group 14 days after the first intraperitoneal injection and into each of the remaining 2 guinea pigs 21 days after the injection, and inject 0.20 mL of horse serum intravenously in the same manner into each guinea pig of the second group. Observe the signs of respiratory distress, collapse or death of the animals for 30 minutes after each intravenous injection and 24 hours later: the animals of the first group exhibit not signs.

All the animals of the second group exhibit symptoms of respiratory distress or collapse and not less than 3 animals are killed.

**Pyrogen** <4.04> Dissolve 6.0 g of Dextran 70 in isotonic sodium chloride solution to make 100 mL, and perform the test: this solution meets the requirement.

**Assay** Weigh accurately about 3 g of Dextran 70, previously dried, dissolve in water to make exactly 50 mL, and use this solution as the sample solution. Determine the optical rotation  $\alpha_D$  as directed under Optical Rotation Determination <2.49> in a 100-mL cell at  $20 \pm 1^\circ\text{C}$ .

$$\text{Amount (mg) of dextran 70} \times \alpha_D = 253.8$$

**Containers and storage** Containers—Tight containers.

## Dextran Sulfate Sodium Sulfur 5

デキストラン硫酸エステルナトリウム イオウ 5

Dextran Sulfate Sodium Sulfur 5 is a sodium salt of sulfate ester obtained by sulfation of partial decomposition products of dextran, which is produced by fermentation of sucrose with *Leuconostoc mesenteroides* Van Tieghem (*Lactobacillaceae*).

**Description** Dextran Sulfate Sodium Sulfur 5 occurs as a white to light yellowish white powder. It is odorless, and has a saline taste.

It is freely soluble in water and practically insoluble in ethanol (95) and in diethyl ether.

It is hygroscopic.

**Identification** (1) To 10 mL of a solution of toluidine blue (1 in 100,000) add 0.05 mL of a solution of Dextran Sulfate Sodium Sulfur 5 (3 in 50) dropwise: a color of the solution changes from blue to red-purple.

(2) To 1 mL of a solution of Dextran Sulfate Sodium Sulfur 5 (1 in 1500) add 2 mL of anthrone TS: a blue-green color develops, which turns dark blue-green gradually. Then, add 1 mL of diluted sulfuric acid (1 in 2) or 1 mL of acetic acid (100) to this solution: the solution remains dark blue-green.

(3) A solution of Dextran Sulfate Sodium Sulfur 5 (1 in 100) responds to the Qualitative Tests <1.09> (1) for sodium salt.

**Optical rotation** <2.49>  $[\alpha]_D^{20}$ :  $+135.0 - +155.0^\circ$  (calculated

ed on the dried basis, 1.5 g, water, 25 mL, 100 mm).

**pH** <2.54> Dissolve 1.0 g of Dextran Sulfate Sodium Sulfur 5 in 20 mL of water: the pH of this solution is between 5.5 and 7.5.

**Purity** (1) Clarity of solution—Dissolve 2.5 g of Dextran Sulfate Sodium Sulfur 5 in 50 mL of water: the solution is clear. And, determine the absorbance of the solution at 420 nm as directed under Ultraviolet-visible Spectrophotometry <2.24>: not more than 0.090.

(2) Chloride <1.03>—Perform the test with 0.10 g of Dextran Sulfate Sodium Sulfur 5. Prepare the control solution with 0.30 mL of 0.01 mol/L hydrochloric acid (not more than 0.106%).

(3) Sulfate <1.14>—Dissolve 0.10 g of Dextran Sulfate Sodium Sulfur 5 in 6 mL of water, add 0.6 mL of barium chloride TS, and heat in a water bath for 4 minutes. After cooling, add 1 mL of dilute hydrochloric acid and water to make 50 mL, allow to stand for 10 minutes, and observe: the turbidity of the solution is not more intense than that of the control solution. Prepare the control solution as follows: to 0.50 mL of 0.005 mol/L sulfuric acid add 6 mL of water, and proceed in the same manner (not more than 0.240%).

(4) Heavy metals <1.07>—Proceed with 1.0 g of Dextran Sulfate Sodium Sulfur 5 according to Method 2, and perform the test. Prepare the control solution with 2.0 mL of Standard Lead Solution (not more than 20 ppm).

(5) Arsenic <1.11>—Prepare the test solution with 1.0 g of Dextran Sulfate Sodium Sulfur 5 according to Method 3, and perform the test (not more than 2 ppm).

**Sulfur content** Weigh accurately about 1.0 g of Dextran Sulfate Sodium Sulfur 5, dissolve in 5 mL of water, add 1.5 mL of hydrochloric acid, and heat in a water bath for 1 hour. After cooling, add water to make exactly 100 mL, and use this solution as the sample solution. To exactly 10 mL of the sample solution add exactly 20 mL of 0.02 mol/L barium chloride VS, add 5 mL of methanol, and heat in a water bath for 30 minutes. After cooling, neutralize with sodium hydroxide TS, and add 70 mL of water, 10 mL of a solution of zinc disodium ethylenediamine tetraacetate tetrahydrate (1 in 20), 3 mL of ammonium chloride TS and 7 mL of strong ammonium water, and titrate <2.50> with 0.02 mol/L disodium dihydrogen ethylenediamine tetraacetate VS until the color of the solution changes from red to light blue (indicator: 5 drops of eriochrome black T TS). Perform a blank determination. Amount of sulfur (S: 32.07), calculated on the dried basis, is between 3.0 and 6.0%.

$$\begin{aligned} \text{Each mL of 0.02 mol/L barium chloride VS} \\ = 0.6413 \text{ mg of S} \end{aligned}$$

**Loss on drying** <2.41> Not more than 10.0% (0.5 g, in vacuum, phosphorus (V) oxide,  $60^\circ\text{C}$ , 4 hours).

**Viscosity** <2.53> Weigh accurately about 1.5 g of Dextran Sulfate Sodium Sulfur 5, calculated on the dried basis, dissolve in a solution of sodium chloride (29 in 500) to make exactly 100 mL, and use this solution as the sample solution. Perform the test with the sample solution and a solution of sodium chloride (29 in 500) at  $25 \pm 0.02^\circ\text{C}$  as directed: the intrinsic viscosity is between 0.030 and 0.040.

**Containers and storage** Containers—Tight containers.

## Dextran Sulfate Sodium Sulfur 18

デキストラン硫酸エステルナトリウム イオウ 18

Dextran Sulfate Sodium Sulfur 18 is a sodium salt of sulfate ester obtained by sulfation of partial decomposition products of dextran, which is produced by fermentation of sucrose with *Leuconostoc mesenteroides* Van Tieghem (*Lactobacillaceae*).

**Description** Dextran Sulfate Sodium Sulfur 18 occurs as a white to light yellowish white powder. It is odorless, and has a saline taste.

It is freely soluble in water and practically insoluble in ethanol (95) and in diethyl ether.

It is hygroscopic.

**Identification (1)** To 10 mL of a solution of toluidine blue (1 in 100,000) add 0.05 mL of a solution of Dextran Sulfate Sodium Sulfur 18 (3 in 50) dropwise: a color of the solution changes from blue to red-purple.

**(2)** To 1 mL of a solution of Dextran Sulfate Sodium Sulfur 18 (1 in 1500) add 2 mL of anthrone TS: a blue-green color develops, which turns dark blue-green gradually. Then, add 1 mL of diluted sulfuric acid (1 in 2) or 1 mL of acetic acid (100) to this solution: the solution remains dark blue-green.

**(3)** A solution of Dextran Sulfate Sodium Sulfur 18 (1 in 100) responds to the Qualitative Tests <1.09> (1) for sodium salt.

**Optical rotation** <2.49>  $[\alpha]_D^{20}$ : +90.0 – +110.0° (calculated on the dried basis, 1.5 g, water, 25 mL, 100 mm).

**pH** <2.54> Dissolve 1.0 g of Dextran Sulfate Sodium Sulfur 18 in 20 mL of water: the pH of this solution is between 5.5 and 7.5.

**Purity (1)** Chloride <1.03>—Perform the test with 0.10 g of Dextran Sulfate Sodium Sulfur 18. Prepare the control solution with 0.30 mL of 0.01 mol/L hydrochloric acid (not more than 0.106%).

**(2)** Sulfate <1.14>—Dissolve 0.10 g of Dextran Sulfate Sodium Sulfur 18 in 6 mL of water, add 0.6 mL of barium chloride TS, and heat in a water bath for 4 minutes. After cooling, add 1 mL of dilute hydrochloric acid and water to make 50 mL, allow to stand for 10 minutes, and observe: the turbidity of the solution is not more intense than that of the control solution. Prepare the control solution as follows: to 1.0 mL of 0.005 mol/L sulfuric acid add 6 mL of water, and proceed in the same manner (not more than 0.480%).

**(3)** Heavy metals <1.07>—Proceed with 1.0 g of Dextran Sulfate Sodium Sulfur 18 according to Method 2, and perform the test. Prepare the control solution with 2.0 mL of Standard Lead Solution (not more than 20 ppm).

**(4)** Arsenic <1.11>—Prepare the test solution with 1.0 g of Dextran Sulfate Sodium Sulfur 18 according to Method 3, and perform the test (not more than 2 ppm).

**Sulfur content** Weigh accurately about 0.5 g of Dextran Sulfate Sodium Sulfur 18, dissolve in 5 mL of water, add 1.5 mL of hydrochloric acid, and heat in a water bath for 1 hour. After cooling, add water to make exactly 100 mL, and use this solution as the sample solution. To exactly 10 mL of the

sample solution add exactly 20 mL of 0.02 mol/L barium chloride VS, add 5 mL of methanol, and heat in a water bath for 30 minutes. After cooling, neutralize with sodium hydroxide TS, and add 70 mL of water, 10 mL of a solution of zinc disodium ethylenediamine tetraacetate tetrahydrate (1 in 20), 3 mL of ammonium chloride TS and 7 mL of strong ammonium water, and titrate <2.50> with 0.02 mol/L disodium dihydrogen ethylenediamine tetraacetate VS until the color of the solution changes from red to light blue (indicator: 5 drops of eriochrome black T TS). Perform a blank determination. Amount of sulfur (S: 32.07), calculated on the dried basis, is between 15.0 and 20.0%.

Each mL of 0.02 mol/L barium chloride VS  
= 0.6413 mg of S

**Loss on drying** <2.41> Not more than 10.0% (0.5 g, in vacuum, phosphorus (V) oxide, 60°C, 4 hours).

**Viscosity** <2.53> Weigh accurately about 1.5 g of Dextran Sulfate Sodium Sulfur 18, calculated on the dried basis, dissolve in a solution of sodium chloride (29 in 500) to make exactly 100 mL, and use this solution as the sample solution. Perform the test with the sample solution and a solution of sodium chloride (29 in 500) at  $25 \pm 0.02^\circ\text{C}$  as directed: the intrinsic viscosity is between 0.020 and 0.032.

**Containers and storage** Containers—Tight containers.

## Dextrin

デキストリン

**Description** Dextrin occurs as a white or light yellow, amorphous powder or granules. It has a slight, characteristic odor and a sweet taste. It does not irritate the tongue. Dextrin is freely soluble in boiling water, soluble in water, and practically insoluble in ethanol (95) and in diethyl ether.

**Identification** To 0.1 g of Dextrin add 100 mL of water, shake, and filter if necessary. To 5 mL of the filtrate add 1 drop of iodine TS: a light red-brown or light red-purple color develops.

**Purity (1)** Clarity and color of solution—Take 2.0 g of Dextrin in a Nessler tube, add 40 mL of water, dissolve by heating, cool, and add water to make 50 mL: the solution is colorless or light yellow. It is clear, and even if turbid, the turbidity is not more than that of the following control solution.

Control solution: To 1.0 mL of 0.005 mol/L sulfuric acid VS add 1 mL of dilute hydrochloric acid, 46 mL of water and 2 mL of barium chloride TS, allow to stand for 10 minutes, and shake before use.

**(2)** Acidity—To 1.0 g of Dextrin add 5 mL of water, dissolve by heating, cool, and add 1 drop of phenolphthalein TS and 0.50 mL of 0.1 mol/L sodium hydroxide VS: a red color develops.

**(3)** Chloride <1.03>—To 2.0 g of Dextrin add 80 mL of water, dissolve by heating, cool, add water to make 100 mL, and filter. Take 40 mL of the filtrate, and add 6 mL of dilute nitric acid and water to make 50 mL. Perform the test using this solution as the test solution. Prepare the control solution with 0.30 mL of 0.01 mol/L hydrochloric acid VS (not more

than 0.013%).

(4) Sulfate <1.14>—To 45 mL of the filtrate obtained in (3) add 1 mL of dilute hydrochloric acid and water to make 50 mL, and perform the test using this solution as the test solution. Prepare the control solution with 0.35 mL of 0.005 mol/L sulfuric acid VS (not more than 0.019%).

(5) Oxalate—To 1.0 g of Dextrin add 20 mL of water, dissolve by heating, cool, add 1 mL of acetic acid (31), and filter. To 5 mL of the filtrate add 5 drops of calcium chloride TS: no turbidity is produced immediately.

(6) Calcium—To a 5-mL portion of the filtrate obtained in (5) add 5 drops of ammonium oxalate TS: no turbidity is immediately produced.

(7) Heavy metals <1.07>—Proceed with 0.5 g of Dextrin according to Method 2, and perform the test. Prepare the control solution with 2.5 mL of Standard Lead Solution (not more than 50 ppm).


**Loss on drying** <2.41> Not more than 10% (0.5 g, 105°C, 4 hours).

**Residue on ignition** <2.44> Not more than 0.5% (0.5 g).

**Containers and storage** Containers—Well-closed containers.

## Dextromethorphan Hydrobromide Hydrate

デキストロメトルファン臭化水素酸塩水和物



C<sub>18</sub>H<sub>25</sub>NO.HBr.H<sub>2</sub>O: 370.32  
(9*S*,13*S*,14*S*)-3-Methoxy-17-methylmorphinan  
monohydrobromide monohydrate [6700-34-1]

Dextromethorphan Hydrobromide Hydrate contains not less than 98.0% of dextromethorphan hydrobromide (C<sub>18</sub>H<sub>25</sub>NO.HBr: 352.31), calculated on the anhydrous basis.

**Description** Dextromethorphan Hydrobromide Hydrate occurs as white crystals or crystalline powder.

It is very soluble in methanol, freely soluble in ethanol (95) and in acetic acid (100), and sparingly soluble in water.

Melting point: about 126°C (Insert the capillary tube into the bath preheated to 116°C, and continue the heating so that the temperature rises at a rate of about 3°C per minute.)

**Identification** (1) Determine the absorption spectrum of a solution of Dextromethorphan Hydrobromide Hydrate (1 in 10,000) as directed under Ultraviolet-visible Spectrophotometry <2.24>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wavelengths.

(2) Determine the infrared absorption spectrum of Dextromethorphan Hydrobromide Hydrate as directed in the potassium bromide disk method under Infrared Spec-

trophotometry <2.25>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wave numbers.

(3) To 50 mL of a solution of Dextromethorphan Hydrobromide Hydrate (1 in 100) add 2 drops of phenolphthalein TS and sodium hydroxide TS until a red color develops. Add 50 mL of chloroform, shake, and add 5 mL of dilute nitric acid to 40 mL of the water layer. This solution responds to the Qualitative Tests <1.09> for bromide.

**Optical rotation** <2.49>  $[\alpha]_D^{20}$ : +26 – +30° (0.34 g, calculated on the anhydrous basis, water, 20 mL, 100 mm).

**pH** <2.54> Dissolve 1.0 g of Dextromethorphan Hydrobromide Hydrate in 100 mL of water: the pH of this solution is between 5.2 and 6.5.

**Purity** (1) Clarity and color of solution—Dissolve 0.20 g of Dextromethorphan Hydrobromide Hydrate in 20 mL of water: the solution is clear and colorless.

(2) *N,N*-dimethylaniline—To 0.50 g of Dextromethorphan Hydrobromide Hydrate add 20 mL of water, and dissolve by heating on a water bath. After cooling, add 2 mL of dilute acetic acid, 1 mL of sodium nitrite TS and water to make 25 mL: the solution has no more color than the following control solution.

Control solution: Dissolve 0.10 g of *N,N*-dimethylaniline in 400 mL of water by warming on a water bath, cool, and add water to make 500 mL. Pipet 5 mL of this solution, and add water to make 200 mL. To 1.0 mL of this solution add 2 mL of dilute acetic acid, 1 mL of sodium nitrite TS and water to make 25 mL.

(3) Heavy metals <1.07>—Proceed with 1.0 g of Dextromethorphan Hydrobromide Hydrate according to Method 4, and perform the test. Prepare the control solution with 2.0 mL of Standard Lead Solution (not more than 20 ppm).

(4) Phenolic compounds—Dissolve 5 mg of Dextromethorphan Hydrobromide Hydrate in 1 drop of dilute hydrochloric acid and 1 mL of water, add 2 drops of iron (III) chloride TS and 2 drops of potassium hexacyanoferrate (III) TS, shake, and allow to stand for 15 minutes: no blue-green color develops.

(5) Related substances—Dissolve 0.25 g of Dextromethorphan Hydrobromide Hydrate in 10 mL of methanol, and use this solution as the sample solution. Pipet 1 mL of the sample solution, add methanol to make exactly 200 mL, and use this solution as the standard solution. Perform the test with these solutions as directed under Thin-layer Chromatography <2.03>. Spot 5 μL each of the sample solution and standard solution on a plate of silica gel for thin-layer chromatography. Develop the plate with a mixture of toluene, ethyl acetate, methanol, dichloromethane and 13.5 mol/L ammonia TS (55:20:13:10:2) to a distance of about 15 cm, and air-dry the plate. Spray evenly bismuth potassium iodide TS on the plate, and then spray evenly hydrogen peroxide TS on the plate: the spots other than the principal spot from the sample solution are not more intense than the spot from the standard solution.

**Water** <2.48> 4.0 – 5.5% (0.2 g, back titration).

**Residue on ignition** <2.44> Not more than 0.1% (1 g).

**Assay** Weigh accurately about 0.5 g of Dextromethorphan Hydrobromide Hydrate, dissolve in 10 mL of acetic acid (100) and add 40 mL of acetic anhydride. Titrate <2.50> with

0.1 mol/L perchloric acid VS (potentiometric titration). Perform a blank determination, and make any necessary correction.

Each mL of 0.1 mol/L perchloric acid VS  
= 35.23 mg of  $C_{18}H_{25}NO \cdot HBr$

**Containers and storage** Containers—Well-closed containers.

## Diastase

ジアスターゼ

Diastase is an enzyme drug mainly prepared from malt.

It has amylolytic activity.

It contains not less than 440 starch saccharifying activity units per g.

It is usually diluted with suitable diluents.

**Description** Diastase occurs as a light yellow to light brown powder.

It is hygroscopic.

**Purity** Rancidity—Diastase has no unpleasant or rancid odor, and has no unpleasant or rancid taste.

**Loss on drying** <2.41> Not more than 4.0% (1 g, 105°C, 5 hours).

**Assay** (i) Substrate solution—Use potato starch TS for amylolytic activity test.

(ii) Sample solution—Weigh accurately about 0.1 g of Diastase, and dissolve in water to make exactly 100 mL.

(iii) Procedure—Proceed as directed in (i) Measurement of starch saccharifying activity of (1) Assay for starch digestive activity under Digestion Test <4.03>.

**Containers and storage** Containers—Tight containers.

Storage—Not exceeding 30°C.

## Diastase and Sodium Bicarbonate Powder

ジアスターゼ・重曹散

### Method of preparation

|                                |       |
|--------------------------------|-------|
| Diastase                       | 200 g |
| Sodium Bicarbonate             | 300 g |
| Precipitated Calcium Carbonate | 400 g |
| Magnesium Oxide                | 100 g |
| To make 1000 g                 |       |

Prepare before use as directed under Powders, with the above ingredients.

**Description** Diastase and Sodium Bicarbonate Powder occurs as a light yellow powder. It has a characteristic, salty taste.

**Containers and storage** Containers—Well-closed containers.

## Compound Diastase and Sodium Bicarbonate Powder

複方ジアスターゼ・重曹散

### Method of preparation

|                    |       |
|--------------------|-------|
| Diastase           | 200 g |
| Sodium Bicarbonate | 600 g |
| Magnesium Oxide    | 150 g |
| Powdered Gentian   | 50 g  |
| To make 1000 g     |       |


Prepare before use as directed under Powders, with the above ingredients.

**Description** Compound Diastase and Sodium Bicarbonate Powder occurs as a slightly brownish, light yellow powder. It has a characteristic odor and a bitter taste.

**Containers and storage** Containers—Well-closed containers.

## Diazepam

ジアゼパム



$C_{16}H_{13}ClN_2O$ : 284.74

7-Chloro-1-methyl-5-phenyl-1,3-dihydro-2H-1,4-benzodiazepin-2-one [439-14-5]

Diazepam, when dried, contains not less than 98.0% of  $C_{16}H_{13}ClN_2O$ .

**Description** Diazepam occurs as a white to light yellow, crystalline powder. It is odorless, and has a slightly bitter taste.

It is freely soluble in acetone, soluble in acetic anhydride and in ethanol (95), sparingly soluble in diethyl ether, slightly soluble in ethanol (99.5), and practically insoluble in water.

**Identification** (1) Dissolve 0.01 g of Diazepam in 3 mL of sulfuric acid, and observe under ultraviolet light (main wavelength: 365 nm): the solution shows a yellow-green fluorescence.

(2) Dissolve 2 mg of Diazepam in 200 mL of a solution of sulfuric acid in ethanol (99.5) (3 in 1000). Determine the absorption spectrum of the solution as directed under Ultraviolet-visible Spectrophotometry <2.24>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wavelengths.

(3) Perform the test with Diazepam as directed under Flame Coloration Test <1.04> (2): a blue to blue-green color appears.

**Absorbance** <2.24>  $E_{1\text{cm}}^{1\%}$  (285 nm): 425 – 445 [after drying, 2 mg, a solution of sulfuric acid in ethanol (99.5) (3 in 1000), 200 mL].

**Melting point** <2.60> 130 – 134°C

**Purity** (1) Clarity of solution—Dissolve 0.10 g of Diazepam in 20 mL of ethanol (95); the solution is clear.

(2) Chloride <1.03>—To 1.0 g of Diazepam add 50 mL of water, allow to stand for 1 hour, with occasional shaking, and filter. To 25 mL of the filtrate add 6 mL of dilute nitric acid and water to make 50 mL. Perform the test using this solution as the test solution. Prepare the control solution with 0.20 mL of 0.01 mol/L hydrochloric acid VS (not more than 0.014%).

(3) Heavy metals <1.07>—Proceed with 1.0 g of Diazepam according to Method 2, and perform the test. Prepare the control solution with 2.0 mL of Standard Lead Solution (not more than 20 ppm).

(4) Related substances—Dissolve 1.0 g of Diazepam in 10 mL of acetone, and use this solution as the sample solution. Pipet 1 mL of the sample solution, and add acetone to make exactly 100 mL. Pipet 1 mL of this solution, add acetone to make exactly 10 mL, and use this solution as the standard solution. Perform the test with these solutions as directed under Thin-layer Chromatography <2.03>. Spot 5  $\mu\text{L}$  each of the sample solution and standard solution on a plate of silica gel with fluorescent indicator for thin-layer chromatography. Develop the plate with a mixture of ethyl acetate and hexane (1:1) to a distance of about 12 cm, and air-dry the plate. Examine under ultraviolet light (main wavelength: 254 nm): the spots other than the principal spot from the sample solution are not more intense than the spot from the standard solution.

**Loss on drying** <2.41> Not more than 0.5% (1 g, 105°C, 2 hours).

**Residue on ignition** <2.44> Not more than 0.1% (1 g).

**Assay** Weigh accurately about 0.6 g of Diazepam, previously dried, dissolve in 60 mL of acetic anhydride, and titrate <2.50> with 0.1 mol/L perchloric acid VS (potentiometric titration). Perform a blank determination, and make any necessary correction.

Each mL of 0.1 mol/L perchloric acid VS  
= 28.48 mg of  $\text{C}_{16}\text{H}_{13}\text{ClN}_2\text{O}$

**Containers and storage** Containers—Tight containers.  
Storage—Light-resistant.

## Dibasic Sodium Phosphate Hydrate

リン酸水素ナトリウム水和物

$\text{Na}_2\text{HPO}_4 \cdot 12\text{H}_2\text{O}$ : 358.14

Dibasic Sodium Phosphate Hydrate, when dried, contains not less than 98.0% of disodium hydrogen phosphate ( $\text{Na}_2\text{HPO}_4$ : 141.96)

**Description** Dibasic Sodium Phosphate Hydrate occurs as colorless or white crystals. It is odorless.

It is freely soluble in water, and practically insoluble in ethanol (95) and in diethyl ether.

It effloresces in warm, dry air.

**Identification** A solution of Dibasic Sodium Phosphate Hydrate (1 in 10) responds to the Qualitative Tests <1.09> (1) and (2) for sodium salt, and the Qualitative Tests <1.09> for phosphate.

**pH** <2.54> Dissolve 1.0 g of Dibasic Sodium Phosphate Hydrate in 50 mL of water: the pH of this solution is between 9.0 and 9.4.

**Purity** (1) Clarity and color of solution—Dissolve 1.0 g of Dibasic Sodium Phosphate Hydrate in 20 mL of water: the solution is clear and colorless.

(2) Chloride <1.03>—Dissolve 1.0 g of Dibasic Sodium Phosphate Hydrate in 7 mL of dilute nitric acid and water to make 50 mL. Perform the test using this solution as the test solution. Prepare the control solution with 0.40 mL of 0.01 mol/L hydrochloric acid VS (not more than 0.014%).

(3) Sulfate <1.14>—Dissolve 0.5 g of Dibasic Sodium Phosphate Hydrate in 2 mL of dilute hydrochloric acid and water to make 50 mL. Perform the test using this solution as the test solution. Prepare the control solution with 0.40 mL of 0.005 mol/L sulfuric acid VS (not more than 0.038%).

(4) Carbonate—To 2.0 g of Dibasic Sodium Phosphate Hydrate add 5 mL of water, boil, and add 2 mL of hydrochloric acid after cooling: the solution does not effervesce.

(5) Heavy metals <1.07>—Dissolve 2.0 g of Dibasic Sodium Phosphate Hydrate in 4 mL of acetic acid (31) and water to make 50 mL. Perform the test using this solution as the test solution. Prepare the control solution with 2.0 mL of Standard Lead Solution by adding 2 mL of dilute acetic acid and water to make 50 mL (not more than 10 ppm).

(6) Arsenic <1.11>—Prepare the test solution with 1.0 g of Dibasic Sodium Phosphate Hydrate according to Method 1, and perform the test (not more than 2 ppm).

**Loss on drying** <2.41> 57.0 – 61.0% (10 g, at 40°C for 3 hours at first and then at 105°C for 5 hours).

**Assay** Dissolve about 3 g of Dibasic Sodium Phosphate Hydrate, previously dried and accurately weighed, in 50 mL of water. Titrate <2.50> it with 0.5 mol/L sulfuric acid VS at 15°C until the green color of the solution changes to dark-greenish red-purple (indicator: 3 to 4 drops of methyl orange-xylene cyanol FF TS).


Each mL of 0.5 mol/L sulfuric acid VS  
= 142.0 mg of  $\text{Na}_2\text{HPO}_4$

**Containers and storage** Containers—Tight containers.



## Dibekacin Sulfate

ジベカシン硫酸塩



$C_{18}H_{37}N_5O_8 \cdot xH_2SO_4$

3-Amino-3-deoxy- $\alpha$ -D-glucopyranosyl-(1 $\rightarrow$ 6)-[2,6-diamino-2,3,4,6-tetra-deoxy- $\alpha$ -D-erythro-hexopyranosyl-(1 $\rightarrow$ 4)]-2-deoxy-D-streptamine sulfate [58580-55-5]

Dibekacin Sulfate is the sulfate of a derivative of bekanamycin.

It contains not less than 640  $\mu$ g (potency) and not more than 740  $\mu$ g (potency) per mg, calculated on the dried basis. The potency of Dibekacin Sulfate is expressed as mass (potency) of dibekacin ( $C_{18}H_{37}N_5O_8$ : 451.52).

**Description** Dibekacin Sulfate occurs as a white to yellowish white powder.

It is very soluble in water, and practically insoluble in ethanol (99.5).

**Identification** (1) Dissolve 20 mg each of Dibekacin Sulfate and Dibekacin Sulfate Reference Standard in 1 mL of water, and use these solutions as the sample solution and the standard solution. Perform the test with these solutions as directed under Thin-layer Chromatography <2.03>. Spot 5  $\mu$ L each of the sample solution and standard solution on a plate of silica gel for thin-layer chromatography. Develop the plate with a mixture of ammonia solution (28) and methanol (1:1) to a distance of about 10 cm, and air-dry the plate. Spray evenly 0.2% ninhydrin-water saturated 1-butanol TS, and heat at 100°C for 10 minutes: the principal spots obtained from the sample solution and standard solution show a purple-brown color and the same  $R_f$  value.

(2) To 5 mL of a solution of Dibekacin Sulfate (1 in 50) add 1 drop of barium chloride TS: a white precipitate is produced.

**Optical rotation** <2.49>  $[\alpha]_D^{20}$ : +96 – +106° (0.25 g calculated on the dried basis, water, 25 mL, 100 mm).

**pH** <2.54> The pH of a solution obtained by dissolving 1.0 g of Dibekacin Sulfate in 20 mL of water is between 6.0 and 8.0.

**Purity** (1) Clarity and color of solution—Dissolve 1.0 g of Dibekacin Sulfate in 10 mL of water: the solution is clear and colorless to pale yellow.

(2) Heavy metals <1.07>—Proceed with 1.0 g of Dibekacin Sulfate according to Method 1, and perform the test. Prepare the control solution with 2.0 mL of Standard Lead Solution (not more than 20 ppm).

**Loss on drying** <2.41> Not more than 5.0% (1 g, reduced pressure not exceeding 0.67 kPa, 60°C, 3 hours).

**Assay** Perform the test according to the Cylinder-plate method as directed under Microbial Assay for Antibiotics <4.02> according to the following conditions.

(i) Test organism—*Bacillus subtilis* ATCC 6633

(ii) Culture medium—Use the medium i in 1) Medium for test organism [5] under (1) Agar media for seed and base layer having pH 6.5 to 6.6 after sterilization.

(iii) Standard solutions—Weigh accurately an amount of Dibekacin Sulfate Reference Standard, previously dried, equivalent to about 20 mg (potency), dissolve in diluted phosphate buffer solution, pH 6.0 (1 in 2) to make exactly 50 mL, and use this solution as the standard stock solution. Keep the standard stock solution at 5 to 15°C and use within 30 days. Take exactly a suitable amount of the standard stock solution before use, add 0.1 mol/L phosphate buffer solution, pH 8.0 to make solutions so that each mL contains 20  $\mu$ g (potency) and 5  $\mu$ g (potency), and use these solutions as the high concentration standard solution and low concentration standard solution, respectively.


(iv) Sample solutions—Weigh accurately an amount of Dibekacin Sulfate, equivalent to about 20 mg (potency), and dissolve in water to make exactly 50 mL. Take exactly a suitable amount of this solution, add 0.1 mol/L phosphate buffer solution, pH 8.0 to make solutions so that each mL contains 20  $\mu$ g (potency) and 5  $\mu$ g (potency), and use these solutions as the high concentration sample solution and low concentration sample solution, respectively.

**Containers and storage** Containers—Tight containers.

## Dibucaine Hydrochloride

### Cinchocaine Hydrochloride

ジブカイン塩酸塩



$C_{20}H_{29}N_3O_2 \cdot HCl$ : 379.92

2-Butyloxy-N-(2-diethylaminoethyl)-4-quinolinecarboxamide monohydrochloride [61-12-1]

Dibucaine Hydrochloride, when dried, contains not less than 98.0% of  $C_{20}H_{29}N_3O_2 \cdot HCl$ .

**Description** Dibucaine Hydrochloride occurs as white crystals or crystalline powder.

It is very soluble in water, in ethanol (95) and in acetic acid (100), freely soluble in acetic anhydride, and practically insoluble in diethyl ether.

It is hygroscopic.

**Identification (1)** Determine the absorption spectrum of a solution of Dibucaine Hydrochloride in 1 mol/L hydrochloric acid TS (1 in 100,000) as directed under Ultraviolet-visible Spectrophotometry <2.24>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wavelengths.

(2) Determine the infrared absorption spectrum of Dibucaine Hydrochloride, previously dried, as directed in the potassium bromide disk method under Infrared Spectrophotometry <2.25>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wave numbers.

(3) A solution of Dibucaine Hydrochloride (1 in 10) responds to the Qualitative Tests <1.09> for chloride.

**Melting point** <2.60> 95 – 100 °C Charge Dibucaine Hydrochloride into a capillary tube for melting point determination, and dry in vacuum over phosphorus (V) oxide at 80 °C for 5 hours. Seal immediately the open end of the tube, and determine the melting point.

**pH** <2.54> Dissolve 1.0 g of Dibucaine Hydrochloride in 50 mL of water: the pH of this solution is between 5.0 and 6.0.

**Purity (1)** Clarity and color of solution—Dissolve 1.0 g of Dibucaine Hydrochloride in 20 mL of water: the solution is clear and colorless. Determine the absorbance of this solution at 430 nm as directed under Ultraviolet-visible Spectrophotometry <2.24>, using water as the blank: it is not more than 0.03.

(2) Sulfate <1.14>—Perform the test with 0.30 g of Dibucaine Hydrochloride. Prepare the control solution with 0.35 mL of 0.005 mol/L sulfuric acid VS (not more than 0.056%).

(3) Heavy metals <1.07>—Proceed with 1.0 g of Dibucaine Hydrochloride according to Method 1, and perform the test. Prepare the control solution with 2.0 mL of Standard Lead Solution (not more than 20 ppm).

(4) Related substances—Dissolve 0.20 g of Dibucaine Hydrochloride in 5 mL of ethanol (95), and use this solution as the sample solution. Pipet 1 mL of the sample solution, add ethanol (95) to make exactly 20 mL, then pipet 2 mL of this solution, add ethanol (95) to make exactly 20 mL, and use this solution as the standard solution. Perform the test with these solutions as directed under Thin-layer Chromatography <2.03>. Spot 5  $\mu$ L each of the sample solution and standard solution on a plate of silica gel for thin-layer chromatography. Develop the plate with a mixture of ethyl acetate, water and acetic acid (100) (3:1:1) to a distance of about 10 cm, and air-dry the plate. Examine under ultraviolet (main wavelength: 254 nm): the spots other than the principal spot from the sample solution are not more intense than the spot from the standard solution.

**Loss on drying** <2.41> Not more than 2.0% (1 g, in vacuum, phosphorus (V) oxide, 80 °C, 5 hours).

**Residue on ignition** <2.44> Not more than 0.1% (1 g).


**Assay** Weigh accurately about 0.3 g of Dibucaine Hydrochloride, previously dried, dissolve in 50 mL of a mixture of acetic anhydride and acetic acid (100) (7:3), and titrate <2.50> with 0.1 mol/L perchloric acid VS (potentiometric titration). Perform a blank determination, and make any necessary correction.

Each mL of 0.1 mol/L perchloric acid VS  
= 19.00 mg of  $C_{20}H_{29}N_3O_2 \cdot HCl$

**Containers and storage** Containers—Tight containers.

## Diclofenac Sodium

ジクロフェナクナトリウム



$C_{14}H_{10}Cl_2NNaO_2$ : 318.13

Monosodium 2-(2,6-dichlorophenylamino)phenylacetate  
[15307-79-6]

Diclofenac Sodium, when dried, contains not less than 98.5% of  $C_{14}H_{10}Cl_2NNaO_2$ .

**Description** Diclofenac Sodium occurs as white to pale yellowish white crystals or crystalline powder.

It is freely soluble in methanol and in ethanol (95), sparingly soluble in water and in acetic acid (100), and practically insoluble in diethyl ether.

It is hygroscopic.

**Identification (1)** To 1 mL of a solution of Diclofenac Sodium in methanol (1 in 250) add 1 mL of nitric acid: a dark red color develops.

(2) Perform the test with 5 mg of Diclofenac Sodium as directed under Flame Coloration Test <1.04> (2): a light green color appears.

(3) Determine the infrared absorption spectrum of Diclofenac Sodium, previously dried, as directed in the potassium bromide disk method under Infrared Spectrophotometry <2.25>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wave numbers.

(4) A solution of Diclofenac Sodium (1 in 100) responds to the Qualitative Tests <1.09> for sodium salt.

**Purity (1)** Heavy metals <1.07>—Proceed with 2.0 g of Diclofenac Sodium according to Method 2, and perform the test. Prepare the control solution with 2.0 mL of Standard Lead Solution (not more than 10 ppm).

(2) Arsenic <1.11>—Prepare the test solution with 1.0 g of Diclofenac Sodium according to Method 3, and perform the test (not more than 2 ppm).

(3) Related substances—Dissolve 0.05 g of Diclofenac Sodium in 50 mL of the mobile phase, and use this solution as the sample solution. Pipet 2 mL of the sample solution, and add the mobile phase to make exactly 50 mL. Pipet 5 mL of this solution, add the mobile phase to make exactly 100 mL, and use this solution as the standard solution. Perform the test with exactly 20  $\mu$ L each of these solutions as directed under Liquid Chromatography <2.01>. Determine each peak area of these solutions by the automatic integration method: the area of each peak other than the peak of diclofenac from the sample solution is not larger than the peak area of the standard solution.

**Operating conditions—**

**Detector:** An ultraviolet absorption photometer (wavelength: 240 nm).

**Column:** A stainless steel column 4.6 mm in inside diameter and 25 cm in length, packed with octadecylsilanized silica gel for liquid chromatography (7  $\mu$ m in particle diameter).

**Column temperature:** A constant temperature of about 40°C.

**Mobile phase:** A mixture of methanol and diluted acetic acid (100) (3 in 2500) (4:3).

**Flow rate:** Adjust the flow rate so that the retention time of diclofenac is about 20 minutes.

**Time span of measurement:** About twice as long as the retention time of diclofenac beginning after the solvent peak.

**System suitability—**

**System performance:** Dissolve 35 mg of ethyl parahydroxybenzoate and 0.05 g of propyl parahydroxybenzoate in 100 mL of the mobile phase. To 1 mL of this solution add the mobile phase to make 50 mL. When the procedure is run with 20  $\mu$ L of this solution under the above operating conditions, ethyl parahydroxybenzoate and propyl parahydroxybenzoate are eluted in this order with the resolution between these peaks being not less than 5.

**System repeatability:** When the test is repeated 6 times with 20  $\mu$ L of the standard solution under the above operating conditions, the relative standard deviation of the peak areas of diclofenac is not more than 2.0%.

**Loss on drying** <2.41> Not more than 0.5% (1 g, 105°C, 3 hours).


**Assay** Weigh accurately about 0.5 g of Diclofenac Sodium, previously dried, dissolve with 40 mL of water in a separator, add 2 mL of dilute hydrochloric acid, and extract the precipitate formed with 50 mL of chloroform. Extract again with two 20-mL portions of chloroform, and filter the extract each time through a pledget of absorbent cotton moistened with chloroform. Wash the tip of the separator and the absorbent cotton with 15 mL of chloroform, combine the washing with the extracts, add 10 mL of a solution of 1 mol/L hydrochloric acid TS in ethanol (99.5) (1 in 100), and titrate <2.50> with 0.1 mol/L potassium hydroxide-ethanol VS from the first equivalent point to the second equivalent point (potentiometric titration).

Each mL of 0.1 mol/L potassium hydroxide-ethanol VS  
= 31.81 mg of  $C_{14}H_{10}Cl_2NNaO_2$

**Containers and storage** Containers—Tight containers.

**Diclofenamide****Dichlorophenamide**

ジクロフェナミド



$C_6H_4Cl_2N_2O_4S_2$ : 305.16

4,5-Dichlorobenzene-1,3-disulfonamide [120-97-8]

Diclofenamide, when dried, contains not less than 98.0% of  $C_6H_4Cl_2N_2O_4S_2$ .

**Description** Diclofenamide occurs as a white, crystalline powder.

It is very soluble in *N,N*-dimethylformamide, soluble in ethanol (95), and very slightly soluble in water.

It dissolves in sodium hydroxide TS.

**Identification (1)** Dissolve 0.01 g of Diclofenamide in 100 mL of 0.01 mol/L sodium hydroxide TS. To 10 mL of the solution add 0.1 mL of hydrochloric acid. Determine the absorption spectrum of the solution as directed under Ultraviolet-visible Spectrophotometry <2.24>, and compare the spectrum with the Reference Spectrum or the spectrum of a solution of Diclofenamide Reference Standard prepared in the same manner as the sample solution: both spectra exhibit similar intensities of absorption at the same wavelengths.

**(2)** Determine the infrared absorption spectrum of Diclofenamide, as directed in the potassium bromide disk method under Infrared Spectrophotometry <2.25>, and compare the spectrum with the Reference Spectrum or the spectrum of Diclofenamide Reference Standard: both spectra exhibit similar intensities of absorption at the same wave numbers.

**Melting point** <2.60> 237 – 240°C

**Purity (1)** Chloride < 1.03 > —Dissolve 0.10 g of Diclofenamide in 10 mL of *N,N*-dimethylformamide, and add 6 mL of dilute nitric acid and water to make 50 mL. Perform the test using this solution as the test solution. Prepare the control solution as follows: to 0.45 mL of 0.01 mol/L hydrochloric acid VS add 10 mL of *N,N*-dimethylformamide, 6 mL of dilute nitric acid and water to make 50 mL (not more than 0.160%).

**(2)** Selenium—To 0.10 g of Diclofenamide add 0.5 mL of a mixture of perchloric acid and sulfuric acid (1:1) and 2 mL of nitric acid, and heat on a water bath until no more brown gas evolves and the solution becomes to be a light yellow clear solution. After cooling, add 4 mL of nitric acid to this solution, then add water to make exactly 50 mL, and use this solution as the sample solution. Separately, pipet 3 mL of Standard Selenium Solution, add 0.5 mL of a mixture of perchloric acid and sulfuric acid (1:1) and 6 mL of nitric acid, then add water to make exactly 50 mL, and use this solution as the standard solution. Perform the test with the sample solution and standard solution as directed under Atomic Ab-

sorption Spectrophotometry <2.23> according to the following conditions, and determine constant absorbances,  $A_T$  and  $A_S$ , obtained on a recorder after rapid increasing of the absorption:  $A_T$  is smaller than  $A_S$  (not more than 30 ppm).

Perform the test by using a hydride generating system and a thermal absorption cell.

Lamp: An selenium hollow cathode lamp

Wavelength: 196.0 nm

Temperature of sample atomizer: When an electric furnace is used, about 1000°C.

Carrier gas: Nitrogen or Argon

(3) Heavy metals <1.07>—Proceed with 2.0 g of Diclofenamide according to Method 2, and perform the test. Prepare the control solution with 2.0 mL of Standard Lead Solution (not more than 10 ppm).

(4) Related substances—Dissolve 0.10 g of Diclofenamide in 50 mL of the mobile phase, and use this solution as the sample solution. Pipet 2 mL of the sample solution, add the mobile phase to make exactly 100 mL, and use this solution as the standard solution. Perform the test with exactly 10  $\mu$ L each of the sample solution and standard solution as directed under Liquid Chromatography <2.01> according to the following conditions, and determine each peak area by the automatic integration method: the total area of the peaks other than the peak of diclofenamide from the sample solution is not larger than the peak area of diclofenamide from the standard solution.

**Operating conditions—**

Detector, column, column temperature, mobile phase, and flow rate: Proceed as directed in the operating conditions in the Assay.

Time span of measurement: About 5 times as long as the retention time of diclofenamide.

**System suitability—**

Test for required detection: To exactly 5 mL of the standard solution add the mobile phase to make exactly 100 mL. Confirm that the peak area of diclofenamide obtained from 10  $\mu$ L of this solution is equivalent to 3.5 to 6.5% of that of diclofenamide obtained from 10  $\mu$ L of the standard solution.

System performance: Proceed as directed in the system suitability in the Assay.

System repeatability: When the test is repeated 6 times with 10  $\mu$ L of the standard solution under the above operating conditions, the relative standard deviation of the peak area of diclofenamide is not more than 1.0%.

**Loss on drying** <2.41> Not more than 1.0% (1 g, in vacuum at a pressure not exceeding 0.67 kPa, 100°C, 5 hours).

**Residue on ignition** <2.44> Not more than 0.1% (1 g).

**Assay** Weigh accurately about 50 mg each of Diclofenamide and Diclofenamide Reference Standard, previously dried, and dissolve each in 30 mL of the mobile phase. To each add exactly 10 mL of the internal standard solution and the mobile phase to make 50 mL, and use these solutions as the sample solution and the standard solution. Perform the test with 10  $\mu$ L each of the sample solution and standard solution as directed under Liquid Chromatography <2.01> according to the following conditions, and calculate the ratios,  $Q_T$  and  $Q_S$ , of the peak area of diclofenamide to that of the internal standard, respectively.

$$\text{Amount (mg) of } C_6H_6Cl_2N_2O_4S_2 = W_S \times (Q_T/Q_S)$$

$W_S$ : Amount (mg) of Diclofenamide Reference Standard

**Internal standard solution—**A solution of butyl parahydroxy benzoate in the mobile phase (3 in 5000).

**Operating conditions—**

Detector: An ultraviolet absorption photometer (wavelength: 280 nm).

Column: A stainless steel column 4 mm in inside diameter and 30 cm in length, packed with octadecylsilanized silica gel for liquid chromatography (10  $\mu$ m in particle diameter).

Column temperature: A constant temperature of about 25°C.

Mobile phase: A mixture of sodium phosphate TS and acetonitrile (1:1).

Flow rate: Adjust the flow rate so that the retention time of diclofenamide is about 7 minutes.

**System suitability—**

System performance: When the procedure is run with 10  $\mu$ L of the standard solution under the above operating conditions, diclofenamide and the internal standard are eluted in this order with the resolution between these peaks being not less than 9.

System repeatability: When the test is repeated 6 times with 10  $\mu$ L of the standard solution under the above operating conditions, the relative standard deviation of the ratios of the peak area of diclofenamide to that of the internal standard is not more than 1.0%.

**Containers and storage** Containers—Tight containers.

## Diclofenamide Tablets

### Dichlorphenamide Tablets

ジクロフェナミド錠

Diclofenamide Tablets contain not less than 92% and not more than 108% of the labeled amount of diclofenamide ( $C_6H_6Cl_2N_2O_4S_2$ ; 305.16).

**Method of preparation** Prepare as directed under Tablets, with Diclofenamide.

**Identification** To a quantity of powdered Diclofenamide Tablets, equivalent to 0.2 g of Diclofenamide according to the labeled amount, add 20 mL of methanol, shake, and filter. Evaporate the filtrate on a water bath to dryness, and dissolve 0.01 g of the residue in 100 mL of 0.01 mol/L sodium hydroxide TS. To 10 mL of this solution add 0.1 mL of hydrochloric acid TS, and determine the absorption spectrum of this solution as directed under Ultraviolet-visible Spectrophotometry <2.24>: it exhibits maxima between 284 nm and 288 nm, and between 293 nm and 297 nm.

**Dissolution** <6.10> Perform the test according to the following method: it meets the requirement.

Perform the test with 1 tablet of Diclofenamide Tablets at 50 revolutions per minute according to the Paddle method, using 900 mL of water as the test solution. Take 20 mL or more of the dissolved solution 60 minutes after starting the test, and filter through a membrane filter with a pore size not exceeding 0.8  $\mu$ m. Discard the first 10 mL of the filtrate, and use the subsequent as the sample solution. Separately, weigh accurately about 55 mg of Diclofenamide Reference Stan-

dard, previously dried in vacuum at a pressure not exceeding 0.67 kPa at 100°C for 5 hours, dissolve in 10 mL of methanol, and add water to make exactly 100 mL. Pipet 10 mL of this solution, add water to make exactly 100 mL, and use this solution as the standard solution. Determine the absorbances,  $A_T$  and  $A_S$ , of the sample solution and the standard solution at 285 nm as directed under Ultraviolet-visible Spectrophotometry <2.24>. The dissolution rate of Diclofenamide Tablets in 60 minutes is not less than 70%.

Dissolution rate (%) with respect to the labeled amount of diclofenamide ( $C_6H_6Cl_2N_2O_4S_2$ )

$$= W_S \times (A_T/A_S) \times (1/C) \times 90$$

$W_S$ : Amount (mg) of Diclofenamide Reference Standard.

$C$ : Labeled amount (mg) of diclofenamide ( $C_6H_6Cl_2N_2O_4S_2$ ) in 1 tablet.

**Assay** Weigh accurately, and powder not less than 20 tablets of Diclofenamide Tablets. Weigh accurately a portion of the powder, equivalent to about 50 mg of diclofenamide ( $C_6H_6Cl_2N_2O_4S_2$ ), add exactly 25 mL of the mobile phase, shake for 15 minutes, and centrifuge. Pipet 10 mL of the supernatant liquid, add exactly 4 mL of the internal standard solution and the mobile phase to make 20 mL, and use this solution as the sample solution. Separately, weigh accurately about 50 mg of Diclofenamide Reference Standard, previously dried at 100°C in vacuum at a pressure not exceeding 0.67 kPa for 5 hours, dissolve in 30 mL of the mobile phase, add exactly 10 mL of the internal standard solution and the mobile phase to make 50 mL, and use this solution as the standard solution. Proceed as directed in the Assay under Diclofenamide.

Amount (mg) of diclofenamide ( $C_6H_6Cl_2N_2O_4S_2$ )

$$= W_S \times (Q_T/Q_S)$$


$W_S$ : Amount (mg) of Diclofenamide Reference Standard

**Internal standard solution**—A solution of butyl parahydroxybenzoate in the mobile phase (3 in 5000).

**Containers and storage** Containers—Well-closed containers.

## Dicloxacillin Sodium Hydrate

ジクロキサシリンナトリウム水和物



$C_{19}H_{16}Cl_2N_3NaO_5S \cdot H_2O$ : 510.32

Monosodium (2*S*,5*R*,6*R*)-6-{[3-(2,6-dichlorophenyl)-5-methylisoxazole-4-carbonyl]amino}-3,3-dimethyl-7-oxo-4-thia-1-azabicyclo[3.2.0]heptane-2-carboxylate monohydrate [13412-64-1]

Dicloxacillin Sodium Hydrate contains not less than 910  $\mu$ g (potency) and not more than 1020  $\mu$ g (potency) per mg, calculated on the anhydrous basis. The potency of Dicloxacillin Sodium Hydrate is expressed as

mass (potency) of dicloxacillin ( $C_{19}H_{17}Cl_2N_3O_5S$ : 470.33).

**Description** Dicloxacillin Sodium Hydrate occurs as a white to light yellowish white crystalline powder.

It is freely soluble in water and in methanol, and soluble in ethanol (95).

**Identification** (1) Determine the absorption spectrum of a solution of Dicloxacillin Sodium Hydrate (1 in 2500) as directed under Ultraviolet-visible Spectrophotometry <2.24>, and compare the spectrum with the Reference Spectrum or the spectrum of a solution of Dicloxacillin Sodium Reference Standard prepared in the same manner as the sample solution: both spectra exhibit similar intensities of absorption at the same wavelengths.

(2) Determine the infrared absorption spectrum of Dicloxacillin Sodium Hydrate as directed in the potassium bromide disk method under Infrared Spectrophotometry <2.25>, and compare the spectrum with the Reference Spectrum or the spectrum of Dicloxacillin Sodium Reference Standard: both spectra exhibit similar intensities of absorption at the same wave numbers.

(3) Dicloxacillin Sodium Hydrate responds to the Quantitative Tests <1.09> (1) for sodium salt.

**Water** <2.48> Not less than 3.0% and not more than 4.5% (0.1 g, volumetric titration, direct titration).

**Assay** Perform the test according to the Cylinder-plate method as directed under Microbial Assay for Antibiotics <4.02> according to the following conditions.

(i) Test organism—*Bacillus subtilis* ATCC 6633

(ii) Culture medium—Use the medium i in 1) Medium for test organism [5] under (1) Agar media for seed and base layer. Adjust the pH of the medium so that it will be 6.5 to 6.6 after sterilization.


(iii) Standard solutions—Weigh accurately an amount of Dicloxacillin Sodium Reference Standard equivalent to about 50 mg (potency), dissolve in phosphate buffer solution, pH 6.0 to make exactly 50 mL, and use this solution as the standard stock solution. Keep the standard stock solution at 5°C or below and use within 24 hours. Take exactly a suitable amount of the standard stock solution before use, add phosphate buffer solution, pH 6.0 to make solutions so that each mL contains 10  $\mu$ g (potency) and 2.5  $\mu$ g (potency), and use these solutions as the high concentration standard solution and low concentration standard solution, respectively.

(iv) Sample solutions—Weigh accurately an amount of Dicloxacillin Sodium Hydrate equivalent to about 50 mg (potency), dissolve in phosphate buffer solution, pH 6.0 to make exactly 50 mL. Take exactly a suitable amount of the solution, add phosphate buffer solution, pH 6.0 to make solutions so that each mL contains 10  $\mu$ g (potency) and 2.5  $\mu$ g (potency), and use these solutions as the high concentration sample solution and low concentration sample solution, respectively.

**Containers and storage** Containers—Tight containers.

## Diethylcarbamazine Citrate

ジエチルカルバマジンクエン酸塩



$C_{10}H_{21}N_3O \cdot C_6H_8O_7$ ; 391.42

*N,N*-Diethyl-4-methylpiperazine-1-carboxamide  
monocitrate [1642-54-2]

Diethylcarbamazine Citrate, when dried, contains not less than 98.0% of  $C_{10}H_{21}N_3O \cdot C_6H_8O_7$ .

**Description** Diethylcarbamazine Citrate occurs as a white, crystalline powder. It is odorless, and has an acid and bitter taste.

It is very soluble in water, soluble in ethanol (95), and practically insoluble in acetone, in chloroform and in diethyl ether.

A solution of Diethylcarbamazine Citrate (1 in 20) is acid. It is hygroscopic.

**Identification (1)** Dissolve 0.5 g of Diethylcarbamazine Citrate in 2 mL of water, add 10 mL of sodium hydroxide TS, and extract with four 5-mL portions of chloroform. Wash the combined chloroform extracts with 10 mL of water, and evaporate the chloroform on a water bath. Add 1 mL of iodoethane to the residue, and boil gently under a reflux condenser for 5 minutes. Evaporate the excess iodoethane with the aid of a current of air, and dissolve the residue in 4 mL of ethanol (95). Cool the ethanol solution in an ice bath, with continuous stirring, add diethyl ether until precipitates are formed, and stir until crystallization is evident. Allow to stand in the ice bath for 30 minutes, and collect the precipitate. Dissolve the precipitate in 4 mL of ethanol (95), repeat the recrystallization in the same manner, then dry at 105°C for 4 hours: the crystals so obtained melt <2.60> between 151°C and 155°C.

(2) Neutralize the remaining aqueous layer obtained in (1) with dilute sulfuric acid: the solution responds to the Qualitative Tests <1.09> (2) and (3) for citrate.

**Melting point** <2.60> 135.5 – 138.5°C

**Purity** Heavy metals <1.07>—Proceed with 2.0 g of Diethylcarbamazine Citrate according to Method 4, and perform the test. Prepare the control solution with 4.0 mL of Standard Lead Solution (not more than 20 ppm).

**Loss on drying** <2.41> Not more than 1.0% (2 g, 105°C, 4 hours).

**Residue on ignition** <2.44> Not more than 0.1% (1 g).

**Assay** Weigh accurately about 0.75 g of Diethylcarbamazine Citrate, previously dried, dissolve in 50 mL of acetic acid (100) by warming, cool, and titrate <2.50> with 0.1 mol/L perchloric acid VS (potentiometric titration). Perform a blank determination, and make any necessary correction.

Each mL of 0.1 mol/L perchloric acid VS  
= 39.14 mg of  $C_{10}H_{21}N_3O \cdot C_6H_8O_7$

**Containers and storage** Containers—Tight containers.

## Diethylcarbamazine Citrate Tablets

ジエチルカルバマジンクエン酸塩錠

Diethylcarbamazine Citrate Tablets contain not less than 95% and not more than 105% of the labeled amount of diethylcarbamazine citrate ( $C_{10}H_{21}N_3O \cdot C_6H_8O_7$ ; 391.42).

**Method of preparation** Prepare as directed under Tablets, with Diethylcarbamazine Citrate.

**Identification (1)** To a quantity of powdered Diethylcarbamazine Citrate Tablets, equivalent to 0.5 g of Diethylcarbamazine Citrate according to the labeled amount, add 10 mL of water, shake, and filter. Add 10 mL of sodium hydroxide TS to the filtrate, and proceed as directed in the Identification (1) under Diethylcarbamazine Citrate.

(2) To a quantity of powdered Diethylcarbamazine Citrate Tablets, equivalent to 0.8 g of diethylcarbamazine citrate according to the labeled amount, add 10 mL of water, shake, centrifuge, and filter the supernatant liquid. To 5 mL of the filtrate add 5 mL of sodium hydroxide TS, and extract with two 20-mL portions of chloroform. Separate the aqueous layer, and neutralize with dilute hydrochloric acid: the solution responds to the Qualitative Tests <1.09> (2) and (3) for citrate.

**Assay** Weigh accurately and powder not less than 20 Diethylcarbamazine Citrate Tablets. Weigh accurately a portion of the powder, equivalent to about 50 mg of diethylcarbamazine citrate ( $C_{10}H_{21}N_3O \cdot C_6H_8O_7$ ), add 10 mL of water, shake well, add 5 mL of sodium hydroxide TS, then add exactly 20 mL of the internal standard solution, and shake vigorously for 10 minutes. Centrifuge, discard the aqueous layer, and use the chloroform layer as the sample solution. Separately, weigh accurately about 50 mg of Diethylcarbamazine Citrate Reference Standard, previously dried at 105°C for 4 hours, dissolve in 10 mL of water, add 5 mL of sodium hydroxide TS, proceed in the same manner as the preparation of the sample solution, and use the chloroform layer as the standard solution. Perform the test with 2  $\mu$ L of the sample solution and standard solution as directed under Gas Chromatography <2.02> according to the following conditions, and calculate the ratios,  $Q_T$  and  $Q_S$ , of the peak area of diethylcarbamazine to that of the internal standard, respectively.

$$\begin{aligned} &\text{Amount (mg) of diethylcarbamazine citrate} \\ & (C_{10}H_{21}N_3O \cdot C_6H_8O_7) \\ & = W_S \times (Q_T / Q_S) \end{aligned}$$

$W_S$ : Amount (mg) of Diethylcarbamazine Citrate Reference Standard

**Internal standard solution**—A solution of *n*-octadecane in chloroform (1 in 1250).

**Operating conditions**—

Detector: A hydrogen flame-ionization detector.

Column: A glass tube 3 mm in inside diameter and 1 m in length, packed with silanized siliceous earth for gas chro-

matography (180 to 250  $\mu\text{m}$  in particle diameter) coated with 35% methylphenyldimethyl silicone polymer for gas chromatography in the ratio of 3%.

Column temperature: A constant temperature of about 145  $^{\circ}\text{C}$ .

Carrier gas: Nitrogen

Flow rate: Adjust the flow rate so that the retention time of diethylcarbamazine is about 4 minutes.

System suitability—


System performance: When the procedure is run with 2  $\mu\text{L}$  of the standard solution under the above operating conditions, diethylcarbamazine and the internal standard are eluted in this order with the resolution between these peaks being not less than 5.

System repeatability: When the test is repeated 6 times with 2  $\mu\text{L}$  of the standard solution under the above operating conditions, the relative standard deviation of the ratios of the peak area of diethylcarbamazine to that of the internal standard is not more than 1.5%.

**Containers and storage** Containers—Well-closed containers.

## Difenidol Hydrochloride

ジフェニドール塩酸塩



$\text{C}_{21}\text{H}_{27}\text{NO} \cdot \text{HCl}$ : 345.91

1,1-Diphenyl-4-piperidin-1-ylbutan-1-ol  
monohydrochloride [3254-89-5]

Difenidol Hydrochloride, when dried, contains not less than 98.5% of  $\text{C}_{21}\text{H}_{27}\text{NO} \cdot \text{HCl}$ .

**Description** Difenidol Hydrochloride occurs as white crystals or crystalline powder. It is odorless.

It is freely soluble in methanol, soluble in ethanol (95), sparingly soluble in water and in acetic acid (100), and practically insoluble in diethyl ether.

Melting point: about 217 $^{\circ}\text{C}$  (with decomposition).

**Identification** (1) Dissolve 0.01 g of Difenidol Hydrochloride in 1 mL of sulfuric acid: an orange-red color develops. To this solution add carefully 3 drops of water: the solution becomes yellowish brown, and colorless on the addition of 10 mL of water.

(2) To 5 mL of a solution of Difenidol Hydrochloride (1 in 100) add 2 mL of Reinecke salt TS: a light red precipitate is formed.

(3) To 10 mL of a solution of Difenidol Hydrochloride (1 in 100) add 2 mL of sodium hydroxide TS, and extract with two 15-mL portions of chloroform. Combine the extracts, wash with three 10-mL portions of water, evaporate the chloroform on a water bath, and dry the residue in a desiccator (in vacuum, silica gel, 55 $^{\circ}\text{C}$ ) for 5 hours: the residue melts <2.60> between 103 $^{\circ}\text{C}$  and 106 $^{\circ}\text{C}$ .

(4) A solution of Difenidol Hydrochloride (1 in 100)

responds to the Qualitative Tests <1.09> for chloride.

**pH** <2.54> Dissolve 1.0 g of Difenidol Hydrochloride in 100 mL of freshly boiled and cooled water: the pH of this solution is between 4.7 and 6.5.

**Purity** (1) Clarity and color of solution—Dissolve 1.0 g of Difenidol Hydrochloride in 10 mL of methanol: the solution is clear and colorless.

(2) Heavy metals <1.07>—Proceed with 1.0 g of Difenidol Hydrochloride according to Method 2, and perform the test. Prepare the control solution with 2.0 mL of Standard Lead Solution (not more than 20 ppm).

(3) Arsenic <1.11>—Prepare the test solution with 2.0 g of Difenidol Hydrochloride according to Method 3, and perform the test (not more than 1 ppm).

(4) Related substances—Dissolve 0.10 g of Difenidol Hydrochloride in methanol to make exactly 10 mL, and use this solution as the sample solution. Separately, dissolve 10 mg of 1,1-diphenyl-4-piperidino-1-butene hydrochloride for thin-layer chromatography in methanol to make exactly 20 mL, pipet 1 mL of this solution, add methanol to make exactly 10 mL, and use this solution as the standard solution. Perform the test with these solutions as directed under Thin-layer Chromatography <2.03>. Spot 5  $\mu\text{L}$  each of the sample solution and standard solution on a plate of silica gel with fluorescent indicator for thin-layer chromatography. Develop the plate with a mixture of toluene, methanol and acetic acid (100) (10:2:1) to a distance of about 15 cm, and air-dry the plate. Examine under ultraviolet light (main wavelength: 254 nm): the spots other than the principal spot from the sample solution are not more intense than the spot from the standard solution.

**Loss on drying** <2.41> Not more than 0.5% (1 g, in vacuum, silica gel, 5 hours).

**Residue on ignition** <2.44> Not more than 0.1% (1 g).


**Assay** Weigh accurately about 0.35 g of Difenidol Hydrochloride, previously dried, dissolve in 30 mL of acetic acid (100) by warming if necessary, cool, add 30 mL of acetic anhydride, and titrate <2.50> with 0.05 mol/L perchloric acid VS (potentiometric titration). Perform a blank determination, and make any necessary correction.

Each mL of 0.05 mol/L perchloric acid VS  
= 17.30 mg of  $\text{C}_{21}\text{H}_{27}\text{NO} \cdot \text{HCl}$

**Containers and storage** Containers—Well-closed containers.

## Digitoxin

ジギトキシン


 $C_{41}H_{64}O_{13}$ : 764.94

3 $\beta$ -[2,6-Dideoxy- $\beta$ -D-ribo-hexopyranosyl-(1 $\rightarrow$ 4)-2,6-dideoxy- $\beta$ -D-ribo-hexopyranosyl-(1 $\rightarrow$ 4)-2,6-dideoxy- $\beta$ -D-ribo-hexopyranosyloxy]-14-hydroxy-5 $\beta$ ,14 $\beta$ -card-20(22)-enolide [71-63-6]

Digitoxin, when dried, contains not less than 90.0% of  $C_{41}H_{64}O_{13}$ .

**Description** Digitoxin occurs as a white to light yellowish white, crystalline powder. It is odorless.

It is soluble in chloroform, sparingly soluble in methanol and in ethanol (95), and practically insoluble in water and in diethyl ether.

**Identification** (1) Transfer 1 mg of Digitoxin to a small test tube about 10 mm in inside diameter, dissolve in 1 mL of a solution of iron (III) chloride hexahydrate in acetic acid (100) (1 in 10,000), and underlay gently with 1 mL of sulfuric acid: at the zone of contact of the two liquids a brown ring free from a reddish color is produced, and the color of the upper layer near the contact zone changes to green through purple. Finally the color of the entire acetic acid layer changes to green through deep blue.

(2) To 2 mg of Digitoxin add 25 mL of a freshly prepared solution of 1,3-dinitrobenzene in ethanol (95) (1 in 100), and dissolve by shaking. Take 2 mL of this solution, add 2 mL of a solution of tetramethylammonium hydroxide in ethanol (95) (1 in 200), and mix: a red-purple color develops slowly, and then fades.

(3) Dissolve 1 mg each of Digitoxin and Digitoxin Reference Standard in a mixture of chloroform and ethanol (95) (1:1) to make 50 mL, and use these solutions as the sample solution and the standard solution, respectively. Perform the test with these solutions as directed under Thin-layer Chromatography <2.03>. Spot 20  $\mu$ L each of the sample solution and standard solution on a plate of silica gel for thin-layer chromatography. Develop the plate with a mixture of dichloromethane, methanol and water (84:15:1) to a distance of about 10 cm, and air-dry the plate. Spray evenly dilute sulfuric acid upon the plate, and heat at 110°C for 10 minutes: the spot from the sample solution shows the same  $R_f$  value as

the spot from the standard solution.

**Optical rotation** <2.49>  $[\alpha]_D^{20}$ : +16 – +18° (after drying. 0.5 g, chloroform, 20 mL, 200 mm).

**Purity** Digitonin—Dissolve 10 mg of Digitoxin in 2 mL of ethanol (95) in a test tube, having the inner walls which are free from scratches, add 2 mL of a solution of cholesterol in ethanol (95) (1 in 200), mix gently, and allow to stand for 10 minutes: no turbidity is produced.

**Loss on drying** <2.41> Not more than 1.5% (0.5 g, in vacuum, 100°C, 2 hours).

**Residue on ignition** <2.44> Not more than 0.5% (0.1 g).

**Assay** Dissolve about 20 mg each of Digitoxin and Digitoxin Reference Standard, previously dried and accurately weighed, in methanol to make exactly 200 mL. Pipet 5 mL each of these solutions, add exactly 10 mL of the internal standard solution to each solution, add 12.5 mL of water, then add methanol to make 50 mL, and use these solutions as the sample solution and standard solution, respectively. Perform the test with 50  $\mu$ L each of the sample solution and standard solution as directed under Liquid Chromatography <2.01> according to the following conditions, and calculate the ratios,  $Q_T$  and  $Q_S$ , of the peak area of digitoxin to that of the internal standard, respectively.

$$\begin{aligned} \text{Amount (mg) of } C_{41}H_{64}O_{13} \\ = W_S \times (Q_T/Q_S) \end{aligned}$$

$W_S$ : Amount (mg) of Digitoxin Reference Standard

**Internal standard solution**—A solution of acenaphthene in methanol (3 in 1,000,000).

**Operating conditions**—

**Detector**: An ultraviolet absorption photometer (wavelength: 230 nm).

**Column**: A stainless steel column about 4 mm in inside diameter and 15 to 20 cm in length, packed with octadecylsilanized silica gel for liquid chromatography (5  $\mu$ m in particle diameter).

**Column temperature**: Room temperature.

**Mobile phase**: A mixture of methanol and water (3:1).

**Flow rate**: Adjust the flow rate so that the retention time of digitoxin is about 5 minutes.

**Selection of column**: Proceed with 50  $\mu$ L of the standard solution under the above operating conditions, and calculate the resolution. Use a column giving elution of digitoxin and the internal standard in this order with the resolution between these peaks being not less than 6.

**Containers and storage** Containers—Tight containers.

Storage—Light-resistant.

## Digitoxin Tablets

ジギトキシン錠

Digitoxin Tablets contain not less than 90% and not more than 110% of the labeled amount of digitoxin ( $C_{41}H_{64}O_{13}$ : 764.94).

**Method of preparation** Prepare as directed under Tablets, with Digitoxin.



**Identification (1)** Place a portion of powdered Digitoxin Tablets, equivalent to 2 mg of digitoxin ( $C_{41}H_{64}O_{13}$ ) according to the labeled amount, in a separator, shake with 30 mL of water, and shake vigorously with 30 mL of chloroform. Filter the chloroform extract with a funnel on which a small amount of anhydrous sodium sulfate is placed, and transfer to a round-bottomed flask connected by a universal joint. Evaporate the solution to dryness by warming under reduced pressure, and dissolve the residue in 10 mL of chloroform. Transfer 5 mL of this solution to a small test tube about 10 mm in inside diameter, and evaporate to dryness on a water bath with the aid of a current of air. Proceed with the residue as directed in the Identification (1) under Digitoxin.

**(2)** Evaporate 4 mL of the chloroform solution obtained in (1) to dryness, by warming under reduced pressure, add a freshly prepared solution of 1,3-dinitrobenzene in ethanol (95) (1 in 100) to the residue, and dissolve by shaking. Proceed with 2 mL of this solution as directed in the Identification (2) under Digitoxin.

**Uniformity of dosage units <6.02>** Perform the test according to the following method: it meets the requirement of the Content uniformity test.

Transfer 1 tablet of Digitoxin Tablets to a 50-mL beaker, add 0.5 mL of water to disintegrate the tablet, add 5 mL of acetonitrile, and warm on a water bath for 5 minutes, covering the beaker with a watch glass. After cooling, transfer the solution to separator A, rinse the beaker with 30 mL of chloroform and then with 20 mL of water, transfer the rinsings to separator A, and extract by vigorous shaking. Transfer the chloroform extract to separator B containing 5 mL of a solution of sodium hydrogen carbonate (1 in 100), and shake to wash. Filter the chloroform layer through a pledget of absorbent cotton, previously moistened with chloroform. Extract the water layer in separator A with two 30-mL portions of chloroform, wash the chloroform extract with a solution of sodium hydrogen carbonate (1 in 100) in separator B, filter in the same manner, and combine the filtrate with the first one. Evaporate this filtrate to dryness under reduced pressure by warming, add diluted ethanol (95) (4 in 5) to make exactly  $V$  mL of a solution containing  $5 \mu\text{g}$  of digitoxin ( $C_{41}H_{64}O_{13}$ ) per mL. Shake vigorously for 20 minutes to dissolve, and use this solution as the sample solution. Separately, weigh accurately about 10 mg of Digitoxin Reference Standard, previously dried at  $100^\circ\text{C}$  for 2 hours, and dissolve in diluted ethanol (95) (4 in 5) to make exactly 100 mL. Pipet 5 mL of this solution, add diluted ethanol (95) (4 in 5) to make exactly 100 mL, and use this solution as the standard solution. Pipet 2 mL each of the sample solution, the standard solution and diluted ethanol (95) (4 in 5) into brown glass-stoppered test tubes T, S and B. Add exactly 10 mL each of 0.02 w/v% L-ascorbic acid-hydrochloric acid TS, shake well, and immediately add exactly 1 mL each of dilute hydrogen peroxide TS. Shake vigorously, and allow to stand at a constant temperature between  $25^\circ\text{C}$  and  $30^\circ\text{C}$  for 45 minutes. Determine the fluorescence intensities,  $F_T$ ,  $F_S$  and  $F_B$ , of these solutions at 400 nm of the excitation wavelength and at about 570 nm of the fluorescence wavelength as directed under Fluorometry <2.22>, respectively.

$$\begin{aligned} &\text{Amount (mg) of digitoxin } (C_{41}H_{64}O_{13}) \\ &= W_S \times \{(F_T - F_B)/(F_S - F_B)\} \times (V/2000) \end{aligned}$$

$W_S$ : Amount (mg) of Digitoxin Reference Standard

**Dissolution <6.10>** Perform the test according to the following method: it meets the requirement.

Take 1 tablet of Digitoxin Tablets, and perform the test using 500 mL of diluted hydrochloric acid (3 in 500), degassed by a suitable method, as the test solution at 100 revolutions per minute as directed in the Basket method. At 30 minutes after starting the test, take  $a + 15$  mL of the dissolved solution, and immediately add the same volume of fresh test solution, previously warmed at  $37 \pm 0.5^\circ\text{C}$ , to the vessel carefully. Filter  $a + 15$  mL of the dissolved solution through a membrane filter (less than  $0.8 \mu\text{m}$  in pore size). Discard the first 10 mL of the filtrate, and use the subsequent filtrate as the sample solution. Measure exactly  $a$  mL of the sample solution, equivalent to about  $2 \mu\text{g}$  of digitoxin ( $C_{41}H_{64}O_{13}$ ) according to the labeled amount, transfer to a glass-stoppered centrifuge tube  $T_{30}$ , and warm at  $37 \pm 0.5^\circ\text{C}$  for 30 minutes. Further, at 60 minutes after starting the test, take  $a + 15$  mL of the dissolved solution, proceed in the same manner, measure exactly  $a$  mL of the sample solution, and transfer to a glass-stoppered centrifuge tube  $T_{60}$ . Separately, weigh accurately 100 times the labeled amount of Digitoxin Reference Standard, previously dried under reduced pressure at  $100^\circ\text{C}$  for 2 hours, and dissolve in ethanol (95) to make exactly 100 mL. Measure exactly 1 mL of this solution, add the test solution to make exactly 500 mL, warm at  $37 \pm 0.5^\circ\text{C}$  for 60 minutes, and filter through a membrane filter (less than  $0.8 \mu\text{m}$  in pore size). Discard the first 10 mL of the filtrate, and use the subsequent filtrate as the standard solution. Measure exactly  $a$  mL each of the standard solution and the test solution, transfer to glass-stoppered centrifuge tubes  $T_S$  and  $T_B$ , respectively. Add exactly 7 mL of chloroform to each of the glass-stoppered centrifuge tubes  $T_{30}$ ,  $T_{60}$ ,  $T_S$  and  $T_B$ , shake vigorously for 10 minutes and centrifuge. Discard the aqueous layer, measure exactly 5 mL of the chloroform layer, transfer to brown test tubes  $T'_{30}$ ,  $T'_{60}$ ,  $T'_S$  and  $T'_B$ , evaporate the chloroform, add exactly 4 mL each of 0.05 w/v% L-ascorbic acid-hydrochloric acid TS, shake well, and allow to stand for 10 minutes. Then add exactly 0.5 mL each of dilute hydrogen peroxide TS, shake well, and allow to stand at a constant temperature between  $25^\circ\text{C}$  and  $30^\circ\text{C}$  for 45 minutes. Determine the fluorescence intensities,  $F_{30}$ ,  $F_{60}$ ,  $F_S$  and  $F_B$ , of these solutions at about 395 nm of the excitation wavelength and at about 560 nm of the fluorescence wavelength as directed under Fluorometry <2.22>, respectively. Dissolution rates of Digitoxin Tablets after 30 minutes and 60 minutes should be not less than 60% and 85%, respectively.

No retest requirement is applied to Digitoxin Tablets.

$$\begin{aligned} &\text{Dissolution rate (\%)} \text{ with respect to the labeled} \\ &\text{amount of digitoxin } (C_{41}H_{64}O_{13}) \text{ for 30 minutes} \\ &= W_S \times \{(F_{30} - F_B)/(F_S - F_B)\} \times (1/C) \end{aligned}$$

$$\begin{aligned} &\text{Dissolution rate (\%)} \text{ with respect to the labeled} \\ &\text{amount of digitoxin } (C_{41}H_{64}O_{13}) \text{ for 60 minutes} \\ &= W_S \times \left( \frac{F_{60} - F_B}{F_S - F_B} + \frac{F_{30} - F_B}{F_S - F_B} \times \frac{a + 15}{500} \right) \\ &\quad \times \frac{1}{C} \end{aligned}$$

$W_S$ : Amount (mg) of Digitoxin Reference Standard.

$C$ : The labeled amount (mg) of digitoxin ( $C_{41}H_{64}O_{13}$ ) in 1 tablet.

$a + 15$ : Measured volume (mL) of dissolved solution at the

specified time.

**Assay** Weigh accurately and powder not less than 20 Digitoxin Tablets. Weigh accurately a portion of the powder, equivalent to about 0.5 mg of digitoxin ( $C_{41}H_{64}O_{13}$ ), and shake with 12.5 mL of water for 10 minutes. Add exactly 10 mL of the internal standard solution, shake for 20 minutes, and add methanol to make 50 mL. Centrifuge this solution, and use the supernatant liquid as the sample solution. Separately, weigh accurately about 20 mg of Digitoxin Reference Standard, previously dried in vacuum at 100°C for 2 hours, dissolve in methanol to make exactly 200 mL. Pipet 5 mL of the solution, add exactly 10 mL of the internal standard solution, add 12.5 mL of water, then methanol to make 50 mL, and use this solution as the standard solution. Proceed as directed in the Assay under Digitoxin.

$$\begin{aligned} &\text{Amount (mg) of digitoxin } (C_{41}H_{64}O_{13}) \\ &= W_S \times (Q_T/Q_S) \times 0.025 \end{aligned}$$


$W_S$ : Amount (mg) of Digitoxin Reference Standard

**Internal standard solution**—A solution of acenaphthene in methanol (3 in 1,000,000).

**Containers and storage** Containers—Tight containers.  
Storage—Light-resistant.

## Digoxin

ジゴキシン



$C_{41}H_{64}O_{14}$ : 780.94

3 $\beta$ -[2,6-Dideoxy- $\beta$ -D-ribo-hexopyranosyl-(1 $\rightarrow$ 4)-2,6-dideoxy- $\beta$ -D-ribo-hexopyranosyl-(1 $\rightarrow$ 4)-2,6-dideoxy- $\beta$ -D-ribo-hexopyranosyloxy]-12 $\beta$ ,14-dihydroxy-5 $\beta$ ,14 $\beta$ -card-20(22)-enolide [20830-75-5]

Digoxin, when dried, contains not less than 96.0% and not more than 106.0% of  $C_{41}H_{64}O_{14}$ .

**Description** Digoxin occurs as colorless or white crystals or a white crystalline powder.

It is freely soluble in pyridine, slightly soluble in ethanol (95), very slightly soluble in acetic acid (100), and practically insoluble in water.

**Identification** (1) Transfer 1 mg of Digoxin to a small test tube about 10 mm in inside diameter, dissolve in 1 mL of a

solution of iron (III) chloride hexahydrate in acetic acid (100) (1 in 10,000), and underlay gently with 1 mL of sulfuric acid: at the zone of contact of the two liquids a brown ring free from a reddish color is produced, and the color of the upper layer near the contact zone changes to green through purple. Finally the entire acetic acid layer shows a green color through a deep blue color.

(2) Determine the infrared absorption spectrum of Digoxin, previously dried, as directed in the potassium bromide disk method under Infrared Spectrophotometry <2.25>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wave numbers.

**Optical rotation** <2.49>  $[\alpha]_D^{20}$ : +10.0 – +13.0° (after drying, 0.20 g, dehydratead pyridine, 10 mL, 100 mm).

**Purity** (1) Clarity and color of solution—Dissolve 0.10 g of Digoxin in 15 mL of diluted ethanol (95) (4 in 5) by warming: the solution is clear and colorless.

(2) Related substances—Dissolve exactly 25.0 mg of Digoxin in 50 mL of warm ethanol (95), cool, and add ethanol (95) to make exactly 100 mL. Pipet 10 mL of this solution, add 10 mL of water and dilute ethanol to make 50 mL, and use this solution as the sample solution. Separately, dissolve exactly 5.0 mg of Gitoxin Reference Standard, previously dried under reduced pressure at 105°C for 1 hour, in a mixture of acetonitrile and water (7:3) to make exactly 200 mL. Pipet 2 mL of this solution, add dilute ethanol to make 50 mL, and use this solution as the standard solution. Perform the test with exactly 10  $\mu$ L each of the sample solution and standard solution as directed under Liquid Chromatography <2.01> according to the following conditions, and determine the peak areas,  $A_T$  and  $A_S$ , of gitoxin:  $A_T$  is not larger than  $A_S$ , and the total of the areas of the peaks other than digitoxin and gitoxin, obtained by the area percentage method, is not more than 3%.

**Operating conditions**—

Detector, column, column temperature, mobile phase, and flow rate: Proceed as directed in the operating conditions in the Assay.

Time span of measurement: About 4 times as long as the retention time of digoxin beginning after the solvent peak.

**System suitability**—

Test for required detectability: Pipet 1 mL of the sample solution, add the mobile phase to make exactly 100 mL, and use this solution as the solution for system suitability test. Pipet 1 mL of the solution, and add the mobile phase to make exactly 10 mL. Confirm that the peak area of digoxin obtained from 10  $\mu$ L of this solution is equivalent to 7 to 13% of that from the solution for system suitability test.

System performance: Dissolve 25 mg of digoxin in 50 mL of warm ethanol (95), cool, and add ethanol (95) to make exactly 100 mL. Pipet 10 mL of this solution, add exactly 5 mL of a solution of propyl parahydroxybenzoate in ethanol (95) (1 in 4000), 10 mL of water and dilute ethanol to make 50 mL. When the procedure is run with 10  $\mu$ L of this solution under the above operating conditions, digoxin and propyl parahydroxybenzoate are eluted in this order with the resolution between these peaks being not less than 5.

System repeatability: When the test is repeated 6 times with 10  $\mu$ L of the solution for system suitability test under the above operating conditions, the relative standard deviation of the peak area of digoxin is not more than 1.0%.

**Loss on drying** <2.41> Not more than 1.0% (0.5 g, in vacuum, 105°C, 1 hour).

**Residue on ignition** <2.44> Not more than 0.5% (0.1 g).

**Assay** Weigh accurately about 25 mg each of Digoxin and Digoxin Reference Standard, previously dried, dissolve in 50 mL of warm ethanol (95), cool, and add ethanol (95) to make exactly 100 mL. Pipet 10 mL of these solutions, add exactly 5 mL of the internal standard solution, 10 mL of water and dilute ethanol to make 50 mL, and use these solutions as the sample solution and standard solution. Perform the test with 10  $\mu$ L each of the sample solution and standard solution as directed under Liquid Chromatography <2.01> according to the following conditions, and determine the ratios,  $Q_T$  and  $Q_S$ , of the peak area of digoxin to that of the internal standard.

$$\text{Amount (mg) of } C_{41}H_{64}O_{14} = W_S \times (Q_T/Q_S)$$

$W_S$ : Amount (mg) of Digoxin Reference Standard

**Internal standard solution**—A solution of propyl parahydroxybenzoate in ethanol (95) (1 in 4000).

**Operating conditions**—

**Detector**: An ultraviolet absorption photometer (wavelength: 220 nm).

**Column**: A stainless steel column 4.6 mm in inside diameter and 25 cm in length, packed with octadecylsilanized silica gel for liquid chromatography (5  $\mu$ m in particle diameter).

**Column temperature**: A constant temperature of about 30°C.

**Mobile phase**: A mixture of water and acetonitrile (7:3).

**Flow rate**: Adjust the flow rate so that the retention time of digoxin is about 10 minutes.

**System suitability**—

**System performance**: When the procedure is run with 10  $\mu$ L of the standard solution under the above operating conditions, digoxin and the internal standard are eluted in this order with the resolution between these peaks being not less than 5.

**System repeatability**: When the test is repeated 6 times with 10  $\mu$ L of the standard solution under the above operating conditions, the relative standard deviation of the ratio of the peak area of digoxin to that of the internal standard is not more than 1.0%.

**Containers and storage** Containers—Tight containers.

Storage—Light-resistant.

## Digoxin Injection

ジゴキシン注射液

Digoxin Injection is an aqueous solution for injection.

It contains not less than 90.0% and not more than 105.0% of the labeled amount of digoxin ( $C_{41}H_{64}O_{14}$ ; 780.94).

**Method of preparation** Prepare as directed under Injections, with a solution of Digoxin in 5 to 50 vol% ethanol.

**Description** Digoxin Injection is a clear, colorless liquid.

**Identification** Dilute Digoxin Injection, if necessary, with methanol so that each mL contains about 0.25 mg of Digoxin according to the labeled amount, and use this solution as the sample solution. In case where ingredients are suspected to affect the test, remove them by means of a solid-phase extraction. Separately, dissolve 0.5 mg of Digoxin Reference Standard in 2 mL of methanol, and use this solution as the standard solution. Perform the test with these solutions as directed under Thin-layer Chromatography <2.03>. Spot 10  $\mu$ L each of the sample solution and standard solution on a plate of octadecylsilanized silica gel for thin-layer chromatography. Develop the plate with a mixture of methanol and water (7:3) to a distance of about 10 cm, and air-dry the plate. Spray evenly a mixture of a solution of trichloroacetic acid in ethanol (99.5) (1 in 4) and a freshly prepared solution of sodium toluenesulfonchloramide trihydrate (3 in 100) (4:1) on the plate, heat at 110°C for 10 minutes, and examine under ultraviolet light (main wavelength: 366 nm): the  $R_f$  values of the principal spots with the sample solution and the standard solution are not different each other.

**Bacterial endotoxins** <4.01> Less than 200 EU/mg.

**Extractable volume** <6.05> It meets the requirements.

**Foreign insoluble matter** <6.06> Perform the test according to Method 1: it meets the requirement.

**Insoluble particulate matter** <6.07> Perform the test according to Method 1: it meets the requirement.

**Sterility** <4.06> Perform the test according to the Membrane filtration method: it meets the requirement.

**Assay** To an exact volume of Digoxin Injection, equivalent to about 2.5 mg of digoxin ( $C_{41}H_{64}O_{14}$ ), add exactly 5 mL of the internal standard solution and dilute ethanol to make 50 mL, and use this solution as the sample solution. Separately, weigh accurately about 25 mg of Digoxin Reference Standard, previously dried under reduced pressure at 105°C for 1 hour, dissolve in 50 mL of warm ethanol (95), cool, and add ethanol (95) to make exactly 100 mL. Pipet 10 mL of this solution, add exactly 5 mL of the internal standard solution, 10 mL of water and dilute ethanol to make 50 mL, and use this solution as the standard solution. Perform the test with 10  $\mu$ L each of the sample solution and standard solution as directed under Liquid Chromatography <2.01> according to the following conditions, and determine the ratios,  $Q_T$  and  $Q_S$ , of the peak area of digoxin to that of the internal standard.

$$\begin{aligned} \text{Amount (mg) of digoxin (} C_{41}H_{64}O_{14} \text{)} \\ = W_S \times (Q_T/Q_S) \times (1/10) \end{aligned}$$

$W_S$ : Amount (mg) of Digoxin Reference Standard

**Internal standard solution**—A solution of propyl parahydroxybenzoate in ethanol (95) (1 in 4000).

**Operating conditions**—

**Detector**: An ultraviolet absorption photometer (wavelength: 220 nm).

**Column**: A stainless steel column 4.6 mm in inside diameter and 25 cm in length, packed with octadecylsilanized silica gel for liquid chromatography (5  $\mu$ m in particle diameter).

**Column temperature**: A constant temperature of about 30°C.

Mobile phase: A mixture of water and acetonitrile (7:3).

Flow rate: Adjust the flow rate so that the retention time of digoxin is about 10 minutes.

*System suitability*—

System performance: When the procedure is run with 10  $\mu$ L of the standard solution under the above operating conditions, digoxin and the internal standard are eluted in this order with the resolution between these peaks being not less than 5.

System repeatability: When the test is repeated 6 times with 10  $\mu$ L of the standard solution under the above operating conditions, the relative standard deviation of the ratio of the peak area of digoxin to that of the internal standard is not more than 1.0%.

**Containers and storage** Containers—Hermetic containers, and colored containers may be used.

Storage—Light-resistant.

## Digoxin Tablets

ジゴキシン錠

Digoxin Tablets contain not less than 90.0% and not more than 105.0% of the labeled amount of digoxin ( $C_{41}H_{64}O_{14}$ ; 780.94).

**Method of preparation** Prepare as directed under Tablets, with Digoxin.

**Identification** To an amount of pulverized Digoxin Tablets, equivalent to 0.5 mg of Digoxin according to the labeled amount, add 2 mL of methanol, shake for 10 minutes, filter, and use the filtrate as the sample solution. Separately, dissolve 0.5 mg of Digoxin Reference Standard in 2 mL of methanol, and use this as the standard solution. Perform the test with these solutions as directed under Thin-layer Chromatography <2.03>. Spot 10  $\mu$ L each of the sample solution and standard solution on a plate of octadecylsilanized silica gel for thin-layer chromatography. Develop the plate with a mixture of methanol and water (7:3) to a distance of about 10 cm, and air-dry the plate. Spray evenly a mixture of a solution of trichloroacetic acid in ethanol (99.5) (1 in 4) and a freshly prepared solution of sodium toluenesulfonchloramide trihydrate (3 in 100) (4:1) on the plate, heat at 110°C for 10 minutes, and examine under ultraviolet light (main wavelength: 366 nm): the  $R_f$  values of the principal spots with the sample solution and the standard solution are not different each other.

**Uniformity of dosage units** <6.02> Perform the test according to the following method: it meets the requirement of the Content uniformity test.

To 1 tablet of Digoxin Tablets add 0.5 mL of water to disintegrate, then add exactly 0.5 mL of the internal standard solution, and add  $V$  mL of dilute ethanol so that each mL contains about 21  $\mu$ g of digoxin ( $C_{41}H_{64}O_{14}$ ). Exposure this solution to ultrasonic waves for 20 minutes, shake for 5 minutes, filter, and use the filtrate as the sample solution. Separately, weigh accurately about 25 mg of Digoxin Reference Standard, previously dried under reduced pressure at 105°C for 1 hour, dissolve in 50 mL of warm ethanol (95), cool, and add ethanol (95) to make exactly 100 mL. Pipet

10 mL of this solution, and add ethanol (95) to make exactly 20 mL. Pipet 1 mL of this solution, add exactly 0.5 mL of the internal standard solution, then add 1.5 mL of water and ( $V - 2$ ) mL of dilute ethanol, and use this as the standard solution. Proceed with the sample solution and standard solution as directed in the Assay.

$$\begin{aligned} &\text{Amount (mg) of digoxin (C}_{41}\text{H}_{64}\text{O}_{14}) \\ &= W_S \times (Q_T/Q_S) \times (1/200) \end{aligned}$$

$W_S$ : Amount (mg) of Digoxin Reference Standard

*Internal standard solution*—A solution of propyl parahydroxybenzoate in ethanol (95) (1 in 40,000/ $V$ ).

**Dissolution** <6.10> Perform the test according to the following method: it meets the requirement.

Perform the test with 1 tablet of Digoxin Tablets, using 500 mL of diluted hydrochloric acid (3 in 500) at 100 revolutions per minute according to the Basket method as the dissolution medium. Withdraw 30 mL or more of the dissolved solution 60 minutes after starting the test, and filter through a membrane filter (less than 0.8  $\mu$ m in pore size). Discard the first 10 mL of the filtrate, and use the subsequent filtrate as the sample solution. Separately, weigh accurately about 25 mg of Digoxin Reference Standard, previously dried in vacuum at 105°C for 1 hour, dissolve in a small portion of ethanol (95), and add a mixture of ethanol (95) and water (4:1) to make exactly 500 mL. Pipet 5 mL of this solution, add the dissolution medium to make exactly 500 mL, and use this solution as the standard solution. Pipet 2 mL each of the sample solution, the standard solution and the dissolution medium, and transfer to brown glass-stoppered test tubes. Add exactly 10 mL of 0.012 g/dL L-ascorbic acid-hydrochloric acid TS to these tubes, and shake. Immediately add exactly 1 mL of dilute hydrogen peroxide TS, shake well, and allow to stand at a constant temperature between 25°C and 30°C for 45 minutes. Determine the fluorescence intensities,  $F_T$ ,  $F_S$ , and  $F_B$ , of these solutions at 360 nm of the excitation wavelength and at 485 nm of the fluorescence wavelength as directed under Fluorometry <2.22>, respectively: the dissolution rate in 60 minutes is not less than 65%. No retest requirement is applied to Digoxin Tablets.

Dissolution rate (%) with respect to the labeled amount of digoxin ( $C_{41}H_{64}O_{14}$ )

$$= W_S \times \{(F_T - F_B)/(F_S - F_B)\} \times (1/C)$$

$W_S$ : Amount (mg) of Digoxin Reference Standard

$C$ : The labeled amount (mg) of digoxin ( $C_{41}H_{64}O_{14}$ ) in 1 tablet

**Assay** Weigh accurately the mass of not less than 20 Digoxin Tablets, and powder. Weigh accurately a portion of the powder, equivalent to about 2.5 mg of digoxin ( $C_{41}H_{64}O_{14}$ ), add 30 mL of dilute ethanol, exposure to ultrasonic waves for 20 minutes, and shake for 5 minutes. Add exactly 5 mL of the internal standard solution and dilute ethanol to make 50 mL, centrifuge, and use the supernatant liquid as the sample solution. Separately, weigh accurately about 25 mg of Digoxin Reference Standard, previously dried under reduced pressure at 105°C for 1 hour, dissolve in 50 mL of warm ethanol (95), cool, and add ethanol (95) to make exactly 100 mL. Pipet 10 mL of this solution, add exactly 5 mL of the internal standard solution, 10 mL of water and dilute ethanol to make 50 mL, and use this solution as the standard solution. Perform

the test with 10  $\mu$ L each of the sample solution and standard solution as directed under Liquid Chromatography <2.01> according to the following conditions, and determine the ratios,  $Q_T$  and  $Q_S$ , of the peak area of digoxin to that of the internal standard.

$$\begin{aligned} \text{Amount (mg) of digoxin (C}_{41}\text{H}_{64}\text{O}_{14}) \\ = W_S \times (Q_T/Q_S) \times (1/10) \end{aligned}$$

$W_S$ : Amount (mg) of Digoxin Reference Standard

**Internal standard solution**—A solution of propyl parahydroxybenzoate in ethanol (95) (1 in 4000).

**Operating conditions**—

**Detector:** An ultraviolet absorption photometer (wavelength: 220 nm).

**Column:** A stainless steel column 4.6 mm in inside diameter and 25 cm in length, packed with octadecylsilanized silica gel for liquid chromatography (5  $\mu$ m in particle diameter).

**Column temperature:** A constant temperature of about 30°C.

**Mobile phase:** A mixture of water and acetonitrile (7:3).

**Flow rate:** Adjust the flow rate so that the retention time of digoxin is about 10 minutes.

**System suitability**—

**System performance:** When the procedure is run with 10  $\mu$ L of the standard solution under the above operating conditions, digoxin and the internal standard are eluted in this order with the resolution between these peaks being not less than 5.


**System repeatability:** When the test is repeated 6 times with 10  $\mu$ L of the standard solution under the above operating conditions, the relative standard deviation of the ratio of the peak area of digoxin to that of the internal standard is not more than 1.0%.

**Containers and storage** Containers—Tight containers.

Storage—Light-resistant.

## Dihydrocodeine Phosphate

ジヒドロコデインリン酸塩



$\text{C}_{18}\text{H}_{23}\text{NO}_3 \cdot \text{H}_3\text{PO}_4$ : 399.38

(5*R*,6*S*)-4,5-Epoxy-3-methoxy-17-methylmorphinan-6-ol monophosphate [24204-13-5]

Dihydrocodeine Phosphate contains not less than 98.0% of  $\text{C}_{18}\text{H}_{23}\text{NO}_3 \cdot \text{H}_3\text{PO}_4$ , calculated on the dried basis.

**Description** Dihydrocodeine Phosphate occurs as a white to yellowish white, crystalline powder.

It is freely soluble in water and in acetic acid (100), slightly soluble in ethanol (95), and practically insoluble in diethyl ether.

The pH of a solution of Dihydrocodeine Phosphate (1 in 10) is between 3.0 and 5.0.

It is affected by light.

**Identification (1)** Determine the absorption spectrum of a solution of Dihydrocodeine Phosphate (1 in 10,000) as directed under Ultraviolet-visible Spectrophotometry <2.24>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wavelengths.

**(2)** Determine the infrared spectrum of Dihydrocodeine Phosphate, previously dried, as directed in the potassium bromide disk method under Infrared Spectrophotometry <2.25>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wave numbers.

**(3)** A solution of Dihydrocodeine Phosphate (1 in 20) responds to the Qualitative Tests <1.09> (1) for phosphate.

**Purity (1) Chloride <1.03>**—Perform the test with 0.5 g of Dihydrocodeine Phosphate. Prepare the control solution with 0.30 mL of 0.01 mol/L hydrochloric acid (not more than 0.021%).

**(2) Sulfate <1.14>**—Perform the test with 0.20 g of Dihydrocodeine Phosphate. Prepare the control solution with 1.0 mL of 0.005 mol/L sulfuric acid (not more than 0.240%).

**(3) Related substances**—Dissolve 0.20 g of Dihydrocodeine Phosphate in 10 mL of diluted ethanol (95) (1 in 2), and use this solution as the sample solution. Pipet 1 mL of the sample solution, add diluted ethanol (95) (1 in 2) to make exactly 50 mL, and use this solution as the standard solution. Perform the test with these solutions as directed under Thin-layer Chromatography <2.03>. Spot 10  $\mu$ L of the sample solution and standard solution on a plate of silica gel with fluorescent indicator for thin-chromatography. Develop the plate with a mixture of ethanol (99.5), toluene, acetone and ammonia solution (28) (14:14:7:1) to a distance of about 15 cm, and air-dry the plate. Examine under ultraviolet light (main wavelength: 254 nm): the spots other than the principal spot from the sample solution are not more intense than the spot from the standard solution.

**Loss on drying <2.41>** Not more than 1.0% (0.5 g, 105°C, 4 hours).

**Assay** Weigh accurately about 0.5 g of Dihydrocodeine Phosphate, dissolve in 70 mL of acetic acid (100), and titrate <2.50> with 0.1 mol/L perchloric acid VS until the color of the solution changes from purple through blue to greenish blue (indicator: 3 drops of crystal violet TS). Perform a blank determination, and make any necessary correction.

$$\begin{aligned} \text{Each mL of 0.1 mol/L perchloric acid VS} \\ = 39.94 \text{ mg of } \text{C}_{18}\text{H}_{23}\text{NO}_3 \cdot \text{H}_3\text{PO}_4 \end{aligned}$$

**Containers and storage** Containers—Tight containers.

Storage—Light-resistant.

## 1% Dihydrocodeine Phosphate Powder

ジヒドロコデインリン酸塩散 1%

1% Dihydrocodeine Phosphate Powder contains not less than 0.90% and not more than 1.10% of dihydrocodeine phosphate ( $C_{18}H_{23}NO_3 \cdot H_3PO_4$ ; 399.38).

### Method of preparation

|                          |                       |
|--------------------------|-----------------------|
| Dihydrocodeine Phosphate | 10 g                  |
| Lactose Hydrate          | a sufficient quantity |
| To make                  | 1000 g                |

Prepare as directed under Powders, with the above ingredients.

**Identification** Determine the absorption spectrum of a solution of 1% Dihydrocodeine Phosphate Powder (1 in 100) as directed under Ultraviolet-visible Spectrophotometry <2.24>: it exhibits a maximum between 281 nm and 285 nm.

**Assay** Weigh accurately about 5 g of 1% Dihydrocodeine Phosphate Powder, dissolve in water to make exactly 100 mL, then pipet 10 mL of this solution, add exactly 10 mL of the internal standard solution, and use this solution as the sample solution. Separately, weigh accurately about 50 mg of dihydrocodeine phosphate for assay (previously determine the loss on drying <2.41> (105°C, 4 hours)), dissolve in water to make exactly 100 mL, then pipet 10 mL of this solution, add exactly 10 mL of the internal standard solution, and use this solution as the standard solution. Perform the test with 20  $\mu$ L each of the sample solution and standard solution as directed under Liquid Chromatography <2.01> according to the following conditions, and calculate the ratios,  $Q_T$  and  $Q_S$ , of the peak area of dihydrocodeine to that of the internal standard.

$$\begin{aligned} &\text{Amount (mg) of dihydrocodeine phosphate} \\ & (C_{18}H_{23}NO_3 \cdot H_3PO_4) \\ & = W_S \times (Q_T/Q_S) \end{aligned}$$

$W_S$ : Amount (mg) of dihydrocodeine phosphate for assay, calculated on the dried basis

**Internal standard solution**—A solution of ethylefuran hydrochloride (3 in 10,000).

### Operating conditions—

**Detector**: An ultraviolet absorption photometer (wavelength: 280 nm).

**Column**: A stainless steel column 4.6 mm in inside diameter and 15 cm in length, packed with octadecylsilanized silica gel for liquid chromatography (5  $\mu$ m in particle diameter).

**Column temperature**: A constant temperature of about 40°C.

**Mobile phase**: Dissolve 1.0 g of sodium lauryl sulfate in 500 mL of diluted phosphoric acid (1 in 1000), and adjust the pH to 3.0 with sodium hydroxide TS. To 240 mL of this solution add 70 mL of tetrahydrofuran.

**Flow rate**: Adjust the flow rate so that the retention time of dihydrocodeine is about 9 minutes.

### System suitability—

**System performance**: When the procedure is run with 20  $\mu$ L of the standard solution under the above operating conditions, dihydrocodeine and the internal standard are eluted in this order with the resolution between these peaks being not less than 4.

**System repeatability**: When the test is repeated 5 times with 20  $\mu$ L of the standard solution under the above operating conditions, the relative standard deviation of the ratios of the peak area of dihydrocodeine to that of the internal standard is not more than 1.0%.

**Containers and storage** Containers—Tight containers.

## 10% Dihydrocodeine Phosphate Powder

ジヒドロコデインリン酸塩散 10%

10% Dihydrocodeine Phosphate Powder contains not less than 9.3% and not more than 10.7% of dihydrocodeine phosphate ( $C_{18}H_{23}NO_3 \cdot H_3PO_4$ ; 399.38).

### Method of preparation

|                          |                       |
|--------------------------|-----------------------|
| Dihydrocodeine Phosphate | 100 g                 |
| Lactose Hydrate          | a sufficient quantity |
| To make                  | 1000 g                |

Prepare as directed under Powders, with the above ingredients.

**Identification** Determine the absorption spectrum of a solution of 10% Dihydrocodeine Phosphate Powder (1 in 1000) as directed under Ultraviolet-visible Spectrophotometry <2.24>: it exhibits a maximum between 281 nm and 285 nm.

**Assay** Weigh accurately about 2.5 g of 10% Dihydrocodeine Phosphate Powder, dissolve in water to make exactly 100 mL, then pipet 2 mL of this solution, add exactly 10 mL of the internal standard solution and water to make 20 mL, and use this solution as the sample solution. Separately, weigh accurately about 50 mg of dihydrocodeine phosphate for assay, (previously determine the loss on drying <2.41> (105°C, 4 hours)), dissolve in water to make exactly 100 mL, then pipet 10 mL of this solution, add exactly 10 mL of the internal standard solution, and use this solution as the standard solution. Perform the test with 20  $\mu$ L each of the sample solution and standard solution as directed under Liquid Chromatography <2.01> according to the following conditions, and calculate the ratios,  $Q_T$  and  $Q_S$ , of the peak area of dihydrocodeine to that of the internal standard.

$$\begin{aligned} &\text{Amount (mg) of dihydrocodeine phosphate} \\ & (C_{18}H_{23}NO_3 \cdot H_3PO_4) \\ & = W_S \times (Q_T/Q_S) \times 5 \end{aligned}$$

$W_S$ : Amount (mg) of dihydrocodeine phosphate for assay, calculated on the dried basis

**Internal standard solution**—A solution of ethylefrine hydrochloride (3 in 10,000).

### Operating conditions—

**Detector**: An ultraviolet absorption photometer (wave-

length: 280 nm).

Column: A stainless steel column 4.6 mm in inside diameter and 15 cm in length, packed with octadecylsilanized silica gel for liquid chromatography (5  $\mu$ m in particle diameter).

Column temperature: A constant temperature of about 40°C.

Mobile phase: Dissolve 1.0 g of sodium lauryl sulfate in 500 mL of diluted phosphoric acid (1 in 1000), and adjust the pH to 3.0 with sodium hydroxide TS. To 240 mL of this solution add 70 mL of tetrahydrofuran.

Flow rate: Adjust the flow rate so that the retention time of dihydrocodeine is about 9 minutes.

System suitability—


System performance: When the procedure is run with 20  $\mu$ L of the standard solution under the above operating conditions, dihydrocodeine and the internal standard are eluted in this order with the resolution between these peaks being not less than 4.

System repeatability: When the test is repeated 5 times with 20  $\mu$ L of the standard solution under the above operating conditions, the relative standard deviation of the ratios of the peak area of dihydrocodeine to that of the internal standard is not more than 1.0%.

**Containers and storage** Containers—Tight containers.

## Dihydroergotamine Mesilate

ジヒドロエルゴタミンメシル酸塩



$C_{33}H_{37}N_5O_5 \cdot CH_4O_3S$ : 679.78  
(5'S,10R)-5'-Benzyl-12'-hydroxy-2'-methyl-9,10-dihydroergotaman-3',6',18-trione monomethanesulfonate  
[6190-39-2]

Dihydroergotamine Mesilate contains not less than 97.0% of  $C_{33}H_{37}N_5O_5 \cdot CH_4O_3S$ , calculated on the dried basis.

**Description** Dihydroergotamine Mesilate occurs as a white to yellowish white or grayish white to reddish white powder.

It is freely soluble in acetic acid (100), sparingly soluble in methanol and in chloroform, slightly soluble in water and in ethanol (95), and practically insoluble in acetic anhydride and in diethyl ether.

It is gradually colored by light.

Melting point: about 214°C (with decomposition).

**Identification (1)** Dissolve 1 mg of Dihydroergotamine Mesilate in 5 mL of a solution of L-tartaric acid (1 in 100). To 1 mL of this solution add 2 mL of 4-dimethylaminobenzaldehyde-ferric chloride TS, and shake: a blue color develops.

(2) To 0.1 g of Dihydroergotamine Mesilate add 0.4 g of sodium hydroxide, stir well, and incinerate by gradual ignition. After cooling, add 10 mL of water to the residue, heat to boiling, cool, and filter. To the filtrate add 0.5 mL of hydrochloric acid: the solution responds to the Qualitative Tests <1.09> for sulfate. Separately, to 0.1 g of Dihydroergotamine Mesilate add 5 mL of dilute hydrochloric acid, shake for 5 minutes, filter, and to the filtrate add 1 mL of barium chloride TS: the solution is clear.

(3) Determine the absorption spectrum of a solution of Dihydroergotamine Mesilate in methanol (1 in 20,000) as directed under Ultraviolet-visible Spectrophotometry <2.24>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wavelengths.

(4) Determine the infrared absorption spectrum of Dihydroergotamine Mesilate, previously dried, as directed in the potassium bromide disk method under Infrared Spectrophotometry <2.25>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wave numbers.

**Optical rotation** <2.49>  $[\alpha]_D^{20}$ :  $-16.7 - -22.7^\circ$  [0.5 g, calculated on the dried basis, a mixture of ethanol (99.5), chloroform and ammonia solution (28) (10:10:1), 20 mL, 100 mm].

**pH** <2.54> Dissolve 0.05 g of Dihydroergotamine Mesilate in 50 mL of water: the pH of this solution is between 4.4 and 5.4.

**Purity (1)** Clarity and color of solution—Dissolve 0.10 g of Dihydroergotamine Mesilate in 0.1 mL of a solution of methanesulfonic acid (7 in 100) and 50 mL of water: the solution is clear, and has no more color than the following control solutions [1] or [2].

Control solution [1]: Pipet 0.6 mL of Ferric Chloride Stock CS and 0.15 mL of Cobaltous Chloride Stock CS, mix, and add diluted hydrochloric acid (1 in 40) to make exactly 100 mL.

Control solution [2]: Pipet 0.6 mL of Ferric Chloride Stock CS, 0.25 mL of Cobaltous Chloride Stock CS and 0.1 mL of copper (II) sulfate CS, mix, and add diluted hydrochloric acid (1 in 40) to make exactly 100 mL.

(2) Related substances—Conduct this procedure without exposure to daylight, using light-resistant vessels. Dissolve 0.10 g of Dihydroergotamine Mesilate in 5 mL of a mixture of chloroform and methanol (9:1), and use this solution as the sample solution. Pipet 1 mL of the sample solution, add a mixture of chloroform and methanol (9:1) to make exactly 200 mL, and use this solution as the standard solution (1). Pipet 10 mL of the standard solution (1), add a mixture of chloroform and methanol (9:1) to make exactly 25 mL, and use this solution as the standard solution (2). Perform the test with these solutions as directed under Thin-layer Chromatography <2.03>. Spot 5  $\mu$ L each of the sample solution and standard solutions (1) and (2) on a plate of silica gel for thin-layer chromatography. Develop the plate with a mixture of dichloromethane, ethyl acetate, methanol and ammonia solution (28) (50:50:6:1) to a distance of about 15 cm, and dry the plate with cold wind within 1 minute. Develop the plate again immediately with a freshly prepared mixture of dichloromethane, ethyl acetate, methanol and ammonia solution (28) (50:50:6:1) to a distance of about 15 cm, and air-dry

the plate. Spray evenly 4-dimethylaminobenzaldehyde TS for spraying on the plate, and dry the plate with warm wind: the spots other than the principal spot from the sample solution are not more intense than the spot from the standard solution (1), and the spots, which are more intense than the spot from the standard solution (2), are not more than two.

**Loss on drying** <2.41> Not more than 4.0% (0.5 g, in vacuum at a pressure not exceeding 0.67 kPa, 100°C, 6 hours).



**Assay** Weigh accurately about 0.2 g of Dihydroergotamine Mesilate, dissolve in 170 mL of a mixture of acetic anhydride and acetic acid (100) (10:1), and titrate <2.50> with 0.02 mol/L perchloric acid VS (potentiometric titration). Perform a blank determination, and make any necessary correction.

Each mL of 0.02 mol/L perchloric acid VS  
= 13.60 mg of  $C_{33}H_{37}N_5O_5 \cdot CH_4O_3S$

**Containers and storage** Containers—Tight containers.  
Storage—Light-resistant.

## Dihydroergotoxine Mesilate

ジヒドロエルゴトキシンメシル酸塩



### Dihydroergocornine Mesilate

$C_{31}H_{41}N_5O_5 \cdot CH_4O_3S$ : 659.79

(5'S,10R)-12'-Hydroxy-2',5'-bis(1-methylethyl)-9,10-dihydroergotaman-3',6',18-trione monomethanesulfonate

### Dihydro- $\alpha$ -ergocryptine Mesilate

$C_{32}H_{43}N_5O_5 \cdot CH_4O_3S$ : 673.82

(5'S,10R)-12'-Hydroxy-2'-(1-methylethyl)-5'-(2-methylpropyl)-9,10-dihydroergotaman-3',6',18-trione monomethanesulfonate

### Dihydro- $\beta$ -ergocryptine Mesilate

$C_{32}H_{43}N_5O_5 \cdot CH_4O_3S$ : 673.82

(5'S,10R)-12'-Hydroxy-2'-(1-methylethyl)-5'-(1-methylpropyl)-9,10-dihydroergotaman-3',6',18-trione monomethanesulfonate

### Dihydroergocristine Mesilate

$C_{35}H_{41}N_5O_5 \cdot CH_4O_3S$ : 707.84

(5'S,10R)-5'-Benzyl-12'-hydroxy-2'-(1-methylethyl)-9,10-dihydroergotaman-3',6',18-trione

monomethanesulfonate

[8067-24-1, Dihydroergotoxine Mesilate]

Dihydroergotoxine Mesilate contains not less than 97.0% and not more than 103.0% of dihydroergotoxine mesilate [as a mixture of dihydroergocornine mesilate ( $C_{31}H_{41}N_5O_5 \cdot CH_4O_3S$ ), dihydro- $\alpha$ -ergocryptine mesilate ( $C_{32}H_{43}N_5O_5 \cdot CH_4O_3S$ ), dihydro- $\beta$ -ergocryptine mesilate ( $C_{32}H_{43}N_5O_5 \cdot CH_4O_3S$ ) and dihydroergocristine mesilate ( $C_{35}H_{41}N_5O_5 \cdot CH_4O_3S$ )], calculated on the anhydrous basis. The relative contents of dihydroergocornine mesilate ( $C_{31}H_{41}N_5O_5 \cdot CH_4O_3S$ ), dihydroergocryptine mesilate ( $C_{32}H_{43}N_5O_5 \cdot CH_4O_3S$ ) and dihydroergocristine mesilate ( $C_{35}H_{41}N_5O_5 \cdot CH_4O_3S$ ) are 30.3–36.3% each, and the content ratio of dihydro- $\alpha$ -ergocryptine mesilate and dihydro- $\beta$ -ergocryptine mesilate is 1.5–2.5:1.

**Description** Dihydroergotoxine Mesilate occurs as a white to pale yellow powder.

It is soluble in methanol, sparingly soluble in ethanol (95), slightly soluble in water, in acetonitrile and in chloroform, and practically insoluble in diethyl ether.

**Identification** Determine the infrared absorption spectrum of Dihydroergotoxine Mesilate as directed in the potassium bromide disk method under Infrared Spectrophotometry <2.25>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wave numbers.

**Optical rotation** <2.49>  $[\alpha]_D^{20}$ : +11.0 – +15.0° (0.2 g, calculated on the anhydrous basis, dilute ethanol, 20 mL, 100 mm).

**Purity** (1) Clarity and color of solution—Dissolve 0.10 g of Dihydroergotoxine Mesilate in 20 mL of water: the solution is clear and the color of the solution is not more intense than that of the following control solution.

Control solution: To a mixture of 1.0 mL of Cobaltous Chloride Stock CS, 0.4 mL of Cupric Sulfate Stock CS and 2.4 mL of Ferric Chloride Stock CS add diluted hydrochloric acid (1 in 40) to make exactly 200 mL.

(2) Heavy metals <1.07>—Proceed with 1.0 g of Dihydroergotoxine Mesilate according to Method 2, and perform the test. Prepare the control solution with 2.0 mL of Standard Lead Solution (not more than 20 ppm).

(3) Related substances—Weigh accurately 0.100 g of Dihydroergotoxine Mesilate, dissolve it in a mixture of chloroform and methanol (9:1) to make exactly 5 mL, and use this solution as the sample solution. Separately, weigh accurately 10 mg of dihydroergocristine mesilate for thin-layer chromatography, and dissolve in a mixture of chloroform and methanol (9:1) to make exactly 100 mL. Pipet 6 mL, 4 mL and 2 mL of this solution, add a mixture of chloroform and methanol (9:1) to make exactly 10 mL, respectively, and use these solutions as the standard solutions (1), (2) and (3), respectively. Perform the test with these solutions as directed under Thin-layer Chromatography <2.03> without putting the filter paper in the developing vessel. Spot 5  $\mu$ L each of the sample solution and standard solutions (1), (2) and (3) on a plate of silica gel for thin-layer chromatography. Develop the plate with a mixture of dichloromethane, ethyl acetate, methanol and ammonia solution (28) (50:50:3:1) to a distance of about 15 cm, and dry the plate with the aid of a cool air



stream. Immediately after that, develop the plate again with a newly prepared mixture of dichloromethane, ethyl acetate, methanol and ammonia solution (28) (50:50:3:1) to a distance of about 15 cm, and dry the plate within 1 minute with the aid of a cool air stream. Spray evenly *p*-dimethylaminobenzaldehyde-hydrochloric acid TS on the plate, dry the plate within 2 minutes with the aid of a cool air stream, and heat it at 40°C for 15 minutes: the spots other than the principal spot from the sample solution are not more intense than the spot from the standard solution (1), not more than 2 spots are more intense than that from the standard solution (2), and not more than 4 spots are more intense than that from the standard solution (3).

**Water** <2.48> Not more than 5.0% (0.2 g, direct titration).

**Residue on ignition** <2.44> Not more than 0.1% (1 g).

**Assay** (1) Dihydroergotoxine mesilate—Weigh accurately about 30 mg each of Dihydroergotoxine Mesilate and Dihydroergotoxine Mesilate Reference Standard, and dissolve them separately in a suitable amount of a mixture of water and acetonitrile (3:1). To these solutions add exactly 10 mL of the internal standard solution and an amount of a mixture of water and acetonitrile (3:1) to make 50 mL, and use these solutions as the sample solution and standard solution. Perform the test with 20  $\mu$ L of the sample solution and standard solution as directed under Liquid Chromatography according to the following conditions, and calculate the ratios of the peak areas of dihydroergocornine, dihydro- $\alpha$ -ergocryptine, dihydroergocristine and dihydro- $\beta$ -ergocryptine to the peak area of the internal standard of these solutions.

$$\begin{aligned} &\text{Amount (mg) of dihydroergotoxine mesilate} \\ &= W_S \times \{(M_{TA} + M_{TB} + M_{TC} + M_{TD}) / \\ &\quad (M_{SA} + M_{SB} + M_{SC} + M_{SD})\} \end{aligned}$$

$W_S$ : Amount (mg) of Dihydroergotoxine Mesilate Reference Standard, calculated on the anhydrous basis

$M_{TA}$ : Ratio of the peak area of dihydroergocornine to that of the internal standard of the sample solution  $\times 659.80$

$M_{TB}$ : Ratio of the peak area of dihydro- $\alpha$ -ergocryptine to that of the internal standard of the sample  $\times 673.83$

$M_{TC}$ : Ratio of the peak area of dihydroergocristine to that of the internal standard of the sample solution  $\times 707.85$

$M_{TD}$ : Ratio of the peak area of dihydro- $\beta$ -ergocryptine to that of the internal standard of the sample solution  $\times 673.83$

$M_{SA}$ : Ratio of the peak area of dihydroergocornine to that of the internal standard of the standard solution  $\times 659.80$

$M_{SB}$ : Ratio of the peak area of dihydro- $\alpha$ -ergocryptine to that of the internal standard of the standard solution  $\times 673.83$

$M_{SC}$ : Ratio of the peak area of dihydroergocristine to that of the internal standard of the standard solution  $\times 707.85$

$M_{SD}$ : Ratio of the peak area of dihydro- $\beta$ -ergocryptine to that of the internal standard of the standard solution  $\times 673.83$

**Internal standard solution**—Dissolve 0.04 g of chloramphenicol in a mixture of water and acetonitrile (3:1) to make 250 mL.

#### Operating conditions—

**Detector:** An ultraviolet absorption photometer (wavelength: 280 nm).

**Column:** A stainless steel column 4.6 mm in inside diameter and 15 cm in length, packed with octadecylsilanized silica gel for liquid chromatography (5  $\mu$ m in particle diameter).

**Column temperature:** A constant temperature of about 25°C.

**Mobile phase:** A mixture of water, acetonitrile and triethylamine (30:10:1).

**Flow rate:** Adjust the flow rate so that the retention time of chloramphenicol is about 5 minutes.

#### System suitability—

**System performance:** When the procedure is run with 20  $\mu$ L of the standard solution under the above operating conditions, the internal standard, dihydroergocornine, dihydro- $\alpha$ -ergocryptine, dihydroergocristine and dihydro- $\beta$ -ergocryptine are eluted in this order with the resolution between the peaks of dihydro- $\alpha$ -ergocryptine and dihydroergocristine being not less than 1.5.

**System repeatability:** When the test is repeated 6 times with 20  $\mu$ L of the standard solution under the above operating conditions, the relative standard deviation of the ratios of the peak area of dihydroergocornine, dihydro- $\alpha$ -ergocryptine, dihydroergocristine and dihydro- $\beta$ -ergocryptine to that of the internal standard is not more than 0.5%.

(2) Relative contents of dihydroergocornine mesilate, dihydroergocryptine mesilate and dihydroergocristine mesilate—Calculate the relative amounts of dihydroergocornine mesilate, dihydroergocryptine mesilate (dihydro- $\alpha$ -ergocryptine mesilate and dihydro- $\beta$ -ergocryptine mesilate) and dihydroergocristine mesilate from the chromatogram obtained in Assay (1) for the sample solution using the following equations:

$$\begin{aligned} &\text{Relative amount (\%)} \text{ of dihydroergocornine mesilate} \\ &= \{M_{TA} / (M_{TA} + M_{TB} + M_{TC} + M_{TD})\} \times 100 \end{aligned}$$

$$\begin{aligned} &\text{Relative amount (\%)} \text{ of dihydroergocryptine mesilate} \\ &= \{(M_{TB} + M_{TD}) / (M_{TA} + M_{TB} + M_{TC} + M_{TD})\} \times 100 \end{aligned}$$

$$\begin{aligned} &\text{Relative amount (\%)} \text{ of dihydroergocristine mesilate} \\ &= \{M_{TC} / (M_{TA} + M_{TB} + M_{TC} + M_{TD})\} \times 100 \end{aligned}$$

(3) Ratio of the content of dihydro- $\alpha$ -ergocryptine mesilate to dihydro- $\beta$ -ergocryptine mesilate—Calculate the ratio of the amount of dihydro- $\alpha$ -ergocryptine mesilate to dihydro- $\beta$ -ergocryptine mesilate from the chromatogram obtained in the Assay (1) for the sample solution using the following equations:


$$\begin{aligned} &\text{Ratio of the content of dihydro-}\alpha\text{-ergocryptine} \\ &\text{mesilate to dihydro-}\beta\text{-ergocryptine mesilate} \\ &= (M_{TB} / M_{TD}) \end{aligned}$$

**Containers and storage** Containers—Well-closed containers.

Storage—Light-resistant.

## Dilazep Hydrochloride Hydrate

ジラゼプ塩酸塩水和物



$C_{31}H_{44}N_2O_{10} \cdot 2HCl \cdot H_2O$ : 695.63

3,3'-(1,4-Diazepane-1,4-diyl)dipropyl bis(3,4,5-trimethoxybenzoate) dihydrochloride monohydrate [20153-98-4, anhydride]

Dilazep Hydrochloride Hydrate contains not less than 98.0% of dilazep hydrochloride ( $C_{31}H_{44}N_2O_{10} \cdot 2HCl$ : 677.62), calculated on the dried basis.

**Description** Dilazep Hydrochloride Hydrate occurs as a white, crystalline powder. It is odorless.

It is freely soluble in acetic acid (100) and in chloroform, soluble in water, slightly soluble in ethanol (95) and in acetic anhydride, and practically insoluble in diethyl ether.

Melting point: 200 – 204°C Immerse the sample in a bath of 110°C, and raise the temperature at the rate of about 3°C per minute from 140°C to 150°C, about 10°C per minute from 160°C to 195°C and about 1°C per minute from 195°C.

**Identification (1)** To 1 mL of a solution of Dilazep Hydrochloride Hydrate (1 in 100) add 0.1 mL of a solution of hydroxylammonium chloride (1 in 10) and 0.1 mL of 8 mol/L potassium hydroxide TS, and warm in a water bath of 70°C for 10 minutes. After cooling, add 0.5 mL of dilute hydrochloric acid and 0.1 mL of iron (III) chloride TS: a purple color develops.

**(2)** To 5 mL of a solution of Dilazep Hydrochloride Hydrate (3 in 500) add 0.3 mL of Reinecke salt TS: a light red precipitate is formed.

**(3)** Determine the absorption spectrum of a solution of Dilazep Hydrochloride Hydrate (1 in 50,000) as directed under Ultraviolet-visible Spectrophotometry <2.24>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wavelengths.

**(4)** Determine the infrared absorption spectrum of Dilazep Hydrochloride Hydrate as directed in the potassium chloride disk method under Infrared Spectrophotometry <2.25>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wave numbers.

**pH** <2.54> Dissolve 1.0 g of Dilazep Hydrochloride Hydrate in 100 mL of water: the pH of this solution is between 3.0 and 4.0.

**Purity (1)** Clarity and color of solution—Dissolve 1.0 g of Dilazep Hydrochloride Hydrate in 20 mL of water: the solution is clear and colorless.

**(2)** Sulfate <1.14>—Perform the test with 0.5 g of Dilazep Hydrochloride Hydrate. Prepare the control solution with

0.50 mL of 0.005 mol/L sulfuric acid VS (not more than 0.048%).

**(3)** Heavy metals <1.07>—Proceed with 2.0 g of Dilazep Hydrochloride Hydrate according to Method 1, and perform the test. Prepare the control solution with 2.0 mL of Standard Lead Solution (not more than 10 ppm).

**(4)** Arsenic <1.11>—Prepare the test solution with 1.0 g of Dilazep Hydrochloride Hydrate according to Method 3, and perform the test (not more than 2 ppm).

**(5)** Related substances—Dissolve 0.40 g of Dilazep Hydrochloride Hydrate in 10 mL of chloroform, and use this solution as the sample solution. Pipet 1 mL of the sample solution, add chloroform to make exactly 200 mL, and use this solution as the standard solution. Perform the test with these solutions as directed under Thin-layer Chromatography <2.03>. Spot 5  $\mu$ L each of the sample solution and standard solution on a plate of silica gel for thin-layer chromatography. Develop the plate with a mixture of methanol, ethyl acetate, dichloromethane and hydrochloric acid (500:200:100:1) to a distance of about 10 cm, and air-dry the plate. Spray evenly Dragendorff's TS for spraying on the plate: the spots other than the principal spot from the sample solution are not more intense than the spot from the standard solution.

**Loss on drying** <2.41> 2.0 – 3.0% (1 g, 105°C, 3 hours).

**Residue on ignition** <2.44> Not more than 0.1% (1 g).


**Assay** Weigh accurately about 0.3 g of Dilazep Hydrochloride Hydrate, dissolve in 40 mL of a mixture of acetic anhydride and acetic acid (100) (7:3), and titrate <2.50> with 0.1 mol/L perchloric acid VS (potentiometric titration). Perform a blank determination, and make any necessary correction.

Each mL of 0.1 mol/L perchloric acid VS  
= 33.88 mg of  $C_{31}H_{44}N_2O_{10} \cdot 2HCl$

**Containers and storage** Containers—Tight containers.

## Diltiazem Hydrochloride

ジルチアゼム塩酸塩



$C_{22}H_{26}N_2O_4 \cdot S \cdot HCl$ : 450.98

(2S,3S)-5-[2-(Dimethylamino)ethyl]-2-(4-methoxyphenyl)-4-oxo-2,3,4,5-tetrahydro-1,5-benzothiazepin-3-yl acetate monohydrochloride [33286-22-5]

Diltiazem Hydrochloride, when dried, contains not less than 98.5% of  $C_{22}H_{26}N_2O_4 \cdot S \cdot HCl$ .

**Description** Diltiazem Hydrochloride occurs as white crystals or crystalline powder. It is odorless.

It is very soluble in formic acid, freely soluble in water, in methanol and in chloroform, sparingly soluble in acetonitrile, slightly soluble in acetic anhydride and in ethanol (99.5), and practically insoluble in diethyl ether.

**Identification (1)** Dissolve 0.05 g of Diltiazem Hydrochloride in 1 mL of 1 mol/L hydrochloric acid TS, add 2 mL of ammonium thiocyanate-cobaltous nitrate TS and 5 mL of chloroform, shake well, and allow to stand: a blue color develops in the chloroform layer.

**(2)** Proceed as directed under Oxygen Flask Combustion Method <1.06> with 0.03 g of Diltiazem Hydrochloride, using 20 mL of water as the absorbing liquid, and prepare the test solution: the test solution responds to the Qualitative Tests <1.09> (1) for sulfate.

**(3)** Dissolve 0.01 g of Diltiazem Hydrochloride in 0.01 mol/L hydrochloric acid TS to make 100 mL. To 2 mL of the solution add 0.01 mol/L hydrochloric acid TS to make 20 mL. Determine the absorption spectrum of the solution as directed under Ultraviolet-visible Spectrophotometry <2.24>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wavelengths.

**(4)** Determine the infrared absorption spectrum of Diltiazem Hydrochloride, previously dried, as directed in the potassium bromide disk method under Infrared Spectrophotometry <2.25>: it exhibits absorption at the wave numbers of about  $1741\text{ cm}^{-1}$ ,  $1678\text{ cm}^{-1}$ ,  $1252\text{ cm}^{-1}$  and  $1025\text{ cm}^{-1}$ .

**(5)** A solution of Diltiazem Hydrochloride (1 in 50) responds to the Qualitative Tests <1.09> (2) for chloride.

**Optical rotation** <2.49>  $[\alpha]_D^{20}$ :  $+115 - +120^\circ$  (after drying, 0.20 g, water, 20 mL, 100 mm).

**Melting point** <2.60>  $210 - 215^\circ\text{C}$  (with decomposition).

**pH** <2.54> Dissolve 1.0 g of Diltiazem Hydrochloride in 100 mL of water: the pH of this solution is between 4.3 and 5.3.

**Purity (1)** Clarity and color of solution—Dissolve 1.0 g of Diltiazem Hydrochloride in 20 mL of water: the solution is clear and colorless.

**(2)** Sulfate <1.14>—Perform the test with 1.0 g of Diltiazem Hydrochloride. Prepare the control solution with 0.50 mL of 0.005 mol/L sulfuric acid VS (not more than 0.024%).

**(3)** Heavy metals <1.07>—Proceed with 2.0 g of Diltiazem Hydrochloride according to Method 2, and perform the test. Prepare the control solution with 2.0 mL of Standard Lead Solution (not more than 10 ppm).

**(4)** Arsenic <1.11>—Place 1.0 g of Diltiazem Hydrochloride in a decomposition flask, add 5 mL of nitric acid and 2 mL of sulfuric acid, put a small funnel on the neck of the flask, and heat cautiously until white fumes are evolved. After cooling, add 2 mL of nitric acid, heat, and repeat this procedure twice, add several 2-mL portions of hydrogen peroxide (30), and heat until the solution becomes colorless to pale yellow. After cooling, add 2 mL of saturated solution of ammonium oxalate monohydrate, and heat again until white fumes are evolved. After cooling, add water to make 5 mL, use this solution as the test solution, and perform the test: the test solution has no more color than the following control solution (not more than 2 ppm).

Control solution: Proceed in the same manner as the test solution without Diltiazem Hydrochloride, add 2.0 mL of

Standard Arsenic Solution and water to make 5 mL, and proceed in the same manner as the test solution.

**(5)** Related substances—Dissolve 50 mg of Diltiazem Hydrochloride in 50 mL of diluted ethanol (99.5) (4 in 5), and use this solution as the sample solution. Measure exactly 1 mL of the sample solution, add diluted ethanol (99.5) (4 in 5) to make exactly 200 mL, and use this solution as the standard solution. Perform the test with exactly 20  $\mu\text{L}$  each of the sample solution and standard solution as directed under Liquid Chromatography <2.01> according to the following conditions. Determine each peak area of both solutions by automatic integration method: the total peak area of peaks other than the peak of diltiazem obtained from the sample solution is not more than 3/5 times the peak area of diltiazem obtained from the standard solution.

**Operating conditions—**

Detector: An ultraviolet absorption photometer (wavelength: 240 nm).

Column: A stainless steel column 4.6 mm in inside diameter and 15 cm in length, packed with octadecylsilanized silica gel for liquid chromatography (5  $\mu\text{m}$  in particle diameter).

Column temperature: A constant temperature of about  $50^\circ\text{C}$ .

Mobile phase: Dissolve 8 g of sodium acetate trihydrate and 1.5 g of *d*-camphorsulfonic acid in 500 mL of water, and filter using a membrane filter (0.4  $\mu\text{m}$  in pore size). Add 250 mL each of acetonitrile and methanol to the filtrate, and adjust the solution to a pH of 6.6 by adding sodium acetate trihydrate.

Flow rate: Adjust the flow rate so that the retention time of diltiazem is about 9 minutes.

Time span of measurement: About twice as long as the retention time of diltiazem beginning after the solvent peak.

**System suitability—**

Test for required detection: To exactly 2 mL of the standard solution add diluted ethanol (99.5) (4 in 5) to make exactly 10 mL. Confirm that the peak area of diltiazem obtained from 20  $\mu\text{L}$  of this solution is equivalent to 15 to 25% of that of diltiazem obtained from 20  $\mu\text{L}$  of the standard solution.

System performance: Dissolve 0.03 g of Diltiazem Hydrochloride, 0.02 g of *d*-3-hydroxy-*cis*-2,3-dihydro-5-[2-(dimethylamino)ethyl]-2-(4-methoxyphenyl)-1,5-benzothiazepin-4(5*H*)-one hydrochloride and 0.02 g of phenylbenzoate in 160 mL of ethanol (99.5), and add water to make 200 mL. Perform the test with 20  $\mu\text{L}$  of this solution as directed under Liquid Chromatography under the above operating conditions: *d*-3-hydroxy-*cis*-2,3-dihydro-5-[2-(dimethylamino)ethyl]-2-(4-methoxyphenyl)-1,5-benzothiazepin-4(5*H*)-one, diltiazem and phenyl benzoate are eluted in this order with the resolutions between the peaks of *d*-3-hydroxy-*cis*-2,3-dihydro-5-[2-(dimethylamino)ethyl]-2-(4-methoxyphenyl)-1,5-benzothiazepin-4(5*H*)-one and diltiazem and between the peaks of diltiazem and phenyl benzoate being not less than 2.5, respectively.

System repeatability: When the test is repeated 6 times with 20  $\mu\text{L}$  of the standard solution under the above operating conditions, the relative standard deviation of the peak area of diltiazem is not more than 2.0%.

**Loss on drying** <2.41> Not more than 0.5% (1 g,  $105^\circ\text{C}$ , 2 hours).

**Residue on ignition** <2.44> Not more than 0.1% (1 g).


**Assay** Weigh accurately about 0.7 g of Diltiazem Hydrochloride, previously dried, dissolve in 2.0 mL of formic acid, add 60 mL of acetic anhydride, and titrate <2.50> with 0.1 mol/L perchloric acid VS (potentiometric titration). Perform a blank determination, and make any necessary correction.

Each mL of 0.1 mol/L perchloric acid VS  
= 45.10 mg of  $C_{22}H_{26}N_2O_4S \cdot HCl$

**Containers and storage** Containers—Tight containers.  
Storage—Light-resistant.

## Dimemorfan Phosphate

ジメモルファンリン酸塩



$C_{18}H_{25}N \cdot H_3PO_4$ : 353.39  
(9*S*,13*S*,14*S*)-3,17-Dimethylmorphinan monophosphate  
[36304-84-4]

Dimemorfan Phosphate, when dried, contains not less than 98.5% of  $C_{18}H_{25}N \cdot H_3PO_4$ .

**Description** Dimemorfan Phosphate occurs as white to pale yellowish white crystals or crystalline powder.

It is freely soluble in acetic acid (100), sparingly soluble in water and in methanol, slightly soluble in ethanol (95), and practically insoluble in diethyl ether.

Melting point: about 265°C (with decomposition).

**Identification (1)** Determine the absorption spectrum of a solution of Dimemorfan Phosphate (1 in 5000) as directed under Ultraviolet-visible Spectrophotometry <2.24>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wavelengths.

(2) Determine the infrared absorption spectrum of Dimemorfan Phosphate, previously dried, as directed in the potassium bromide disk method under Infrared Spectrophotometry <2.25>, and compare the spectrum with the Reference Spectrum: both spectra exhibits similar intensities of absorption at the same wave numbers.

(3) To 2 mL of a solution of Dimemorfan Phosphate (1 in 100) add 2 to 3 drops of silver nitrate TS: a yellow precipitate is formed, and it dissolves on the addition of dilute nitric acid.

**Optical rotation** <2.49>  $[\alpha]_D^{20}$ : +25 – +27° (after drying, 1 g, methanol, 100 mL, 100 mm).

**pH** <2.54> Dissolve 1.0 g of Dimemorfan Phosphate in 100 mL of water: the pH of this solution is between 4.0 and 5.0.

**Purity (1)** Heavy metals <1.07>—Proceed with 1.0 g of Dimemorfan Phosphate according to Method 1, and perform the test. Prepare the control solution with 2.0 mL of Standard Lead Solution (not more than 20 ppm).

(2) Arsenic <1.11>—Prepare the test solution with 1.0 g of Dimemorfan Phosphate according to Method 3, and perform the test. Use 10 mL of a solution of magnesium nitrate hexahydrate in ethanol (95) (1 in 10) (not more than 2 ppm).

(3) Related substances—Dissolve 0.10 g of Dimemorfan Phosphate in 10 mL of methanol, and use this solution as the sample solution. Pipet 1 mL of the sample solution, add methanol to make exactly 100 mL, and use this solution as the standard solution. Perform the test with these solutions as directed under Thin-layer Chromatography <2.03>. Spot 10  $\mu$ L each of the sample solution and standard solution on a plate of silica gel for thin-layer chromatography. Develop the plate with a mixture of methanol, chloroform and ammonia solution (28) (150:150:1) to a distance of about 10 cm, and air-dry the plate. Spray evenly the plate with Dragendorff's TS for spraying: the spots other than the principal spot from the sample solution are not more intense than the spot from the standard solution.

**Loss on drying** <2.41> Not more than 0.5% (1 g, 105°C, 3 hours).


**Assay** Weigh accurately about 0.6 g of Dimemorfan Phosphate, previously dried, dissolve in 100 mL of acetic acid (100), and titrate <2.50> with 0.1 mol/L perchloric acid VS (potentiometric titration). Perform a blank determination, and make any necessary correction.

Each mL of 0.1 mol/L perchloric acid VS  
= 35.34 mg of  $C_{18}H_{25}N \cdot H_3PO_4$

**Containers and storage** Containers—Tight containers.

## Dimenhydrinate

ジメンヒドリナート



$C_{17}H_{21}NO \cdot C_7H_7ClN_4O_2$ : 469.96  
2-(Diphenylmethoxy)-*N,N*-dimethylethylamine—  
8-chloro-1,3-dimethyl-1*H*-purine-2,6(3*H*,7*H*)-dione  
(1/1) [523-87-5]

Dimenhydrinate, when dried, contains not less than 53.0% and not more than 55.5% of diphenhydramine ( $C_{17}H_{21}NO$ : 255.36), and not less than 44.0% and not more than 47.0% of 8-chlorotheophylline ( $C_7H_7ClN_4O_2$ : 214.61).

**Description** Dimenhydrinate occurs as a white, crystalline powder. It is odorless, and has a bitter taste.

It is very soluble in chloroform, freely soluble in ethanol (95), and slightly soluble in water and in diethyl ether.

**Identification (1)** Dissolve 0.5 g of Dimenhydrinate in 30 mL of dilute ethanol, add 30 mL of water, and use this solution as the sample solution. Transfer 30 mL of the sample solution to a separator, and add 2 mL of ammonia solution (28). Extract with two 10-mL portions of diethyl ether, com-

bine the diethyl ether extracts, wash the combined extracts with 5 mL of water, and then extract the combined extracts with 15 mL of diluted hydrochloric acid (1 in 100). With this acid extract perform the following tests.

(i) To 5 mL of this acid extract add 5 drops of Reinecke salt TS: a light red precipitate is produced.

(ii) To 10 mL of this acid extract add 10 mL of 2,4,6-trinitrophenol TS dropwise, and allow to stand for 30 minutes. Collect the precipitate by filtrating, recrystallize from dilute ethanol, and dry at 105°C for 30 minutes: the crystals melt <2.60> between 128°C and 133°C.

(2) To 30 mL of the sample solution obtained in the Identification (1) add 2 mL of dilute sulfuric acid, and cool for 30 minutes. Scratch the inside wall of the container frequently to facilitate crystallization. Filter, and wash the white crystals with a small amount of ice-cooled water. Dry the crystals for 1 hour at 105°C: the crystals melt <2.60> between 300°C and 305°C with decomposition.

(3) To 0.01 g of the crystals obtained in the Identification (2) add 10 drops of hydrogen peroxide TS and 1 drop of hydrochloric acid, and evaporate on a water bath to dryness: the residue shows a yellow-red color. When the dish containing the residue is held over a vessel containing 2 to 3 drops of ammonia TS, the color changes to red-purple, which is discharged on the addition of 2 to 3 drops of sodium hydroxide TS.

(4) Mix well 0.05 g of the crystals obtained in the Identification (2) with 0.5 g of sodium peroxide in a nickel crucible, and heat until the mass melts. Cool, dissolve the melted mass in 20 mL of water, and acidify with dilute nitric acid: the solution responds to the Qualitative Tests for chloride <1.09>.

**Melting point** <2.60> 102 – 107°C

**Purity** (1) Chloride <1.03>—Transfer 50 mL of the filtrate obtained in the Assay (2) to a Nessler tube, add 1 mL of nitric acid, and allow to stand for 5 minutes: the turbidity of the solution is not greater than that of the following control solution.

Control solution: Dilute 0.25 mL of 0.01 mol/L hydrochloric acid VS with 6 mL of dilute nitric acid and with water to make 50 mL, add 1 mL of silver nitrate TS, and allow to stand for 5 minutes (not more than 0.044%).

(2) Bromide and iodide—Place 0.10 g of Dimenhydrinate in a glass-stoppered test tube, and add 0.05 g of sodium nitrite, 10 mL of chloroform and 10 mL of dilute hydrochloric acid. Stopper, shake well, and allow to stand: the chloroform layer remains colorless.

**Loss on drying** <2.41> Not more than 0.5% (3 g, in vacuum, phosphorus (V) oxide, 24 hours).

**Residue on ignition** <2.44> Not more than 0.3% (1 g).

**Assay** (1) Diphenhydramine—Weigh accurately about 0.5 g of Dimenhydrinate, previously dried, transfer to a 250-mL separator, and add 50 mL of water, 3 mL of ammonia TS and 10 g of sodium chloride. Extract with six 15-mL portions of diethyl ether with shaking, combine the diethyl ether extracts, and wash the combined diethyl ether extracts with three 50-mL portions of water. To the diethyl ether extracts add exactly 25 mL of 0.05 mol/L sulfuric acid VS, and add 25 mL of water. Shake thoroughly, and evaporate the diethyl ether gently. Cool, and titrate the excess sulfuric acid with 0.1 mol/L sodium hydroxide VS (indicator: 3 drops of methyl

red TS). Perform a blank determination, and make any necessary correction.

Each mL of 0.05 mol/L sulfuric acid VS  
= 25.54 mg  $C_{17}H_{21}NO$

(2) 8-Chlorotheophylline—Weigh accurately about 0.8 g of Dimenhydrinate, previously dried, transfer to a 200-mL volumetric flask, add 50 mL of water, 3 mL of ammonia TS and 6 mL of a solution of ammonium nitrate (1 in 10), and heat on a water bath for 5 minutes. Add exactly 25 mL of 0.1 mol/L silver nitrate VS, heat on a water bath for 15 minutes with occasional shaking, cool, and add water to make exactly 200 mL. Allow to stand overnight to settle the precipitate, and filter through a dry filter paper, discarding the first 20 mL of the filtrate. Measure exactly 100 mL of the subsequent filtrate, acidify with nitric acid, add 3 mL of nitric acid, and titrate the excess silver nitrate with 0.1 mol/L ammonium thiocyanate VS (indicator: 2 mL of ammonium iron (III) sulfate TS). Perform a blank determination, and make any necessary correction.

Each mL of 0.1 mol/L silver nitrate VS  
= 21.46 mg of  $C_7H_7ClN_4O_2$

**Containers and storage** Containers—Well-closed containers.

## Dimenhydrinate Tablets

ジメンヒドリナート錠

Dimenhydrinate Tablets contain not less than 95% and not more than 105% of the labeled amount of dimenhydrinate ( $C_{17}H_{21}NO \cdot C_7H_7ClN_4O_2$ ; 469.96).

**Method of preparation** Prepare as directed under Tablets, with Dimenhydrinate.

**Identification** (1) Triturate a quantity of powdered Dimenhydrinate Tablets, equivalent to 0.5 g of Dimenhydrinate according to the labeled amount, with 25 mL of warm ethanol (95), and filter. Dilute the filtrate with 40 mL of water, and filter again. Use the filtrate as the sample solution. Transfer 30 mL of the sample solution to a separator, and proceed as directed in the Identification (1) under Dimenhydrinate.

(2) With 30 mL of the sample solution obtained in (1), proceed as directed in the Identification (2), (3) and (4) under Dimenhydrinate.

**Assay** Weigh accurately, and powder not less than 20 Dimenhydrinate Tablets. Weigh accurately a portion of the powder, equivalent to about 0.5 g of dimenhydrinate ( $C_{17}H_{21}NO \cdot C_7H_7ClN_4O_2$ ), transfer to a flask, add 40 mL of ethanol (95), and heat with swirling on a water bath until the solution just boils. Continue to heat for 30 seconds, and filter through a glass filter (G4). Wash the filter with warm ethanol (95), transfer the filtrate and washings to a flask, and evaporate the ethanol on a water bath to make 5 mL. Add 50 mL of water, 3 mL of ammonia TS and 6 mL of a solution of ammonium nitrate (1 in 10), heat the mixture on a water bath for 5 minutes, add exactly 25 mL of 0.1 mol/L silver nitrate VS, and heat on a water bath for 15 minutes with occasional shaking. Transfer the mixture to a 200-mL volumetric flask,


using water to rinse the flask, cool, add water to make exactly 200 mL, and proceed as directed in the Assay (2) under Dimenhydrinate.

Each mL of 0.1 mol/L silver nitrate VS  
= 47.00 mg of  $C_{17}H_{21}NO \cdot C_7H_7ClN_4O_2$

**Containers and storage** Containers—Well-closed containers.

## Dimercaprol

ジメルカプロール



$C_3H_8OS_2$ : 124.23  
(*2RS*)-2,3-Bisulfanylpropan-1-ol [59-52-9]

Dimercaprol contains not less than 98.5% and not more than 101.5% of  $C_3H_8OS_2$ .

**Description** Dimercaprol is a colorless or pale yellow liquid. It has a mercaptan-like, disagreeable odor.

It is miscible with methanol and with ethanol (99.5)

It is soluble in peanut oil, and sparingly soluble in water.

It shows no optical rotation.

**Identification (1)** Add 1 drop of Dimercaprol to a mixture of 1 drop of a solution of cobalt (II) chloride hexahydrate (1 in 200) and 5 mL of water: a yellow-brown color develops.

(2) Determine the infrared absorption spectrum of Dimercaprol as directed in the liquid film method under Infrared Spectrophotometry <2.25>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wave numbers.

**Refractive index** <2.45>  $n_D^{20}$ : 1.570 – 1.575

**Specific gravity** <2.56>  $d_{20}^{20}$ : 1.238 – 1.248

**Purity (1)** Clarity and color of solution—Dissolve 1.0 mL of Dimercaprol in 20 mL of peanut oil: the solution is clear and colorless to pale yellow.

(2) Bromide—To 2.0 g of Dimercaprol add 25 mL of dilute potassium hydroxide-ethanol TS, and heat in a water bath under a reflux condenser for 2 hours. Evaporate the ethanol in a current of warm air, add 20 mL of water, and cool. Add a mixture of 10 mL of strong hydrogen peroxide and 40 mL of water, boil gently under a reflux condenser for 10 minutes, and filter rapidly after cooling. Wash the residue with two 10-mL portions of water, combine the washings with the filtrate, add 10 mL of dilute nitric acid and exactly 5 mL of 0.1 mol/L silver nitrate VS, and titrate <2.50> the excess silver nitrate with 0.1 mol/L ammonium thiocyanate VS (indicator: 2 mL of ammonium iron (III) sulfate TS). Perform a blank determination: not more than 1.0 mL of 0.1 mol/L silver nitrate VS is consumed.

(3) Heavy metals <1.07>—Proceed with 1.0 g of Dimercaprol according to Method 2, and perform the test. Prepare the control solution with 2.0 mL of Standard Lead Solution (not more than 20 ppm).

**Assay** Weigh accurately about 0.15 g of Dimercaprol into a glass-stoppered flask, dissolve in 10 mL of methanol, and ti-

trate <2.50> immediately with 0.05 mol/L iodine VS until a pale yellow color is produced. Perform a blank determination, and make any necessary correction.

Each mL of 0.05 mol/L iodine VS = 6.211 mg of  $C_3H_8OS_2$

**Containers and storage** Containers—Tight containers.  
Storage—Not exceeding 5°C.

## Dimercaprol Injection

ジメルカプロール注射液

Dimercaprol Injection is an oily solution for injection.

It contains not less than 95.0% and not more than 105.0% of the labeled amount of dimercaprol ( $C_3H_8OS_2$ : 124.23).

**Method of preparation** Prepare as directed under Injections, with Dimercaprol. Benzyl Benzoate or Benzyl Alcohol may be added to increase the solubility.

**Description** Dimercaprol Injection is a clear, colorless or light yellow liquid. It has an unpleasant odor.

**Identification** Measure a volume of Dimercaprol Injection, equivalent to 30 mg of Dimercaprol according to the labeled amount, and proceed as directed in the Identification (1).

**Extractable volume** <6.05> It meets the requirement.

**Foreign insoluble matter** <6.06> Perform the test according to Method 1: it meets the requirement.

**Insoluble particulate matter** <6.07> Perform the test according to Method 2: it meets the requirement.

**Sterility** <4.06> Perform the test according to the Membrane filtration method: it meets the requirement.


**Assay** Pipet a volume of Dimercaprol Injection, equivalent to about 0.1 g of dimercaprol ( $C_3H_8OS_2$ ), into a flask, and rinse the pipet several times with a mixture of methanol and diethyl ether (3:1), adding the rinsings to the flask. Add the mixture of methanol and diethyl ether (3:1) to make 50 mL, and titrate <2.50> with 0.05 mol/L iodine VS until a yellow color persists. Perform a blank determination, and make any necessary correction.

Each mL of 0.05 mol/L iodine VS  
= 6.211 mg of  $C_3H_8OS_2$

**Containers and storage** Containers—Hermetic containers.  
Storage—In a cold place.

## Dimorpholamine

ジモルホラミン



$C_{20}H_{38}N_4O_4$ : 398.54

*N,N'*-Ethylenebis(*N*-butylmorpholine-4-carboxamide)

[119-48-2]

Dimorpholamine, when dried, contains not less than 98.0% and not more than 101.0% of  $C_{20}H_{38}N_4O_4$ .

**Description** Dimorpholamine is a white to light yellow, crystalline powder, masses or syrupy liquid.

It is very soluble in ethanol (99.5) and in acetic anhydride, and soluble in water.

The pH of a solution prepared by dissolving 1.0 g of Dimorpholamine in 10 mL of water is between 6.0 and 7.0.

It is hygroscopic.

**Identification (1)** Determine the absorption spectrum of a solution of Dimorpholamine (1 in 50,000) as directed under Ultraviolet-visible Spectrophotometry <2.24>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wavelengths.

**(2)** Determine the infrared absorption spectrum of Dimorpholamine, previously dried, as directed in the potassium bromide disk method under the Infrared Spectrophotometry <2.25>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wave numbers.

**Purity (1)** Clarity and color of solution—Dissolve 1.0 g of Dimorpholamine in 50 mL of water: the solution is clear and colorless to pale yellow.

**(2)** Chloride <1.03>—To 20 mL of the solution obtained in (1) add 6 mL of dilute nitric acid and water to make 50 mL. Perform the test using this solution as the test solution. Prepare the control solution with 0.40 mL of 0.01 mol/L hydrochloric acid VS (not more than 0.036%).

**(3)** Sulfate <1.14>—To 10 mL of the solution obtained in (1) add 1 mL of dilute hydrochloric acid and water to make 50 mL. Perform the test using this solution as the test solution. Prepare the control solution with 0.40 mL of 0.005 mol/L sulfuric acid VS (not more than 0.096%).

**(4)** Heavy metals <1.07>—Proceed with 2.0 g of Dimorpholamine according to Method 2, and perform the test. Prepare the control solution with 2.0 mL of Standard Lead Solution (not more than 10 ppm).

**(5)** Related substances—Dissolve 0.20 g of Dimorpholamine in 10 mL of ethanol (99.5), and use this solution as the sample solution. Pipet 1 mL of the sample solution, add ethanol (99.5) to make exactly 100 mL, and use this solution as the standard solution. Perform the test with these solutions as directed under Thin-layer Chromatography <2.03>.

Spot 10  $\mu$ L each of the sample solution and standard solution on a plate of silica gel for thin-layer chromatography. Develop the plate with a mixture of ethanol (99.5) and water (4:1) to a distance of about 10 cm, and air-dry the plate. Allow the plate to stand in iodine vapor for 10 minutes: the spot other than the principal spot from the sample solution is not more intense than the spot from the standard solution.

**Loss on drying** <2.41> Not more than 0.5% (1 g, in vacuum, phosphorus (V) oxide, 8 hours).

**Residue on ignition** <2.44> Not more than 0.1% (1 g).

**Assay** Weigh accurately about 0.6 g of Dimorpholamine, previously dried, dissolve in 50 mL of acetic anhydride, and titrate <2.50> with 0.1 mol/L perchloric acid VS (potentiometric titration). Perform a blank determination, and make any necessary correction.

Each mL of 0.1 mol/L perchloric acid VS  
= 39.85 mg of  $C_{20}H_{38}N_4O_4$

**Containers and storage** Containers—Tight containers.  
Storage—Light-resistant.

## Dimorpholamine Injection

ジモルホラミン注射液

Dimorpholamine Injection is an aqueous solution for injection.

It contains not less than 95.0% and not more than 105.0% of the labeled amount of dimorpholamine ( $C_{20}H_{38}N_4O_4$ : 398.54).

**Method of preparation** Prepare as directed under Injections, with Dimorpholamine.

**Description** Dimorpholamine Injection is a clear, colorless liquid.

pH: 3.0 – 5.5

**Identification (1)** To a volume of Dimorpholamine Injection, equivalent to 0.1 g of Dimorpholamine according to the labeled amount, add 3 drops of Dragendorff's TS: an orange color develops.

**(2)** To a volume of Dimorpholamine Injection, equivalent to 50 mg of Dimorpholamine according to the labeled amount, add 1 mL of dilute hydrochloric acid, and evaporate on a water bath to dryness. Dissolve the residue in 2 mL of hydrochloric acid, boil for 10 minutes under a reflux condenser, and evaporate to dryness on a water bath. Dissolve the residue with 1 mL of water, neutralize with sodium hydroxide TS, and add 0.2 mL of a solution of acetaldehyde (1 in 20), 0.1 mL of sodium pentacyanonitrosyl ferrate (III) TS and 0.5 mL of sodium carbonate TS: a blue color develops.

**Bacterial endotoxins** <4.01> Less than 5.0 EU/mg. Perform the test with the sample diluted to 0.15 w/v% with water for bacterial endotoxins test.

**Extractable volume** <6.05> It meets the requirement.

**Foreign insoluble matter** <6.06> Perform the test according to Method 1: it meets the requirement.

**Insoluble particulate matter** <6.07> Perform the test according to Method 1: it meets the requirement.

**Sterility** <4.06> Perform the test according to the Membrane filtration method: it meets the requirement.

**Assay** Measure exactly a volume of Dimorpholamine Injection, equivalent to about 30 mg of dimorpholamine ( $C_{20}H_{38}N_4O_4$ ), and add water to make exactly 200 mL. Pipet 1 mL of this solution, shake with exactly 4 mL of the internal standard solution for 5 minutes, and use this solution as the sample solution. Separately, weigh accurately about 0.15 g of dimorpholamine for assay, previously dried in a desiccator (in vacuum, phosphorus (V) oxide) for 8 hours, and dissolve in water to make exactly 1000 mL. Pipet 1 mL of this solution, shake with exactly 4 mL of the internal standard solution for 5 minutes, and use this solution as the standard solution. Perform the test with 10  $\mu$ L each of the sample solution and standard solution as directed under Liquid Chromatography <2.01> according to the following conditions, and calculate the ratios,  $Q_T$  and  $Q_S$ , of the peak area of dimorpholamine to that of the internal standard.

$$\begin{aligned} \text{Amount (mg) of dimorpholamine (C}_{20}\text{H}_{38}\text{N}_4\text{O}_4) \\ = W_S \times (Q_T/Q_S) \times (1/5) \end{aligned}$$

$W_S$ : Amount (mg) of dimorpholamine for assay

**Internal standard solution**—A solution of butyl parahydroxybenzoate in acetonitrile (1 in 25,000).

**Operating conditions**—

**Detector:** An ultraviolet absorption photometer (wavelength: 216 nm).

**Column:** A stainless steel column 4.6 mm in inside diameter and 15 cm in length, packed with octadecylsilanized silica gel for liquid chromatography (5  $\mu$ m in particle diameter).

**Column temperature:** A constant temperature of about 40°C.

**Mobile phase:** A mixture of water and acetonitrile (1:1).

**Flow rate:** Adjust the flow rate so that the retention time of dimorpholamine is about 4 minutes.

**System suitability**—

**System performance:** When the procedure is run with 10  $\mu$ L of the standard solution under the above operating conditions, dimorpholamine and the internal standard are eluted in this order with the resolution between these peaks being not less than 2.0.


**System repeatability:** When the test is repeated 6 times with 10  $\mu$ L of the standard solution under the above operating conditions, the relative standard deviation of the ratio of the peak area of dimorpholamine to that of the internal standard is not more than 1.0%.

**Containers and storage** Containers—Hermetic containers.

## Dinoprost

### Prostaglandin $F_{2\alpha}$

ジノプロスト



$C_{20}H_{34}O_5$ : 354.48

(5Z)-7-[(1R,2R,3R,5S)-3,5-Dihydroxy-2-[(1E,3S)-3-hydroxyoct-1-en-1-yl]cyclopentyl]hept-5-enoic acid  
[551-11-1]

Dinoprost contains not less than 98.5% of  $C_{20}H_{34}O_5$ , calculated on the anhydrous basis.

**Description** Dinoprost occurs as white, waxy masses or powder, or a clear, colorless to light yellow and viscous liquid. It is odorless.

It is very soluble in *N,N*-dimethylformamide, freely soluble in methanol, in ethanol (99.5) and in diethyl ether, and very slightly soluble in water.

**Identification** (1) To 5 mg of Dinoprost add 2 mL of sulfuric acid, and dissolve by shaking for 5 minutes: a dark red color develops. To this solution add 30 mL of sulfuric acid: an orange color develops with a green fluorescence.

(2) Dissolve 1 mg of Dinoprost in 50 mL of diluted sulfuric acid (7 in 10), and warm in a water bath heated at 50°C for 40 minutes. After cooling, determine the absorption spectrum of the solution as directed under Ultraviolet-visible Spectrophotometry <2.24>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wavelengths.

(3) Warm Dinoprost at 40°C to effect a liquid, and determine the infrared absorption spectrum of the liquid as directed in the liquid film method under Infrared Spectrophotometry <2.25>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wave numbers.

**Optical rotation** <2.49>  $[\alpha]_D^{20}$ : +24 – +31° (0.2 g, ethanol (99.5), 10 mL, 100 mm).

**Purity** (1) Clarity and color of solution—Dissolve 0.20 g of Dinoprost in 5 mL of ethanol (99.5): the solution is clear and colorless to pale yellow.

(2) Related substances—Dissolve 10 mg of Dinoprost in 2 mL of methanol, add water to make 10 mL, and use this solution as the sample solution. Pipet 3 mL of the sample solution, add diluted methanol (1 in 5) to make exactly 100 mL, and use this solution as the standard solution. Perform the test with exactly 10  $\mu$ L each of the sample solution and standard solution as directed under Liquid Chromatography <2.01> according to the following conditions. Determine each peak area of these solutions by the automatic integration method: the total area of the peaks other than the peak of dinoprost from the sample solution is not larger than the peak area of dinoprost from the standard solution.



**Operating conditions—**

**Detector:** An ultraviolet absorption photometer (wavelength: 205 nm).

**Column:** A stainless steel column about 5 mm in inside diameter and about 15 cm in length, packed with octadecylsilanized silica gel for liquid chromatography (5  $\mu$ m in particle diameter).

**Column temperature:** A constant temperature of about 25°C.

**Mobile phase:** A mixture of 0.02 mol/L potassium dihydrogenphosphate TS and acetonitrile (5:2).

**Flow rate:** Adjust the flow rate so that the retention time of dinoprost is about 20 minutes.

**Selection of column:** Dissolve 0.01 g each of isopropyl parahydroxybenzoate and propyl parahydroxybenzoate in 2 mL of methanol, and add water to make 10 mL. To 1 mL of this solution add diluted methanol (1 in 5) to make 30 mL, proceed with 10  $\mu$ L of this solution under the above operating conditions, and calculate the resolution. Use a column giving elution of isopropyl parahydroxybenzoate and propyl parahydroxybenzoate in this order with the resolution between these peaks being not less than 2.5.

**Detection sensitivity:** Adjust the detection sensitivity so that the peak height of dinoprost from the standard solution composes 5% to 15% of the full scale.

**Time span of measurement:** About 1.5 times as long as the retention time of dinoprost beginning after the solvent peak.

**Water** <2.48> Not more than 0.5% (0.3 g, direct titration).

**Assay** Weigh accurately about 50 mg of Dinoprost, dissolve in 30 mL of *N,N*-dimethylformamide, and titrate <2.50> with 0.02 mol/L tetramethylammonium hydroxide VS under a stream of nitrogen (potentiometric titration). Perform a blank determination, and make any necessary correction.

Each mL of 0.02 mol/L tetramethylammonium hydroxide VS


$$= 7.090 \text{ mg of } C_{20}H_{34}O_5$$

**Containers and storage** Containers—Tight containers.

**Storage**—Light-resistant, and in a place not exceeding 5°C.

## Diphenhydramine

ジフェンヒドรามイン



$C_{17}H_{21}NO$ : 255.35

2-(Diphenylmethoxy)-*N,N*-dimethylethylamine [58-73-1]

Diphenhydramine contains not less than 96.0% of  $C_{17}H_{21}NO$ .

**Description** Diphenhydramine is a clear, light yellow to yellow liquid. It has a characteristic odor, and has a burning taste at first, followed by a slight sensation of numbness on the tongue.

It is miscible with acetic anhydride, with acetic acid (100), with ethanol (95) and with diethyl ether.

It is very slightly soluble in water.

Boiling point: about 162°C (in vacuum, 0.67 kPa).

Refractive index  $n_D^{20}$ : about 1.55

It is gradually affected by light.

**Identification** (1) To 0.05 g of Diphenhydramine add 2 mL of sulfuric acid: an orange-red precipitate is produced immediately, and its color changes to red-brown on standing. Add carefully 2 mL of water to this solution: the intensity of the color changes, but the color tone does not change.

(2) Dissolve 0.1 g of Diphenhydramine in 10 mL of dilute ethanol, add an excess of a saturated solution of 2,4,6-trinitrophenol in dilute ethanol with stirring, and cool in ice. Collect the produced crystals, recrystallize from dilute ethanol, and dry at 105°C for 30 minutes: the crystals melt between 128°C and 133°C.

**Specific gravity** <2.56>  $d_{20}^{20}$ : 1.013 – 1.020

**Purity** (1)  $\beta$ -Dimethylaminoethanol—Dissolve 1.0 g of Diphenhydramine in 20 mL of diethyl ether, and extract with two 10-mL portions of water with thorough shaking. Combine the water extracts, and add 2 drops of phenolphthalein TS and 1.0 mL of 0.05 mol/L sulfuric acid VS: no red color develops.

(2) Benzohydrol—Transfer 1.0 g of Diphenhydramine to a separator, dissolve in 20 mL of diethyl ether, and extract with two 25-mL portions of diluted hydrochloric acid (1 in 15) with thorough shaking. Separate the diethyl ether layer, evaporate slowly on a water bath, and dry in a desiccator (in vacuum, silica gel) for 2 hours: the mass of the residue is not more than 20 mg.

(3) Heavy metals <1.07>—Proceed with 1.0 g of Diphenhydramine according to Method 2, and perform the test. Prepare the control solution with 2.0 mL of Standard Lead Solution (not more than 20 ppm).

**Residue on ignition** <2.44> Not more than 0.1% (1 g).

**Assay** Weigh accurately about 0.5 g of Diphenhydramine, dissolve in 50 mL of a mixture of acetic anhydride and acetic acid (100) (7:3), and titrate <2.50> with 0.1 mol/L perchloric acid VS (potentiometric titration). Perform a blank determination, and make any necessary correction.

Each mL of 0.1 mol/L perchloric acid VS  
= 25.54 mg of  $C_{17}H_{21}NO$

**Containers and storage** Containers—Tight containers.

**Storage**—Light-resistant, and almost well-filled.

## Diphenhydramine and Bromovalerylurea Powder

ジフェンヒドรามイン・バレリル尿素散

### Method of preparation

|   |                       |
|---|-----------------------|
| Diphenhydramine Tannate                   | 90 g                  |
| Bromovalerylurea                          | 500 g                 |
| Starch, Lactose Hydrate, or their mixture | a sufficient quantity |

To make 1000 g

Prepare as directed under Powders, with the above ingredients.

**Description** Diphenhydramine and Bromovalerylurea Powder occurs as a slightly grayish white powder.

**Identification (1)** To 0.1 g of Diphenhydramine and Bromovalerylurea Powder add 5 mL of dilute hydrochloric acid, 1 mL of ethanol (95) and 10 mL of water, shake, and filter. To the filtrate add 10 mL of sodium hydroxide TS, and extract with 10 mL of chloroform. Separate the chloroform layer, add 1 mL of bromophenol blue TS, and shake: a yellow color develops in the chloroform layer (diphenhydramine tannate).


(2) Shake 0.02 g of Diphenhydramine and Bromovalerylurea Powder with 10 mL of diethyl ether, filter, and evaporate the filtrate on a water bath. Dissolve the residue in 2 mL of sodium hydroxide TS, and add 5 mL of dimethylglyoxime-thiosemicarbazide TS, and heat on a water bath for 30 minutes: a red color develops (bromovalerylurea).

(3) Shake 0.3 g of Diphenhydramine and Bromovalerylurea with 5 mL of methanol, filter, and use the filtrate as the sample solution. Dissolve 0.15 g of bromovalerylurea and 0.03 g of diphenhydramine tannate in 5 mL each of methanol, and use the solutions as standard solution (1) and standard solution (2). Perform the test as directed under Thin-layer Chromatography <2.03> with these solutions. Spot 5  $\mu$ L each of the sample solution and standard solutions (1) and (2) on a plate of silica gel with fluorescent indicator for thin-layer chromatography. Develop the plate in a mixture of ethyl acetate, ethanol (99.5) and ammonia solution (28) (50:5:1) to a distance of about 10 cm. Air-dry the plate, and examine under ultraviolet light (main wavelength: 254 nm): 3 spots from the sample solution and the corresponding spot from standard solutions (1) and (2) show the same *R<sub>f</sub>* value. Spray Dragendorff's TS for spraying evenly on the plate: the spot from the standard solution (2) and the corresponding spot from the sample solution reveal an orange color.

**Containers and storage** Containers—Well-closed containers.

## Diphenhydramine Hydrochloride

ジフェンヒドラミン塩酸塩



$C_{17}H_{21}NO \cdot HCl$ : 291.82

2-(Diphenylmethoxy)-*N,N*-dimethylethylamine monohydrochloride [147-24-0]

Diphenhydramine Hydrochloride, when dried, contains not less than 98.0% of  $C_{17}H_{21}NO \cdot HCl$ .

**Description** Diphenhydramine Hydrochloride occurs as white crystals or crystalline powder. It is odorless, and has a

bitter taste, followed by a sensation of numbness on the tongue.

It is very soluble in methanol and in acetic acid (100), freely soluble in water and in ethanol (95), sparingly soluble in acetic anhydride, and practically insoluble in diethyl ether.

It is gradually affected by light.

**Identification (1)** Determine the absorption spectrum of a solution of Diphenhydramine Hydrochloride in methanol (1 in 2000) as directed under Ultraviolet-visible Spectrophotometry <2.24>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wavelengths.

(2) Determine the infrared absorption spectrum of Diphenhydramine Hydrochloride as directed in the potassium chloride disk method under Infrared Spectrophotometry <2.25>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wave numbers.

(3) A solution of Diphenhydramine Hydrochloride (1 in 50) responds to the Qualitative Tests <1.09> for chloride.

**pH** <2.54> Dissolve 1.0 g of Diphenhydramine Hydrochloride in 10 mL of water: the pH of this solution is between 4.0 and 5.0.

**Melting point** <2.60> 166 – 170°C

**Purity (1)** Clarity and color of solution—Dissolve 1.0 g of Diphenhydramine Hydrochloride in 10 mL of water: the solution is clear and colorless.

(2) Heavy metals <1.07>—Proceed with 1.0 g of Diphenhydramine Hydrochloride according to Method 4, and perform the test. Prepare the control solution with 2.0 mL of Standard Lead Solution (not more than 20 ppm).

(3) Related substances—Dissolve 0.20 g of Diphenhydramine Hydrochloride in 10 mL of methanol, and use this solution as the sample solution. Pipet 1 mL of the sample solution, add methanol to make exactly 200 mL, and use this solution as the standard solution. Perform the test with these solutions as directed under Thin-layer Chromatography <2.03>. Spot 5  $\mu$ L each of the sample solution and standard solution on a plate of silica gel for thin-layer chromatography. Develop the plate with a mixture of hexane, ethyl acetate, methanol and ammonia solution (28) (10:4:2:1) to a distance of about 10 cm, and air-dry the plate. Spray evenly iodine TS on the plate: the spots other than the principal spot from the sample solution and the spot on the original point are not more intense than the spot from the standard solution.

**Loss on drying** <2.41> Not more than 0.5% (2 g, 105°C, 3 hours).

**Residue on ignition** <2.44> Not more than 0.1% (1 g).

**Assay** Weigh accurately about 0.4 g of Diphenhydramine Hydrochloride, previously dried, dissolve in 50 mL of a mixture of acetic anhydride and acetic acid (100) (7:3). Titrate <2.50> with 0.1 mol/L perchloric acid VS (potentiometric titration). Perform a blank determination, and make any necessary correction.

Each mL of 0.1 mol/L perchloric acid VS  
= 29.18 mg of  $C_{17}H_{21}NO \cdot HCl$

**Containers and storage** Containers—Tight containers.

Storage—Light-resistant.

## Diphenhydramine, Phenol and Zinc Oxide Liniment

ジフェンヒドラミン・フェノール・亜鉛華リニメント

### Method of preparation

|                                |       |
|--------------------------------|-------|
| Diphenhydramine                | 20 g  |
| Phenol and Zinc Oxide Liniment | 980 g |
| To make 1000 g                 |       |

Dissolve and mix the above ingredients.

**Description** Diphenhydramine, Phenol and Zinc Oxide Liniment is a white to whitish, pasty mass. It has a slight odor of phenol.

**Identification (1)** To 3 g of Diphenhydramine, Phenol and Zinc Oxide Liniment add 20 mL of hexane, shake well, and separate the hexane layer. Shake thoroughly the hexane solution with 10 mL of 0.2 mol/L hydrochloric acid. Separate the aqueous layer, and adjust with sodium hydroxide TS to a pH of 4.6. Add 1 mL of bromophenol blue-potassium biphthalate TS and 10 mL of chloroform, and shake: a yellow color develops in the chloroform layer (diphenhydramine).

**(2)** Place 1 g of Diphenhydramine, Phenol and Zinc Oxide Liniment in a porcelain crucible, gradually raise the temperature by heating until the mass is charred, and ignite strongly: a yellow color is produced, and disappears on cooling. To the residue add 10 mL of water and 5 mL of dilute hydrochloric acid, shake well, and filter. Add 2 to 3 drops of potassium hexacyanoferrate (II) TS to the filtrate: a white precipitate is produced (zinc oxide).

**(3)** Shake 0.5 g of Diphenhydramine, Phenol and Zinc Oxide Liniment with 1 mL of water and 5 mL of chloroform, filter, and use the filtrate as the sample solution. Dissolve 0.01 g each of diphenhydramine and phenol in 5 mL each of chloroform, and use these solutions as standard solution (1) and standard solution (2). Perform the test as directed under Thin-layer Chromatography <2.03> with the sample solution and the standard solutions. Spot 5  $\mu$ L each of the sample solution and standard solutions (1) and (2) on a plate of silica gel for thin-layer chromatography. Develop the plate with a mixture of ethyl acetate, ethanol (99.5) and ammonia solution (28) (50:5:1) to a distance of about 10 cm, and air-dry the plate. Allow the plate to stand in iodine vapor: two spots from the sample solution and each spot from standard solution (1) and standard solution (2) show the same  $R_f$  value. Sublime iodine, and spray Dragendorff's TS evenly upon the plate: the spot from standard solution (1) and the corresponding spot from the sample solution reveal an orange color.

**Containers and storage** Containers—Tight containers.

Storage—Light-resistant.

## Diphenhydramine Tannate

タンニン酸ジフェンヒドラミン

Diphenhydramine Tannate is a compound of diphenhydramine and tannic acid, and contains not less than 25.0% and not more than 35.0% of diphenhydramine ( $C_{17}H_{21}NO$ : 255.35).

**Description** Diphenhydramine Tannate occurs as a grayish white to light brown powder. It is odorless or has a slight, characteristic odor. It is tasteless.

It is slightly soluble in ethanol (95), and practically insoluble in water and in diethyl ether.

**Identification (1)** To 1 g of Diphenhydramine Tannate add 15 mL of water and 0.3 mL of dilute hydrochloric acid, shake thoroughly for 1 minute, filter, and use this filtrate as the sample solution. Transfer 10 mL of the sample solution to a separator, extract with two 20-mL portions of chloroform, combine the chloroform extracts, and evaporate on a water bath to dryness. To 5 mL of a solution of the residue (1 in 100) add 5 drops of Reinecke salt TS: a light red precipitate is produced.

**(2)** To 10 mL of a solution of the residue obtained in (1) (1 in 100) add 10 mL of 2,4,6-trinitrophenol TS dropwise, and allow to stand for 30 minutes. Collect the precipitate by filtration, recrystallize from dilute ethanol, and dry at 105°C for 30 minutes: the crystals melt <2.60> between 128°C and 133°C.

**(3)** To 1 mL of the sample solution obtained in (1) add 1 drop of iron (III) chloride TS: a dark blue-purple color develops.

**Purity** Heavy metals <1.07>—Proceed with 1.0 g of Diphenhydramine Tannate according to Method 2, and perform the test. Prepare the control solution with 2.0 mL of Standard Lead Solution (not more than 20 ppm).

**Loss on drying** <2.41> Not more than 7.0% (1 g, 105°C, 5 hours).

**Residue on ignition** <2.44> Not more than 1.0% (1 g).

**Assay** Transfer about 1.7 g of Diphenhydramine Tannate, accurately weighed, to a separator, dissolve in 20 mL of water and 3.0 mL of dilute hydrochloric acid with thorough shaking, add 20 mL of a solution of sodium hydroxide (1 in 10) and exactly 25 mL of isooctane, shake vigorously for 5 minutes, dissolve 2 g of sodium chloride with shaking, and allow to stand. To 20 mL of the isooctane layer add exactly 80 mL of acetic acid (100), and titrate <2.50> with 0.1 mol/L perchloric acid VS (potentiometric titration). Perform a blank determination, and make any necessary correction.

Each mL of 0.1 mol/L perchloric acid VS  
= 25.54 mg of  $C_{17}H_{21}NO$

**Containers and storage** Containers—Tight containers.

Storage—Light-resistant.

## Freeze-dried Diphtheria Antitoxin, Equine

乾燥ジフテリアウマ抗毒素

Freeze-dried Diphtheria Antitoxin, Equine, is a preparation for injection which is dissolved before use.

It contains diphtheria antitoxin in immunoglobulin of horse origin.

It conforms to the requirements of Freeze-dried Diphtheria Antitoxin, Equine, in the Minimum Requirements for Biological Products.

**Description** Freeze-dried Diphtheria Antitoxin, Equine, becomes a colorless or light yellow-brown, clear liquid or a slightly whitish turbid liquid on addition of solvent.

## Diphtheria Toxoid

ジフテリアトキソイド

Diphtheria Toxoid is a liquid for injection containing diphtheria toxoid prepared by treating diphtheria toxin with formaldehyde by a method involving no appreciable loss of the immunogenicity.

It conforms to the requirements of Diphtheria Toxoid in the Minimum Requirements for Biological Products.

**Description** Diphtheria Toxoid is a clear, colorless to light yellow-brown liquid.

## Adsorbed Diphtheria Toxoid for Adult Use

成人用沈降ジフテリアトキソイド

Adsorbed Diphtheria Toxoid for Adult Use is a liquid for injection containing diphtheria toxoid prepared by treating diphtheria toxin with formaldehyde by a method involving no appreciable loss of the immunogenicity and very few antigenic substances other than toxoid, and rendered insoluble with aluminum salt.

It conforms to the requirements of Adsorbed Diphtheria Toxoid for Adult Use in the Minimum Requirements of Biological Products.

**Description** Adsorbed Diphtheria Toxoid for Adult Use becomes a homogeneous, whitish turbid liquid on shaking.

## Adsorbed Diphtheria-Purified Pertussis-Tetanus Combined Vaccine

沈降精製百日せきジフテリア破傷風混合ワクチン

Adsorbed Diphtheria-Purified Pertussis-Tetanus Combined Vaccine is a liquid for injection consisting of a liquid containing the protective antigen of *Bordetella pertussis*, Diphtheria Toxoid and a liquid containing tetanus toxoid obtained by detoxifying the tetanus toxin with formaldehyde solution without impairing its immunogenicity, to which aluminum is added to make the antigen and the toxoids insoluble.

It conforms to the requirements of Adsorbed Diphtheria-Purified Pertussis-Tetanus Combined Vaccine in the Minimum Requirements for Biological Products.

**Description** Adsorbed Diphtheria-Purified Pertussis-Tetanus Combined Vaccine becomes a homogeneous, white turbid liquid on shaking.

## Diphtheria-Tetanus Combined Toxoid

ジフテリア破傷風混合トキソイド

Diphtheria-Tetanus Combined Toxoid is a liquid for injection containing diphtheria toxoid and tetanus toxoid which are prepared by treating diphtheria toxin and tetanus toxin, respectively, with formaldehyde by a method involving no appreciable loss of the immunogenicity.

It conforms to the requirements of Diphtheria-Tetanus Combined Toxoid in the Minimum Requirements of Biological Products.

**Description** Diphtheria-Tetanus Combined Toxoid is a colorless or light yellow-brown, clear liquid.

## Adsorbed Diphtheria-Tetanus Combined Toxoid

沈降ジフテリア破傷風混合トキソイド

Adsorbed Diphtheria-Tetanus Combined Toxoid is a liquid for injection containing diphtheria toxoid and tetanus toxoid which are prepared by treating diphtheria toxin and tetanus toxin, respectively, with formaldehyde by a method involving no appreciable loss of the immunogenicity and rendered insoluble by adding aluminum salt.


It conforms to the requirements of Adsorbed

Diphtheria-Tetanus Combined Toxoid in the Minimum Requirements for Biological Products.

**Description** Adsorbed Diphtheria-Tetanus Combined Toxoid becomes a homogeneous, whitish turbid liquid on shaking.

## Dipyridamole

ジピリダモール



$C_{24}H_{40}N_8O_4$ : 504.63

2,2',2'',2'''-[4,8-Di(piperidin-1-yl)pyrimido[5,4-d]pyrimidine-2,6-diyl]dinitrilo}tetraethanol [58-32-2]

Dipyridamole, when dried, contains not less than 98.5% of  $C_{24}H_{40}N_8O_4$ .

**Description** Dipyridamole occurs as yellow crystals or crystalline powder. It is odorless, and has a slightly bitter taste.

It is freely soluble in chloroform, sparingly soluble in methanol and in ethanol (99.5), and practically insoluble in water and in diethyl ether.

**Identification (1)** Dissolve 5 mg of Dipyridamole in 2 mL of sulfuric acid, add 2 drops of nitric acid, and shake: a deep purple color develops.

**(2)** Determine the absorption spectrum of a solution of Dipyridamole in a mixture of methanol and hydrochloric acid (99:1) (1 in 100,000) as directed under Ultraviolet-visible Spectrophotometry <2.24>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wavelengths.

**(3)** Determine the infrared absorption spectrum of Dipyridamole, previously dried, as directed in the potassium bromide disk method under Infrared Spectrophotometry <2.25>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wave numbers.

**Melting point** <2.60> 165 – 169°C

**Purity (1)** Clarity and color of solution—Dissolve 0.5 g of Dipyridamole in 10 mL of chloroform: the solution is clear, and shows a yellow color.

**(2)** Heavy metals <1.07>—Proceed with 2.0 g of Dipyridamole according to Method 2, and perform the test. Prepare the control solution with 2.0 mL of Standard Lead Solution (not more than 10 ppm).

**(3)** Arsenic <1.11>—Prepare the test solution with 1.0 g of Dipyridamole according to Method 3, and perform the test (not more than 2 ppm).

**(4)** Related substances—Dissolve 50 mg of Dipyridamole in 50 mL of the mobile phase, and use this solution as the sample solution. Pipet 0.5 mL of the sample solution, add

the mobile phase to make exactly 100 mL, and use this solution as the standard solution. Perform the test with exactly 20  $\mu$ L each of the sample solution and standard solution as directed under Liquid Chromatography <2.01> according to the following conditions, and determine each peak area by the automatic integration method: the total area of the peaks other than the peak of dipyridamole from the sample solution is not larger than the peak area of dipyridamole from the standard solution.

**Operating conditions—**

**Detector:** An ultraviolet absorption photometer (wavelength: 280 nm).

**Column:** A stainless steel column 4 mm in inside diameter and 15 cm in length, packed with octylsilanized silica gel for liquid chromatography (5  $\mu$ m in particle diameter).

**Column temperature:** A constant temperature of about 40°C.

**Mobile phase:** Dissolve 0.2 g of potassium dihydrogen phosphate in 200 mL of water, and add 800 mL of methanol.

**Flow rate:** Adjust the flow rate so that the retention time of dipyridamole is about 4 minutes.

**Time span of measurement:** About 5 times as long as the retention time of dipyridamole.

**System suitability—**

**Test for required detection:** To exactly 5 mL of the standard solution add the mobile phase to make exactly 25 mL. Confirm that the peak area of dipyridamole obtained from 20  $\mu$ L of this solution is equivalent to 15 to 25% of that of dipyridamole obtained from 20  $\mu$ L of the standard solution.

**System performance:** Dissolve 7 mg of Dipyridamole and 3 mg of terphenyl in 50 mL of methanol. When the procedure is run with 20  $\mu$ L of this solution under the above operating conditions, dipyridamole and terphenyl are eluted in this order with the resolution between these peaks being not less than 5.

**System repeatability:** When the test is repeated 6 times with 20  $\mu$ L of the standard solution under the above operating conditions, the relative standard deviation of the peak area of dipyridamole is not more than 1.0%.

**Loss on drying** <2.41> Not more than 0.2% (1 g, 105°C, 3 hours).

**Residue on ignition** <2.44> Not more than 0.1% (1 g).

**Assay** Weigh accurately about 0.6 g of Dipyridamole, previously dried, dissolve in 70 mL of methanol, and titrate <2.50> with 0.1 mol/L perchloric acid VS (potentiometric titration). Perform a blank determination, and make any necessary correction.

Each mL of 0.1 mol/L perchloric acid VS  
= 50.46 mg of  $C_{24}H_{40}N_8O_4$


**Containers and storage** Containers—Well-closed containers.

Storage—Light-resistant.

## Disodium Edetate Hydrate

### EDTA Sodium Hydrate

エデト酸ナトリウム水和物



$C_{10}H_{14}N_2Na_2O_8 \cdot 2H_2O$ : 372.24

Disodium dihydrogen ethylenediaminetetraacetate dihydrate [6381-92-6]

Disodium Edetate Hydrate contains not less than 99.0% of  $C_{10}H_{14}N_2Na_2O_8 \cdot 2H_2O$ .

**Description** Disodium Edetate Hydrate occurs as white crystals or crystalline powder. It is odorless and has a slight, acid taste.

It is soluble in water, and practically insoluble in ethanol (95) and in diethyl ether.

**Identification (1)** Dissolve 0.01 g of Disodium Edetate Hydrate in 5 mL of water, add 2 mL of a solution of potassium chromate (1 in 200) and 2 mL of arsenic (III) trioxide TS, and heat in a water bath for 2 minutes: a purple color develops.

**(2)** Dissolve 0.5 g of Disodium Edetate Hydrate in 20 mL of water, and add 1 mL of dilute hydrochloric acid: a white precipitate is produced. Collect the precipitate, wash with 50 mL of water, and dry at 105°C for 1 hour: the precipitate melts <2.60> between 240°C and 244°C (with decomposition).

**(3)** A solution of Disodium Edetate Hydrate (1 in 20) responds to the Qualitative Tests <1.09> (1) for sodium salt.

**pH <2.54>** Dissolve 1 g of Disodium Edetate Hydrate in 100 mL of water: the pH of this solution is between 4.3 and 4.7.

**Purity (1)** Clarity and color of solution—Dissolve 1.0 g of Disodium Edetate Hydrate in 50 mL of water: the solution is clear and colorless.

**(2)** Cyanide—Transfer 1.0 g of Disodium Edetate Hydrate to a round-bottomed flask, dissolve in 100 mL of water, add 10 mL of phosphoric acid, and distil. Place 15 mL of 0.5 mol/L sodium hydroxide VS in a 100-mL measuring cylinder, which is used as a receiver, and immerse the bottom end of the condenser into the solution. Distil the mixture until the distillate measures 100 mL, and use this solution as the sample solution. Transfer 20 mL of the sample solution to a glass-stoppered test tube, add 1 drop of phenolphthalein TS, neutralize with dilute acetic acid, and add 5 mL of phosphate buffer solution, pH 6.8, and 1.0 mL of diluted sodium toluensulfoncholoramide TS (1 in 5). Immediately stopper the tube, mix gently, and allow to stand for a few minutes. Mix well with 5 mL of pyridine-pyrazolone TS, and allow to stand between 20°C and 30°C for 50 minutes: the solution has no more color than the following control solution.

Control solution: Pipet 1.0 mL of Standard Cyanide Solution, add 15 mL of 0.5 mol/L sodium hydroxide VS and water to make exactly 1000 mL, transfer 20 mL of this solution to a glass-stoppered test tube, and proceed as directed

for the sample solution.

**(3)** Heavy metals <1.07>—Proceed with 2.0 g of Disodium Edetate Hydrate according to Method 2, and perform the test. Prepare the control solution with 2.0 mL of Standard Lead Solution (not more than 10 ppm).

**(4)** Arsenic <1.11>—Prepare the test solution with 1.0 g of Disodium Edetate Hydrate according to Method 1, and perform the test (not more than 2 ppm).

**Residue on ignition <2.44>** 37.0 – 39.0% (1 g).


**Assay** Weigh accurately about 1 g of Disodium Edetate Hydrate, dissolve in 50 mL of water, add 2 mL of ammonia-ammonium chloride buffer solution, pH 10.7, and 0.04 g of eriochrome black T-sodium chloride indicator, and titrate <2.50> with 0.1 mol/L zinc VS until the color of the solution changes from blue to red.

Each mL of 0.1 mol/L zinc VS  
= 37.22 mg of  $C_{10}H_{14}N_2Na_2O_8 \cdot 2H_2O$

**Containers and storage** Containers—Well-closed containers.

## Disopyramide

ジソピラミド



$C_{21}H_{29}N_3O$ : 339.47

(2*RS*)-4-Bis(1-methylethyl)amino-2-phenyl-2-(pyridin-2-yl)butanamide [3737-09-5]

Disopyramide contains not less than 98.5% of  $C_{21}H_{29}N_3O$ , calculated on the dried basis.

**Description** Disopyramide occurs as white crystals or crystalline powder.

It is very soluble in methanol and in ethanol (95), freely soluble in acetic anhydride, in acetic acid (100) and in diethyl ether, and slightly soluble in water.

**Identification (1)** To 1 mL of a solution of Disopyramide in ethanol (95) (1 in 20) add 10 mL of 2,4,6-trinitrophenol TS, and warm: a yellow precipitate is formed. Filter this precipitate, wash with water, and dry at 105°C for 1 hour: the residue melts <2.60> between 172°C and 176°C.

**(2)** Determine the absorption spectrum of a solution of Disopyramide in 0.05 mol/L sulfuric acid-methanol TS (1 in 25,000) as directed under Ultraviolet-visible Spectrophotometry <2.24>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wavelengths.

**(3)** Determine the infrared absorption spectrum of Disopyramide, previously dried, as directed in the potassium bromide disk method under Infrared Spectrophotometry <2.25>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wave numbers.

**Absorbance** <2.24>  $E_{1\text{cm}}^{1\%}$  (269 nm): 194 – 205 (10 mg, 0.05 mol/L sulfuric acid-methanol TS, 500 mL).

**Purity** (1) Heavy metals <1.07>—Dissolve 1.0 g of Disopyramide in 10 mL of ethanol (95), and add 2 mL of dilute acetic acid (31) and water to make 50 mL. Perform the test using this solution as the test solution. Prepare the control solution as follows: to 2.0 mL of Standard Lead Solution add 10 mL of ethanol (95), 2 mL of dilute acetic acid and water to make 50 mL (not more than 20 ppm).

(2) Arsenic <1.11>—Prepare the test solution with 1.0 g of Disopyramide according to Method 3, and perform the test (not more than 2 ppm).

(3) Related substances—Dissolve 0.40 g of Disopyramide in 10 mL of methanol, and use this solution as the sample solution. Pipet 1 mL of the sample solution, add methanol to make exactly 400 mL, and use this solution as the standard solution. Perform the test with these solutions as directed under Thin-layer Chromatography <2.03>. Spot 10  $\mu\text{L}$  each of the sample solution and standard solution on a plate of silica gel with fluorescent indicator for thin-layer chromatography. Develop the plate with a mixture of 1-butanol, water and ammonia solution (28) (45:4:1) to a distance of about 10 cm, and air-dry the plate. Examine under ultraviolet light (main wavelength: 254 nm): the spots other than the principal spot from the sample solution are not more intense than the spot from the standard solution.

**Loss on drying** <2.41> Not more than 0.5% (0.5 g, in vacuum, 80°C, 2 hours).

**Residue on ignition** <2.44> Not more than 0.20% (1 g).


**Assay** Weigh accurately about 0.25 g of Disopyramide, dissolve in 30 mL of acetic acid (100), and titrate <2.50> with 0.1 mol/L perchloric acid VS (potentiometric titration). Perform a blank determination, and make any necessary correction.

Each mL of 0.1 mol/L perchloric acid VS  
= 16.97 mg of  $\text{C}_{21}\text{H}_{29}\text{N}_3\text{O}$

**Containers and storage** Containers—Tight containers.

## Distigmine Bromide

ジスチグミン臭化物



$\text{C}_{22}\text{H}_{32}\text{Br}_2\text{N}_4\text{O}_4$ ; 576.32

3,3'-[Hexamethylenebis(methyliminocarbonyloxy)]bis(1-methylpyridinium) dibromide [15876-67-2]

Distigmine Bromide contains not less than 98.5% of  $\text{C}_{22}\text{H}_{32}\text{Br}_2\text{N}_4\text{O}_4$ , calculated on the anhydrous basis.

**Description** Distigmine Bromide occurs as a white, crystalline powder.

It is very soluble in water, freely soluble in methanol, in

ethanol (95) and in acetic acid (100), and slightly soluble in acetic anhydride.

The pH of a solution of Distigmine Bromide (1 in 100) is between 5.0 and 5.5.

It is slightly hygroscopic.

It is gradually colored by light.

Melting point: about 150°C (with decomposition).

**Identification** (1) Determine the absorption spectrum of a solution of Distigmine Bromide (1 in 25,000) as directed under Ultraviolet-visible Spectrophotometry <2.24>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wavelengths.

(2) Determine the infrared absorption spectrum of Distigmine Bromide as directed in the potassium bromide disk method under Infrared Spectrophotometry <2.25>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wave numbers.

(3) To 5 mL of a solution of Distigmine Bromide (1 in 10) add 2 mL of dilute nitric acid: the solution responds to the Qualitative Tests <1.09> (1) for bromide.

**Purity** (1) Clarity and color of solution—Dissolve 0.25 g of Distigmine Bromide in 5 mL of water: the solution is clear and colorless.

(2) Sulfate <1.14>—Perform the test with 0.40 g of Distigmine Bromide. Prepare the control solution with 0.40 mL of 0.005 mol/L sulfuric acid VS (not more than 0.048%).

(3) Heavy metals <1.07>—Proceed with 2.0 g of Distigmine Bromide according to Method 2, and perform the test. Prepare the control solution with 2.0 mL of Standard Lead Solution (not more than 10 ppm).

(4) Related substances—Dissolve 40 mg of Distigmine Bromide in 10 mL of methanol, and use this solution as the sample solution. Pipet 1 mL of this solution, add methanol to make exactly 200 mL, and use this solution as the standard solution. Perform the test with these solutions as directed under Thin-layer Chromatography <2.03>. Spot 10  $\mu\text{L}$  each of the sample solution and standard solution on a plate of cellulose with fluorescent indicator for thin-layer chromatography. Develop the plate with a mixture of 1-butanol, water, ethanol (99.5) and acetic acid (100) (8:3:2:1) to a distance of about 13 cm, and air-dry the plate. Examine under ultraviolet light (main wavelength: 254 nm): the spots other than the principal spot from the sample solution are not more intense than the spot from the standard solution. Spray evenly Dragendorff's TS for spraying on the plate: the spots other than the principal spot from the sample solution are not more intense than the spot from the standard solution.

**Water** <2.48> Not more than 1.0% (1 g, direct titration).

**Residue on ignition** <2.44> Not more than 0.1% (1 g).

**Assay** Weigh accurately about 0.4 g of Distigmine Bromide, dissolve in 60 mL of a mixture of acetic anhydride and acetic acid (100) (8:1), and titrate <2.50> with 0.1 mol/L perchloric acid VS (potentiometric titration with platinum electrode). Perform a blank determination, and make any necessary correction.

Each mL of 0.1 mol/L perchloric acid VS  
= 28.82 mg of  $\text{C}_{22}\text{H}_{32}\text{Br}_2\text{N}_4\text{O}_4$

**Containers and storage** Containers—Tight containers.

Storage—Light-resistant.

## Distigmine Bromide Tablets

ジスチグミン臭化物錠

Distigmine Bromide Tablets contain not less than 95% and not more than 105% of the labeled amount of distigmine bromide ( $C_{22}H_{32}Br_2N_4O_4$ ; 576.32).

**Method of preparation** Prepare as directed under Tablets, with Distigmine Bromide.

**Identification** Determine the absorption spectrum of the solution obtained in the Assay, as directed under Ultraviolet-visible Spectrophotometry <2.24>: it exhibits a maximum between 268 nm and 272 nm, and a minimum between 239 nm and 243 nm.

**Assay** Weigh accurately and powder not less than 20 tablets of Distigmine Bromide Tablets. Weigh accurately a portion of the powder, equivalent to about 15 mg of Distigmine Bromide ( $C_{22}H_{32}Br_2N_4O_4$ ), add 30 mL of 0.1 mol/L hydrochloric acid TS, shake for 1 hour, add 0.1 mol/L hydrochloric acid TS to make exactly 50 mL, and filter. Discard the first 20 mL of the filtrate, pipet 10 mL of the subsequent filtrate, add 0.1 mol/L hydrochloric acid TS to make exactly 100 mL, and use this solution as the sample solution. Separately, weigh accurately about 30 mg of distigmine bromide for assay (previously determine the water), and dissolve in 0.1 mol/L hydrochloric acid TS to make exactly 100 mL. Pipet 10 mL of this solution, add 0.1 mol/L hydrochloric acid TS to make exactly 100 mL, and use this solution as the standard solution. Determine the absorbances of the sample solution and standard solution,  $A_{T2}$  and  $A_{S2}$ , at 270 nm and,  $A_{T1}$  and  $A_{S1}$ , at 241 nm as directed under Ultraviolet-visible Spectrophotometry <2.24>, respectively.


$$\text{Amount (mg) of distigmine bromide } (C_{22}H_{32}Br_2N_4O_4) \\ = W_S \times \{(A_{T2} - A_{T1}) / (A_{S2} - A_{S1})\} \times (1/2)$$

$W_S$ : Amount (mg) of distigmine bromide for assay, calculated on the anhydrous basis

**Containers and storage** Containers—Tight containers.

## Disulfiram

ジスルフィラム



$C_{10}H_{20}N_2S_4$ ; 296.54

Tetraethylthiuram disulfide [97-77-8]

Disulfiram, when dried, contains not less than 99.0% of  $C_{10}H_{20}N_2S_4$ .

**Description** Disulfiram occurs as white to yellowish white crystals or crystalline powder.

It is freely soluble in acetone and in toluene, sparingly soluble in methanol and in ethanol (95), and practically insoluble in water.

**Identification** (1) Determine the absorption spectrum of a solution of Disulfiram in ethanol (95) (1 in 100,000) as directed under Ultraviolet-visible Spectrophotometry <2.24>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wavelengths.

(2) Determine the infrared absorption spectrum of Disulfiram, previously dried, as directed in the potassium bromide disk method under Infrared Spectrophotometry <2.25>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wave numbers.

**Melting point** <2.60> 70–73°C

**Purity** (1) Heavy metals <1.07>—Proceed with 2.0 g of Disulfiram according to Method 2, and perform the test. Prepare the control solution with 2.0 mL of Standard Lead Solution (not more than 10 ppm).

(2) Arsenic <1.11>—Prepare the test solution with 1.0 g of Disulfiram according to Method 4, and perform the test (not more than 2 ppm).

(3) Diethyldithiocarbamic acid—Dissolve 0.10 g of Disulfiram in 10 mL of toluene, and shake with 10 mL of diluted sodium carbonate TS (1 in 20). Discard the toluene layer, wash the water layer with 10 mL of toluene, shake with 5 drops of a solution of cupric sulfate (1 in 250) and 2 mL of toluene, and allow to stand: no light yellow color develops in the toluene layer.

(4) Related substances—Dissolve 50 mg of Disulfiram in 40 mL of methanol, add water to make 50 mL, and use this solution as the sample solution. Pipet 1 mL of the sample solution, add the mobile phase to make exactly 200 mL, and use this solution as the standard solution. Perform the test with exactly 10  $\mu$ L each of the sample solution and the standard solution as directed under Liquid Chromatography <2.01> according to the following conditions. Determine each peak area of both solutions by the automatic integration method: the total area of the peaks other than the peak of disulfiram from the sample solution is not larger than the peak area of disulfiram from the standard solution.

**Operating conditions—**

Detector: An ultraviolet absorption photometer (wavelength: 210 nm).

Column: A stainless steel column about 5 mm in inside diameter and about 15 cm in length, packed with octadecylsilylated silica gel for liquid chromatography (5  $\mu$ m in particle diameter).

Column temperature: A constant temperature of about 25°C.

Mobile phase: A mixture of methanol and water (7:3).

Flow rate: Adjust the flow rate so that the retention time of disulfiram is about 8 minutes.

Selection of column: Dissolve 50 mg of Disulfiram and 50 mg of benzophenone in 40 mL of methanol, and add water to make 50 mL. To 1 mL of this solution add the mobile phase to make 200 mL. Proceed with 10  $\mu$ L of this solution under the above operating conditions, and calculate the resolution. Use a column giving elution of benzophenone and disulfiram in this order with the resolution between these peaks being



not less than 4.

**Detection sensitivity:** Adjust the detection sensitivity so that the peak height of disulfiram obtained from 10  $\mu\text{L}$  of the standard solution is 15 – 30 mm.

**Time span of measurement:** About 3.5 times of the retention time of disulfiram.

**Loss on drying** <2.41> Not more than 0.20% (2 g, silica gel, 24 hours).

**Residue on ignition** <2.44> Not more than 0.1% (2 g).


**Assay** Weigh accurately about 0.2 g of Disulfiram, previously dried, in an iodine bottle, dissolve in 20 mL of acetone, add 1.5 mL of water and 1.0 g of potassium iodide, and dissolve by shaking thoroughly. To this solution add 3.0 mL of hydrochloric acid, stopper the bottle tightly, shake, and allow to stand in a dark place for 3 minutes. Add 70 mL of water, and titrate <2.50> with 0.1 mol/L sodium thiosulfate VS (potentiometric titration). Perform a blank determination, and make any necessary correction.

Each mL of 0.1 mol/L sodium thiosulfate VS  
= 14.83 mg of  $\text{C}_{10}\text{H}_{20}\text{N}_2\text{S}_4$

**Containers and storage** Containers—Tight containers.

## Dobutamine Hydrochloride

ドブタミン塩酸塩



$\text{C}_{18}\text{H}_{23}\text{NO}_3 \cdot \text{HCl}$ : 337.84

4-{2-[(1*RS*)-3-(4-Hydroxyphenyl)-1-methylpropylamino]ethyl}benzene-1,2-diol monohydrochloride [49745-95-1]

Dobutamine Hydrochloride, when dried, contains not less than 98.0% of  $\text{C}_{18}\text{H}_{23}\text{NO}_3 \cdot \text{HCl}$ .

**Description** Dobutamine Hydrochloride occurs as white to very pale orange crystalline powder or grains.

It is freely soluble in methanol, sparingly soluble in water and in ethanol (95), and practically insoluble in diethyl ether.

A solution of Dobutamine Hydrochloride (1 in 100) shows no optical rotation.

**Identification (1)** Determine the infrared absorption spectra of Dobutamine Hydrochloride, previously dried, as directed in the potassium bromide disk method under Infrared Spectrophotometry <2.25>, and compare the spectrum with the Reference Spectrum or the spectrum of dried Dobutamine Hydrochloride Reference Standard: both spectra exhibit similar intensities of absorption at the same wave numbers.

**(2)** A solution of Dobutamine Hydrochloride (1 in 50) responds to the Qualitative Tests <1.09> (2) for chloride.

**pH** <2.54> Dissolve 1.0 g of Dobutamine Hydrochloride in 100 mL of water: the pH of this solution is between 4.0 and 5.5.

**Melting point** <2.60> 188 – 191°C

**Purity (1)** Clarity and color of solution—Dissolve 0.5 g of Dobutamine Hydrochloride in 30 mL of water: the solution is clear and colorless.

**(2)** Heavy metals <1.07>—Dissolve 1.0 g of Dobutamine Hydrochloride in 40 mL of water by warming, cool, and add 2 mL of dilute acetic acid and water to make 50 mL. Perform the test using this solution as the test solution. Prepare the control solution as follows: to 2.0 mL of Standard Lead Solution add water to make 50 mL (not more than 20 ppm).

**(3)** Related substances—Dissolve 0.10 g of Dobutamine Hydrochloride in 10 mL of methanol, and use this solution as the sample solution. Pipet 1 mL of the sample solution, add methanol to make exactly 200 mL, and use this solution as the standard solution. Perform the test with these solutions as directed under Thin-layer Chromatography <2.03>. Spot 10  $\mu\text{L}$  each of the sample solution and standard solution on a plate of silica gel for thin-layer chromatography. Develop the plate with a mixture of chloroform, methanol and formic acid (78:22:5) to a distance of about 12 cm, and air-dry the plate. Allow the plate to stand for 5 minutes in iodine vapor: the spots other than the principal spot from the sample solution are not more intense than the spot from the standard solution.

**Loss on drying** <2.41> Not more than 0.30% (1 g, 105°C, 3 hours).

**Residue on ignition** <2.44> Not more than 0.1% (1 g).

**Assay** Weigh accurately about 0.1 g each of Dobutamine Hydrochloride and Dobutamine Hydrochloride Reference Standard, each previously dried, dissolve separately in exactly 10 mL of the internal standard solution, add diluted methanol (1 in 2) to make 50 mL, and use these solutions as the sample solution and standard solution respectively. Perform the test with 5  $\mu\text{L}$  each of the sample solution and standard solution as directed under Liquid Chromatography <2.01> according to the following conditions, and calculate the ratios,  $Q_T$  and  $Q_S$ , of the peak area of dobutamine to that of the internal standard, respectively.

Amount (mg) of  $\text{C}_{18}\text{H}_{23}\text{NO}_3 \cdot \text{HCl}$  =  $W_S \times (Q_T/Q_S)$

$W_S$ : Amount (mg) of Dobutamine Hydrochloride Reference Standard

**Internal standard solution**—A solution of salicylamide in diluted methanol (1 in 2) (1 in 125).

**Operating conditions**—

**Detector:** An ultraviolet absorption photometer (wavelength: 280 nm).

**Column:** A stainless steel column 4.0 mm in inside diameter and 15 cm in length, packed with octadecylsilanized silica gel for liquid chromatography (7  $\mu\text{m}$  in particle diameter).

**Column temperature:** A constant temperature of about 25°C.

**Mobile phase:** A mixture of tartrate buffer solution, pH 3.0 and methanol (7:3).

**Flow rate:** Adjust the flow rate so that the retention time of dobutamine is about 7 minutes.

**System suitability**—

**System performance:** When the procedure is run with 5  $\mu\text{L}$  of the standard solution under the above operating conditions, dobutamine and internal standard are eluted in this


order with the resolution between these peaks being not less than 5.

**System repeatability:** When the test is repeated 6 times with 5  $\mu$ L of the standard solution under the above operating conditions, the relative standard deviation of the ratios of the peak area of dobutamine to that of the internal standard is not more than 1.0%.

**Containers and storage** Containers—Tight containers.

## Dopamine Hydrochloride

ドパミン塩酸塩



$C_8H_{11}NO_2 \cdot HCl$ : 189.64

4-(2-Aminoethyl)benzene-1,2-diol monohydrochloride  
[62-31-7]

Dopamine Hydrochloride, when dried, contains not less than 98.5% of  $C_8H_{11}NO_2 \cdot HCl$ .

**Description** Dopamine Hydrochloride occurs as white crystals or crystalline powder.

It is freely soluble in water and in formic acid, and sparingly soluble in ethanol (95).

Melting point: about 248°C (with decomposition).

**Identification** (1) Determine the absorption spectrum of a solution of Dopamine Hydrochloride in 0.1 mol/L hydrochloric acid TS (1 in 25,000) as directed under Ultraviolet-visible Spectrophotometry <2.24>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wavelengths.

(2) Determine the infrared absorption spectrum of Dopamine Hydrochloride as directed in the potassium chloride disk method under Infrared Spectrophotometry <2.25>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wave numbers.

(3) A solution of Dopamine Hydrochloride (1 in 50) responds to the Qualitative Tests <1.09> (1) for chloride.

**pH** <2.54> Dissolve 1.0 g of Dopamine Hydrochloride in 50 mL of water: the pH of this solution is between 4.0 and 5.5.

**Purity** (1) Clarity and color of solution—Dissolve 1.0 g of Dopamine Hydrochloride in 10 mL of water: the solution is clear and colorless.

(2) Sulfate <1.14>—Perform the test with 0.8 g of Dopamine Hydrochloride. Prepare the control solution with 0.35 mL of 0.005 mol/L sulfuric acid VS (not more than 0.021%).

(3) Heavy metals <1.07>—Proceed with 1.0 g of Dopamine Hydrochloride according to Method 1, and perform the test. Prepare the control solution with 2.0 mL of Standard Lead Solution (not more than 20 ppm).

(4) Arsenic <1.11>—Prepare the test solution with 1.0 g of Dopamine Hydrochloride according to Method 1, and perform the test (not more than 2 ppm).

(5) Related substances—Dissolve 0.10 g of Dopamine Hydrochloride in 10 mL of water, and use this solution as the

sample solution. Pipet 1 mL of the sample solution, add water to make exactly 250 mL, and use this solution as the standard solution. Perform the test with these solutions as directed under Thin-layer Chromatography <2.03>. Spot 5  $\mu$ L each of the sample solution and standard solution on a plate of cellulose with fluorescent indicator for thin-layer chromatography. Develop the plate with a mixture of 1-propanol, water and acetic acid (100) (16:8:1) to a distance of about 10 cm, and air-dry the plate. Spray evenly a solution of ninhydrin in acetone (1 in 50) on the plate, and heat at 90°C for 10 minutes: the spots other than the principal spot from the sample solution are not more intense than the spot from the standard solution.

**Loss on drying** <2.41> Not more than 0.5% (1 g, 105°C, 3 hours).

**Residue on ignition** <2.44> Not more than 0.1% (1 g).

**Assay** Weigh accurately about 0.2 g of Dopamine Hydrochloride, previously dried, dissolve in 5 mL of formic acid, add exactly 15 mL of 0.1 mol/L perchloric acid VS, and heat on a water bath for 15 minutes. After cooling, add 50 mL of acetic acid (100), and titrate <2.50> the excess perchloric acid with 0.1 mol/L sodium acetate VS (potentiometric titration). Perform a blank determination.

Each mL of 0.1 mol/L perchloric acid VS  
= 18.96 mg of  $C_8H_{11}NO_2 \cdot HCl$

**Containers and storage** Containers—Tight containers.

## Dopamine Hydrochloride Injection

ドパミン塩酸塩注射液

Dopamine Hydrochloride Injection is an aqueous solution for injection.

It contains not less than 97% and not more than 103% of the labeled amount of dopamine hydrochloride ( $C_8H_{11}NO_2 \cdot HCl$ : 189.64).

**Method of preparation** Prepare as directed under Injections, with Dopamine Hydrochloride.

**Description** Dopamine Hydrochloride Injection occurs as a clear, colorless liquid.

**Identification** To a volume of Dopamine Hydrochloride Injection, equivalent to 0.04 g of Dopamine Hydrochloride according to the labeled amount, add 0.1 mol/L hydrochloric acid TS to make 100 mL. To 5 mL of this solution add 0.1 mol/L hydrochloric acid TS to make 50 mL. Determine the absorption spectrum of this solution as directed under Ultraviolet-visible Spectrophotometry <2.24>: it exhibits a maximum between 278 nm and 282 nm.

**pH** <2.54> 3.0 – 5.0

**Bacterial endotoxins** <4.01> Less than 4.2 EU/mg.

**Extractable volume** <6.05> It meets the requirement.

**Assay** To an exact volume of Dopamine Hydrochloride Injection, equivalent to about 30 mg of dopamine hydrochloride ( $C_8H_{11}NO_2 \cdot HCl$ ), add the mobile phase to make exactly 50 mL. Pipet 2.5 mL of this solution, add exactly 2.5 mL of

the internal standard solution and the mobile phase to make 20 mL, and use this solution as the sample solution. Separately, weigh accurately about 30 mg of dopamine hydrochloride for assay, previously dried at 105°C for 3 hours, dissolve in the mobile phase to make exactly 50 mL. Pipet 2.5 mL of this solution, add exactly 2.5 mL of the internal standard solution and the mobile phase to make 20 mL, and use this solution as the standard solution. Perform the test with 10  $\mu$ L each of the sample solution and standard solution as directed under Liquid Chromatography <2.01> according to the following conditions, and calculate the ratios,  $Q_T$  and  $Q_S$ , of the peak area of dopamine to that of the internal standard.

$$\text{Amount (mg) of dopamine hydrochloride (C}_8\text{H}_{11}\text{NO}_2\cdot\text{HCl)} = W_S \times (Q_T/Q_S)$$

$W_S$ : Amount (mg) of dopamine hydrochloride for assay

**Internal standard solution**—A solution of uracil in the mobile phase (3 in 10,000).

**Operating conditions**—

**Detector**: An ultraviolet absorption photometer (wavelength: 280 nm).

**Column**: A stainless steel column 4.6 mm in inside diameter and 25 cm in length, packed with octadecylsilanized silica gel for liquid chromatography (5  $\mu$ m in particle diameter).

**Column temperature**: A constant temperature of about 25°C.

**Mobile phase**: Disodium hydrogen phosphate-citric acid buffer solution, pH 3.0

**Flow rate**: Adjust the flow rate so that the retention time of dopamine is about 10 minutes.

**System suitability**—

**System performance**: When the procedure is run with 10  $\mu$ L of the standard solution under the above operating conditions, the internal standard and dopamine are eluted in this order with the resolution between these peaks being not less than 10.


**System repeatability**: When the test is repeated 6 times with 10  $\mu$ L of the standard solution under the above operating conditions, the relative standard deviation of the ratios of peak area of dopamine to that of the internal standard is not more than 1.0%.

**Containers and storage** Containers—Hermetic containers.

Plastic containers for aqueous injections may be used.

## Doxapram Hydrochloride Hydrate

ドキサプラム塩酸塩水和物



$\text{C}_{24}\text{H}_{30}\text{N}_2\text{O}_2\cdot\text{HCl}\cdot\text{H}_2\text{O}$ : 432.98  
(4*RS*)-1-Ethyl-4-[2-(morpholin-4-yl)ethyl]-3,3-diphenylpyrrolidin-2-one monohydrochloride

monohydrate [7081-53-0]

Doxapram Hydrochloride Hydrate contains not less than 98.0% of doxapram hydrochloride ( $\text{C}_{24}\text{H}_{30}\text{N}_2\text{O}_2\cdot\text{HCl}$ : 414.97), calculated on the anhydrous basis.

**Description** Doxapram Hydrochloride Hydrate occurs as white crystals or crystalline powder.

It is freely soluble in methanol and in acetic acid (100), sparingly soluble in water, in ethanol (95) and in acetic anhydride, and practically insoluble in diethyl ether.

**Identification** (1) Determine the absorption spectrum of a solution of Doxapram Hydrochloride Hydrate (1 in 2500) as directed under Ultraviolet-visible Spectrophotometry <2.24>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wavelengths.

(2) Determine the infrared absorption spectrum of Doxapram Hydrochloride Hydrate as directed in the potassium bromide disk method under Infrared Spectrophotometry <2.25>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wave numbers.

(3) A solution of Doxapram Hydrochloride Hydrate (1 in 50) responds to the Qualitative Tests <1.09> for chloride.

**pH** <2.54> Dissolve 1.0 g of Doxapram Hydrochloride Hydrate in 50 mL of water: the pH of this solution is between 3.5 and 5.0.

**Melting point** <2.60> 218 – 222°C

**Purity** (1) Clarity and color of solution—Dissolve 1.0 g of Doxapram Hydrochloride Hydrate in 50 mL of water: the solution is clear and colorless.

(2) Sulfate <1.14>—Perform the test with 1.0 g of Doxapram Hydrochloride Hydrate. Prepare the control solution with 0.50 mL of 0.005 mol/L sulfuric acid VS (not more than 0.024%).

(3) Heavy metals <1.07>—Proceed with 2.0 g of Doxapram Hydrochloride Hydrate according to Method 2, and perform the test. Prepare the control solution with 2.0 mL of Standard Lead Solution (not more than 10 ppm).

(4) Arsenic <1.11>—Prepare the test solution with 1.0 g of Doxapram Hydrochloride Hydrate according to Method 3, and perform the test (not more than 2 ppm).

(5) Related substances—Dissolve 0.5 g of Doxapram Hydrochloride Hydrate in 10 mL of methanol, and use this solution as the sample solution. Pipet 3 mL of the sample solution, and add methanol to make exactly 100 mL. Pipet 5 mL of this solution, add methanol to make exactly 50 mL, and use this solution as the standard solution. Perform the test with these solutions as directed under Thin-layer Chromatography <2.03>. Spot 6  $\mu$ L each of the sample solution and standard solution on a plate of silica gel for thin-layer chromatography. Develop the plate with a mixture of chloroform, formic acid, ethyl formate and methanol (8:3:3:2) to a distance of about 10 cm, and air-dry the plate. Allow the plate to stand in iodine vapor: the spots other than the principal spot from the sample solution are not more intense than the spot from the standard solution.

**Water** <2.48> 3.5 – 4.5% (0.5 g, direct titration).

**Residue on ignition** <2.44> Not more than 0.3% (1 g).


**Assay** Weigh accurately about 0.8 g of Doxapram Hydrochloride, dissolve in 50 mL of a mixture of acetic anhydride and acetic acid (100) (7:3), and titrate <2.50> with 0.1 mol/L perchloric acid VS (potentiometric titration). Perform a blank determination, and make any necessary correction.

Each mL of 0.1 mol/L perchloric acid VS  
= 41.50 mg of  $C_{24}H_{30}N_2O_2 \cdot HCl$

**Containers and storage** Containers—Tight containers.

## Doxifluridine

ドキシフルリジン



$C_9H_{11}FN_2O_5$ : 246.19  
5'-Deoxy-5-fluorouridine  
[3094-09-5]

Doxifluridine, when dried, contains not less than 98.5% and not more than 101.0% of  $C_9H_{11}FN_2O_5$ .

**Description** Doxifluridine occurs as a white crystalline powder.

It is freely soluble in *N,N*-dimethylformamide, soluble in water and in methanol, and slightly soluble in ethanol (99.5).

It dissolves in 0.1 mol/L hydrochloric acid TS and in 0.01 mol/L sodium hydroxide TS.

Melting point: about 191°C (with decomposition).

**Identification (1)** Determine the absorption spectrum of a solution of Doxifluridine in 0.1 mol/L hydrochloric acid TS (1 in 50,000) as directed under Ultraviolet-visible Spectrophotometry <2.24>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wavelengths.

(2) Determine the infrared absorption spectrum of Doxifluridine, previously dried, as directed in the potassium bromide disk method under Infrared Spectrophotometry <2.25>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wave numbers.

**Optical rotation** <2.49>  $[\alpha]_D^{20}$ : +160 – +174° (after drying, 0.1 g, water, 10 mL, 100 mm).

**pH** <2.54> The pH of a solution obtained by dissolving 0.10 g of Doxifluridine in 10 mL of water is between 4.2 and 5.2.

**Purity (1) Fluoride**—Dissolve 0.10 g of Doxifluridine in 10.0 mL of diluted 0.01 mol/L sodium hydroxide TS (1 in 20). Transfer 5.0 mL of this solution into a 20-mL volumetric flask, add 5 mL of a mixture of acetone and lanthanum-alizarin complexone TS (2:1) and water to make 20 mL, allow to stand for 1 hour, and use this solution as the sample solution. Separately, put 1.0 mL of Standard Fluorine Solution in a

20-mL volumetric flask, add 5.0 mL of diluted 0.01 mol/L sodium hydroxide TS (1 in 20) and 5 mL of the mixture of acetone and alizarin complexone TS (2:1), then proceed in the same manner as for preparation of the sample solution, and use the solution so obtained as the standard solution. Determine the absorbances,  $A_T$  and  $A_S$ , of the sample solution and standard solution at 620 nm as directed under Ultraviolet-visible Spectrophotometry <2.24>, using a solution obtained in the same way with 5.0 mL of diluted 0.01 mol/L sodium hydroxide TS (1 in 20) as a blank:  $A_T$  is not larger than  $A_S$ .

(2) Chloride <1.03>—Perform the test with 0.30 g of Doxifluridine. Prepare the control solution with 0.30 mL of 0.01 mol/L hydrochloric acid VS (not more than 0.035%).

(3) Heavy metals <1.07>—Proceed with 1.0 g of Doxifluridine according to Method 1, and perform the test. Prepare the control solution with 2.0 mL of Standard Lead Solution (not more than 20 ppm).

(4) Related substances—Dissolve 20 mg of Doxifluridine in 2 mL of methanol, and use this solution as the sample solution. Pipet 1 mL of the sample solution, and add methanol to make exactly 25 mL. Pipet 5 mL of this solution, add methanol to make exactly 50 mL, and use this solution as the standard solution. Perform the test with these solutions as directed under Thin-layer Chromatography <2.03>. Spot 10  $\mu$ L each of the sample solution and standard solution on a plate of silica gel with fluorescent indicator for thin-layer chromatography. Develop the plate with a mixture of ethyl acetate, acetic acid (100) and water (17:2:1) to a distance of about 12 cm, and air-dry the plate. Examine under ultraviolet light (main wavelength: 254 nm): the number of the spot other than the principal spot is not more than three, and they are not more intense than the spot with the standard solution.

**Loss on drying** <2.41> Not more than 0.5% (1 g, 105°C, 4 hours).

**Residue on ignition** <2.44> Not more than 0.1% (1 g, platinum crucible)

**Assay** Weigh accurately about 0.25 g of Doxifluridine, previously dried, dissolve in 50 mL of *N,N*-dimethylformamide, and titrate <2.50> with 0.1 mol/L tetramethylammonium hydroxide VS (potentiometric titration). Perform a blank determination in the same manner, and make any necessary correction.

Each mL of 0.1 mol/L tetramethylammonium hydroxide VS  
= 24.62 mg of  $C_9H_{11}FN_2O_5$

**Containers and storage** Containers—Tight containers.

## Doxifluridine Capsules

ドキシフルリジンカプセル

Doxifluridine Capsules contain not less than 95.0% and not more than 105.0% of doxifluridine ( $C_9H_{11}FN_2O_5$ : 246.19).

**Method of preparation** Prepare as directed under Capsules, with Doxifluridine.

**Identification (1)** Dissolve an amount of the contents of Doxifluridine Capsules, equivalent to 20 mg of Doxifluridine

according to the labeled amount, in 0.1 mol/L hydrochloric acid TS to make 100 mL, and filter. To 1 mL of the filtrate add 0.1 mol/L hydrochloric acid TS to make 20 mL, and determine the absorption spectrum of this solution as directed under Ultraviolet-visible Spectrophotometry <2.24>, using 0.1 mol/L hydrochloric acid TS as the blank: it exhibits a maximum between 267 nm and 271 nm.

(2) To an amount of powdered contents of Doxifluridine Capsules, equivalent to 20 mg of Doxifluridine according to the labeled amount, add 2 mL of methanol, shake, centrifuge, and use the supernatant liquid as the sample solution. Separately, dissolve 20 mg of doxifluridine in 2 mL of methanol, and use this solution as the standard solution. Perform the test with these solutions as directed under Thin-layer Chromatography <2.03>. Spot 10  $\mu$ L each of the sample solution and standard solution on a plate of silica gel with fluorescent indicator for thin-layer chromatography. Develop the plate with a mixture of ethyl acetate, acetic acid (100) and water (17:2:1) to a distance of about 12 cm, and air-dry the plate. Examine under ultraviolet light (main wavelength: 254 nm): the principal spot with the sample solution and the spot with the standard solution show a dark purple color and these R<sub>f</sub> values are the same.

**Uniformity of dosage units** <6.02> It meets the requirement of the Mass variation test.

**Dissolution** <6.10> Perform the test according to the following method: it meets the requirement.

Perform the test with one capsule of Doxifluridine Capsules at 50 revolutions per minute according to the Paddle method, using 900 mL of water as the dissolution medium, using the sinker. Withdraw 20 mL or more of the dissolution medium 30 minutes after starting the test, and filter through a membrane filter with pore size of not more than 0.45  $\mu$ m. Discard the first 10 mL of the filtrate, pipet the subsequent  $V$  mL, add water to make exactly  $V'$  mL so that each mL contains about 13  $\mu$ g of doxifluridine ( $C_9H_{11}FN_2O_5$ ) according to the labeled amount, and use this solution as the sample solution. Separately, weigh accurately about 26 mg of doxifluridine for assay, previously dried at 105°C for 4 hours, and dissolve in water to make exactly 100 mL. Pipet 5 mL of this solution, add water to make exactly 100 mL, and use this solution as the standard solution. Determine the absorbances,  $A_T$  and  $A_S$ , of these solutions at 269 nm as directed under Ultraviolet-visible Spectrophotometry <2.24>. The dissolution rate in 30 minutes is not less than 85%.

Dissolution rate (%) with respect to the labeled amount of doxifluridine ( $C_9H_{11}FN_2O_5$ )

$$= W_S \times (A_T/A_S) \times (V/V') \times (1/C) \times 9$$

$W_S$ : Amount (mg) of doxifluridine for assay

$C$ : Labeled amount (mg) of doxifluridine ( $C_9H_{11}FN_2O_5$ ) in 1 capsule

**Assay** Weigh accurately the mass and powder the contents of not less than 20 Doxifluridine Capsules. Weigh accurately a portion of the powder, equivalent to about 50 mg of doxifluridine ( $C_9H_{11}FN_2O_5$ ) according to the labeled amount, add 40 mL of water, shake for about 10 minutes, add water to make exactly 50 mL, and filter. Discard the first 10 mL of the filtrate, pipet 5 mL of the subsequent filtrate, add exactly 10 mL of the internal standard solution, then add a mixture of water and methanol (5:3) to make 100 mL, and use this solu-

tion as the sample solution. Separately, weigh accurately about 50 mg of doxifluridine for assay, previously dried at 105°C for 4 hours, and dissolve in water to make exactly 50 mL. Pipet 5 mL of this solution, add exactly 10 mL of the internal standard solution, then add the mixture of water and methanol (5:3) to make exactly 100 mL, and use this solution as the standard solution. Perform the test with 10  $\mu$ L each of the sample solution and standard solution as directed under Liquid Chromatography <2.01> according to the following conditions, and determine the ratios,  $Q_T$  and  $Q_S$ , of the peak height of doxifluridine to that of the internal standard.

$$\begin{aligned} &\text{Amount (mg) of doxifluridine (C}_9\text{H}_{11}\text{FN}_2\text{O}_5\text{)} \\ &= W_S \times (Q_T/Q_S) \end{aligned}$$

$W_S$ : Amount (mg) of doxifluridine for assay

**Internal standard solution**—A solution of anhydrous caffeine (1 in 1000).

**Operating conditions**—

**Detector:** An ultraviolet absorption photometer (wavelength: 254 nm).

**Column:** A stainless steel column 6 mm in inside diameter and 15 cm in length, packed with octadecylsilanized silica gel for liquid chromatography (5  $\mu$ m in particle diameter).

**Column temperature:** A constant temperature of about 25°C.

**Mobile phase:** A mixture of water and methanol (13:7).

**Flow rate:** Adjust the flow rate so that the retention time of doxifluridine is about 2.5 minutes.

**System suitability**—


**System performance:** When the procedure is run with 10  $\mu$ L of the standard solution under the above operating conditions, doxifluridine and the internal standard are eluted in this order with the resolution between these peaks being not less than 5.

**System repeatability:** When the test is repeated 6 times with 10  $\mu$ L of the standard solution under the above operating conditions, the relative standard deviation of the ratio of the peak height of doxifluridine to that of the internal standard is not more than 1.0%.

**Containers and storage** Containers—Tight containers.

## Doxorubicin Hydrochloride

ドキシルビシン塩酸塩



$C_{27}H_{29}NO_{11} \cdot HCl$ : 579.98

(2*S*,4*S*)-4-(3-Amino-2,3,6-trideoxy- $\alpha$ -L-xylohexopyranosyloxy)-2,5,12-trihydroxy-2-hydroxyacetyl-7-methoxy-1,2,3,4-tetrahydrotetracene-6,11-dione monohydrochloride [25316-40-9]

Doxorubicin Hydrochloride is the hydrochloride of

a derivative of daunorubicin.

It contains not less than 980  $\mu\text{g}$  (potency) and not more than 1080  $\mu\text{g}$  (potency) per mg, calculated on the anhydrous basis. The potency of Doxorubicin Hydrochloride is expressed as mass (potency) of doxorubicin hydrochloride ( $\text{C}_{27}\text{H}_{29}\text{NO}_{11}\cdot\text{HCl}$ ).

**Description** Doxorubicin Hydrochloride occurs as a red-orange crystalline powder.

It is sparingly soluble in water, slightly soluble in methanol, very slightly soluble in ethanol (99.5), and practically insoluble in acetonitrile.

**Identification (1)** Determine the absorption spectrum of a solution of Doxorubicin Hydrochloride in methanol (1 in 100,000) as directed under Ultraviolet-visible Spectrophotometry <2.24>, and compare the spectrum with the Reference Spectrum or the spectrum of a solution of Doxorubicin Hydrochloride Reference Standard prepared in the same manner as the sample solution: both spectra exhibit similar intensities of absorption at the same wavelengths.

**(2)** Determine the infrared absorption spectrum of Doxorubicin Hydrochloride as directed in the potassium chloride disk method under Infrared Spectrophotometry <2.25>, and compare the spectrum with the Reference Spectrum or the spectrum of Doxorubicin Hydrochloride Reference Standard: both spectra exhibit similar intensities of absorption at the same wave numbers.

**(3)** A solution of Doxorubicin Hydrochloride (1 in 200) responds to the Qualitative Tests <1.09> (1) for chloride.

**Optical rotation** <2.49>  $[\alpha]_{\text{D}}^{20}$ : +240 – +290° (20 mg calculated on the anhydrous basis, methanol, 20 mL, 100 mm).

**pH** <2.54> The pH of a solution obtained by dissolving 50 mg of Doxorubicin Hydrochloride in 10 mL of water is between 4.0 and 5.5.

**Purity (1)** Clarity and color of solution—Dissolve 50 mg of Doxorubicin Hydrochloride in 10 mL of water: the solution is clear and red.

**(2)** Related substances—Dissolve 25 mg of Doxorubicin Hydrochloride in 100 mL of the mobile phase, and use this solution as the sample solution. Pipet 2 mL of the sample solution, add the mobile phase to make exactly 100 mL, and use this solution as the standard solution. Perform the test with exactly 20  $\mu\text{L}$  each of the sample solution and standard solution as directed under Liquid Chromatography <2.01> according to the following conditions, and determine each peak area by the automatic integration method: the area of the peak other than doxorubicin obtained from the sample solution is not more than 1/4 times the peak area of doxorubicin from the standard solution, and the total area of the peaks other than doxorubicin is not more than the peak area of doxorubicin from the standard solution.

**Operating conditions—**

**Detector:** An ultraviolet absorption photometer (wavelength: 254 nm).

**Column:** A stainless steel column 4.6 mm in inside diameter and 25 cm in length, packed with octadecylsilanized silica gel for liquid chromatography (5  $\mu\text{m}$  in particle diameter).

**Column temperature:** A constant temperature of about 25°C.

**Mobile phase:** Dissolve 3 g of sodium lauryl sulfate in 1000

mL of diluted phosphoric acid (7 in 5000), and add 1000 mL of acetonitrile.

**Flow rate:** Adjust the flow rate so that the retention time of doxorubicin is about 8 minutes.

**Time span of measurement:** About 3 times as long as the retention time of doxorubicin.

**System suitability—**

**Test for required detectability:** Measure 1 mL of the standard solution, and add the mobile phase to make exactly 20 mL. Confirm that the peak area of doxorubicin obtained from 20  $\mu\text{L}$  of this solution is equivalent to 3.5 to 6.5% of that from 20  $\mu\text{L}$  of the standard solution.

**System performance:** Dissolve 5 mg of Doxorubicin Hydrochloride in 20 mL of water, add 1.5 mL of phosphoric acid, and allow to stand at room temperature for 30 minutes. Adjust the pH of this solution to 2.5 with 2 mol/L sodium hydroxide TS. When the procedure is run with 20  $\mu\text{L}$  of this solution under the above operating conditions, doxorubicinone, having the relative retention time of about 0.6 with respect to doxorubicin, and doxorubicin are eluted in this order with the resolution between these peaks being not less than 5.

**System repeatability:** When the test is repeated 6 times with 20  $\mu\text{L}$  of the standard solution under the above operating conditions, the relative standard deviation of the peak area of doxorubicin is not more than 2.0%.

**Water** <2.48> Not more than 3.0% (0.3 g, volumetric titration, direct titration).

**Assay** Weigh accurately an amount of Doxorubicin Hydrochloride and Doxorubicin Hydrochloride Reference Standard, equivalent to about 10 mg (potency), dissolve each in water to make exactly 25 mL. Pipet 5 mL each of these solutions, add water to make exactly 100 mL, and use these solutions as the sample solution and the standard solution, respectively. Determine the absorbances at 495 nm,  $A_{\text{T}}$  and  $A_{\text{S}}$ , of the sample solution and standard solution as directed under Ultraviolet-visible Spectrophotometry <2.24>.


$$\begin{aligned} &\text{Amount } [\mu\text{g (potency)}] \text{ of } \text{C}_{27}\text{H}_{29}\text{NO}_{11}\cdot\text{HCl} \\ &= W_{\text{S}} \times (A_{\text{T}}/A_{\text{S}}) \times 1000 \end{aligned}$$

$W_{\text{S}}$ : Amount [mg (potency)] of Doxorubicin Hydrochloride Reference Standard

**Containers and storage** Containers—Tight containers.

## Doxycycline Hydrochloride Hydrate

ドキシサイクリン塩酸塩水和物



$\text{C}_{22}\text{H}_{24}\text{N}_2\text{O}_8\cdot\text{HCl}\cdot\frac{1}{2}\text{C}_2\text{H}_6\text{O}\cdot\frac{1}{2}\text{H}_2\text{O}$ : 512.94  
(4*S*,4*aR*,5*S*,5*aR*,6*R*,12*aS*)-4-Dimethylamino-3,5,10,12,12*a*-pentahydroxy-6-methyl-1,11-dioxo-1,4,4*a*,5,5*a*,6,11,12*a*-octahydrotetracene-2-carboxamide monohydrochloride hemiethanolate hemihydrate [564-25-0, Doxycycline]

Doxycycline Hydrochloride Hydrate is the hydrochloride of a derivative of oxytetracycline.

It contains not less than 880  $\mu\text{g}$  (potency) and not more than 943  $\mu\text{g}$  (potency) per mg, calculated on the anhydrous basis and corrected by the amount of ethanol. The potency of Doxycycline Hydrochloride Hydrate is expressed as mass (potency) of doxycycline ( $\text{C}_{22}\text{H}_{24}\text{N}_2\text{O}_8$ : 444.43).

**Description** Doxycycline Hydrochloride Hydrate occurs as yellow to dark yellow, crystals or crystalline powder.

It is freely soluble in water and in methanol, and slightly soluble in ethanol (99.5).

**Identification (1)** Determine the infrared absorption spectrum of Doxycycline Hydrochloride Hydrate as directed in the potassium bromide disk method under Infrared Spectrophotometry <2.25>, and compare the spectrum with the Reference Spectrum or the spectrum of Doxycycline Hydrochloride Reference Standard: both spectra exhibit similar intensities of absorption at the same wave numbers.

(2) Dissolve 10 mg of Doxycycline Hydrochloride Hydrate in 10 mL of water, and add silver nitrate TS: a white turbidity is produced.

**Absorbance** <2.24>  $E_{1\text{ cm}}^{1\%}$  (349 nm): 285 – 315 (10 mg, 0.01 mol/L hydrochloric acid-methanol TS, 500 mL).

**Optical rotation** <2.49>  $[\alpha]_{\text{D}}^{20}$ :  $-105 - -120^\circ$  (0.25 g calculated on the anhydrous basis and corrected by the amount of ethanol, 0.01 mol/L hydrochloric acid-methanol TS, 25 mL, 100 mm). Determine within 5 minutes after the sample solution is prepared.

**Purity (1)** Heavy metals <1.07>—Proceed with 1.0 g of Doxycycline Hydrochloride Hydrate according to Method 2, and perform the test. Prepare the control solution with 5.0 mL of Standard Lead Solution (not more than 50 ppm).

(2) Related substance—Dissolve 20 mg of Doxycycline Hydrochloride Hydrate in 0.01 mol/L hydrochloric acid TS to make exactly 25 mL, and use this solution as the sample solution. Separately, dissolve 20 mg of 6-epidoxycycline hydrochloride in 0.01 mol/L hydrochloric acid TS to make exactly 25 mL, and use this solution as 6-epidoxycycline hydrochloride stock solution. Separately, dissolve 20 mg of metacycline hydrochloride in 0.01 mol/L hydrochloric acid TS to make exactly 25 mL, and use this solution as metacycline hydrochloride stock solution. Pipet 2 mL each of 6-epidoxycycline hydrochloride stock solution and metacycline hydrochloride stock solution, add 0.01 mol/L hydrochloric acid TS to make exactly 100 mL, and use this solution as the standard solution. Perform the test with exactly 20  $\mu\text{L}$  each of the sample solution and standard solution as directed under Liquid Chromatography <2.01> according to the following conditions, and determine each peak area by the automatic integration method: the peak areas of metacycline and 6-epidoxycycline obtained from the sample solution are not more than the peak areas of them obtained from the standard solution, respectively, and the areas of the two peaks, appeared between the solvent peak and metacycline and behind of doxycycline, obtained from the sample solution are not more than 1/4 of the peak area of 6-epidoxycycline from the standard solution.

*Operating conditions—*

Detector: An ultraviolet absorption photometer (wavelength: 254 nm).

Column: A stainless steel column 4.6 mm in inside diameter and 25 cm in length, packed with styrene-divinylbenzene copolymer for liquid chromatography (8  $\mu\text{m}$  in particle diameter).

Column temperature: A constant temperature of about 60°C.

Mobile phase: Mix 125 mL of 0.2 mol/L potassium dihydrogen phosphate TS, 117 mL of 0.2 mol/L sodium hydroxide TS, and add water to make 500 mL. To 400 mL of this solution add 50 mL of a solution of tetrabutylammonium hydrogensulfate (1 in 100), 10 mL of a solution of disodium dihydrogen ethylenediamine tetraacetate dihydrate (1 in 25), 60 g of *t*-butanol and 200 mL of water, adjust the pH to 8.0 with 2 mol/L sodium hydroxide TS, and add water to make 1000 mL.

Flow rate: Adjust the flow rate so that the retention time of doxycycline is about 19 minutes.

Time span of measurement: About 2.4 times as long as the retention time of doxycycline beginning after the solvent peak.

*System suitability—*

Test for required detectability: Measure exactly 1 mL of the standard solution, and add 0.01 mol/L hydrochloric acid TS to make exactly 20 mL. Confirm that the peak areas of 6-epidoxycycline and metacycline obtained from 20  $\mu\text{L}$  of this solution are equivalent to 3.5 to 6.5% of them obtained from 20  $\mu\text{L}$  of the standard solution, respectively.

System performance: To 8 mL of the sample solution, 3 mL of 6-epidoxycycline hydrochloride stock solution and 2 mL of metacycline hydrochloride stock solution add 0.01 mol/L hydrochloric acid TS to make 50 mL. When the procedure is run with 20  $\mu\text{L}$  of this solution under the above operating conditions, metacycline, 6-epidoxycycline and doxycycline are eluted in this order with the resolutions between the peaks, metacycline and 6-epidoxycycline, and 6-epidoxycycline and doxycycline, being not less than 1.3 and not less than 2.0, respectively, and the symmetry factor of the peak of doxycycline is not more than 1.3.

System repeatability: When the test is repeated 6 times with 20  $\mu\text{L}$  of the standard solution under the above operating conditions, the relative standard deviations of the peak area of metacycline and 6-epidoxycycline are not more than 3.0% and not more than 2.0%, respectively.

**Ethanol** Weigh accurately about 0.1 g of Doxycycline Hydrochloride Hydrate, dissolve in the internal standard solution to make exactly 10 mL, and use this solution as the sample solution. Separately, weigh accurately about 0.4 g of ethanol (99.5), and add the internal standard solution to make exactly 100 mL. Pipet 1 mL of this solution, add the internal standard solution to make exactly 10 mL, and use this solution as the standard solution. Perform the test with 1  $\mu\text{L}$  each of the sample solution and standard solution as directed under Gas Chromatography <2.02> according to the following conditions, and determine the ratios,  $Q_T$  and  $Q_S$ , of the peak area of ethanol to that of the internal standard: the amount of ethanol is not less than 4.3% and not more than 6.0%.

$$\text{Amount (\%)} \text{ of ethanol} = (W_S/W_T) \times (Q_T/Q_S)$$

$W_S$ : Amount (mg) of ethanol (99.5)

$W_T$ : Amount (mg) of the sample

**Internal standard solution**—A solution of 1-propanol (1 in 2000).

**Operating conditions**—

Detector: An hydrogen flame-ionization detector.

Column: A glass column 3.2 mm in inside diameter and 1.5 m in length, packed with porous ethylvinylbenzene-divinylbenzene copolymer for gas chromatography (0.0075  $\mu$ m in average pore size, 500 – 600 m<sup>2</sup>/g in specific surface area) (150 – 180  $\mu$ m in particle diameter).

Column temperature: A constant temperature of about 135°C.

Carrier gas: Nitrogen

Flow rate: Adjust the flow rate so that the retention time of ethanol is about 5 minutes.

**System suitability**—

System performance: When the procedure is run with 1  $\mu$ L of the standard solution under the above operating conditions, ethanol and the internal standard are eluted in this order with the resolution between these peaks being not less than 2.0.

System repeatability: When the test is repeated 5 times with 1  $\mu$ L of the standard solution under the above operating conditions, the relative standard deviation of the ratios of the peak area of ethanol to that of the internal standard is not more than 2.0%.

**Water** <2.48> Not less than 1.4% and not more than 2.8% (0.6 g, volumetric titration, direct titration).

**Residue on ignition** <2.44> Not more than 0.3% (1 g).

**Assay** Weigh accurately an amount of Doxycycline Hydrochloride Hydrate and Doxycycline Hydrochloride Reference Standard, equivalent to about 50 mg (potency), dissolve each in water to make exactly 50 mL, and use these solutions as the sample solution and standard solution. Perform the test with exactly 10  $\mu$ L each of the sample solution and standard solution as directed under Liquid Chromatography <2.01> according to the following conditions, and determine the peak areas,  $A_T$  and  $A_S$ , of doxycycline.

$$\begin{aligned} &\text{Amount } [\mu\text{g (potency)}] \text{ of doxycycline (C}_{22}\text{H}_{24}\text{N}_2\text{O}_8) \\ &= W_S \times (A_T/A_S) \times 1000 \end{aligned}$$

$W_S$ : Amount [mg (potency)] of Doxycycline Hydrochloride Reference Standard

**Operating conditions**—

Detector: An ultraviolet absorption photometer (wavelength: 280 nm).

Column: A stainless steel column 3.9 mm in inside diameter and 30 cm in length, packed with octadecylsilanized silica gel for liquid chromatography (10  $\mu$ m in particle diameter).

Column temperature: A constant temperature of about 30°C.

Mobile phase: Dissolve 7.0 g of sodium dihydrogen phosphate dihydrate in 450 mL of water, add 553 mL of a mixture of methanol and *N,N*-dimethyl-*n*-octylamine (550:3), and adjust the pH to 8.0 with a solution of sodium hydroxide (43 in 200).

Flow rate: Adjust the flow rate so that the retention time of doxycycline is about 6 minutes.

**System suitability**—

System performance: When the procedure is run with 10  $\mu$ L of the standard solution under the above operating conditions, the theoretical plates and the symmetry factor of the peak of doxycycline are not less than 1000 and not more than 2.0, respectively.


System repeatability: When the test is repeated 6 times with 10  $\mu$ L of the standard solution under the above operating conditions, the relative standard deviation of the peak area of doxycycline is not more than 1.0%.

**Containers and storage** Containers—Tight containers.

Storage—Light-resistant.

## Droperidol

ドロペリドール



$\text{C}_{22}\text{H}_{22}\text{FN}_3\text{O}_2$ : 379.43

1-[1-[4-(4-Fluorophenyl)-4-oxobutyl]-1,2,3,6-tetrahydropyridin-4-yl]-1,3-dihydro-2H-benzimidazol-2-one [548-73-2]

Droperidol, when dried, contains not less than 98.0% of  $\text{C}_{22}\text{H}_{22}\text{FN}_3\text{O}_2$ .

**Description** Droperidol occurs as a white to light yellow powder.

It is freely soluble in acetic acid (100), soluble in dichloromethane, slightly soluble in ethanol (99.5), and practically insoluble in water.

It is gradually colored by light.

**Identification** (1) Put 30 mg of Droperidol in a brown volumetric flask, and dissolve in 10 mL of 0.1 mol/L hydrochloric acid TS and ethanol (95) to make 100 mL. Transfer 5 mL of the solution to a brown volumetric flask, and add 10 mL of 0.1 mol/L hydrochloric acid TS and ethanol (95) to make 100 mL. Determine the absorption spectrum of the solution as directed under Ultraviolet-visible Spectrophotometry <2.24>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wavelengths.

(2) Determine the infrared absorption spectrum of Droperidol, previously dried, as directed in the potassium bromide disk method under Infrared Spectrophotometry <2.25>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wave numbers. If any difference appears between the spectra, dissolve Droperidol in acetone, evaporate the acetone, dry the residue in a desiccator (in vacuum, silica gel, 70°C) for 4 hours, and perform the test with the residue.

**Purity** (1) Heavy metals <1.07>—Proceed with 1.0 g of Droperidol in a platinum crucible according to Method 2, and perform the test. Prepare the control solution with 2.0



mL of Standard Lead Solution (not more than 20 ppm).

(2) Related substances—Conduct this procedure without exposure to daylight, using light-resistant vessels. Dissolve 50 mg of Droperidol in 5 mL of dichloromethane, and use this solution as the sample solution. Pipet 1 mL of the sample solution, add dichloromethane to make exactly 100 mL, and use this solution as the standard solution. Perform the test with these solutions as directed under Thin-layer Chromatography <2.03>. Spot 10  $\mu$ L each of the sample solution and standard solution on a plate of silica gel with fluorescent indicator for thin-layer chromatography. Develop the plate with a mixture of ethyl acetate, chloroform, methanol and acetic acid-sodium acetate buffer solution, pH 4.7, (54:23:18:5) to a distance of about 15 cm, and air-dry the plate. Examine under ultraviolet light (main wavelength: 254 nm): the spots other than the principal spot from the sample solution are not more intense than the spot from the standard solution.

**Loss on drying** <2.41> Not more than 3.0% (0.5 g, in vacuum, silica gel, 70°C, 4 hours).

**Residue on ignition** <2.44> Not more than 0.2% (1 g, platinum crucible).


**Assay** Weigh accurately about 0.5 g of Droperidol, previously dried, dissolve in 50 mL of acetic acid (100), and titrate <2.50> with 0.1 mol/L perchloric acid VS (potentiometric titration). Perform a blank determination, and make any necessary correction.

Each mL of 0.1 mol/L perchloric acid VS  
= 37.94 mg of  $C_{22}H_{22}FN_3O_2$

**Containers and storage** Containers—Tight containers.  
Storage—Light-resistant.

## Dydrogesterone

ジドロゲステロン



$C_{21}H_{28}O_2$ : 312.45  
9 $\beta$ ,10 $\alpha$ -Pregna-4,6-diene-3,20-dione [152-62-5]

Dydrogesterone, when dried, contains not less than 98.0% and not more than 102.0% of  $C_{21}H_{28}O_2$ .

**Description** Dydrogesterone occurs as white to light yellowish white crystals or crystalline powder. It is odorless.

It is freely soluble in chloroform, soluble in acetonitrile, sparingly soluble in methanol and in ethanol (95), slightly soluble in diethyl ether, and practically insoluble in water.

**Identification** (1) To 5 mg of Dydrogesterone add 5 mL of 4-methoxybenzaldehyde-acetic acid TS and 2 to 3 drops of sulfuric acid, and heat in a water bath for 2 minutes: an orange-red color develops.

(2) Determine the absorption spectrum of a solution of Dydrogesterone in methanol (1 in 200,000) as directed under

Ultraviolet-visible Spectrophotometry <2.24>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wavelengths.

(3) Determine the infrared absorption spectrum of Dydrogesterone, previously dried, as directed in the potassium bromide disk method under Infrared Spectrophotometry <2.25>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wave numbers.

**Optical rotation** <2.49>  $[\alpha]_D^{20}$ :  $-470 - -500^\circ$  (after drying, 0.1 g, chloroform, 10 mL, 100 mm).

**Melting point** <2.60> 167 – 171°C

**Purity** (1) Heavy metals <1.07>—Proceed with 1.0 g of Dydrogesterone according to Method 2, and perform the test. Prepare the control solution with 2.0 mL of Standard Lead Solution (not more than 20 ppm).

(2) Related substances—Dissolve 10 mg of Dydrogesterone in 200 mL of the mobile phase, and use this solution as the sample solution. Pipet 1 mL of the sample solution, add the mobile phase to make exactly 100 mL, and use this solution as the standard solution. Perform the test with exactly 10  $\mu$ L each of the sample solution and standard solution as directed under Liquid Chromatography <2.01> according to the following conditions. Determine each peak area of these solutions by the automatic integration method: the total area of peaks other than the peak of dydrogesterone from the sample solution is not larger than the peak area of dydrogesterone from the standard solution.

**Operating conditions**—

**Detector:** An ultraviolet absorption photometer (wavelength: 280 nm).

**Column:** A stainless steel column about 4 mm in inside diameter and about 15 cm in length, packed with octadecylsilylated silica gel for liquid chromatography (3  $\mu$ m in particle diameter).

**Column temperature:** A constant temperature of about 40°C.

**Mobile phase:** A mixture of water, ethanol (95) and acetonitrile (53:26:21).

**Flow rate:** Adjust the flow rate so that the retention time of dydrogesterone is about 12 minutes.

**Selection of column:** Dissolve 1 mg each of Dydrogesterone and progesterone in 20 mL of the mobile phase. Proceed with 10  $\mu$ L each of these solutions under the above operating conditions, and calculate the resolution. Use a column giving elution of dydrogesterone and progesterone in this order with the resolution between these peaks being not less than 8. Wavelength is 265 nm.

**Detection sensitivity:** Adjust the detection sensitivity so that the peak height of dydrogesterone obtained from 10  $\mu$ L of the standard solution is between 5 mm and 10 mm.

**Time span of measurement:** About twice as long as the retention time of dydrogesterone beginning after the solvent peak.

**Loss on drying** <2.41> Not more than 0.5% (0.5 g, in vacuum, phosphorus (V) oxide, 24 hours).

**Residue on ignition** <2.44> Not more than 0.1% (1 g).

**Assay** Weigh accurately about 50 mg of Dydrogesterone, previously dried, and dissolve in methanol to make exactly

100 mL. Pipet 1 mL of this solution, and add methanol to make exactly 100 mL. Determine the absorbance  $A$  of this solution at the wavelength of maximum absorption at about 286 nm as directed under Ultraviolet-visible Spectrophotometry <2.24>.

$$\text{Amount (mg) of } C_{21}H_{28}O_2 = (A/845) \times 100,000$$

**Containers and storage** Containers—Tight containers.

## Dydrogesterone Tablets

ジドロゲステロン錠

Dydrogesterone Tablets contain not less than 95% and not more than 105% of the labeled amount of dydrogesterone ( $C_{21}H_{28}O_2$ : 312.45).

**Method of preparation** Prepare as directed under Tablets, with Dydrogesterone.

**Identification (1)** To a quantity of powdered Dydrogesterone Tablets, equivalent to 0.05 g of Dydrogesterone according to the labeled amount, add 50 mL of methanol, shake well, and filter. Evaporate 5 mL of the filtrate on a water bath to dryness. Proceed with the residue as directed in the Identification (1) under Dydrogesterone.

**(2)** To 1 mL of the filtrate obtained in (1) add methanol to make 200 mL. Determine the absorption spectrum of this solution as directed under Ultraviolet-visible Spectrophotometry <2.24>: it exhibits a maximum between 284 nm and 288 nm.

**Dissolution <6.10>** Perform the test according to the following method: it meets the requirement.

Perform the test with 1 tablet of Dydrogesterone Tablets at 50 revolutions per minute according to the Paddle method, using 900 mL of water as the dissolution medium. Take 20 mL or more of the dissolved solution 30 minutes after starting the test, and filter. Discard the first 10 mL of the filtrate, and use the subsequent as the sample solution. Separately, weigh accurately about 50 mg of dydrogesterone for assay, previously dried in a desiccator (in vacuum, phosphorus (V) oxide) for 24 hours, and dissolve in methanol to make exactly 100 mL. Pipet 1 mL of this solution, add water to make exactly 100 mL, and use this solution as the standard solution. Determine the absorbances,  $A_T$  and  $A_S$ , of the sample solution and standard solution at 296 nm as directed under Ultraviolet-visible Spectrophotometry <2.24>.

The dissolution rate of Dydrogesterone Tablets in 30 minutes is not less than 80%.

$$\begin{aligned} &\text{Dissolution rate (\%)} \text{ with respect to} \\ &\text{the labeled amount of dydrogesterone } (C_{21}H_{28}O_2) \\ &= W_S \times (A_T/A_S) \times (1/C) \times 9 \end{aligned}$$

$W_S$ : Amount (mg) of dydrogesterone for assay

$C$ : Labeled amount (mg) of dydrogesterone ( $C_{21}H_{28}O_2$ ) in 1 tablet.

**Assay** Weigh accurately and powder not less than 20 Dydrogesterone Tablets. Weigh accurately a portion of the powder, equivalent to about 10 mg of dydrogesterone ( $C_{21}H_{28}O_2$ ), shake well with 50 mL of methanol, and add methanol to make exactly 100 mL. Filter this solution, dis-

card the first 20 mL of the filtrate, pipet the subsequent 5 mL, add methanol to make exactly 100 mL, and use this solution as the sample solution. Separately, weigh accurately about 0.01 g of dydrogesterone for assay, previously dried in a desiccator (in vacuum, phosphorus (V) oxide) for 24 hours, proceed in the same manner as the preparation of the sample solution, and use the solution as the standard solution. Determine the absorbances,  $A_T$  and  $A_S$ , of the sample solution and standard solution at 286 nm as directed under Ultraviolet-visible Spectrophotometry <2.24>.


$$\begin{aligned} &\text{Amount (mg) of dydrogesterone } (C_{21}H_{28}O_2) \\ &= W_S \times (A_T/A_S) \end{aligned}$$

$W_S$ : Amount (mg) of dydrogesterone for assay

**Containers and storage** Containers—Tight containers.

## Ecothiopate Iodide

エコチオパートヨウ化物



$C_9H_{23}INO_3PS$ : 383.23

2-(Diethoxyphosphorylsulfanyl)-*N,N,N*-trimethylethylaminium iodide [513-10-0]

Ecothiopate Iodide contains not less than 95.0% of  $C_9H_{23}INO_3PS$ , calculated on the dried basis.

**Description** Ecothiopate Iodide occurs as white crystals or crystalline powder.

It is very soluble in water, freely soluble in methanol, slightly soluble in ethanol (95), and practically insoluble in diethyl ether.

**Identification (1)** Dissolve 0.1 g of Ecothiopate Iodide in 2 mL of water, and add 1 mL of nitric acid: a brown precipitate is formed. To 1 drop of the turbid solution containing this precipitate add 1 mL of hexane, and shake: a light red color develops in the hexane layer.

**(2)** Heat the suspension of the precipitate obtained in (1) until it becomes colorless, cool, add 10 mL of water, and use this solution as the sample solution. Two mL of the sample solution responds to the Qualitative Tests <1.09> (2) for phosphate.

**(3)** Two mL of the sample solution obtained in (2) responds to the Qualitative Tests <1.09> for sulfate.

**pH <2.54>** Dissolve 0.1 g of Ecothiopate Iodide in 40 mL of water: the pH of this solution is between 3.0 and 5.0.

**Melting point <2.60>** 116 – 122°C

**Purity (1)** Clarity and color of solution—Dissolve 0.5 g of Ecothiopate Iodide in 5 mL of water: the solution is clear and colorless.

**(2)** Heavy metals <1.07>—To 1.0 g of Ecothiopate Iodide in a Kjeldahl flask add 5 mL of nitric acid and 2 mL of sulfuric acid, put a small funnel on the mouth of the flask, and heat carefully until white fumes are evolved. After cooling, add 2 mL of nitric acid, and heat. Repeat this procedure twice, add several 2-mL portions of hydrogen peroxide (30), and heat

until the solution becomes colorless, and white fumes are evolved. After cooling, transfer the solution together with a small quantity of water to a Nessler tube, and add water to make about 20 mL. Adjust the solution with ammonia solution (28) and ammonia TS to a pH between 3.0 and 3.5, add water to make 50 mL, and use this solution as the test solution. Prepare the control solution as follows: proceed in the same manner as the preparation of the test solution, and add 2.0 mL of Standard Lead Solution and water to make 50 mL (not more than 20 ppm).

(3) Related substances—Dissolve 0.20 g of Ecothiopate Iodide in 10 mL of methanol, and use this solution as the sample solution. Pipet 3 mL of the sample solution, add methanol to make exactly 200 mL, and use this solution as the standard solution. Perform the test with these solutions as directed under Thin-layer Chromatography <2.03>. Spot 10  $\mu$ L each of the sample solution and standard solution on a plate of cellulose for thin-layer chromatography. Develop the plate with a mixture of 1-butanol, water and acetic acid (100) (4:2:1) to a distance of about 10 cm, and air-dry the plate. Spray evenly Dragendorff's TS for spraying on the plate: the spots other than the principal spot from the sample solution are not more intense than the spot from the standard solution.

**Loss on drying** <2.41> Not more than 1.0% (1 g, in vacuum, phosphorus (V) oxide, 50°C, 3 hours).

**Assay** Weigh accurately about 0.125 g of Ecothiopate Iodide, and dissolve in water to make exactly 100 mL. Pipet 10 mL of this solution, add 30 mL of water, then add exactly 10 mL of phosphate buffer solution, pH 12, stopper the container, and allow to stand at  $25 \pm 3^\circ\text{C}$  for 20 minutes. To this solution add quickly 2 mL of acetic acid (100), and titrate <2.50> with 0.002 mol/L iodine VS (potentiometric titration). Perform the test in the same manner without phosphate buffer solution, pH 12, and make any necessary correction.


Each mL of 0.002 mol/L iodine VS  
= 1.533 mg of  $\text{C}_9\text{H}_{23}\text{INO}_3\text{PS}$

**Containers and storage** Containers—Tight containers.

Storage—Light-resistant, and not exceeding 0°C.

## Edrophonium Chloride

エドロホニウム塩化物



$\text{C}_{10}\text{H}_{16}\text{ClNO}$ : 201.69

*N*-Ethyl-3-hydroxy-*N,N*-dimethylanilinium chloride  
[116-38-1]

Edrophonium Chloride, when dried, contains not less than 98.0% of  $\text{C}_{10}\text{H}_{16}\text{ClNO}$ .

**Description** Edrophonium Chloride occurs as white crystals or crystalline powder. It is odorless.

It is very soluble in water, freely soluble in ethanol (95) and

in acetic acid (100), and practically insoluble in acetic anhydride and in diethyl ether.

It is hygroscopic.

It is gradually colored by light.

**Identification** (1) To 5 mL of a solution of Edrophonium Chloride (1 in 100) add 1 drop of iron (III) chloride TS: a light red-purple color develops.

(2) Determine the absorption spectrum of a solution of Edrophonium Chloride in 0.1 mol/L hydrochloric acid TS (1 in 20,000) as directed under Ultraviolet-visible Spectrophotometry <2.24>, and compare the spectrum with the Reference Spectrum or the spectrum of a solution of Edrophonium Chloride Reference Standard prepared in the same manner as the sample solution: both spectra exhibit similar intensities of absorption at the same wavelengths.

(3) A solution of Edrophonium Chloride (1 in 50) responds to the Qualitative Tests <1.09> for chloride.

**pH** <2.54> Dissolve 1.0 g of Edrophonium Chloride in 10 mL of water: the pH of this solution is between 3.5 and 5.0.

**Melting point** <2.60> 166 – 171°C (with decomposition).

**Purity** (1) Clarity and color of solution—Dissolve 1.0 g of Edrophonium Chloride in 10 mL of water: the solution is clear and colorless.

(2) Heavy metals <1.07>—Proceed with 1.0 g of Edrophonium Chloride according to Method 1, and perform the test. Prepare the control solution with 2.0 mL of Standard Lead Solution (not more than 20 ppm).

(3) Arsenic <1.11>—Prepare the test solution with 1.0 g of Edrophonium Chloride according to Method 1, and perform the test (not more than 2 ppm).

(4) Related substances—Dissolve 0.50 g of Edrophonium Chloride in 10 mL of ethanol (95), and use this solution as the sample solution. Pipet 1 mL of the sample solution, and add ethanol (95) to make exactly 100 mL. Pipet 3 mL of this solution, add ethanol (95) to make exactly 10 mL, and use this solution as the standard solution. Perform the test with these solutions as directed under Thin-layer Chromatography <2.03>. Spot 10  $\mu$ L each of the sample solution and standard solution on a plate of silica gel with fluorescent indicator for thin-layer chromatography. Develop the plate with a mixture of methanol, chloroform and ammonia solution (28) (16:4:1) to a distance of about 10 cm, and air-dry the plate. Examine under ultraviolet light (main wavelength: 254 nm): the spots other than the principal spot from the sample solution are not more intense than the spot from the standard solution.

**Loss on drying** <2.41> Not more than 0.20% (1 g, in vacuum, phosphorus (V) oxide, 3 hours).

**Residue on ignition** <2.44> Not more than 0.1% (1 g).

**Assay** Weigh accurately about 0.2 g of Edrophonium Chloride, previously dried, and dissolve in 100 mL of a mixture of acetic anhydride and acetic acid (100) (7:3). Titrate <2.50> with 0.1 mol/L perchloric acid VS (potentiometric titration). Perform a blank determination, and make any necessary correction.

Each mL of 0.1 mol/L perchloric acid VS  
= 20.17 mg of  $\text{C}_{10}\text{H}_{16}\text{ClNO}$

**Containers and storage** Containers—Tight containers.

Storage—Light-resistant.

## Edrophonium Chloride Injection

エドロホニウム塩化物注射液

Edrophonium Chloride Injection is an aqueous solution for injection. It contains not less than 95% and not more than 105% of the labeled amount of edrophonium chloride ( $C_{10}H_{16}ClNO$ ; 201.69).

**Method of preparation** Prepare as directed under Injections, with Edrophonium Chloride.

**Description** Edrophonium Chloride Injection is a clear and colorless liquid.

**Identification (1)** To a volume of Edrophonium Chloride Injection, equivalent to 0.04 g of Edrophonium Chloride according to the labeled amount, add 4 mL of barium nitrate TS, shake, and filter. Proceed with the filtrate as directed in the Identification (1) under Edrophonium Chloride.

**(2)** Determine the absorption spectrum of the sample solution obtained in the Assay as directed under Ultraviolet-visible Spectrophotometry <2.24>: it exhibits a maximum between 272 nm and 276 nm.

**pH** <2.54> 6.5 – 8.0

**Extractable volume** <6.05> It meets the requirement.

**Assay** Conduct this procedure without exposure to daylight, using light-resistant containers. Measure exactly a volume of Edrophonium Chloride Injection, equivalent to about 50 mg of edrophonium chloride ( $C_{10}H_{16}ClNO$ ), place in a chromatographic column prepared by pouring 10 mL of weakly basic DEAE-bridged dextran anion exchanger (Cl type) (50 to 150  $\mu$ m in particle diameter) into a chromatographic tube about 2 cm in inside diameter and about 10 cm in length, add 25 mL of water, and elute at the flow rate of 1 to 2 mL per minute. Wash the column with two 25-mL portions of water at the flow rate of 1 to 2 mL per minute. Combine the washings with above effluent solutions, and add water to make exactly 100 mL. Measure exactly 10 mL of this solution, and add 10 mL of phosphate buffer solution, pH 8.0, and 5 g of sodium chloride. Wash this solution with four 20-mL portions of a mixture of diethyl ether and hexane (1:1), collect the water layer, add 0.1 mol/L hydrochloric acid TS to make exactly 100 mL, and use this solution as the sample solution. Separately, weigh accurately about 50 mg of Edrophonium Chloride Reference Standard, previously dried in a desiccator (in vacuum, phosphorus (V) oxide) for 3 hours, and dissolve in water to make exactly 100 mL. Measure exactly 10 mL of this solution, and prepare the standard solution in the same manner as the sample solution. Determine the absorbances,  $A_T$  and  $A_S$ , of the sample solution and standard solution at 273 nm as directed under Ultraviolet-visible Spectrophotometry <2.24>.

$$\begin{aligned} \text{Amount (mg) of edrophonium chloride (C}_{10}\text{H}_{16}\text{ClNO)} \\ = W_S \times (A_T/A_S) \end{aligned}$$

$W_S$ : Amount (mg) of Edrophonium Chloride Reference Standard


**Containers and storage** Containers—Hermetic containers,

and colored containers may be used.

Storage—Light-resistant.

## Elcatonin

エルカトニン



$C_{148}H_{244}N_{42}O_{47}$ ; 3363.77  
[60731-46-6]

Elcatonin contains not less than 5000 Elcatonin Units and not more than 7000 Elcatonin Units per 1 mg of peptide, calculated on the dehydrated and de-acetic acid basis.

**Description** Elcatonin is a white powder.

It is very soluble in water, freely soluble in ethanol (95), and practically insoluble in acetonitrile.

It is hygroscopic.

The pH of its solution (1 in 500) is between 4.5 and 7.0.

**Identification** Dissolve 5 mg of Elcatonin in 5 mL of water. Determine the absorption spectrum of the solution as directed under Ultraviolet-visible Spectrophotometry <2.24>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wavelengths.

**Constituent amino acids** Put about 1 mg of Elcatonin in a test tube for hydrolysis, add phenol-hydrochloric acid TS to dissolve, replace the air inside with Nitrogen, seal the tube under reduced pressure, and heat at  $110 \pm 2^\circ\text{C}$  for 24 hours. After cooling, open the tube, evaporate the hydrolyzate to dryness under reduced pressure, dissolve the residue in 1 mL of 0.02 mol/L hydrochloric acid TS, and use this solution as the sample solution. Separately, weigh exactly 1.33 mg of L-aspartic acid, 1.19 mg of L-threonine, 1.05 mg of L-serine, 1.47 mg of L-glutamic acid, 1.15 mg of L-proline, 0.75 mg of glycine, 0.89 mg of L-alanine, 1.17 mg of L-valine, 1.89 mg of L-2-aminosuberic acid, 1.31 mg of L-leucine, 1.81 mg of L-tyrosine, 1.83 mg of L-lysine hydrochloride, 2.10 mg of L-histidine hydrochloride monohydrate and 2.11 mg of L-arginine hydrochloride, dissolve them in 0.02 mol/L hydrochloric acid TS to make exactly 50 mL, and use this solution as the standard solution. Perform the test with exactly 10  $\mu$ L each of the sample solution and standard solution as directed under Liquid Chromatography <2.01> according to the following conditions: 14 peaks of amino acids appear on the chromatogram obtained from the sample solution, and their respective molar ratios against alanine are 1.7 – 2.2 for aspartic acid, 3.5 – 4.2 for threonine, 2.4 – 3.0 for serine, 2.7 – 3.2 for glutamic acid, 1.7 – 2.2 for proline, 2.7 – 3.2 for glycine, 1.6 – 2.2 for valine, 0.8 – 1.2 for 2-aminosuberic acid, 4.5 – 5.2 for leucine, 0.7 – 1.2 for tyrosine, 1.7 – 2.2 for lysine, 0.8 – 1.2 for histidine and 0.7 – 1.2 for arginine.

**Operating conditions—**

Detector: A visible spectrophotometer (wavelength: 440

nm and 570 nm).

Column: A stainless steel column about 4 mm in inside diameter and about 8 cm in length, packed with strongly acidic ion-exchange resin for liquid chromatography composed with a sulfonated styrene-divinylbenzene copolymer (3  $\mu$ m in particle diameter).

Column temperature: Varied between 50°C and 65°C.

Chemical reaction vessel temperature: A constant temperature of about 130°C.

Color developing time: About 1 minute.

Mobile phase: Buffer solutions A, B, C and D, with sodium ion concentrations of 0.10 mol/L, 0.135 mol/L, 1.26 mol/L and 0.20 mol/L, respectively. The ion concentration of the mobile phase is changed stepwise from 0.10 mol/L to 1.26 mol/L by using these buffer solutions.

| Buffer solution: | Components of buffer solutions |                     |                     |                     |
|------------------|--------------------------------|---------------------|---------------------|---------------------|
|                  | A                              | B                   | C                   | D                   |
| Citric acid      | 8.85 g                         | 7.72 g              | 6.10 g              | —                   |
| Sodium citrate   | 3.87 g                         | 10.05 g             | 26.67 g             | —                   |
| Sodium hydroxide | —                              | —                   | 2.50 g              | 8.00 g              |
| Sodium chloride  | 3.54 g                         | 1.87 g              | 54.35 g             | —                   |
| Ethanol          | 60.0 mL                        | —                   | —                   | 60.0 mL             |
| Thiodiglycol     | 5.0 mL                         | 5.0 mL              | —                   | —                   |
| Purified water   | a sufficient amount            | a sufficient amount | a sufficient amount | a sufficient amount |
| Total amount     | 1000 mL                        | 1000 mL             | 1000 mL             | 1000 mL             |

Reaction reagent: Mix 407 g of lithium acetate dihydrate, 245 mL of acetic acid (100) and 801 mL of 1-methoxy-2-propanol, add water to make 2000 mL, stir for about 20 minutes while passing Nitrogen, and use this solution as solution A. Separately, to 1957 mL of 1-methoxy-2-propanol add 77 g of ninhydrin and 0.134 g of sodium borohydride, stir for about 20 minutes while passing Nitrogen, and use this solution as solution B. Mix solution A and solution B before use.

Flow rate of mobile phase: Adjust the flow rate so that the retention time of arginine is about 75 minutes.

Flow rate of reaction reagent: About 0.2 mL per minute.

Selection of column: Proceed with 10  $\mu$ L of the standard solution under the above operating conditions. Use a column from which aspartic acid, threonine, serine, glutamic acid, proline, glycine, alanine, valine, 2-aminosuberic acid, leucine, tyrosine, lysine, histidine and arginine are eluted in this order, with complete separation of each peak.

**Purity (1)** Acetic acid—Weigh accurately 3–6 mg of Elcatonin quickly under conditions of  $25 \pm 2^\circ\text{C}$  and  $50 \pm 5\%$  relative humidity, add exactly 1 mL of the internal standard solution to dissolve it, and use this solution as the sample solution. Separately, weigh accurately about 0.5 g of acetic acid (100), and add the internal standard solution to make exactly 100 mL. Pipet 5 mL of this solution, add the internal standard solution to make exactly 100 mL, and use this solution as the standard solution. Perform the test with 20  $\mu$ L each of the sample solution and standard solution as directed under Liquid Chromatography <2.01> according to the following conditions, and calculate the ratios,  $Q_T$  and  $Q_S$ , of the peak area of acetic acid to that of the internal standard: the amount of acetic acid is not more than 7.0%.

Amount (%) of acetic acid ( $\text{CH}_3\text{COOH}$ )

$$= (W_{\text{ST}}/W_{\text{SA}}) \times (Q_T/Q_S) \times 50$$

$W_{\text{ST}}$ : Amount (g) of acetic acid (100)

$W_{\text{SA}}$ : Amount (mg) of sample

**Internal standard solution**—A solution of citric acid (1 in 4000).

**Operating conditions**—

Detector: An ultraviolet absorption photometer (wavelength: 210 nm).

Column: A stainless steel column about 4 mm in inside diameter and about 15 cm in length, packed with octadecylsilanized silica gel for liquid chromatography (5  $\mu$ m in particle diameter).

Column temperature: A constant temperature of about 25°C.

Mobile phase: Dissolve 13.2 g of diammonium hydrogenphosphate in 900 mL of water, add phosphoric acid to adjust the pH to 2.5, and add water to make 1000 mL.

Flow rate: Adjust the flow rate so that the retention time of acetic acid is about 4 minutes.

Selection of column: Proceed with 20  $\mu$ L of the standard solution under the above operating conditions, and calculate the resolution. Use a column from which acetic acid and citric acid are eluted in this order with the resolution between their peaks being not less than 2.0.

**(2) Related substances**—Dissolve 1.0 mg of Elcatonin in 1 mL of a mixture of trifluoroacetic acid TS and acetonitrile (2:1), and use this solution as the sample solution. Take exactly 0.3 mL of the sample solution, add a mixture of trifluoroacetic acid TS and acetonitrile (2:1) to make exactly 10 mL, and use this solution as the standard solution. Perform the test with exactly 10  $\mu$ L each of the sample solution and standard solution as directed under Liquid Chromatography <2.01> according to the following conditions, and determine each peak area by the automatic integration method: the total of the peak areas other than the peak of elcatonin of the sample solution is not larger than the peak area of elcatonin of the standard solution, and each peak area other than the peak of elcatonin of the sample solution is not larger than 1/3 of the peak area of elcatonin of the standard solution.

**Operating conditions**—

Detector: An ultraviolet absorption photometer (wavelength: 225 nm).

Column: A stainless steel column about 4 mm in inside diameter and about 15 cm in length, packed with octadecylsilanized silica gel for liquid chromatography (5  $\mu$ m in particle diameter).

Column temperature: A constant temperature of about 40°C.

Mobile phase: A mixture of trifluoroacetic acid TS and acetonitrile (change the ratio linearly from 85:15 to 55:45 in 30 minutes).

Flow rate: Adjust the flow rate so that the retention time of elcatonin is about 25 minutes.

Selection of column: Dissolve 2 mg of Elcatonin in 200  $\mu$ L of trypsin TS for test of elcatonin, warm at 37°C for 1 hour, then add 1 drop of acetic acid (100), and heat at 95°C for 1 minute. To 10  $\mu$ L of this solution add 50  $\mu$ L of the sample solution, and mix. Proceed with 10  $\mu$ L of this solution under the above operating conditions, and calculate the resolution. Use a column such that the resolution between the peak of elcatonin and the peak which appears immediately before the

peak of elcatonin is not less than 2.0, and the retention time of elcatonin is about 25 minutes.

**Detection sensitivity:** Adjust the detection sensitivity so that the peak height of elcatonin from 10  $\mu$ L of the standard solution is between 50 mm and 200 mm.

**Time span of measurement:** Continue measurement until the regularly changing base-line of the chromatogram disappears, beginning after the solvent peak.

**Water <2.48>** Weigh accurately 1 – 3 mg of Elcatonin quickly under conditions of  $25 \pm 2^\circ\text{C}$  and  $50 \pm 5\%$  relative humidity, and perform the test as directed in 2. Coulometric titration: not more than 8.0%.

**Nitrogen content** Weigh accurately 0.015 – 0.02 g of Elcatonin quickly under conditions of  $25 \pm 2^\circ\text{C}$  and  $50 \pm 5\%$  relative humidity, and perform the test as directed under Nitrogen Determination <1.08>: it contains not less than 16.1% and not more than 18.7% of nitrogen (N: 14.01) in the peptide, calculated on the dehydrated and de-acetic acid basis.

**Assay** (i) Animals: Select healthy male Sprague-Dawley rats each weighing between 90 g and 110 g. Keep the rats for not less than 3 days before use, providing an appropriate uniform diet and water.

(ii) Diluent for elcatonin: Dissolve 2.72 g of sodium acetate in water to make 200 mL, add 0.2 g of bovine serum albumin, and adjust the pH to 6.0 with acetic acid (100). Prepare before use.

(iii) Standard solution: Dissolve Elcatonin Reference Standard in the diluent for elcatonin to make two standard solutions, one to contain exactly 0.075 Unit in each mL which is designated as the high-dose standard solution,  $S_H$ , and the other to contain exactly 0.0375 Unit in each mL which is designated as the low-dose standard solution,  $S_L$ .

(iv) Sample solution: Weigh accurately 0.5 – 2.0 mg of Elcatonin quickly under conditions of  $25 \pm 2^\circ\text{C}$  and  $50 \pm 5\%$  relative humidity, and dissolve in the diluent for elcatonin to make two sample solutions, the high-dose sample solution,  $T_H$ , which contains the Units per mL equivalent to  $S_H$  and the low-dose sample solution,  $T_L$ , which contains the Units per mL equivalent to  $S_L$ .

(v) Deproteinizing solution for elcatonin: Dissolve 160 g of trichloroacetic acid and 30.6 g of strontium chloride in water to make 3600 mL.

(vi) Procedure: Divide the animals into 4 equal groups of not less than 10 animals each. Withhold all food, but not water, for 18 to 24 hours before the injections, and withhold water during the assay until the final blood sample is taken. Handle the animals with care in order to avoid undue excitement.

Inject exactly 0.2 mL each of the standard solutions and the sample solutions into the tail vein of each animal as indicated in the following design:

|              |       |              |       |
|--------------|-------|--------------|-------|
| First group  | $S_H$ | Third group  | $T_H$ |
| Second group | $S_L$ | Fourth group | $T_L$ |

At 1 hour after the injection, take a sufficient blood sample to perform the test from the carotid artery and vein of each animal under ether anesthesia, centrifuge the blood samples to separate serum, and determine the serum calcium according to the following (vii).

(vii) Serum calcium determination: Take exactly 0.3 mL of the serum, add the deproteinizing solution for elcatonin to

make exactly 3 mL, mix well, centrifuge, and use the supernatant liquid as the sample solution for calcium determination. Separately, pipet 1 mL of Standard Calcium Solution for Atomic Absorption Spectrophotometry <2.23>, and add a solution of sodium chloride (17 in 2000) to make exactly 10 mL. Pipet 5 mL of this solution, add the deproteinizing solution for elcatonin to make exactly 50 mL, and use this solution as the standard solution for calcium determination. Determine the absorbances,  $A_T$  and  $A_S$ , of the sample solution and the standard solution as directed under Atomic Absorption Spectrophotometry <2.23> according to the following conditions. Determine the absorbance,  $A_0$ , of a solution obtained in the same manner used for preparation of the standard solution, but with 1 mL of water instead of the standard solution.

$$\begin{aligned} \text{Amount (mg) of calcium (Ca) in 100 mL of the serum} \\ = 0.01 \times \{(A_T - A_0)/(A_S - A_0)\} \times 10 \times 100 \end{aligned}$$

Gas: Combustible gas—Acetylene

Supporting gas—Air

Lamp: Calcium hollow-cathode lamp

Wavelength: 422.7 nm

(viii) Calculation: Amounts of calcium in 100 mL of the serum obtained with  $S_H$ ,  $S_L$ ,  $T_H$  and  $T_L$  in (vii) are symbolized as  $y_1$ ,  $y_2$ ,  $y_3$  and  $y_4$ , respectively. Sum up individual  $y_1$ ,  $y_2$ ,  $y_3$  and  $y_4$  to obtain  $Y_1$ ,  $Y_2$ ,  $Y_3$  and  $Y_4$ , respectively.

Units per mg of peptide, calculated on the dehydrated and de-acetic acid basis

$$= \text{antilog } M \times (\text{units per mL of } S_H \times (b/a))$$

$$M = 0.3010 \times (Y_a/Y_b)$$

$$Y_a = -Y_1 - Y_2 + Y_3 + Y_4$$

$$Y_b = Y_1 - Y_2 + Y_3 - Y_4$$

$a$ : Amount (mg) of the sample

$$\times \{100 - [\text{water content (\%)} + \text{acetic acid content (\%)}]/100\}$$

$b$ : Total volume (mL) of the high-dose sample solution prepared by dissolving the sample with diluent for elcatonin.

$F'$  computed by the following equation should be smaller than  $F$  shown in the table against  $n$  with which  $s^2$  is calculated. Calculate  $L$  ( $P=0.95$ ) by use of the following equation:  $L$  should be not more than 0.20. If  $F'$  exceeds  $F$ , or if  $L$  exceeds 0.20, repeat the test, increasing the number of animals or arranging the assay conditions so that  $F'$  is not more than  $F$  and  $L$  is not more than 0.20.

$$F' = (-Y_1 + Y_2 + Y_3 - Y_4)^2 / 4fs^2$$

$f$ : Number of the animals of each group.

$$s^2 = \{\Sigma y^2 - (Y/f)\} / n$$

$\Sigma y^2$ : The sum of squares of  $y_1$ ,  $y_2$ ,  $y_3$  and  $y_4$  in each group.

$$Y = Y_1^2 + Y_2^2 + Y_3^2 + Y_4^2$$

$$n = 4(f - 1)$$

$$L = 2\sqrt{(C - 1)(CM^2 + 0.09062)}$$

$$C = Y_b^2 / (Y_b^2 - 4fs^2t^2)$$


$t^2$ : Value shown in the following table against  $n$  used to calculate  $s^2$ .

| <i>n</i> | <i>t</i> <sup>2</sup> = <i>F</i> | <i>n</i> | <i>t</i> <sup>2</sup> = <i>F</i> | <i>n</i> | <i>t</i> <sup>2</sup> = <i>F</i> |
|----------|----------------------------------|----------|----------------------------------|----------|----------------------------------|
| 1        | 161.45                           | 13       | 4.667                            | 25       | 4.242                            |
| 2        | 18.51                            | 14       | 4.600                            | 26       | 4.225                            |
| 3        | 10.129                           | 15       | 4.543                            | 27       | 4.210                            |
| 4        | 7.709                            | 16       | 4.494                            | 28       | 4.196                            |
| 5        | 6.608                            | 17       | 4.451                            | 29       | 4.183                            |
| 6        | 5.987                            | 18       | 4.414                            | 30       | 4.171                            |
| 7        | 5.591                            | 19       | 4.381                            | 40       | 4.085                            |
| 8        | 5.318                            | 20       | 4.351                            | 60       | 4.001                            |
| 9        | 5.117                            | 21       | 4.325                            | 120      | 3.920                            |
| 10       | 4.965                            | 22       | 4.301                            | ∞        | 3.841                            |
| 11       | 4.844                            | 23       | 4.279                            |          |                                  |
| 12       | 4.747                            | 24       | 4.260                            |          |                                  |

**Containers and storage** Containers—Tight containers.  
Storage—Not exceeding 8°C.

## Enflurane

エンフルラン



C<sub>3</sub>H<sub>2</sub>ClF<sub>5</sub>O: 184.49  
(2*RS*)-2-Chloro-1-(difluoromethoxy)-1,1,2-trifluoroethane  
[13838-16-9]

**Description** Enflurane is a clear, colorless liquid.

It is slightly soluble in water.

It is miscible with ethanol (95) and with diethyl ether.

It is a volatile, and not an inflammable.

It shows no optical rotation.

Boiling point: 54 – 57°C

**Identification** (1) Take 50 μL of Enflurane, and prepare the test solution as directed to the Oxygen Flask Combustion Method <1.06> using 40 mL of water as the absorbing liquid. The test solution responds to the Qualitative Tests <1.09> for chloride and fluoride.

(2) Determine the infrared absorption spectrum of Enflurane as directed in the liquid film method under Infrared Spectrophotometry <2.25>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wave numbers.

**Refractive index** <2.45> *n*<sub>D</sub><sup>20</sup>: 1.302 – 1.304

**Specific gravity** <2.56> *d*<sub>20</sub><sup>20</sup>: 1.520 – 1.540

**Purity** (1) Acidity or alkalinity—To 60 mL of Enflurane add 60 mL of freshly boiled and cooled water, shake for 3 minutes, separate the water layer, and use the layer as the sample solution. To 20 mL of the sample solution add one drop of bromocresol purple TS and 0.10 mL of 0.01 mol/L sodium hydroxide TS: the color of the solution is purple. To 20 mL of the sample solution add one drop of bromocresol purple TS and 0.06 mL of 0.01 mol/L hydrochloric acid TS: the color of the solution is yellow.

(2) Chloride <1.03>—To 20 g of Enflurane add 20 mL of water, shake well, and separate the water layer. Take 10 mL

of the water layer add 6 mL of dilute nitric acid and water to make 50 mL, and perform the test using this solution as the test solution. Prepare the control solution with 0.30 mL of 0.01 mol/L hydrochloric acid (not more than 0.001%).

(3) Nonvolatile residue—Evaporate exactly 65 mL of Enflurane on a water bath to dryness, and dry the residue at 105°C for 1 hour: the of the residue is not more than 1.0 mg.

(4) Related substances—Proceed the test with 5 μL of Enflurane as directed under Gas chromatography <2.02> according to the following conditions. Determine each peak area other than the peak of air which appears soon after injection of the sample by the automatic integration method, and calculate the amount of each peak by the area percentage method: the amount of the substances other than enflurane is not more than 0.10%.

**Operating conditions—**

Detector: A thermal conductivity detector.

Column: A column 3 mm in inside diameter and 3 m in length, packed with siliceous earth for gas chromatography, 180 to 250 μm in particle diameter, coated with diethylene glycol succinate ester for gas chromatography in the ratio of 20%.

Column temperature: A constant temperature of about 80°C.

Carrier gas: Helium

Flow rate: Adjust the flow rate so that the retention time of enflurane is about 3 minutes.

Time span of measurement: About three times as long as the retention time of enflurane.

**System suitability—**

Test for required detection: Pipet exactly 1 mL of enflurane add 2-propanol to make exactly 100 mL. To exactly 2 mL of this solution add 2-propanol to make exactly 10 mL, and use this solution as the solution for system suitability test. Pipet 1 mL of the solution, and add 2-propanol to make exactly 10 mL. Confirm that the peak area of enflurane obtained from 5 μL of this solution is equivalent to 7 to 13% of that of enflurane obtained from 5 μL of the solution for system suitability test.

System performance: Mix 5 mL of Enflurane and 5 mL of 2-propanol. When the procedure is run with 5 μL of this mixture under the above operating conditions, enflurane and 2-propanol are eluted in this order with the resolution between these peaks being not less than 2.0.

System repeatability: When the test is repeated 6 times with 5 μL of the solution for system suitability test under the above operating conditions, the relative standard deviation of the peak area of enflurane is not more than 2.0%.


**Water** <2.48> Not more than 0.10% (10 g, direct titration).

**Containers and storage** Containers—Tight containers.

Storage—Not exceeding 30°C.

## Enoxacin Hydrate

エノキサシン水和物



$C_{15}H_{17}FN_4O_3 \cdot 1\frac{1}{2}H_2O$ : 347.34

1-Ethyl-6-fluoro-4-oxo-7-(piperazin-1-yl)-1,4-dihydro-1,8-naphthyridine-3-carboxylic acid sesquihydrate  
[84294-96-2]

Enoxacin Hydrate, when dried, contains not less than 98.5% of enoxacin ( $C_{15}H_{17}FN_4O_3$ : 320.32).

**Description** Enoxacin Hydrate occurs as white to pale yellow-brown crystals or crystalline powder.

It is freely soluble in acetic acid (100), slightly soluble in methanol, very slightly soluble in chloroform, and practically insoluble in water, in ethanol (95) and in diethyl ether.

It dissolves in dilute sodium hydroxide TS.

It is gradually colored by light.

**Identification (1)** Place 0.02 g of Enoxacin Hydrate and 0.05 g of sodium in a test tube, and heat gradually to ignition with precaution. After cooling, add 0.5 mL of methanol and then 5 mL of water, and heat to boiling. To this solution add 2 mL of dilute acetic acid, and filter: the filtrate responds to the Qualitative Tests <1.09> (2) for fluoride.

(2) Dissolve 0.05 g of Enoxacin Hydrate in dilute sodium hydroxide TS to make 100 mL. To 1 mL of the solution add water to make 100 mL. Determine the absorption spectrum of the solution as directed under Ultraviolet-visible Spectrophotometry <2.24>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wavelengths.

(3) Determine the infrared absorption spectrum of Enoxacin Hydrate as directed in the potassium bromide disk method under Infrared Spectrophotometry <2.25>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wave numbers.

**Melting point** <2.60> 225 – 229°C (after drying).

**Purity (1)** Sulfate <1.14>—Dissolve 1.0 g of Enoxacin Hydrate in 50 mL of dilute sodium hydroxide TS, shake with 10 mL of dilute hydrochloric acid, and centrifuge. Filter the supernatant liquid, and to 30 mL of the filtrate add water to make 50 mL. Perform the test using this solution as the test solution. Prepare the control solution as follows: to 0.50 mL of 0.005 mol/L sulfuric acid VS add 25 mL of dilute sodium hydroxide TS, 5 mL of dilute hydrochloric acid TS and water to make 50 mL (not more than 0.048%).

(2) Heavy metals <1.07>—Proceed with 1.0 g of Enoxacin Hydrate according to Method 2, and perform the test. Prepare the control solution with 2.0 mL of Standard Lead Solution (not more than 20 ppm).

(3) Arsenic <1.11>—Prepare the test solution with 1.0 g of Enoxacin Hydrate according to Method 3, and perform

the test (not more than 2 ppm).

(4) Related substances—Dissolve 50 mg of Enoxacin Hydrate in 25 mL of a mixture of chloroform and methanol (7:3), and use this solution as the sample solution. Pipet 1 mL of the sample solution, add a mixture of chloroform and methanol (7:3) to make exactly 200 mL, and use this solution as the standard solution. Perform the test with these solutions as directed under Thin-layer Chromatography <2.03>. Spot 5  $\mu$ L each of the sample solution and standard solution on a plate of silica gel with fluorescent indicator for thin-layer chromatography. Develop the plate with a mixture of 1-butanol, water and acetic acid (100) (3:1:1) to a distance of about 10 cm, and air-dry the plate. Examine under ultraviolet light (main wavelength: 254 nm): the spots other than the principal spot from the sample solution are not more intense than the spot from the standard solution.

**Loss on drying** <2.41> 7.0 – 9.0% (1 g, 105°C, 3 hours).

**Residue on ignition** <2.44> Not more than 0.1% (1 g, platinum crucible).


**Assay** Weigh accurately about 0.3 g of Enoxacin Hydrate, previously dried, dissolve in 30 mL of acetic acid (100), and titrate <2.50> with 0.1 mol/L perchloric acid VS (potentiometric titration). Perform a blank determination, and make any necessary correction.

Each mL of 0.1 mol/L perchloric acid VS  
= 32.03 mg of  $C_{15}H_{17}FN_4O_3$

**Containers and storage** Containers—Tight containers.  
Storage—Light-resistant.

## Enviomycin Sulfate

エンビオマイシン硫酸塩



Tuberactinomycin N Sulfate

$C_{25}H_{43}N_{13}O_{10} \cdot 1\frac{1}{2}H_2SO_4$ : 832.81

Tuberactinomycin O Sulfate

$C_{25}H_{43}N_{13}O_9 \cdot 1\frac{1}{2}H_2SO_4$ : 816.81

Tuberactinomycin N Sulfate

(3*R*,4*R*)-*N*-[(3*S*,9*S*,12*S*,15*S*)-9,12-Bis(hydroxymethyl)-3-[(4*R*)-2-iminohexahydropyrimidin-4-yl]-2,5,8,11,14-penta-oxo-6-(*Z*)-ureidomethylene-1,4,7,10,13-pentaazacyclohexadec-15-yl]-3,6-diamino-4-hydroxyhexanamide sesquisulfate



[33103-22-9, Tuberactinomycin N]

Tuberactinomycin O Sulfate

(3*S*)-*N*-[(3*S*,9*S*,12*S*,15*S*)-9,12-Bis(hydroxymethyl)-3-[(4*R*)-2-imino-6-hydroxypyrimidin-4-yl]-2,5,8,11,14-penta-oxo-6-(*Z*)-ureidomethylene-1,4,7,10,13-pentaazacyclohexadec-15-yl]-3,6-diaminohexanamide sesquisulfate

[33137-73-4, Tuberactinomycin O]

Enviomycin Sulfate is the sulfate of a mixture of peptide substances having antibacterial activity produced by the growth of *Streptomyces griseoverticillatus* var. *tuberacticus*.

It contains not less than 770  $\mu\text{g}$  (potency) per mg, calculated on the dried basis. The potency of Enviomycin Sulfate is expressed as mass (potency) of tuberactinomycin N ( $\text{C}_{25}\text{H}_{43}\text{N}_{13}\text{O}_{10}$ : 685.69).

**Description** Enviomycin Sulfate occurs as a white powder.

It is very soluble in water, and practically insoluble in ethanol (99.5).

**Identification** (1) To 5 mL of a solution of Enviomycin Sulfate (1 in 200) add 1.5 mL of sodium hydroxide TS, and add 1 drop of a mixture of 0.01 mol/L citric acid TS and copper (II) sulfate TS (97:3): a blue-purple color develops.

(2) Determine the absorption spectrum of a solution of Enviomycin Sulfate (1 in 100,000) as directed under Ultraviolet-visible Spectrophotometry <2.24>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wavelengths.

(3) To 2 mL of a solution of Enviomycin Sulfate (1 in 20) add 1 drop of barium chloride TS: a white precipitate is produced.

**Optical rotation** <2.49>  $[\alpha]_{\text{D}}^{20}$ :  $-16$  to  $-22^\circ$  (0.5 g calculated on the dried basis, water, 50 mL, 100 mm).

**pH** <2.54> The pH of a solution obtained by dissolving 2.0 g of Enviomycin Sulfate in 20 mL of water is between 5.5 and 7.5.

**Content ratio of the active principle** Dissolve 0.1 g of Enviomycin Sulfate in water to make 100 mL, and use this solution as the sample solution. Perform the test with 3  $\mu\text{L}$  of the sample solution as directed under Liquid Chromatography <2.01> according to the following conditions, and determine the peak areas,  $A_{\text{T1}}$  and  $A_{\text{T2}}$ , of tuberactinomycin N and tuberactinomycin O, having the relative retention time,  $1.4 \pm 0.4$ , with respect to tuberactinomycin N, by the automatic integration method:  $A_{\text{T2}}/(A_{\text{T1}} + A_{\text{T2}})$  is between 0.090 and 0.150.

**Operating conditions**—

Detector: An ultraviolet absorption photometer (wavelength: 254 nm).

Column: A stainless steel column 4.6 mm in inside diameter and 15 cm in length, packed with silica gel for liquid chromatography (5  $\mu\text{m}$  in particle diameter).

Column temperature: A constant temperature of about 25°C.

Mobile phase: A mixture of ammonium acetate TS, 1,4-dioxane, tetrahydrofuran, water and ammonia solution (28) (100:75:50:23:2).

Flow rate: Adjust the flow rate so that the retention time of tuberactinomycin N is about 9 minutes.

**System suitability**—

**System performance:** When the procedure is run with 3  $\mu\text{L}$  of the sample solution under the above operating conditions, tuberactinomycin N and tuberactinomycin O are eluted in this order with the resolution between these peaks being not less than 1.5.

**System repeatability:** When the test is repeated 6 times with 3  $\mu\text{L}$  of the sample solution under the above operating conditions, the relative standard deviation of the peak area of tuberactinomycin N is not more than 2.0%.

**Purity** (1) Clarity and color of solution—Dissolve 1.0 g of Enviomycin Sulfate in 10 mL of water: the solution is clear and colorless.

(2) Heavy metals <1.07>—Proceed with 2.0 g of Enviomycin Sulfate according to Method 1, and perform the test. Prepare the control solution with 2.0 mL of Standard Lead Solution (not more than 10 ppm).

(3) Arsenic <1.11>—Prepare the test solution with 2.0 g of Enviomycin Sulfate according to Method 1, and perform the test (not more than 1 ppm).

**Loss on drying** <2.41> Not more than 4.0% (0.2 g, in vacuum, phosphorus (V) oxide, 60°C, 3 hours).

**Assay** Perform the test according to the Cylinder-plate method as directed under Microbial Assay for Antibiotics <4.02> according to the following conditions.

(i) Test organism—*Bacillus subtilis* ATCC 6633

(ii) Culture medium—Use the medium i in 1) Medium for test organism [5] under (1) Agar media for seed and base layer.


(iii) Standard solutions—Weigh accurately an amount of Enviomycin Sulfate Reference Standard, equivalent to about 20 mg (potency), dissolve in water to make exactly 20 mL, and use this solution as the standard stock solution. Keep the standard stock solution at a temperature not exceeding 5°C and use within 10 days. Take exactly a suitable amount of the standard stock solution before use, add 0.1 mol/L phosphate buffer solution, pH 8.0 to make solutions so that each mL contains 400  $\mu\text{g}$  (potency) and 100  $\mu\text{g}$  (potency), and use these solutions as the high concentration standard solution and low concentration standard solution, respectively.

(iv) Sample solutions—Weigh accurately an amount of Enviomycin Sulfate, equivalent to about 20 mg (potency), and dissolve in water to make exactly 20 mL. Take exactly a suitable amount of this solution, add 0.1 mol/L phosphate buffer solution, pH 8.0 to make solutions so that each mL contains 400  $\mu\text{g}$  (potency) and 100  $\mu\text{g}$  (potency), and use these solutions as the high concentration sample solution and low concentration sample solution, respectively.

**Containers and storage** Containers—Tight containers.

## Eperisone Hydrochloride

エペリゾン塩酸塩



$C_{17}H_{25}NO \cdot HCl$ : 295.85

(2*RS*)-1-(4-Ethylphenyl)-2-methyl-3-piperidin-1-ylpropan-1-one monohydrochloride [56839-43-1]

Eperisone Hydrochloride contains not less than 98.5% and not more than 101.0% of  $C_{17}H_{25}NO \cdot HCl$ , calculated on the anhydrous basis.

**Description** Eperisone Hydrochloride occurs as a white crystalline powder.

It is freely soluble in water, in methanol and in acetic acid (100), and soluble in ethanol (99.5).

Melting point: about 167°C (with decomposition).

A solution of Eperisone Hydrochloride in methanol (1 in 100) shows no optical rotation.

**Identification (1)** Determine the absorption spectrum of a solution of Eperisone Hydrochloride in methanol (1 in 100,000) as directed under Ultraviolet-visible Spectrophotometry <2.24>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wavelengths.

(2) Determine the infrared absorption spectrum of Eperisone Hydrochloride as directed in the potassium chloride disk method under Infrared Spectrophotometry <2.25>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wave numbers.

(3) A solution of Eperisone Hydrochloride (1 in 50) responds to the Qualitative Tests <1.09> for chloride.

**Purity (1)** Heavy metals <1.07>—Proceed with 1.0 g of Eperisone Hydrochloride according to Method 1, and perform the test. Prepare the control solution with 2.0 mL of Standard Lead Solution (not more than 20 ppm).

(2) Piperidine hydrochloride—Dissolve 1.0 g of Eperisone Hydrochloride in 20 mL of water, add 2.0 mL of diluted hydrochloric acid (1 in 2), 2.0 mL of a solution of copper (II) sulfate pentahydrate (1 in 20) and 1.5 mL of ammonia solution (28), and use this solution as the sample solution. Separately, to 2.0 mL of a solution of piperidine hydrochloride (1 in 1000) add 18 mL of water, 2.0 mL of diluted hydrochloric acid (1 in 2), 2.0 mL of a solution of copper (II) sulfate pentahydrate (1 in 20) and 1.5 mL of ammonia solution (28), and use this solution as the standard solution. To each of the sample solution and standard solution add 10 mL of a mixture of isopropylether and carbon disulfide (3:1), shake for 30 seconds, allow them to stand for 2 minutes, and compare the color of the upper layer: the color obtained from the sample solution is not more darker than that from the standard solution.

(3) Related substances—Dissolve 0.1 g of Eperisone

Hydrochloride in 100 mL of the mobile phase, and use this solution as the sample solution. Pipet 1 mL of the sample solution, add the mobile phase to make exactly 100 mL, and use this solution as the standard solution. Perform the test with exactly 10  $\mu$ L each of the sample solution and standard solution as directed under Liquid Chromatography <2.01> according to the following conditions, and determine each peak area by the automatic integration method: the total area of the peaks other than the peak of eperisone is not more than 1/5 of the peak area of eperisone from the standard solution.

**Operating conditions—**

**Detector:** An ultraviolet absorption photometer (wavelength: 254 nm).

**Column:** A stainless steel column 4.6 mm in inside diameter and 15 cm in length, packed with octadecylsilanized silica gel for liquid chromatography (5  $\mu$ m in particle diameter).

**Column temperature:** A constant temperature of about 30°C.

**Mobile phase:** A mixture of methanol, 0.0375 mol/L sodium 1-decanesulfonate TS and perchloric acid (600:400:1).

**Flow rate:** Adjust the flow rate so that the retention time of eperisone is about 17 minutes.

**Time span of measurement:** About 2 times as long as the retention time of eperisone.

**System suitability—**

**Test for required detectability:** Pipet 1 mL of the standard solution, and add the mobile phase to make exactly 10 mL. Confirm that the peak area of eperisone obtained from 10  $\mu$ L of this solution is equivalent to 7 to 13% of that obtained from 10  $\mu$ L of the standard solution.

**System performance:** When the procedure is run with 10  $\mu$ L of the standard solution under the above operating conditions, the number of theoretical plates and the symmetry factor of the peak of eperisone are not less than 4000 steps and not more than 2.0, respectively.

**System repeatability:** When the test is repeated 6 times with 10  $\mu$ L of the standard solution under the above operating conditions, the relative standard deviation of the peak area of eperisone is not more than 3.0%.

**Water <2.48>** Not more than 0.2% (0.1 g, coulometric titration).

**Residue on ignition <2.44>** Not more than 0.2% (1 g).


**Assay** Weigh accurately about 0.6 g of Eperisone Hydrochloride, dissolve in 20 mL of acetic acid (100), add 80 mL of acetic anhydride, and titrate <2.50> with 0.1 mol/L perchloric acid VS (potentiometric titration). Perform a blank determination in the same manner, and make any necessary correction.

Each mL of 0.1 mol/L perchloric acid VS  
= 29.58 mg of  $C_{17}H_{25}NO \cdot HCl$

**Containers and storage** Containers—Well-closed containers.

## Ephedrine Hydrochloride

エフェドリン塩酸塩



$C_{10}H_{15}NO \cdot HCl$ : 201.69

(1*R*,2*S*)-2-Methylamino-1-phenylpropan-1-ol  
monohydrochloride [50-98-6]

Ephedrine Hydrochloride, when dried, contains not less than 99.0% of  $C_{10}H_{15}NO \cdot HCl$ .

**Description** Ephedrine Hydrochloride occurs as white crystals or crystalline powder.

It is freely soluble in water, soluble in ethanol (95), slightly soluble in acetic acid (100), and practically insoluble in acetonitrile and in acetic anhydride.

**Identification (1)** Determine the absorption spectrum of a solution of Ephedrine Hydrochloride (1 in 2000) as directed under Ultraviolet-visible Spectrophotometry <2.24>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wavelengths.

(2) Determine the infrared absorption spectrum of Ephedrine Hydrochloride, previously dried, as directed in the potassium chloride disk method under Infrared Spectrophotometry <2.25>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wave numbers.

(3) A solution of Ephedrine Hydrochloride (1 in 15) responds to the Qualitative Tests <1.09> for chloride.

**Optical rotation** <2.49>  $[\alpha]_D^{20}$ :  $-33.0 - -36.0^\circ$  (after drying, 1 g, water, 20 mL, 100 mm).

**pH** <2.54> Dissolve 1.0 g of Ephedrine Hydrochloride in 20 mL of water: the pH of this solution is between 4.5 and 6.5.

**Melting point** <2.60> 218 – 222°C

**Purity (1)** Clarity and color of solution—Dissolve 0.5 g of Ephedrine Hydrochloride in 10 mL of water: the solution is clear and colorless.

(2) Sulfate <1.14>—Dissolve 0.05 g of Ephedrine Hydrochloride in 40 mL of water, add 1 mL of dilute hydrochloric acid and 1 mL of barium chloride TS, and allow to stand for 10 minutes: no turbidity is produced.

(3) Heavy metals <1.07>—Proceed with 1.0 g of Ephedrine Hydrochloride according to Method 2, and perform the test. Prepare the control solution with 1.0 mL of Standard Lead Solution (not more than 10 ppm).

(4) Related substances—Dissolve 0.05 g of Ephedrine Hydrochloride in 50 mL of the mobile phase, and use this solution as the sample solution. Pipet 1 mL of the sample solution, add the mobile phase to make exactly 100 mL, and use this solution as the standard solution. Perform the test with exactly 10  $\mu$ L each of the sample solution and standard solution as directed under Liquid Chromatography <2.07> ac-

cording to the following conditions, and calculate the areas of each peak by the automatic integration method: the total area of the peaks other than ephedrine from the sample solution is not larger than the peak area of ephedrine from the standard solution.

**Operating conditions—**

**Detector:** An ultraviolet absorption photometer (wavelength: 210 nm).

**Column:** A stainless steel column 4.6 mm in inside diameter and 15 cm in length, packed with octadecylsilanized silica gel for liquid chromatography (5  $\mu$ m in particle diameter).

**Column temperature:** A constant temperature of about 45°C.

**Mobile phase:** A mixture of a solution of sodium lauryl sulfate (1 in 128), acetonitrile and phosphoric acid (640:360:1).

**Flow rate:** Adjust the flow rate so that the retention time of ephedrine is about 14 minutes.

**Time span of measurement:** About 3 times as long as the retention time of ephedrine beginning after the solvent peak.

**System suitability—**

**Test for required detectability:** To exactly 1 mL of the standard solution add the mobile phase to make exactly 20 mL. Confirm that the peak area of ephedrine obtained from 10  $\mu$ L of this solution is equivalent to 4 to 6% of that from the standard solution.

**System performance:** Dissolve 1 mg of ephedrine hydrochloride for assay and 4 mg of atropine sulfate in 100 mL of diluted methanol (1 in 2). When the procedure is run with 10  $\mu$ L of this solution under the above operating conditions, ephedrine and atropine are eluted in this order with the resolution between these peaks being not less than 1.5.

**System repeatability:** When the test is repeated 6 times with 10  $\mu$ L of the standard solution under the above operating conditions, the relative standard deviation of the peak areas of ephedrine is not more than 2.0%.

**Loss on drying** <2.41> Not more than 0.5% (1 g, 105°C, 3 hours).

**Residue on ignition** <2.44> Not more than 0.1% (1 g).

**Assay** Weigh accurately about 0.4 g of Ephedrine Hydrochloride, previously dried, and dissolve in 50 mL of a mixture of acetic anhydride and acetic acid (100) (7:3) by warming. Cool, and titrate <2.50> with 0.1 mol/L perchloric acid VS (potentiometric titration). Perform a blank determination, and make any necessary correction.

Each mL of 0.1 mol/L perchloric acid VS  
= 20.17 mg of  $C_{10}H_{15}NO \cdot HCl$

**Containers and storage** Containers—Well-closed containers.

## Ephedrine Hydrochloride Injection

エフェドリン塩酸塩注射液

Ephedrine Hydrochloride Injection is aqueous solution for injection. It contains not less than 95% and not more than 105% of the labeled amount of ephed-

drine hydrochloride ( $C_{10}H_{15}NO \cdot HCl$ : 201.69).

**Method of preparation** Prepare as directed under Injections, with Ephedrine Hydrochloride.

**Description** Ephedrine Hydrochloride Injection is a clear, colorless liquid.

pH: 4.5 – 6.5

**Identification** To a volume of Ephedrine Hydrochloride Injection, equivalent to 0.05 g of Ephedrine Hydrochloride according to the labeled amount, add water to make 100 mL, and determine the absorption spectrum of this solution as directed under Ultraviolet-visible Spectrophotometry <2.24>: it exhibits maxima between 249 nm and 253 nm, between 255 nm and 259 nm, and between 261 nm and 265 nm.

**Bacterial endotoxins** <4.01> Less than 7.5 EU/mg.

**Extractable volume** <6.05> It meets the requirement.

**Assay** To an exact volume of Ephedrine Hydrochloride Injection, equivalent to about 40 mg of ephedrine hydrochloride ( $C_{10}H_{15}NO \cdot HCl$ ) according to the labeled amount, add exactly 10 mL of the internal standard solution and water to make 200 mL, and use this solution as the sample solution. Separately, weigh accurately about 40 mg of ephedrine hydrochloride for assay, previously dried at 105°C for 3 hours, add exactly 10 mL of the internal standard solution to dissolve, add water to make 200 mL, and use this solution as the standard solution. Perform the test with 10  $\mu$ L each of the sample solution and standard solution as directed under Liquid Chromatography <2.01> according to the following conditions, and calculate the ratios,  $Q_T$  and  $Q_S$ , of the peak area of ephedrine to that of the internal standard of each solution.

$$\text{Amount (mg) of ephedrine hydrochloride (C}_{10}\text{H}_{15}\text{NO} \cdot \text{HCl}) = W_S \times (Q_T/Q_S)$$

$W_S$ : Amount (mg) of ephedrine hydrochloride for assay

**Internal standard solution**—A solution of etilefrine hydrochloride (1 in 500).

**Operating conditions**—

Detector, column, column temperature, mobile phase, and flow rate: Proceed as directed in the operating conditions in the Purity (4) under Ephedrine Hydrochloride.

**System suitability**—

System performance: When the procedure is run with 10  $\mu$ L of the standard solution under the above operating conditions, the internal standard and ephedrine are eluted in this order with the resolution between these peaks being not less than 15.

System repeatability: When the test is repeated 6 times with 10  $\mu$ L of the standard solution under the above operating conditions, the relative standard deviation of the ratios of the peak area of ephedrine to that of the internal standard is not more than 1.0%.

**Containers and storage** Containers—Hermetic containers.

Storage—Light-resistant.

## 10% Ephedrine Hydrochloride Powder

### Ephedrine Hydrochloride Powder

エフェドリン塩酸塩散 10%

10% Ephedrine Hydrochloride Powder contains not less than 9.3% and not more than 10.7% of ephedrine hydrochloride ( $C_{10}H_{15}NO \cdot HCl$ : 201.69).

#### Method of preparation

|  |                       |
|--|-----------------------|
| Ephedrine Hydrochloride                  | 100 g                 |
| Starch, Lactose Hydrate or their mixture | a sufficient quantity |
| <hr/>                                    |                       |
| To make                                  | 1000 g                |

Prepare as directed under Powders, with the above ingredients.

**Identification** To 0.5 g of 10% Ephedrine Hydrochloride Powder add 100 mL of water, shake for 20 minutes, and filter. Determine the absorption spectrum of the filtrate as directed under Ultraviolet-visible Spectrophotometry <2.24>: it exhibits maxima between 249 nm and 253 nm, between 255 nm and 259 nm, and between 261 nm and 265 nm.

**Assay** Weigh accurately about 0.4 g of 10% Ephedrine Hydrochloride Powder, add 150 mL of water, and extract with the aid of ultrasonicator for 10 minutes with occasional shaking. Shake more for 10 minutes, then add exactly 10 mL of the internal standard solution and water to make 200 mL, centrifuge, and use the supernatant liquid as the sample solution. Separately, weigh accurately about 40 mg of ephedrine hydrochloride for assay, previously dried at 105°C for 3 hours, add exactly 10 mL of the internal standard solution to dissolve, add water to make 200 mL, and use this solution as the standard solution. Perform the test with 10  $\mu$ L each of the sample solution and standard solution as directed under Liquid Chromatography <2.01> according to the following conditions, and calculate the ratios,  $Q_T$  and  $Q_S$ , of the peak area of ephedrine to that of the internal standard of each solution.

$$\text{Amount (mg) of ephedrine hydrochloride (C}_{10}\text{H}_{15}\text{NO} \cdot \text{HCl}) = W_S \times (Q_T/Q_S)$$

$W_S$ : Amount (mg) of ephedrine hydrochloride for assay

**Internal standard solution**—A solution of etilefrine hydrochloride (1 in 500).

**Operating conditions**—

Detector, column, column temperature, mobile phase and flow rate: Perform as directed in the operating conditions in the Purity (4) under Ephedrine Hydrochloride.

**System suitability**—

System performance: When the procedure is run with 10  $\mu$ L of the standard solution under the above operating conditions, the internal standard and ephedrine are eluted in this order with the resolution between these peaks being not less than 15.

System repeatability: When the test is repeated 6 times with

10  $\mu$ L of the standard solution under the above operating conditions, the relative standard deviation of the ratios of the peak area of ephedrine to that of the internal standard is not more than 1.0%.

**Containers and storage** Containers—Well-closed containers.

## Ephedrine Hydrochloride Tablets

エフェドリン塩酸塩錠

Ephedrine Hydrochloride Tablets contain not less than 93% and not more than 107% of the labeled amount of ephedrine hydrochloride ( $C_{10}H_{15}NO \cdot HCl$ ; 201.69).

**Method of preparation** Prepare as directed under Tablets, with Ephedrine Hydrochloride.

**Identification** To an amount of powdered Ephedrine Hydrochloride Tablets, equivalent to 0.05 g of Ephedrine Hydrochloride, add 100 mL of water, shake for 20 minutes, and filter. Determine the absorption spectrum of the filtrate as directed under Ultraviolet-visible Spectrophotometry <2.24>: it exhibits maxima between 249 nm and 253 nm, between 255 nm and 259 nm, and between 261 nm and 265 nm.

**Assay** Weigh accurately not less than 20 tablets of Ephedrine Hydrochloride Tablets, and powder. Weigh accurately an amount of the powder, equivalent to about 40 mg of ephedrine hydrochloride ( $C_{10}H_{15}NO \cdot HCl$ ), add 150 mL of water, and extract with the aid of ultrasonicator for 10 minutes with occasional shaking. Shake more for 10 minutes, then add exactly 10 mL of the internal standard solution and water to make 200 mL, centrifuge, and use the supernatant liquid as the sample solution. Separately, weigh accurately about 40 mg of ephedrine hydrochloride for assay, previously dried at 105°C for 3 hours, add exactly 10 mL of the internal standard solution to dissolve, add water to make 200 mL, and use this solution as the standard solution. Perform the test with 10  $\mu$ L each of the sample solution and standard solution as directed under Liquid Chromatography <2.01> according to the following conditions, and calculate the ratios,  $Q_T$  and  $Q_S$ , of the peak area of ephedrine to that of the internal standard of each solution.

$$\text{Amount (mg) of ephedrine hydrochloride } (C_{10}H_{15}NO \cdot HCl) = W_S \times (Q_T/Q_S)$$

$W_S$ : Amount (mg) of ephedrine hydrochloride for assay

**Internal standard solution**—A solution of etilefrine hydrochloride (1 in 500).

**Operating conditions**—

Detector, column, column temperature, mobile phase, and flow rate: Proceed as directed in the operating conditions in the Purity (4) under Ephedrine Hydrochloride.

**System suitability**—

System performance: When the procedure is run with 10  $\mu$ L of the standard solution under the above operating conditions, the internal standard and ephedrine are eluted in this order with the resolution between these peaks being not less than 15.


System repeatability: When the test is repeated 6 times with 10  $\mu$ L of the standard solution under the above operating conditions, the relative standard deviation of the ratios of the peak area of ephedrine to that of the internal standard is not more than 1.0%.

**Containers and storage** Containers—Well-closed containers.

## Epirizole

メピリゾール

エピリゾール



$C_{11}H_{14}N_4O_2$ : 234.25

4-Methoxy-2-(5-methoxy-3-methyl-1H-pyrazol-1-yl)-6-methylpyrimidine [18694-40-1]

Epirizole, when dried, contains not less than 99.0% of  $C_{11}H_{14}N_4O_2$ .

**Description** Epirizole occurs as white crystals or crystalline powder. It is odorless, and has a bitter taste.

It is very soluble in methanol and in acetic acid (100), freely soluble in ethanol (95), and sparingly soluble in water and in diethyl ether.

It dissolves in dilute hydrochloric acid and in sulfuric acid.

The pH of a solution of Epirizole (1 in 100) is between 6.0 and 7.0.

**Identification (1)** To 0.1 g of Epirizole add 0.1 g of vanillin, 5 mL of water and 2 mL of sulfuric acid, and mix with shaking for a while: a yellow precipitate is formed.

**(2)** Dissolve 0.1 g of Epirizole in 10 mL of water, and add 10 mL of 2,4,6-trinitrophenol TS: a yellow precipitate is produced. Collect the precipitate by filtration, wash with 50 mL of water, and dry at 105°C for 1 hour: it melts <2.60> between 163°C and 169°C.

**(3)** Determine the absorption spectrum of a solution of Epirizole in 0.1 mol/L hydrochloric acid TS (1 in 200,000) as directed under Ultraviolet-visible Spectrophotometry <2.24>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wavelengths.

**Melting point** <2.60> 88 – 91°C

**Purity (1)** Clarity and color of solution—Dissolve 0.20 g of Epirizole in 20 mL of water: the solution is clear and colorless.

**(2)** Chloride <1.03>—Add 0.5 g of Epirizole to a ground mixture of 0.7 g of potassium nitrate and 1.2 g of anhydrous sodium carbonate, mix well, transfer little by little to a platinum crucible, previously heated, and heat until the reaction is completed. After cooling, add 15 mL of dilute sulfuric acid and 5 mL of water to the residue, boil for 5 minutes, filter, wash the insoluble matter with 10 mL of water, and add 6 mL

of dilute nitric acid and water to the combined filtrate and washings to make 50 mL. Perform the test with this solution as the test solution. Prepare the control solution as follows: proceed with the same quantities of the same reagents as directed for the preparation of the test solution, and add 0.25 mL of 0.01 mol/L hydrochloric acid VS and water to make 50 mL (not more than 0.018%).

(3) Heavy metals <1.07>—Proceed with 2.0 g of Epirizole according to Method 2, and perform the test. Prepare the control solution with 2.0 mL of Standard Lead Solution (not more than 10 ppm).

(4) Arsenic <1.11>—Prepare the test solution with 1.0 g of Epirizole according to Method 3, and perform the test (not more than 2 ppm).

(5) Related substances—Dissolve 1.0 g of Epirizole in 10 mL of methanol, and use this solution as the sample solution. Pipet 1 mL of the sample solution, and add methanol to make exactly 50 mL. Pipet 1 mL of this solution, add methanol to make exactly 10 mL, and use this solution as the standard solution. Perform the test with these solutions as directed under Thin-layer Chromatography <2.03>. Spot 2  $\mu$ L each of the sample solution and standard solution on a plate of silica gel with fluorescent indicator for thin-layer chromatography. Develop the plate with a mixture of isopropyl diethyl ether, ethanol (95) and water (23:10:2) to a distance of about 10 cm, and air-dry the plate. Examine under ultraviolet light (main wavelength: 254 nm): the spots other than the principal spot from the sample solution are not more intense than the spot from the standard solution. Place this plate in a chamber filled with iodine vapor: the spots other than the principal spot from the sample solution are not more intense than the spot from the standard solution.

(6) Readily carbonizable substances <1.15>—Perform the test with 0.10 g of Epirizole: the solution has no more color than Matching Fluid A.

**Loss on drying** <2.41> Not more than 0.5% (1 g, silica gel, 4 hours).

**Residue on ignition** <2.44> Not more than 0.1% (1 g).

**Assay** Weigh accurately about 0.5 g of Epirizole, previously dried, dissolve in 40 mL of acetic acid (100) and titrate <2.50> with 0.1 mol/L perchloric acid VS (indicator: 2 drops of crystal violet TS) until the color of the solution changes from purple through blue-green to green.


Perform a blank determination, and make any necessary correction.

Each mL of 0.1 mol/L perchloric acid VS  
= 23.43 mg of  $C_{27}H_{29}NO_{11}$

**Containers and storage** Containers—Well-closed containers.

## Epirubicin Hydrochloride

エピルビシン塩酸塩



$C_{27}H_{29}NO_{11} \cdot HCl$ : 579.98

(2*S*,4*S*)-4-(3-Amino-2,3,6-trideoxy- $\alpha$ -L-arabino-hexopyranosyloxy)-2,5,12-trihydroxy-2-hydroxyacetyl-7-methoxy-1,2,3,4-tetrahydrotetracyclic-6,11-dione monohydrochloride [56390-09-1]

Epirubicin Hydrochloride is the hydrochloride of a derivative of daunorubicin.

It contains not less than 970  $\mu$ g (potency) and not more than 1020  $\mu$ g (potency) per mg, calculated on the anhydrous basis and corrected by the amount of the residual solvent. The potency of Epirubicin Hydrochloride is expressed as mass (potency) of epirubicin hydrochloride ( $C_{27}H_{29}NO_{11} \cdot HCl$ ).

**Description** Epirubicin Hydrochloride occurs as a pale yellowish red to brownish red powder.

It is soluble in water and in methanol, slightly soluble in ethanol (95), and practically insoluble in acetonitrile.

It is hygroscopic.

**Identification** (1) Determine the absorption spectrum of a solution of Epirubicin Hydrochloride in methanol (3 in 200,000) as directed under Ultraviolet-visible Spectrophotometry <2.24>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wavelengths.

(2) Determine the infrared absorption spectrum of Epirubicin Hydrochloride and Epirubicin Hydrochloride Reference Standard as directed in the potassium bromide disk method under Infrared Spectrophotometry <2.25>, and compare these spectra: both spectra exhibit similar intensities of absorption at the same wave numbers.

**Optical rotation** <2.49>  $[\alpha]_D^{20}$ : +310 – +340° (10 mg calculated on the anhydrous basis and collected by the amount of the residual solvent, methanol, 20 mL, 100 mm).

**pH** <2.54> Dissolve 10 mg of Epirubicin Hydrochloride in 2 mL of water: the pH of the solution is between 4.0 and 5.5.

**Purity** (1) Clarity and color of solution—Dissolve 50 mg of Epirubicin Hydrochloride in 5 mL of water: the solution is clear and dark red.

(2) Heavy metals <1.07>—Proceed with 1.0 g of Epirubicin Hydrochloride according to Method 2, and perform the test. Prepare the control solution with 2.0 mL of Standard Lead Solution (not more than 20 ppm).

(3) Related substances—Perform the test with 10  $\mu$ L of

the sample solution obtained in the Assay as directed under Liquid Chromatography <2.01> according to the following conditions, determine each peak area by the automatic integration method, and calculate the sum amount of the peaks other than epirubicin and 2-naphthalenesulfonic acid by the area percentage method: not more than 5.0%.

**Operating conditions—**

Detector, column, column temperature, mobile phase, and flow rate: Proceed as directed in the operating conditions in the Assay.

Time span of measurement: About 3 times as long as the retention time of epirubicin beginning after the solvent peak.

**System suitability—**

Test for required detectability: To 1 mL of the sample solution add the mobile phase to make 100 mL, and use this solution as the solution for system suitability test. Pipet 1 mL of the solution for system suitability test, and add the mobile phase to make exactly 10 mL. Confirm that the peak area of epirubicin obtained from 10  $\mu$ L of this solution is equivalent to 7 to 13% of that obtained from 10  $\mu$ L of the solution for system suitability test.

System performance, and system repeatability: Proceed as directed in the system suitability in the Assay.

(4) Residual solvents <2.46>—Weigh accurately about 0.3 g of Epirubicin Hydrochloride, add exactly 0.6 mL of the internal standard solution, add *N,N*-dimethylformamide to make 6 mL, and use this solution as the sample solution. Separately, pipet 1 mL of methanol, add *N,N*-dimethylformamide to make exactly 25 mL, and use this solution as methanol standard stock solution. Take exactly 125  $\mu$ L of acetone, 30  $\mu$ L of ethanol (99.5), 32  $\mu$ L of 1-propanol and 17  $\mu$ L of the methanol standard stock solution, add exactly 10 mL of the internal standard solution and *N,N*-dimethylformamide to make 100 mL, and use this solution as the standard solution. Perform the test with 1  $\mu$ L each of the sample solution and standard solution as directed under Gas Chromatography <2.02> according to the following condition, and determine the ratios of the peak areas of acetone, ethanol, 1-propanol and methanol to that of the internal standard,  $Q_{TA}$  and  $Q_{SA}$ ,  $Q_{TB}$  and  $Q_{SB}$ ,  $Q_{TC}$  and  $Q_{SC}$ , and  $Q_{TD}$  and  $Q_{SD}$ , respectively. Calculate the amounts of acetone, ethanol, 1-propanol and methanol by the following equations: the amounts of acetone, ethanol, 1-propanol and methanol are not more than 1.5%, not more than 0.5%, not more than 0.5% and not more than 0.1%, respectively.

$$\begin{aligned} \text{Amount (\%)} & \text{ of acetone} \\ &= (1/W_T) \times (Q_{TA}/Q_{SA}) \times 593 \end{aligned}$$

$$\begin{aligned} \text{Amount (\%)} & \text{ of ethanol} \\ &= (1/W_T) \times (Q_{TB}/Q_{SB}) \times 142 \end{aligned}$$

$$\begin{aligned} \text{Amount (\%)} & \text{ of 1-propanol} \\ &= (1/W_T) \times (Q_{TC}/Q_{SC}) \times 154 \end{aligned}$$

$$\begin{aligned} \text{Amount (\%)} & \text{ of methanol} \\ &= (1/W_T) \times (Q_{TD}/Q_{SD}) \times 2.23 \end{aligned}$$

$W_T$ : Amount (mg) of Epirubicin Hydrochloride

**Internal standard solution—**A solution of 1,4-dioxane in *N,N*-dimethylformamide (1 in 100).

**Operating conditions—**

Detector: Hydrogen flame-ionization detector.

Column: A fused silica column 0.53 mm in inside diameter

and 30 m in length, coated with polyethylene glycol for gas-chromatography 1  $\mu$ m in thickness.

Column temperature: 40°C for 11 minutes after injection of the sample, then rise to 90°C at a rate of 10°C per minute. If necessary, rise to 130°C at a rate of 50°C per minute and maintain the temperature for 30 minutes.

Injection port temperature: A constant temperature of about 120°C.

Detector temperature: A constant temperature of about 150°C.

Carrier gas: Helium

Flow rate: Adjust the flow rate so that the retention time of the internal standard is about 8 minutes.

Split ratio: 1:15

**System suitability—**

System performance: When the procedure is run with 1  $\mu$ L of the standard solution under the above operating conditions, acetone, methanol, ethanol, 1-propanol and the internal standard are eluted in this order with the resolution between the peaks of acetone and the internal standard being not less than 30.

System repeatability: When the test is repeated 6 times with 1  $\mu$ L of the standard solution under the above operating conditions, the relative standard deviations of the peak areas of acetone, methanol, ethanol and 1-propanol are not more than 4.0%, respectively.

**Water** Not more than 8.0% (0.1 g, volumetric titration, direct titration).

**Residue on ignition** <2.44> Not more than 0.5% (0.1 g).

**Assay** Weigh accurately an amount of Epirubicin Hydrochloride and Epirubicin Hydrochloride Reference Standard, equivalent to about 50 mg (potency), dissolve each in the internal standard solution to make exactly 50 mL, and use these solutions as the sample solution and standard solution, respectively. Perform the test with 10  $\mu$ L each of the sample solution and standard solution as directed under Liquid Chromatography <2.01> according to the following conditions, and determine the ratios,  $Q_T$  and  $Q_S$ , of the peak area of epirubicin to that of the internal standard.

$$\begin{aligned} \text{Amount } [\mu\text{g (potency)}] & \text{ of } C_{27}H_{29}NO_{11}.HCl \\ &= W_S \times (Q_T/Q_S) \times 1000 \end{aligned}$$

$W_S$ : Amount [mg (potency)] of Epirubicin Hydrochloride Reference Standard

**Internal standard solution—**A solution of sodium 2-naphthalene sulfonate in a mixture of water, acetonitrile, methanol and phosphoric acid (540:290:170:1) (1 in 2000).

**Operating conditions—**

Detector: An ultraviolet absorption photometer (wavelength: 254 nm).

Column: A stainless steel column 4 mm in inside diameter and 25 cm in length, packed with trimethylsilylated silica gel for liquid chromatography (6  $\mu$ m in particle diameter).

Column temperature: A constant temperature of about 35°C.

Mobile phase: Dissolve 2 g of sodium lauryl sulfate in a mixture of water, acetonitrile, methanol and phosphoric acid (540:290:170:1) to make 1000 mL.

Flow rate: Adjust the flow rate so that the retention time of epirubicin is about 9.5 minutes.

**System suitability—**

**System performance:** When the procedure is run with 10  $\mu$ L of the standard solution under the above operating conditions, the internal standard and epirubicin are eluted in this order with the resolution between these peaks being not less than 20.

**System repeatability:** When the test is repeated 5 times with 10  $\mu$ L of the standard solution under the above operating conditions, the relative standard deviation of the ratios of the peak area of epirubicin to that of the internal standard is not more than 1.0%.

**Containers and storage** Containers—Tight containers.


Storage—At a temperature between 0°C and 5°C.

## Ergocalciferol

### Calciferol

### Vitamin D<sub>2</sub>

エルゴカルシフェロール



$C_{28}H_{44}O$ : 396.65  
(3*S*,5*Z*,7*E*,22*E*)-9,10-Secoergosta-5,7,10(19),22-tetraen-3-ol  
[50-14-6]

Ergocalciferol contains not less than 97.0% and not more than 103.0% of  $C_{28}H_{44}O$ .

**Description** Ergocalciferol occurs as white crystals. It is odorless, or has a faint, characteristic odor.

It is freely soluble in ethanol (95), in diethyl ether and in chloroform, sparingly soluble in isooctane, and practically insoluble in water.

It is affected by air and by light.

**Melting point:** 115 – 118°C Transfer Ergocalciferol to a capillary tube, and dry for 3 hours in a desiccator (in vacuum at a pressure not exceeding 2.67 kPa). Immediately fire-seal the capillary tube, put it in a bath fluid, previously heated to a temperature about 10°C below the expected melting point, and heat at a rate of rise of about 3°C per minute, and read the melting point.

**Identification (1)** Dissolve 0.5 mg of Ergocalciferol in 5 mL of chloroform, add 0.3 mL of acetic anhydride and 0.1 mL of sulfuric acid, and shake: a red color is produced, and rapidly changes through purple and blue to green.

**(2)** Determine the infrared absorption spectrum of Ergocalciferol as directed in the potassium bromide disk method under Infrared Spectrophotometry <2.25>, and compare the spectrum with the Reference Spectrum or the spectrum of Ergocalciferol Reference Standard: both spectra exhibit similar intensities of absorption at the same wave numbers.

**Absorbance** <2.24>  $E_{1\text{ cm}}^{1\%}$  (265 nm): 445 – 485 (0.01 g, ethanol (95), 100 mL).

**Optical rotation** <2.49>  $[\alpha]_D^{20}$ : +102 – +107° (0.3 g, ethanol (95), 20 mL, 100 mm). Prepare the solution of Ergocalciferol within 30 minutes after the container has been opened, and determine the rotation within 30 minutes after the solution has been prepared.

**Purity** Ergosterol—Dissolve 10 mg of Ergocalciferol in 2.0 mL of diluted ethanol (95) (9 in 10), add a solution of 20 mg of digitonin in 2.0 mL of diluted ethanol (95) (9 in 10), and allow the mixture to stand for 18 hours: no precipitate is formed.

**Assay** Weigh accurately about 30 mg each of Ergocalciferol and Ergocalciferol Reference Standard, and dissolve each in isooctane to make exactly 50 mL. Pipet 10 mL each of these solutions, add exactly 3 mL each of the internal standard solution, then add the mobile phase to make 50 mL, and use these solutions as the sample solution and the standard solution, respectively. Perform the test with 10 to 20  $\mu$ L each of the sample solution and standard solution as directed under Liquid Chromatography <2.01> according to the following conditions, and calculate the ratios,  $Q_T$  and  $Q_S$ , of the peak area of ergocalciferol to that of the internal standard. Perform the procedure rapidly avoiding contact with air or other oxidizing agents and using light-resistant containers.

$$\text{Amount (mg) of } C_{28}H_{44}O = W_S \times (Q_T/Q_S)$$

$W_S$ : Amount (mg) of Ergocalciferol Reference Standard

**Internal standard solution**—A solution of dimethyl phthalate in isooctane (1 in 100).

**Operating conditions**—

**Detector:** An ultraviolet absorption photometer (wavelength: 254 nm).

**Column:** A stainless steel column 4.6 mm in inside diameter and 25 cm in length, packed with a silica gel for liquid chromatography (10  $\mu$ m particle diameter).

**Column temperature:** A constant temperature of about 20°C.

**Mobile phase:** A mixture of hexane and n-amylalcohol (997:3).

**Flow rate:** Adjust the flow rate so that the retention time of ergocalciferol is about 25 minutes.

**System suitability**—

**System performance:** Dissolve 15 mg of Ergocalciferol Reference Standard in 25 mL of isooctane. Transfer this solution to a flask, heat in an oil bath under a reflux condenser for 2 hours, and cool immediately to room temperature. Transfer the solution to a quartz test tube, and irradiate with a short-wave lamp (main wavelength: 254 nm) and a long-wave lamp (main wavelength: 365 nm) for 3 hours. To 10 mL of this solution add the mobile phase to make 50 mL. When the procedure is run with 10  $\mu$ L of this solution under the above operating conditions. Use a column with the ratios of the retention time of previtamin D<sub>2</sub>, trans-vitamin D<sub>2</sub> and tachysterol<sub>2</sub> to that of ergocalciferol being about 0.5, about 0.6 and about 1.1, respectively, and with resolution between previtamin D<sub>2</sub> and trans-vitamin D<sub>2</sub> being not less than 0.7, and that between ergocalciferol and tachysterol<sub>2</sub> being not less than 1.0.

**System repeatability:** When the test is repeated 6 times with




10  $\mu$ L of the standard solution under the above operating conditions, the relative standard deviation of the ratios of the peak area of ergocalciferol to that of the internal standard is not more than 1.0%.

**Containers and storage** Containers—Hermetic containers.

Storage—Light-resistant, under Nitrogen atmosphere, and in a cold place.

## Ergometrine Maleate

エルゴメトリンマレイン酸塩



$C_{19}H_{23}N_3O_2 \cdot C_4H_4O_4$ : 441.48

(8*S*)-*N*-[(1*S*)-2-Hydroxy-1-methylethyl]-6-methyl-9,10-didehydroergoline-8-carboxamide monomaleate  
[129-51-1]

Ergometrine Maleate, when dried, contains not less than 98.0% of  $C_{19}H_{23}N_3O_2 \cdot C_4H_4O_4$ .

**Description** Ergometrine Maleate occurs as a white to pale yellow, crystalline powder. It is odorless.

It is sparingly soluble in water, slightly soluble in methanol and in ethanol (95), and practically insoluble in diethyl ether.

Melting point: about 185°C (with decomposition).

It gradually changes to yellow in color on exposure to light.

**Identification (1)** Prepare a solution of Ergometrine Maleate (1 in 50): the solution shows a blue fluorescence.

(2) Dissolve 1 mg of Ergometrine Maleate in 5 mL of water. To 1 mL of this solution add 2 mL of 4-dimethylaminobenzaldehyde-ferric chloride TS, shake, and allow to stand for 5 to 10 minutes: a deep blue color develops.

(3) To 5 mL of a solution of Ergometrine Maleate (1 in 500) add 1 drop of potassium permanganate TS: the red color of the solution disappears immediately.

**Optical rotation** <2.49>  $[\alpha]_D^{20} +48 - +57^\circ$  (after drying 0.25 g, water, 25 mL, 100 mm).

**pH** <2.54> Dissolve 0.10 g of Ergometrine Maleate in 10 mL of water. The pH of the solution is between 3.0 and 5.0.

**Purity (1)** Clarity and color of solution—Dissolve 0.10 g of Ergometrine Maleate in 10 mL of water: the solution is clear and colorless to light yellow.

(2) Ergotamine and ergotamine—To 0.02 g of Ergometrine Maleate add 2 mL of a solution of sodium hydroxide (1 in 10), and heat to boiling: the gas evolved does not change moistened red litmus paper to blue.

(3) Related substances—Dissolve 5 mg each of Ergometrine Maleate and Ergometrine Maleate Reference Standard in 1.0 mL of methanol, and use these solutions as the sample solution and standard solution, respectively. Perform the test with these solutions as directed under Thin-layer Chromatography <2.03>.

Spot 10  $\mu$ L each of the sample solution and standard solution on a plate, prepared with silica gel for thin-layer chromatography and dilute sodium hydroxide TS. Develop the plate with a mixture of chloroform and methanol (4:1) to a distance of about 10 cm, and air-dry the plate. Spray evenly 4-dimethylaminobenzaldehyde TS on the plate: the spots obtained from the sample solution and the standard solution show a red-purple color and the same *R<sub>f</sub>* value, and any spot from the sample solution other than that corresponding to the spot from the standard solution does not appear.

**Loss on drying** <2.41> Not more than 2.0% (0.2 g, silica gel, 4 hours).

**Assay** Weigh accurately about 10 mg each of Ergometrine Maleate and Ergometrine Maleate Reference Standard, previously dried in a desiccator (silica gel) for 4 hours, dissolve in water to make exactly 250 mL, and use these solutions as the sample solution and the standard solution, respectively. Pipet 2 mL of each solution into a separate brown glass-stoppered tube. To each tube add 4 mL of 4-dimethylaminobenzaldehyde-iron (III) chloride TS, exactly measured, while cooling in an ice bath, then warm at 45°C for 10 minutes. Allow to stand at room temperature for 20 minutes, and perform the test with these solutions as directed under Ultraviolet-visible Spectrophotometry <2.24>, using a solution, prepared with 2 mL of water in the same manner, as the blank. Determine the absorbances, *A<sub>T</sub>* and *A<sub>S</sub>*, of the subsequent solutions of the sample solution and the standard solution at 550 nm, respectively.

$$\begin{aligned} \text{Amount (mg) of } C_{19}H_{23}N_3O_2 \cdot C_4H_4O_4 \\ = W_S \times (A_T/A_S) \end{aligned}$$

*W<sub>S</sub>*: Amount (mg) of Ergometrine Maleate Reference Standard

**Containers and storage** Containers—Tight containers.

Storage—Light-resistant.

## Ergometrine Maleate Injection

エルゴメトリンマレイン酸塩注射液

Ergometrine Maleate Injection is an aqueous solution for injection.

It contains not less than 90% and not more than 110% of the labeled amount of ergometrine maleate ( $C_{19}H_{23}N_3O_2 \cdot C_4H_4O_4$ : 441.48).

**Method of preparation** Prepare as directed under Injections, with Ergometrine Maleate.

**Description** Ergometrine Maleate Injection is a clear, colorless to pale yellow liquid.

pH: 2.7 – 3.5

**Identification (1)** Measure a volume of Ergometrine Maleate Injection, equivalent to 3 mg of Ergometrine Maleate according to the labeled amount, if necessary, dilute with water or evaporate on a water bath to make 15 mL, and use this solution as the sample solution. The sample solution shows a blue fluorescence.

(2) To 1 mL of the sample solution obtained in (1) add 1

mL of ammonia TS, and extract with 20 mL of diethyl ether. To the diethyl ether extract add 1 mL of dilute sulfuric acid, shake, and warm to remove diethyl ether in a water bath. Cool, to the residue obtained add 2 mL of 4-dimethylaminobenzaldehyde-iron (III) chloride TS, and allow to stand for 5 to 10 minutes: a deep blue color develops.

(3) To 5 mL of the sample solution obtained in (1) add 1 drop of potassium permanganate TS: a red color disappears immediately.

**Extractable volume** <6.05> It meets the requirement.

**Assay** Transfer an exactly measured volume of Ergometrine Maleate Injection, equivalent to about 2 mg of ergometrine maleate ( $C_{19}H_{23}N_3O_2 \cdot C_4H_4O_4$ ), and add sodium chloride in a ratio of 0.3 g to 1 mL of the solution. To this mixture add 20 mL of diethyl ether and 2 mL of ammonia TS, shake, and extract. Further, extract with three 15-mL portions of diethyl ether, combine all the extracts, add 5 g of anhydrous sodium sulfate, filter through a pledget of absorbent cotton, and wash with three 5-mL portions of diethyl ether. Add the washings to the filtrate, shake with 5 mL of dilute sulfuric acid, evaporate the diethyl ether by warming in a current of nitrogen, to the remaining solution add water to make exactly 50 mL, and use this solution as the sample solution. Weigh accurately about 2 mg of Ergometrine Maleate Reference Standard, previously dried in a desiccator (silica gel) for 4 hours, add water to make exactly 50 mL, and use this solution as the standard solution. Transfer 2 mL each of the sample solution and standard solution, accurately measured, to separate glass-stoppered test tubes, and proceed as directed in the Assay under Ergometrine Maleate.

Amount (mg) of ergometrine maleate ( $C_{19}H_{23}N_3O_2 \cdot C_4H_4O_4$ )  
 $= W_S \times (A_T/A_S)$

$W_S$ : Amount (mg) of Ergometrine Maleate Reference Standard

**Containers and storage** Containers—Hermetic containers, and colored containers may be used.

Storage—Light-resistant, and in a cold place.

## Ergometrine Maleate Tablets

エルゴメトリンマレイン酸塩錠

Ergometrine Maleate Tablets contain not less than 90% and not more than 110% of the labeled amount of ergometrine maleate ( $C_{19}H_{23}N_3O_2 \cdot C_4H_4O_4$ : 441.48).

**Method of preparation** Prepare as directed under Tablets, with Ergometrine Maleate.

**Identification** To a quantity of powdered Ergometrine Maleate Tablets, equivalent to 3 mg of Ergometrine Maleate according to the labeled amount, add 15 mL of warm water, shake, and filter: the filtrate shows a blue fluorescence. Proceed with this solution as directed in the Identification (2) and (3) under Ergometrine Maleate.

**Uniformity of dosage units** <6.02> Perform the test according to the following method: it meets the requirement of the Content uniformity test.

Transfer 1 tablet of Ergometrine Maleate Tablets to a

glass-stoppered centrifuge tube, and add a solution of L-tartaric acid (1 in 100) to make exactly  $V$  mL of a solution containing about  $40 \mu\text{g}$  of ergometrine maleate ( $C_{19}H_{23}N_3O_2 \cdot C_4H_4O_4$ ) per mL. Stopper the tube, shake for 30 minutes vigorously, centrifuge, and use the supernatant liquid as the sample solution. Separately, weigh accurately about 4 mg of Ergometrine Maleate Reference Standard, previously dried in a desiccator (silica gel) for 4 hours, dissolve in water to make exactly 100 mL, and use this solution as the standard solution. Pipet 4 mL each of the sample solution and standard solution into separate brown glass-stoppered test tubes, add exactly 8 mL each of 4-dimethylaminobenzaldehyde-iron (III) chloride TS while cooling in an ice bath, after shaking, and allow to stand for 1 hour at ordinary temperature. Perform the test with these solutions as directed under Ultraviolet-visible Spectrophotometry <2.24>, using a solution, prepared with 4 mL of water in the same manner, as the blank. Determine the absorbances,  $A_T$  and  $A_S$ , of the subsequent solutions of the sample solution and the standard solution at 550 nm, respectively.

Amount (mg) of ergometrine maleate ( $C_{19}H_{23}N_3O_2 \cdot C_4H_4O_4$ )  
 $= W_S \times (A_T/A_S) \times (V/100)$

$W_S$ : Amount (mg) of Ergometrine Maleate Reference Standard

**Assay** Weigh accurately, and powder not less than 20 Ergometrine Maleate Tablets. Weigh accurately a portion of the powder, equivalent to about 2 mg of ergometrine maleate ( $C_{19}H_{23}N_3O_2 \cdot C_4H_4O_4$ ), transfer to a glass filter (G4), add 10 mL of a solution of L-tartaric acid (1 in 100), and filter with thorough shaking. Repeat the procedures 3 times, combine the filtrates, add a solution of L-tartaric acid (1 in 100) to make exactly 50 mL, and use this solution as the sample solution. Separately, weigh accurately about 2 mg of Ergometrine Maleate Reference Standard, previously dried in a desiccator (silica gel) for 4 hours, dissolve in a solution of L-tartaric acid (1 in 100) to make exactly 50 mL, and use this solution as the standard solution. Pipet 2 mL each of the sample solution and standard solution, and proceed as directed in the Assay under Ergometrine Maleate.

Amount (mg) of ergometrine maleate ( $C_{19}H_{23}N_3O_2 \cdot C_4H_4O_4$ )  
 $= W_S \times (A_T/A_S)$


$W_S$ : Amount (mg) of Ergometrine Maleate Reference Standard

**Containers and storage** Containers—Well-closed containers.

Storage—Light-resistant.

## Ergotamine Tartrate

エルゴタミン酒石酸塩



$(C_{33}H_{35}N_5O_5)_2 \cdot C_4H_6O_6$ ; 1313.41

(5'S)-5'-Benzyl-12'-hydroxy-2'-methyletergotaman-3',6',18-trione hemitartrate [379-79-3]

Ergotamine Tartrate contains not less than 98.0% of  $(C_{33}H_{35}N_5O_5)_2 \cdot C_4H_6O_6$ , calculated on the dried basis.

**Description** Ergotamine Tartrate occurs as colorless crystals, or a white to pale yellowish white or grayish white, crystalline powder.

It is slightly soluble in water and in ethanol (95).

Melting point: about 180°C (with decomposition).

**Identification (1)** Dissolve 1 mg of Ergotamine Tartrate in 10 mL of a mixture of acetic acid (100) and ethyl acetate (1:1). To 0.5 mL of this solution add slowly 0.5 mL of sulfuric acid, with shaking in cold water, and allow to stand: a purple color develops. To this solution add 0.1 mL of diluted iron (III) chloride TS (1 in 12): the color of the solution changes to blue to blue-purple.

**(2)** Dissolve 1 mg of Ergotamine Tartrate in 5 mL of a solution of L-tartaric acid (1 in 100). To 1 mL of this solution add 2 mL of 4-dimethylaminobenzaldehyde-iron (III) chloride TS, and shake: a blue color develops.

**Optical rotation** <2.49> Ergotamine base  $[\alpha]_D^{20}$ :  $-155$  –  $-165^\circ$ . Dissolve 0.35 g of Ergotamine Tartrate in 25 mL of a solution of L-tartaric acid (1 in 100), add 0.5 g of sodium hydrogen carbonate, shake gently and sufficiently, and extract with four 10-mL portions of ethanol-free chloroform. Filter the extracts successively through a small filter paper, moistened with ethanol-free chloroform, into a 50-mL volumetric flask. Allow the flask to stand in a water bath at 20°C for 10 minutes, and determine the optical rotation in a 100-mm cell. Separately, pipet 25 mL of this solution, evaporate to dryness under reduced pressure at a temperature not higher than 45°C, dissolve the residue in 25 mL of acetic acid (100), and titrate <2.50> with 0.05 mol/L perchloric acid VS (indicator: 1 drop of crystal violet TS). Perform a blank determination, and make any necessary correction. Calculate the specific rotation of the ergotamine base from the consumed volume of 0.05 mol/L perchloric acid VS and the optical rotation.

Each mL of 0.05 mol/L perchloric acid VS  
= 29.08 mg of  $C_{33}H_{35}N_5O_5$

**Purity** Related substances—Conduct this procedure without exposure to daylight, using light-resistant vessels. To 40 mg of Ergotamine Tartrate add 10 mL of a solution of L-

tartaric acid in diluted methanol (1 in 2) (1 in 1000), dissolve with thorough shaking, and use this solution as the sample solution. Pipet 1 mL of this solution, add a solution of L-tartaric acid in diluted methanol (1 in 2) (1 in 1000) to make exactly 50 mL, and use this solution as the standard solution. Perform the test with these solutions as directed under Thin-layer Chromatography <2.03>. Spot 10  $\mu$ L each of the sample solution and standard solution on a plate of silica gel for thin-layer chromatography. Develop the plate with a mixture of chloroform and methanol (9:1) to a distance of about 10 cm, and air-dry the plate. Spray evenly 4-dimethylaminobenzaldehyde TS on the plate: the spots other than the principal spot from the sample solution are not more intense than the spot from the standard solution.

**Loss on drying** <2.41> Not more than 5.0% (0.1 g, in vacuum, 60°C, 4 hours).

**Assay** Weigh accurately about 0.2 g of Ergotamine Tartrate, dissolve in 15 mL of a mixture of acetic acid (100) and acetic anhydride (50:3), and titrate <2.50> with 0.05 mol/L perchloric acid VS (indicator: 1 drop of crystal violet TS). Perform a blank determination, and make any necessary correction.


Each mL of 0.05 mol/L perchloric acid VS  
= 32.84 mg of  $(C_{33}H_{35}N_5O_5)_2 \cdot C_4H_6O_6$

**Containers and storage** Containers—Tight containers.

Storage—Light-resistant, and almost well-filled, or under nitrogen atmosphere, and not exceeding 5°C.

## Erythromycin

エリスロマイシン



$C_{37}H_{67}NO_{13}$ ; 733.93

(2R,3S,4S,5R,6R,8R,10R,11R,12S,13R)-5-(3,4,6-Trideoxy-3-dimethylamino- $\beta$ -D-xylo-hexopyranosyloxy)-3-(2,6-dideoxy-3-C-methyl-3-O-methyl- $\alpha$ -L-ribo-hexopyranosyloxy)-6,11,12-trihydroxy-2,4,6,8,10,12-hexamethyl-9-oxopentadecan-13-olide [114-07-8]

Erythromycin is a macrolide substance having antibacterial activity produced by the growth of *Saccharopolyspora erythraea*.

It contains not less than 930  $\mu$ g (potency) and not more than 1020  $\mu$ g (potency) per mg, calculated on the anhydrous basis. The potency of Erythromycin is expressed as mass (potency) of erythromycin ( $C_{37}H_{67}NO_{13}$ ).

**Description** Erythromycin occurs as a white to light yellow-

ish white powder.

It is very soluble in *N,N*-dimethylformamide, freely soluble in methanol and in ethanol (95), and very slightly soluble in water.

**Identification (1)** Determine the infrared absorption spectrum of Erythromycin as directed in the potassium bromide disk method under Infrared Spectrophotometry <2.25>, and compare the spectrum with the Reference Spectrum or the spectrum of Erythromycin Reference Standard: both spectra exhibit similar intensities of absorption at the same wave numbers.

(2) Dissolve 10 mg each of Erythromycin and Erythromycin Reference Standard in 1 mL of methanol, and use these solutions as the sample solution and standard solution. Perform the test with these solutions as directed under Thin-layer Chromatography <2.03>. Spot 5  $\mu$ L each of the sample solution and standard solution on a plate of silica gel for thin-layer chromatography. Develop the plate with a mixture of methanol and ammonia solution (28) (50:1) to a distance of about 10 cm, and air-dry the plate. Spray evenly 4-methoxybenzaldehyde-sulfuric acid TS on the plate, and heat at 100°C for 15 minutes: the principal spot from the sample solution and the spot from the standard solution are dark purple in color, and their *R<sub>f</sub>* values are the same.

**Optical rotation** <2.49>  $[\alpha]_D^{20}$ :  $-71 - -78^\circ$  (1 g calculated on the anhydrous basis, ethanol (95), 50 mL, 100 mm).

**Purity (1)** Heavy metals <1.07>—Proceed with 1.0 g of Erythromycin according to Method 4, and perform the test. Prepare the control solution with 2.0 mL of Standard Lead Solution (not more than 20 ppm).

(2) Arsenic <1.11>—Prepare the test solution with 1.0 g of Erythromycin according to Method 5 using hydrochloric acid instead of diluted hydrochloric acid (1 in 2), and perform the test (not more than 2 ppm).

(3) Related substances—Dissolve 40 mg of Erythromycin in 2 mL of methanol, add a mixture of phosphate buffer solution, pH 7.0 and methanol (15:1) to make exactly 10 mL, and use this solution as the sample solution. Separately, dissolve 16 mg of Erythromycin Reference Standard in 2 mL of methanol, add a mixture of phosphate buffer solution, pH 7.0 and methanol (15:1) to make exactly 10 mL, and use this solution as the standard stock solution. Dissolve 5 mg each of erythromycin B and erythromycin C in 2 mL of methanol, add exactly 2 mL of the standard stock solution, add a mixture of phosphate buffer solution, pH 7.0 and methanol (15:1) to make exactly 25 mL, and use this solution as the standard solution. Perform the test with exactly 100  $\mu$ L each of the sample solution and standard solution as directed under Liquid Chromatography <2.01> according to the following conditions, and determine each peak area by the automatic integration method: the peak areas of erythromycin B and erythromycin C from the sample solution are not more than those of erythromycin B and erythromycin C from the standard solution, respectively, and each area of the peaks other than erythromycin, erythromycin B and erythromycin C is not more than the area of the peak of erythromycin from the standard solution.

**Operating conditions—**

**Detector:** An ultraviolet absorption photometer (wavelength: 215 nm).

**Column:** A stainless steel column 4.6 mm in inside di-

ameter and 25 cm in length, packed with styrene-divinylbenzene copolymer for liquid chromatography (8  $\mu$ m in particle diameter).

**Column temperature:** A constant temperature of about 70°C.

**Mobile phase:** Dissolve 3.5 g of dipotassium hydrogen phosphate in water to make 100 mL, and adjust the pH to 9.0 with diluted phosphoric acid (1 in 10). To 50 mL of this solution add 190 mL of *t*-butanol, 30 mL of acetonitrile and water to make 1000 mL.

**Flow rate:** Adjust the flow rate so that the retention time of erythromycin is about 20 minutes.

**Time span of measurement:** About 4 times as long as the retention time of erythromycin beginning after the solvent peak.

**System suitability—**

**System performance:** Dissolve 2 mg of *N*-demethylerythromycin in 10 mL of the standard solution. When the procedure is run with 100  $\mu$ L of this solution under the above operating conditions, *N*-demethylerythromycin, erythromycin C, erythromycin and erythromycin B are eluted in this order, with the resolution between the peaks of *N*-demethylerythromycin and erythromycin C being not less than 0.8, and with the resolution between the peaks of *N*-demethylerythromycin and erythromycin being not less than 5.5.

**System repeatability:** When the test is repeated 3 times with 100  $\mu$ L of the standard solution under the above operating conditions, the relative standard deviation of the peak area of erythromycin is not more than 3.0%.

**Water** <2.48> Not more than 10.0% (0.2 g, volumetric titration, direct titration).

**Residue on ignition** <2.44> Not more than 0.2% (1 g).

**Assay** Perform the test according to the Cylinder-plate method as directed under Microbial Assay for Antibiotics <4.02> according to the following conditions.

(i) Test organism—*Staphylococcus aureus* ATCC 6538 P

(ii) Culture medium—Use the medium i in 3) Medium for other organisms under (1) Agar media for seed and base layer. Adjust the pH of the medium so that it will be 7.8 to 8.0 after sterilization.

(iii) Standard solutions—Weigh accurately an amount of Erythromycin Reference Standard, equivalent to about 25 mg (potency), dissolve in 25 mL of methanol, add 0.1 mol/L phosphate buffer solution, pH 8.0 to make exactly 100 mL, and use this solution as the standard stock solution. Keep the standard stock solution at 5°C or below, and use within 7 days. Take exactly a suitable amount of the standard stock solution before use, add 0.1 mol/L phosphate buffer solution, pH 8.0 to make solutions so that each mL contains 20  $\mu$ g (potency) and 5  $\mu$ g (potency), and use these solutions as the high concentration standard solution and low concentration standard solution, respectively.


(iv) Sample solutions—Weigh accurately an amount of Erythromycin, equivalent to about 25 mg (potency), dissolve in 25 mL of methanol, and add 0.1 mol/L phosphate buffer solution, pH 8.0 to make exactly 100 mL. Take exactly a suitable amount of this solution, add 0.1 mol/L phosphate buffer solution, pH 8.0 to make solutions so that each mL contains 20  $\mu$ g (potency) and 5  $\mu$ g (potency), and use these solutions as the high concentration sample solution and low

concentration sample solution, respectively.

**Containers and storage** Containers—Well-closed containers.

## Erythromycin Ethylsuccinate

エリスロマイシンエチルコハク酸エステル



$C_{43}H_{75}NO_{16}$ : 862.05

(2*R*,3*S*,4*S*,5*R*,6*R*,8*R*,10*R*,11*R*,12*S*,13*R*)-5-[3,4,6-Trideoxy-2-*O*-(3-ethoxycarbonylpropanoyl)-3-dimethylamino-β-*D*-xylo-hexopyranosyloxy]-3-(2,6-dideoxy-3-*C*-methyl-3-*O*-methyl-α-*L*-ribo-hexopyranosyloxy)-6,11,12-trihydroxy-2,4,6,8,10,12-hexamethyl-9-oxopentadecan-13-olide [41342-53-4]

Erythromycin Ethylsuccinate is a derivative of erythromycin.

It contains not less than 780 μg (potency) and not more than 900 μg (potency) per mg, calculated on the anhydrous basis. The potency of Erythromycin Ethylsuccinate is expressed as mass (potency) of erythromycin ( $C_{37}H_{67}NO_{13}$ : 733.93).

**Description** Erythromycin Ethylsuccinate occurs as a white powder.

It is freely soluble in methanol and in acetone, soluble in ethanol (95), and practically insoluble in water.

**Identification (1)** Dissolve 3 mg of Erythromycin Ethylsuccinate in 2 mL of acetone, and add 2 mL of hydrochloric acid: an orange color develops and is immediately changed to red to deep purple.

**(2)** Determine the infrared absorption spectrum of Erythromycin Ethylsuccinate, previously dried in a desiccator (reduced pressure, silica gel) for 24 hours, as directed in the potassium bromide disk method under Infrared Spectrophotometry <2.25>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wave numbers.

**Water** <2.48> Not more than 5.0% (0.5 g, volumetric titration, direct titration).

**Assay** Perform the test according to the Cylinder-plate method as directed under Microbial Assay for Antibiotics <4.02> according to the following conditions.

(i) Test organism—*Staphylococcus aureus* ATCC 6538 P

(ii) Culture medium—Use the medium i in 3) Medium for other organisms under (1) Agar media for seed and base layer. Adjust the pH of the medium so that it will be 7.8 to

8.0 after sterilization.


(iii) Standard solutions—Weigh accurately an amount of Erythromycin Reference Standard equivalent to about 50 mg (potency), dissolve in 50 mL of methanol, add 0.1 mol/L phosphate buffer solution, pH 8.0 to make exactly 100 mL, and use this solution as the standard stock solution. Keep the standard stock solution at 5°C or below and use within 7 days. Take exactly a suitable amount of the standard stock solution before use, add 0.1 mol/L phosphate buffer solution, pH 8.0 to make solutions so that each mL contains 20 μg (potency) and 5 μg (potency), and use these solutions as the high concentration standard solution and low concentration standard solution, respectively.

(iv) Sample solutions—Weigh accurately an amount of Erythromycin Ethylsuccinate equivalent to about 50 mg (potency), dissolve in 50 mL of methanol, and add 0.1 mol/L phosphate buffer solution, pH 8.0 to make exactly 100 mL. Take exactly a suitable amount of the solution, add 0.1 mol/L phosphate buffer solution, pH 8.0 to make solutions so that each mL contains 20 μg (potency) and 5 μg (potency), and use these solutions as the high concentration sample solution and low concentration sample solution, respectively.

**Containers and storage** Containers—Tight containers.

## Erythromycin Lactobionate

エリスロマイシンラクトビオン酸塩



$C_{37}H_{67}NO_{13} \cdot C_{12}H_{22}O_{12}$ : 1092.22

(2*R*,3*S*,4*S*,5*R*,6*R*,8*R*,10*R*,11*R*,12*S*,13*R*)-5-(3,4,6-Trideoxy-3-dimethylamino-β-*D*-xylo-hexopyranosyloxy)-3-(2,6-dideoxy-3-*C*-methyl-3-*O*-methyl-α-*L*-ribo-hexopyranosyloxy)-6,11,12-trihydroxy-2,4,6,8,10,12-hexamethyl-9-oxopentadecan-13-olide mono(4-*O*-β-*D*-galactopyranosyl-D-gluconate) [3847-29-8]

Erythromycin Lactobionate is the lactobionate of erythromycin.

It contains not less than 590 μg (potency) and not more than 700 μg (potency) per mg, calculated on the anhydrous basis. The potency of Erythromycin Lactobionate is expressed as mass (potency) of erythromycin ( $C_{37}H_{67}NO_{13}$ : 733.93).

**Description** Erythromycin Lactobionate occurs as a white powder.

It is freely soluble in water, in methanol and in ethanol (99.5), and very slightly soluble in acetone.

**Identification (1)** To 3 mg of Erythromycin Lactobionate add 2 mL of acetone, and add 2 mL of hydrochloric acid: an orange color is produced, and it changes immediately to red

to deep purple.

(2) Transfer about 0.3 g of Erythromycin Lactobionate to a separator, add 15 mL of ammonia TS and 15 mL of chloroform, shake, and take the separated aqueous layer. Wash the aqueous layer with three 15-mL portions of chloroform, and evaporate the aqueous liquid on a water bath to dryness. Dissolve the residue in 10 mL of a mixture of methanol and water (3:2), and use this solution as the sample solution. Separately, dissolve 0.10 g of lactobionic acid in 10 mL of a mixture of methanol and water (3:2), and use this solution as the standard solution. Perform the test with these solutions as directed under Thin-layer Chromatography <2.03>. Spot 10  $\mu$ L each of the sample solution and standard solution on a plate of silica gel for thin-layer chromatograph. Develop the plate with the upper layer obtained from a mixture of water, 1-butanol and acetic acid (100) (3:3:1) to a distance of about 10 cm, and air-dry the plate. Spray evenly dilute sulfuric acid, and heat at 105°C for 20 minutes: the principal spot obtained from the sample solution shows a deep brown and the *R<sub>f</sub>* value which are the same as those of the principal spot from the standard solution.

**pH** <2.54> The pH of a solution obtained by dissolving 0.5 g of Erythromycin Lactobionate in 10 mL of water is between 5.0 and 7.5.

**Water** <2.48> Not more than 5.0% (0.5 g, volumetric titration, direct titration).

**Assay** Perform the test according to the Cylinder-plate method as directed under Microbial Assay for Antibiotics <4.02> according to the following conditions.

(i) Test organism—*Staphylococcus aureus* ATCC 6538 P

(ii) Culture medium—Use the medium i in 3) Medium for other organisms under (1) Agar media for seed and base layer. Adjust the pH of the medium so that it will be 7.8 to 8.0 after sterilization.


(iii) Standard solutions—Weigh accurately an amount of Erythromycin Reference Standard, equivalent to about 50 mg (potency), dissolve in 50 mL of methanol, add 0.1 mol/L phosphate buffer solution, pH 8.0 to make exactly 100 mL, and use this solution as the standard stock solution. Keep the standard stock solution at not exceeding 5°C and use within 7 days. Take exactly a suitable amount of the standard stock solution before use, add 0.1 mol/L phosphate buffer solution, pH 8.0 to make solutions so that each mL contains 20  $\mu$ g (potency) and 5  $\mu$ g (potency), and use these solutions as the high concentration standard solution and low concentration standard solution, respectively.

(iv) Sample solutions—Weigh accurately an amount of Erythromycin Lactobionate, equivalent to about 50 mg (potency), dissolve in 50 mL of methanol, and add 0.1 mol/L phosphate buffer solution, pH 8.0 to make exactly 100 mL. Take exactly a suitable amount of this solution, add 0.1 mol/L phosphate buffer solution, pH 8.0 to make solutions so that each mL contains 20  $\mu$ g (potency) and 5  $\mu$ g (potency), and use these solutions as the high concentration sample solution and low concentration sample solution, respectively.

**Containers and storage** Containers—Tight containers.

## Erythromycin Stearate

エリスロマイシンステアリン酸塩



$C_{37}H_{67}NO_{13} \cdot C_{18}H_{36}O_2$ : 1018.40

(2*R*,3*S*,4*S*,5*R*,6*R*,8*R*,10*R*,11*R*,12*S*,13*R*)-5-(3,4,6-Trideoxy-3-dimethylamino- $\beta$ -D-xyllo-hexopyranosyloxy)-3-(2,6-dideoxy-3-*C*-methyl-3-*O*-methyl- $\alpha$ -L-ribo-hexopyranosyloxy)-6,11,12-trihydroxy-2,4,6,8,10,12-hexamethyl-9-oxopentadecan-13-olide monostearate [643-22-1]

Erythromycin Stearate is the stearate of erythromycin.

It contains not less than 600  $\mu$ g (potency) and not more than 720  $\mu$ g (potency) per mg, calculated on the anhydrous basis. The potency of Erythromycin Stearate is expressed as mass (potency) of erythromycin ( $C_{37}H_{67}NO_{13}$ : 733.93).

**Description** Erythromycin Stearate occurs as a white powder.

It is freely soluble in ethanol (95) and in acetone, soluble in methanol, and practically insoluble in water.

**Identification** (1) Dissolve 3 mg of Erythromycin Stearate in 2 mL of acetone, and add 2 mL of hydrochloric acid: an orange color develops and is immediately changed to red to deep purple.

(2) Determine the infrared absorption spectrum of Erythromycin Stearate, previously dried in a desiccator (reduced pressure, silica gel) for 24 hours, as directed in the potassium bromide disk method under Infrared Spectrophotometry <2.25>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wave numbers.

**Water** <2.48> Not more than 5.0% (0.5 g, volumetric titration, direct titration).

**Assay** Perform the test according to the Cylinder-plate method as directed under Microbial Assay for Antibiotics <4.02> according to the following conditions.

(i) Test organism—*Staphylococcus aureus* ATCC 6538 P

(ii) Culture medium—Use the medium i in 3) Medium for other organisms under (1) Agar media for seed and base layer. Adjust the pH of the medium so that it will be 7.8 to 8.0 after sterilization.

(iii) Standard solutions—Weigh accurately an amount of Erythromycin Reference Standard equivalent to about 50 mg


(potency), dissolve in 50 mL of methanol, add 0.1 mol/L phosphate buffer solution, pH 8.0 to make exactly 100 mL, and use this solution as the standard stock solution. Keep the standard stock solution at 5°C or below and use within 7 days. Take exactly a suitable amount of the standard stock solution before use, add 0.1 mol/L phosphate buffer solution, pH 8.0 to make solutions so that each mL contains 20 µg (potency) and 5 µg (potency), and use these solutions as the high concentration standard solution and low concentration standard solution, respectively.

(iv) Sample solutions—Weigh accurately an amount of Erythromycin Stearate equivalent to about 50 mg (potency), dissolve in 50 mL of methanol, and add 0.1 mol/L phosphate buffer solution, pH 8.0 to make exactly 100 mL. Take exactly a suitable amount of the solution, add 0.1 mol/L phosphate buffer solution, pH 8.0 to make solutions so that each mL contains 20 µg (potency) and 5 µg (potency), and use these solutions as the high concentration sample solution and low concentration sample solution, respectively.

**Containers and storage** Containers—Tight containers.

## Estazolam

エスタゾラム



$C_{16}H_{11}ClN_4$ : 294.74

8-Chloro-6-phenyl-4H-[1,2,4]triazolo[4,3-a][1,4]benzodiazepine [29975-16-4]

Estazolam, when dried, contains not less than 98.5% of  $C_{16}H_{11}ClN_4$ .

**Description** Estazolam occurs as white to pale yellowish white crystals or crystalline powder. It is odorless, and has a bitter taste.

It is soluble in methanol and in acetic anhydride, sparingly soluble in ethanol (95), and practically insoluble in water and in diethyl ether.

**Identification** (1) Dissolve 0.01 g of Estazolam in 3 mL of sulfuric acid: the solution shows a yellow-green fluorescence under ultraviolet light (main wavelength: 365 nm).

(2) Determine the absorption spectrum of a solution of Estazolam in 1 mol/L hydrochloric acid TS (1 in 100,000) as directed under Ultraviolet-visible Spectrophotometry <2.24>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wavelengths.

(3) Perform the test with Estazolam as directed under Flame Coloration Test (2) <1.04>: a green color appears.

**Melting point** <2.60> 229 – 233°C

**Purity** (1) Clarity and color of solution—Dissolve 0.10 g of Estazolam in 10 mL of ethanol (95): the solution is clear

and colorless.

(2) Chloride <1.03>—Dissolve 1.0 g of Estazolam in 10 mL of ethanol (95) by heating, add 40 mL of water, cool with shaking in ice water, allow to stand to attain ordinary temperature, and filter. To 30 mL of the filtrate add 6 mL of dilute nitric acid and water to make 50 mL, and perform the test using this solution as the test solution. Prepare the control solution with 0.25 mL of 0.01 mol/L hydrochloric acid VS and 6 mL of ethanol (95) (not more than 0.015%).

(3) Heavy metals <1.07>—Proceed with 1.0 g of Estazolam according to Method 2, and perform the test. Prepare the control solution with 2.0 mL of Standard Lead Solution (not more than 20 ppm).

(4) Arsenic <1.11>—Prepare the test solution with 1.0 g of Estazolam according to Method 3, and perform the test (not more than 2 ppm).

(5) Related substances—Dissolve 0.20 g of Estazolam in 10 mL of methanol, and use this solution as the sample solution. Pipet 1 mL of the sample solution, add methanol to make exactly 200 mL, and use this solution as the standard solution. Perform the test with these solutions as directed under Thin-layer Chromatography <2.03>. Spot 10 µL each of the sample solution and standard solution on a plate of silica gel with fluorescent indicator for thin-layer chromatography. Develop the plate with a mixture of hexane, chloroform and methanol (5:3:1) to a distance of about 10 cm, and air-dry the plate. Examine the plate under ultraviolet light (main wavelength: 254 nm): the spots other than the principal spot from the sample solution are not more intense than the principal spot from the standard solution.

**Loss on drying** <2.41> Not more than 1.0% (1 g, 105°C, 4 hours).

**Residue on ignition** <2.44> Not more than 0.1% (2 g).


**Assay** Weigh accurately about 0.25 g of Estazolam, previously dried, dissolve in 100 mL of acetic anhydride, and titrate <2.50> with 0.1 mol/L perchloric acid VS (potentiometric titration), until the solution changes to the second equivalence point. Perform a blank determination, and make any necessary correction.

Each mL of 0.1 mol/L perchloric acid VS  
= 14.74 mg of  $C_{16}H_{11}ClN_4$

**Containers and storage** Containers—Well-closed containers.

## Estradiol Benzoate

エストラジオール安息香酸エステル



$C_{25}H_{28}O_3$ : 376.49

Estra-1,3,5(10)-triene-3,17β-diol 3-benzoate [50-50-0]

Estradiol Benzoate, when dried, contains not less

than 97.0% of  $C_{25}H_{28}O_3$ .

**Description** Estradiol Benzoate occurs as a white, crystalline powder. It is odorless.

It is sparingly soluble in acetone, slightly soluble in methanol, in ethanol (95) and in diethyl ether, and practically insoluble in water.

**Identification (1)** To 2 mg of Estradiol Benzoate add 2 mL of sulfuric acid: a yellowish green color with a blue fluorescence is produced, and the color of the solution changes to light orange on the careful addition of 2 mL of water.

(2) Determine the infrared absorption spectrum of Estradiol Benzoate, previously dried, as directed in the potassium bromide disk method under Infrared Spectrophotometry <2.25>, and compare the spectrum with the Reference Spectrum or the spectrum of dried Estradiol Benzoate Reference Standard: both spectra exhibit similar intensities of absorption at the same wave numbers.

**Optical rotation** <2.49>  $[\alpha]_D^{20}$ : +54 – +58° (after drying, 0.1 g, acetone, 10 mL, 100 mm).

**Melting point** <2.60> 191 – 198°C

**Purity (1)** 3,17 $\alpha$ -Estradiol—Dissolve 5.0 mg each of Estradiol Benzoate and Estradiol Benzoate Reference Standard in acetone to make exactly 100 mL, and use these solutions as the sample solution and standard solution, respectively. Place exactly 2 mL each of the sample solution and standard solution in separate glass-stoppered test tube, add boiling stones, evaporate the acetone by heating in a water bath, and dry the residue in a desiccator (in vacuum, phosphorus (V) oxide) for 1 hour. Add 1.0 mL of dilute iron-phenol TS to each test tube. Stopper the test tubes loosely, heat for 30 seconds in a water bath, shake in a water bath for several seconds, and heat for 2 minutes. Cool the solutions in ice for 2 minutes, add 4.0 mL of diluted sulfuric acid (7 in 20), and mix well: the solution obtained from the sample solution has no more color than that from the standard solution.

(2) Related substances—Dissolve 40 mg of Estradiol Benzoate in 2 mL of acetone, and use this solution as the sample solution. Pipet 1 mL of this solution, add acetone to make exactly 100 mL, and use this solution as the standard solution. Perform the test with these solutions as directed under Thin-layer Chromatography <2.03>. Spot 10  $\mu$ L each of the sample solution and standard solution on a plate of silica gel with fluorescent indicator for thin-layer chromatography. Develop the plate with a mixture of chloroform and diethylamine (19:1) to a distance of about 15 cm, and air-dry the plate. Examine under ultraviolet light (main wavelength: 254 nm): the spots other than the principal spot from the sample solution are not more intense than the spot from the standard solution.

**Loss on drying** <2.41> Not more than 0.5% (0.5 g, in vacuum, phosphorus (V) oxide, 4 hours).

**Residue on ignition** <2.44> Not more than 0.2% (0.1 g).

**Assay** Weigh accurately about 10 mg each of Estradiol Benzoate and Estradiol Benzoate Reference Standard, previously dried, and dissolve each in methanol to make exactly 20 mL. Pipet 5 mL each of these solutions, add 5 mL of the internal standard solution, then add methanol to make 20 mL, and use these solutions as the sample solution and standard solu-

tion, respectively. Perform the test with 5  $\mu$ L of the sample solution and standard solution as directed under Liquid Chromatography <2.01> according to the following conditions, and calculate the ratios,  $Q_T$  and  $Q_S$ , of the peak area of estradiol benzoate to that of the internal standard.

$$\text{Amount (mg) of } C_{25}H_{28}O_3 = W_S \times (Q_T/Q_S)$$

$W_S$ : Amount (mg) of Estradiol Benzoate Reference Standard

**Internal standard solution**—A solution of progesterone in methanol (13 in 80,000).

**Operating conditions**—

Detector: An ultraviolet absorption photometer (wavelength: 230 nm).

Column: A stainless steel column 4.6 mm in inside diameter and 15 cm in length, packed with octadecylsilanized silica gel for liquid chromatography (5  $\mu$ m in particle diameter).

Column temperature: A constant temperature of about 35°C.

Mobile phase: A mixture of acetonitrile and water (7:3).

Flow rate: Adjust the flow rate so that the retention time of estradiol benzoate is about 10 minutes.

**System suitability**—

System performance: When the procedure is run with 5  $\mu$ L of the standard solution under the above operating conditions, the internal standard and estradiol benzoate are eluted in this order with the resolution between these peaks being not less than 9.

System repeatability: When the test is repeated 6 times with 5  $\mu$ L of the standard solution under the above operating conditions, the relative standard deviation of the ratios of the peak area of estradiol benzoate to that of the internal standard is not more than 1.0%.

**Containers and storage** Containers—Tight containers.

Storage—Light-resistant.

## Estradiol Benzoate Injection

エストラジオール安息香酸エステル注射液

Estradiol Benzoate Injection is an oily solution for injection.

It contains not less than 90% and not more than 110% of the labeled amount of estradiol benzoate ( $C_{25}H_{28}O_3$ ; 376.49).

**Method of preparation** Prepare as directed under Injections, with Estradiol Benzoate.

**Description** Estradiol Benzoate Injection is a clear, oily liquid.

**Identification** To a volume of Estradiol Benzoate Injection, equivalent to 1 mg of Estradiol Benzoate according to the labeled amount, add chloroform to make 5 mL, and use this solution as the sample solution. Separately dissolve 1 mg of Estradiol Benzoate Reference Standard in 5 mL of chloroform, and use this solution as the standard solution. Perform the test with these solutions as directed under Thin-layer Chromatography <2.03>. Spot 50  $\mu$ L each of the sample so-



lution and standard solution on a plate of silica gel with fluorescent indicator for thin-layer chromatography. Develop the plate with dichloromethane to a distance of about 15 cm, and air-dry the plate. Then develop the plate with a mixture of chloroform and methanol (99:1) to a distance of about 15 cm, and air-dry the plate. Examine under ultraviolet light (main wavelength: 254 nm): the principal spot obtained from the sample solution and the spot obtained from the standard solution show the same *R<sub>f</sub>* value.

**Extractable volume** <6.05> It meets the requirement.

**Assay** Transfer an exactly measured volume of Estradiol Benzoate Injection, equivalent to about 10 mg of estradiol benzoate ( $C_{25}H_{28}O_3$ ), to a separator, add 30 mL of hexane saturated with diluted methanol (9 in 10), and extract with five 15-mL portions of dilute methanol (9 in 10) saturated with hexane. Filter the extract through a filter paper washed with 10 mL of diluted methanol (9 in 10), to the filtrate add methanol to make exactly 200 mL, and use this solution as the sample solution. Separately, weigh accurately about 25 mg of Estradiol Benzoate Reference Standard, previously dried in a desiccator (in vacuum, phosphorus (V) oxide) for 4 hours, and dissolve in methanol to make exactly 100 mL. Pipet 10 mL of this solution, add methanol to make exactly 50 mL, and use this solution as the standard solution. Transfer 2 mL each of the sample solution and standard solution, exactly measured, to light-resistant 20-mL volumetric flasks, and evaporate to dryness on a water bath with the aid of a current of air. Dissolve the residue in 1 mL of methanol, add 10 mL of boric acid-methanol buffer solution, shake, and boil under a reflux condenser for 30 minutes. Cool, add 5 mL of boric acid-methanol buffer solution, shake, and cool with ice. To each solution add 2 mL of ice-cold diazo TS quickly, shake vigorously, add 2 mL of sodium hydroxide TS, then add water to make 20 mL, and filter after shaking. Discard the first 3 mL of the filtrate, and perform the test with the subsequent filtrate as directed under Ultraviolet-visible Spectrophotometry <2.24> using a solution, prepared with 2 mL of methanol in the same manner, as the blank. Determine the absorbances,  $A_T$  and  $A_S$ , of the subsequent solutions obtained from the sample solution and standard solution in a 4-cm cell at 490 nm, respectively.

$$\begin{aligned} &\text{Amount (mg) of estradiol benzoate } (C_{25}H_{28}O_3) \\ &= W_S \times (A_T/A_S) \times (2/5) \end{aligned}$$

$W_S$ : Amount (mg) of Estradiol Benzoate Reference Standard

**Containers and storage** Containers—Hermetic containers.

## Estradiol Benzoate Injection (Aqueous Suspension)

エストラジオール安息香酸エステル水性懸濁注射液

Estradiol Benzoate Injection (Aqueous Suspension) is an aqueous suspension for injection. It contains not less than 90% and not more than 110% of the labeled amount of estradiol benzoate ( $C_{25}H_{28}O_3$ ; 376.49).

**Method of preparation** Prepare as directed under Injection,

with Estradiol Benzoate.

**Description** Estradiol Benzoate Injection (Aqueous Suspension) produces a white turbidity on shaking.

**Identification** Extract a volume of Estradiol Benzoate Injection (Aqueous Suspension), equivalent to 1 mg of Estradiol Benzoate according to the labeled amount, with 5 mL of chloroform, and use this extract as the sample solution. Separately, dissolve 1 mg of Estradiol Benzoate Reference Standard in 5 mL of chloroform, and use this solution as the standard solution. Perform the test with these solutions as directed under Thin-layer Chromatography <2.03>. Spot 50  $\mu$ L each of the sample solution and standard solution on a plate of silica gel with fluorescent indicator for thin-layer chromatography. Develop the plate with a mixture of chloroform and methanol (99:1) to a distance of about 15 cm, and air-dry the plate. Examine under ultraviolet light (main wavelength: 254 nm): the principal spot obtained from the sample solution and the spot obtained from the standard solution show the same *R<sub>f</sub>* value.

**Extractable volume** <6.05> It meets the requirement.

**Assay** Measure exactly a volume of well-mixed Estradiol Benzoate Injection (Aqueous Suspension), equivalent to about 2 mg of estradiol benzoate ( $C_{25}H_{28}O_3$ ), dissolve the crystals with an appropriate quantity of methanol, and add methanol to make exactly 20 mL. Pipet 10 mL of this solution, add exactly 10 mL of the internal standard solution, add methanol to make 100 mL, and use this solution as the sample solution. Separately, weigh accurately about 10 mg of Estradiol Benzoate Reference Standard, previously dried in desiccator (reduced pressure, phosphorus (V) oxide) for 4 hours, and dissolve in methanol to make exactly 100 mL. Pipet 10 mL of this solution, add exactly 10 mL of the internal standard solution and methanol to make 100 mL, and use this solution as the standard solution. Proceed with these solutions as directed in the Assay under Estradiol Benzoate.

$$\begin{aligned} &\text{Amount (mg) of estradiol benzoate } (C_{25}H_{28}O_3) \\ &= W_S \times (Q_T/Q_S) \times (1/5) \end{aligned}$$


$W_S$ : Amount (mg) of Estradiol Benzoate Reference Standard

**Internal standard solution**—A solution of progesterone in methanol (13 in 100,000).

**Containers and storage** Containers—Hermetic containers.

## Estradiol

エストリオール



$C_{18}H_{24}O_3$ ; 288.38

Estra-1,3,5(10)-triene-3,16 $\alpha$ ,17 $\beta$ -triol [50-27-1]

Estradiol, when dried, contains not less than 97.0% and not more than 102.0% of  $C_{18}H_{24}O_3$ .

**Description** Estriol occurs as a white, crystalline powder. It is odorless.

It is sparingly soluble in methanol, slightly soluble in ethanol (95) and in 1,4-dioxane, and practically insoluble in water and in diethyl ether.

**Identification** (1) Dissolve 0.01 g of Estriol in 100 mL of ethanol (95) by warming, and use this solution as the sample solution. Evaporate 1 mL of this solution on a water bath to dryness, add 5 mL of a solution of sodium *p*-phenolsulfonate in diluted phosphoric acid (1 in 50), heat at 150°C for 10 minutes, and cool: a red-purple color develops.

(2) Determine the absorption spectrum of the sample solution obtained in (1) as directed under Ultraviolet-visible Spectrophotometry <2.24>, and compare the spectrum with the Reference Spectrum or the spectrum of a solution of Estriol Reference Standard prepared in the same manner as the sample solution: both spectra exhibit similar intensities of absorption at the same wavelengths.

(3) Determine the infrared absorption spectrum of Estriol, previously dried, as directed in the potassium bromide disk method under Infrared Spectrophotometry <2.25>, and compare the spectrum with the Reference Spectrum or the spectrum of previously dried Estriol Reference Standard: both spectra exhibit similar intensities of absorption at the same wave numbers.

**Optical rotation** <2.49>  $[\alpha]_D^{20}$ : +54 – +62° (after drying, 40 mg, 1,4-dioxane, 10 mL, 100 mm).

**Melting point** <2.60> 281 – 286°C

**Purity** (1) Heavy metals <1.07>—Proceed with 1.0 g of Estriol according to Method 2, and perform the test. Prepare the control solution with 2.0 mL of Standard Lead Solution (not more than 20 ppm).

(2) Related substances—Dissolve 40 mg of Estriol in 10 mL of ethanol (95) by warming, and use this solution as the sample solution. Pipet 1 mL of the sample solution, add ethanol (95) to make exactly 100 mL, and use this solution as the standard solution. Perform the test with these solutions as directed under Thin-layer Chromatography <2.03>. Spot 5  $\mu$ L each of the sample solution and standard solution on a plate of silica gel for thin-layer chromatography. Develop the plate with a mixture of chloroform, methanol, acetone and acetic acid (100) (18:1:1:1) to a distance of about 15 cm, and air-dry the plate. Spray evenly diluted sulfuric acid (1 in 2) on the plate, and heat at 105°C for 15 minutes: the spots other than the principal spot from the sample solution are not more intense than the spot from the standard solution.

**Loss on drying** <2.41> Not more than 0.5% (0.5 g, 105°C, 3 hours).

**Residue on ignition** <2.44> Not more than 0.1% (0.5 g).

**Assay** Weigh accurately about 25 mg each of Estriol and Estriol Reference Standard, previously dried, and dissolve each in methanol to make exactly 50 mL. Pipet 10 mL each of these solutions, add exactly 5 mL of the internal standard solution, add methanol to make 100 mL, and use these solutions as the sample solution and the standard solution, respectively. Perform the test with 10  $\mu$ L each of the sample solution and standard solution as directed under Liquid Chromatography <2.01> according to the following conditions, and calculate the ratios,  $Q_T$  and  $Q_S$ , of the peak area of

estriol to that of the internal standard, respectively.

$$\text{Amount (mg) of } C_{18}H_{24}O_3 = W_S \times (Q_T/Q_S)$$

$W_S$ : Amount (mg) of Estriol Reference Standard

**Internal standard solution**—A solution of methyl benzoate for estriol limit test in methanol (1 in 1000).

**Operating conditions**—

Detector: An ultraviolet absorption photometer (wavelength: 280 nm).

Column: A stainless steel column 4.6 mm in inside diameter and 15 cm in length, packed with octadecylsilanized silica gel for liquid chromatography (5  $\mu$ m in particle diameter).

Column temperature: A constant temperature of about 25°C.

Mobile phase: A mixture of water and methanol (51:49).

Flow rate: Adjust the flow rate so that the retention time of estriol is about 10 minutes.

**System suitability**—

System performance: When the procedure is run with 10  $\mu$ L of the standard solution under the above operating conditions, estriol and the internal standard are eluted in this order with the resolution between these peaks being not less than 8.

System repeatability: When the test is repeated 6 times with 10  $\mu$ L of the standard solution under the above operating conditions, the relative standard deviation of the ratios of the peak area of estriol to that of the internal standard is not more than 1.0%.

**Containers and storage** Containers—Tight containers.

## Estriol Injection (Aqueous Suspension)

エストリオール水性懸濁注射液

Estriol Injection (Aqueous Suspension) is an aqueous suspension for injection. It contains not less than 90% and not more than 110% of the labeled amount of estriol ( $C_{18}H_{24}O_3$ : 288.38).

**Method of preparation** Prepare as directed under Injections, with Estriol.

**Description** Shake Estriol Injection (Aqueous Suspension): a white turbidity is produced.

**Identification** (1) Shake well, take a volume of Estriol Injection (Aqueous Suspension), equivalent to 2 mg of Estriol according to the labeled amount, add ethanol (95) to make 20 mL, and use this solution as the sample solution. Proceed with the sample solution as directed in the Identification (1) under Estriol.

(2) Determine the absorption spectrum of the sample solution obtained in (1) as directed under Ultraviolet-visible Spectrophotometry <2.24>: it exhibits a maximum between 279 nm and 283 nm.

**Extractable volume** <6.05> It meets the requirement.

**Assay** Shake well, pipet a volume of Estriol Injection (Aqueous Suspension), equivalent to about 5 mg of estriol

(C<sub>18</sub>H<sub>24</sub>O<sub>3</sub>), and dissolve in methanol to make exactly 20 mL. Pipet 4 mL of this solution, add exactly 5 mL of the internal standard solution, then add methanol to make 50 mL, and use this solution as the sample solution. Separately, weigh accurately about 25 mg of Estriol Reference Standard, previously dried at 105°C for 3 hours, and dissolve in methanol to make exactly 100 mL. Pipet 4 mL of this solution, add exactly 5 mL of the internal standard solution, then add methanol to make 50 mL, and use this solution as the standard solution. Proceed as directed in the Assay under Estriol.

$$\begin{aligned} &\text{Amount (mg) of estriol (C}_{18}\text{H}_{24}\text{O}_3\text{)} \\ &= W_S \times (Q_T/Q_S) \times (1/5) \end{aligned}$$

$W_S$ : Amount (mg) of Estriol Reference Standard

**Internal standard solution**—A solution of methyl benzoate for estriol limit test in ethanol (95) (1 in 5000).

**Containers and storage** Containers—Hermetic containers.

## Estriol Tablets

エストリオール錠

Estriol Tablets contain not less than 90% and not more than 110% of the labeled amount of estriol (C<sub>18</sub>H<sub>24</sub>O<sub>3</sub>; 288.38).

**Method of preparation** Prepare as directed under Tablets, with Estriol.

**Identification (1)** Weigh a portion of powdered Estriol Tablets, equivalent to 2 mg of Estriol according to the labeled amount, add ethanol (95) to make 20 mL, shake for 10 minutes, centrifuge, and use the supernatant liquid as the sample solution. Proceed with the sample solution as directed in the Identification (1) under Estriol.

(2) Determine the absorption spectrum of the sample solution obtained in (1) as directed under Ultraviolet-visible Spectrophotometry <2.24>: it exhibits a maximum between 279 nm and 283 nm.

**Uniformity of dosage units <6.02>** Perform the test according to the following method: it meets the requirement of the Content uniformity test.

To 1 tablet of Estriol Tablets add exactly 5 mL of water, disperse the fine particles with ultrasonic wave, add exactly 15 mL of methanol, and shake for 15 minutes. Centrifuge this solution for 10 minutes, pipet a definite amount of the supernatant liquid, and add methanol to make exactly a definite amount of solution so that each mL of the solution contains about 5 µg of estriol (C<sub>18</sub>H<sub>24</sub>O<sub>3</sub>). Pipet 5 mL of this solution, add exactly 1 mL of the internal standard solution, and use this solution as the sample solution. Proceed with 20 µL of the sample solution as directed in the Assay under Estriol. Use a solution of methyl benzoate in methanol (1 in 40,000) as the internal standard solution. Calculate the mean value from each ratio of peak areas of 10 samples: the samples conform to the requirements if the deviation (%) of the mean value and each ratio of peak areas is within 15%. If the deviation (%) exceeds 15%, and 1 sample shows deviation within 25%, repeat the test with 20 samples. Calculate the deviation (%) of the mean value from each ratio of peak areas of the 30

samples used in the 2 tests and each ratio of peak areas: the samples conform to the requirements if the deviation exceeds 15%, not more than 1 sample shows deviation within 25%, and no sample shows deviation exceeding 25%.

**Dissolution <6.10>** Perform the test according to the following method: it meets the requirement.

Perform the test with 1 tablet of Estriol Tablets at 50 revolutions per minute according to the Paddle method, using 900 mL of water as the dissolution medium. Take 20 mL or more of the dissolved solution 30 minutes after starting the test, and filter through a membrane filter with pore size of not more than 0.8 µm. Discard the first 10 mL of the filtrate, pipet the subsequent  $V$  mL, add water to make exactly  $V'$  mL so that each mL contains about 0.1 µg of estriol (C<sub>18</sub>H<sub>24</sub>O<sub>3</sub>) according to the labeled amount, and use this solution as the sample solution. Separately, weigh accurately about 10 mg of Estriol Reference Standard, previously dried at 105°C for 3 hours, dissolve in methanol to make exactly 100 mL, then pipet 5 mL of this solution, and add water to make exactly 100 mL. Pipet 2 mL of this solution, add water to make exactly 100 mL, and use this solution as the standard solution. Perform the test with exactly 100 µL each of the sample solution and standard solution according to the operating conditions as directed in the Assay under Estriol, and determine the peak areas of estriol,  $A_T$  and  $A_S$ , from these solutions.

The dissolution rate of Estriol Tablets in 30 minutes is not less than 80%.

$$\begin{aligned} &\text{Dissolution rate (\%)} \text{ with respect to} \\ &\text{the labeled amount of estriol (C}_{18}\text{H}_{24}\text{O}_3\text{)} \\ &= W_S \times (A_T/A_S) \times (V'/V) \times (1/C) \times (9/10) \end{aligned}$$

$W_S$ : Amount (mg) of Estriol Reference Standard

$C$ : Labeled amount (mg) of estriol (C<sub>18</sub>H<sub>24</sub>O<sub>3</sub>) in 1 tablet

**Assay** Weigh accurately and powder not less than 20 Estriol Tablets. Weigh accurately a portion of the powder, equivalent to about 1 mg of estriol (C<sub>18</sub>H<sub>24</sub>O<sub>3</sub>), add exactly 5 mL of water, disperse the fine particles with ultrasonic wave, shake with 25 mL of methanol for 10 minutes, centrifuge, and take the supernatant liquid. Add 25 mL of methanol, repeat the above procedure twice, combine the supernatant liquid, add exactly 5 mL of the internal standard solution, then add methanol to make 100 mL, and use this solution as the sample solution. Separately, weigh accurately about 25 mg of Estriol Reference Standard, previously dried at 105°C for 3 hours, and dissolve in methanol to make exactly 100 mL. Pipet 4 mL of this solution, add exactly 5 mL of the internal standard solution, then add methanol to make 100 mL, and use this solution as the standard solution. Proceed with 20 µL each of the sample solution and standard solution as directed in the Assay under Estriol.

$$\begin{aligned} &\text{Amount (mg) of estriol (C}_{18}\text{H}_{24}\text{O}_3\text{)} \\ &= W_S \times (Q_T/Q_S) \times (1/25) \end{aligned}$$


$W_S$ : Amount (mg) of Estriol Reference Standard

**Internal standard solution**—A solution of methyl benzoate for estriol limit test in methanol (1 in 5000).

**Containers and storage** Containers—Tight containers.

## Etacrynic Acid

エタクリン酸



$C_{13}H_{12}Cl_2O_4$ : 303.14  
[2,3-Dichloro-4-(2-ethylacryloyl)phenoxy]acetic acid  
[58-54-8]

Etacrynic Acid, when dried, contains not less than 98.0% of  $C_{13}H_{12}Cl_2O_4$ .

**Description** Etacrynic Acid occurs as a white, crystalline powder. It is odorless, and has a slightly bitter taste.

It is very soluble in methanol, freely soluble in ethanol (95), in acetic acid (100) and in diethyl ether, and very slightly soluble in water.

**Identification (1)** Dissolve 0.2 g of Etacrynic Acid in 10 mL of acetic acid (100), and to 5 mL of this solution add 0.1 mL of bromine TS: the color of the test solution disappears. To the remaining 5 mL of the solution add 0.1 mL of potassium permanganate TS: the color of the test solution changes to light orange immediately.

(2) To 0.01 g of Etacrynic Acid add 1 mL of sodium hydroxide TS, and heat in a water bath for 3 minutes. After cooling, add 1 mL of disodium chlorotriphosphate TS, and heat in a water bath for 10 minutes: a deep purple color develops.

(3) Determine the absorption spectrum of a solution of Etacrynic Acid in methanol (1 in 20,000) as directed under Ultraviolet-visible Spectrophotometry <2.24>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wavelengths.

(4) Perform the test with Etacrynic Acid as directed under Flame Coloration Test (2) <1.04>: a green color appears.

**Melting point** <2.60> 121 – 125°C

**Purity (1)** Clarity and color of solution—Dissolve 1.0 g of Etacrynic Acid in 10 mL of methanol: the solution is clear and colorless.

(2) Heavy metals <1.07>—Proceed with 1.0 g of Etacrynic Acid according to Method 4, and perform the test. Prepare the control solution with 2.0 mL of Standard Lead Solution (not more than 20 ppm).

(3) Arsenic <1.11>—Prepare the test solution with 1.0 g of Etacrynic Acid according to Method 3, and perform the test. Add 10 mL of a solution of magnesium nitrate hexahydrate in ethanol (95) (1 in 50), then add 1.5 mL of hydrogen peroxide (30), and fire to burn (not more than 2 ppm).

(4) Related substances—Dissolve 0.20 g of Etacrynic Acid in 10 mL of ethanol (95), and use this solution as the sample solution. Pipet 3 mL of the sample solution, add ethanol (95) to make exactly 200 mL, and use this solution as the standard solution. Perform the test with these solutions as directed under Thin-layer Chromatography <2.03>. Spot 10  $\mu$ L each of the sample solution and standard solution on a plate of silica gel with fluorescent indicator for thin-layer

chromatography. Develop the plate with a mixture of chloroform, ethyl acetate and acetic acid (100) (6:5:2) to a distance of about 15 cm, and air-dry the plate. Examine under ultraviolet light (main wavelength: 254 nm): the spots other than the principal spot from the sample solution are not more intense than the spot from the standard solution.

**Loss on drying** <2.41> Not more than 0.25% (1 g, in vacuum, 60°C, 2 hours).

**Residue on ignition** <2.44> Not more than 0.1% (1 g).

**Assay** Weigh accurately about 0.1 g of Etacrynic Acid, previously dried, place in an iodine bottle, dissolve in 20 mL of acetic acid (100), and add exactly 20 mL of 0.05 mol/L bromine VS. To this solution add 3 mL of hydrochloric acid, stopper tightly at once, shake, and allow to stand in a dark place for 60 minutes. Add carefully 50 mL of water and 15 mL of potassium iodide TS, stopper tightly at once, shake well, and titrate <2.50> the liberated iodine with 0.1 mol/L sodium thiosulfate VS (indicator: 1 mL of starch TS). Perform a blank determination.

Each mL of 0.05 mol/L bromine VS  
= 15.16 mg of  $C_{13}H_{12}Cl_2O_4$

**Containers and storage** Containers—Well-closed containers.

## Etacrynic Acid Tablets

エタクリン酸錠

Etacrynic Acid Tablets contain not less than 90% and not more than 110% of the labeled amount of etacrynic acid ( $C_{13}H_{12}Cl_2O_4$ : 303.14).

**Method of preparation** Prepare as directed under Tablets, with Etacrynic Acid.

**Identification (1)** Weigh a quantity of powdered Etacrynic Acid Tablets, equivalent to 0.3 g of Etacrynic Acid according to the labeled amount, add 25 mL of 0.1 mol/L hydrochloric acid TS, and extract with 50 mL of Dichloromethane. Filter the dichloromethane extract, and evaporate the filtrate on a water bath to dryness. Proceed with the residue as directed in the Identification (1), (2) and (4) under Etacrynic Acid.

(2) Prepare a solution of the residue obtained in (1), equivalent to a solution of Etacrynic Acid in methanol (1 in 20,000), and determine the absorption spectrum as directed under Ultraviolet-visible Spectrophotometry <2.24>: it exhibits a maximum between 268 nm and 272 nm.

**Dissolution** <6.10> Perform the test according to the following method: it meets the requirement.

Perform the test with 1 tablet of Etacrynic Acid Tablets at 50 revolutions per minute according to the Paddle method, using 900 mL of water as the test solution. Take 20 mL or more of the dissolved solution 45 minutes after starting the test, and filter through a membrane filter with pore size of not more than 0.8  $\mu$ m. Discard the first 10 mL of the filtrate, and use the subsequent as the sample solution. Separately, weigh accurately about 55 mg of etacrynic acid for assay, previously dried at 60°C for 2 hours, dissolve in 10 mL of methanol, and add water to make exactly 100 mL. Pipet 5 mL of this so-

lution, add water to make exactly 100 mL, and use this solution as the standard solution. Determine the absorbances,  $A_T$  and  $A_S$ , of the sample solution and standard solution at 277 nm as directed under Ultraviolet-visible Spectrophotometry <2.24>.

The dissolution rate of Etacrylic Acid Tablets in 45 minutes is not less than 70%.

$$\begin{aligned} & \text{Dissolution rate (\%)} \text{ with respect to} \\ & \text{the labeled amount of etacrylic acid (C}_{13}\text{H}_{12}\text{Cl}_2\text{O}_4\text{)} \\ & = W_S \times (A_T/A_S) \times (1/C) \times 45 \end{aligned}$$

$W_S$ : Amount (mg) of etacrylic acid for assay

$C$ : Labeled amount (mg) of etacrylic acid (C<sub>13</sub>H<sub>12</sub>Cl<sub>2</sub>O<sub>4</sub>) in 1 tablet


**Assay** Weigh accurately and powder not less than 20 Etacrylic Acid Tablets. Weigh accurately a portion of the powder, equivalent to about 0.1 g of etacrylic acid (C<sub>13</sub>H<sub>12</sub>Cl<sub>2</sub>O<sub>4</sub>), add 25 mL of 0.1 mol/L hydrochloric acid TS, and extract with three 30-mL portions of dichloromethane. Filter the dichloromethane extracts through a pledget of absorbent cotton into an iodine bottle. Wash the pledget of absorbent cotton with a small amount of dichloromethane, and combine the washing with the extracts. Evaporate this solution on a water bath to dryness in a current of air, to the residue add 20 mL of acetic acid (100), and proceed as directed in the Assay under Etacrylic Acid.

$$\begin{aligned} & \text{Each mL of 0.05 mol/L bromine VS} \\ & = 15.16 \text{ mg of C}_{13}\text{H}_{12}\text{Cl}_2\text{O}_4 \end{aligned}$$

**Containers and storage** Containers—Well-closed containers.

## Ethambutol Hydrochloride

エタンブトール塩酸塩



C<sub>10</sub>H<sub>24</sub>N<sub>2</sub>O<sub>2</sub>·2HCl: 277.23

2,2'-(Ethylenediimino)bis[(2S)-butan-1-ol] dihydrochloride  
[1070-11-7]

Ethambutol Hydrochloride, when dried, contains not less than 98.5% of C<sub>10</sub>H<sub>24</sub>N<sub>2</sub>O<sub>2</sub>·2HCl.

**Description** Ethambutol Hydrochloride occurs as white crystals or crystalline powder. It is odorless, and has a bitter taste.

It is very soluble in water, soluble in methanol and in ethanol (95), and practically insoluble in diethyl ether.

The pH of a solution of Ethambutol Hydrochloride (1 in 20) is between 3.4 and 4.0.

**Identification** (1) To 10 mL of a solution of Ethambutol Hydrochloride (1 in 100) add 0.5 mL of copper (II) sulfate TS and 2 mL of sodium hydroxide TS: a deep blue color is produced.

(2) Dissolve 0.1 g of Ethambutol Hydrochloride in 40 mL

of water, add 20 mL of 2,4,6-trinitrophenol TS, and allow to stand for 1 hour. Collect the precipitate, wash with 50 mL of water, and dry at 105°C for 2 hours: the precipitate melts <2.60> between 193°C and 197°C.

(3) A solution of Ethambutol Hydrochloride (1 in 30) responds to the Qualitative Tests <1.09> for chloride.

**Optical rotation** <2.49>  $[\alpha]_D^{20}$ : +5.5 – +6.1° (after drying, 5 g, water, 50 mL, 200 mm).

**Melting point** <2.60> 200 – 204°C

**Purity** (1) Clarity and color of solution—Dissolve 1.0 g of Ethambutol Hydrochloride in 10 mL of water: the solution is clear and colorless.

(2) Heavy metals <1.07>—Proceed with 2.0 g Ethambutol Hydrochloride according to Method 1, and perform the test. Prepare the control solution with 2.0 mL of Standard Lead Solution (not more than 10 ppm).

(3) Arsenic <1.11>—Prepare the test solution with 1.0 g of Ethambutol Hydrochloride according to Method 1, and perform the test (not more than 2 ppm).

(4) 2-Aminobutanol—Dissolve 5.0 g of Ethambutol Hydrochloride in methanol to make exactly 100 mL, and use this solution as the sample solution. Dissolve 0.05 g of 2-amino-1-butanol in methanol to make exactly 100 mL, and use this solution as the standard solution. Perform the test with these solutions as directed under Thin-layer Chromatography <2.03>. Spot 2  $\mu$ L each of the sample solution and standard solution on a plate of silica gel for thin-layer chromatography. Develop the plate with a mixture of ethyl acetate, acetic acid (100), hydrochloric acid and water (11:7:1:1) to a distance of about 10 cm, air-dry the plate, and heat at 105°C for 5 minutes. Cool, spray evenly ninhydrin-L-ascorbic acid TS upon the plate, air-dry the plate, and heat at 105°C for 5 minutes: the spot from the sample solution, corresponding to that from the standard solution, has no more color than that from the standard solution.

**Loss on drying** <2.41> Not more than 0.5% (1 g, 105°C, 3 hours).

**Residue on ignition** <2.44> Not more than 0.1% (1 g).

**Assay** Weigh accurately about 0.2 g of Ethambutol Hydrochloride, previously dried, dissolve in 20 mL of water, and add 1.8 mL of copper (II) sulfate TS. To the solution add 7 mL of sodium hydroxide TS with shaking, add water to make exactly 50 mL, and centrifuge. Pipet 10 mL of the supernatant liquid, add 10 mL of ammonia-ammonium chloride buffer solution, pH 10.0 and 100 mL of water, and titrate <2.50> with 0.01 mol/L disodium dihydrogen ethylenediamine tetraacetate VS until the color of the solution changes from blue-purple through light red to light yellow (indicator: 0.15 mL of Cu-PAN TS). Perform a blank determination, and make any necessary correction.


$$\begin{aligned} & \text{Each mL of 0.01 mol/L disodium dihydrogen} \\ & \text{ethylenediamine tetraacetate VS} \\ & = 2.772 \text{ mg of C}_{10}\text{H}_{24}\text{N}_2\text{O}_2\cdot 2\text{HCl} \end{aligned}$$

**Containers and storage** Containers—Tight containers.

## Ethanol

### Alcohol

エタノール



$\text{C}_2\text{H}_6\text{O}$ : 46.07

Ethanol [64-17-5]

This monograph is harmonized with the European Pharmacopoeia and the U.S. Pharmacopeia. The parts of the text that are not harmonized are marked with symbols (♦ ♦).

Ethanol contains not less than 95.1 vol% and not more than 96.9 vol% (by specific gravity) of  $\text{C}_2\text{H}_6\text{O}$  at 15°C.

♦**Description** Ethanol is a clear, colorless liquid.

It is miscible with water.

It is flammable and burns with a light blue flame on ignition.

It is volatile.♦

**Identification** (1) Determine the infrared absorption spectrum of Ethanol as directed in the liquid film method under Infrared Spectrophotometry <2.25>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wave numbers.

**Specific gravity** <2.56>  $d_{4}^{15}$ : 0.809 – 0.816

**Purity** (1) Clarity and color of solution—Ethanol is clear and colorless. To 1.0 mL of Ethanol add water to make 20 mL, and allow to stand for 5 minutes: the resulting liquid is clear. Control solution: water.

(2) Acidity or alkalinity—To 20 mL of Ethanol add 20 mL of freshly boiled and cooled water and 0.1 mL of a solution prepared by addition of 7.0 mL of ethanol (95) and 2.0 mL of water to 1.0 mL of phenolphthalein TS: no color develops. Add 1.0 mL of 0.01 mol/L sodium hydroxide VS to this solution: a light red color develops.

(3) Volatile impurities—Pipet 500 mL of Ethanol, add 150  $\mu\text{L}$  of 4-methylpentan-2-ol, and use this solution as the sample solution. Separately, to 100  $\mu\text{L}$  of anhydrous methanol add Ethanol to make exactly 50 mL. Pipet 5 mL of this solution, add Ethanol to make exactly 50 mL, and use this solution as the standard solution (1). Separately, to exactly 50  $\mu\text{L}$  each of anhydrous methanol and acetaldehyde add Ethanol to make exactly 50 mL. To exactly 100  $\mu\text{L}$  of this solution add Ethanol to make exactly 10 mL, and use this solution as the standard solution (2). Separately, to exactly 150  $\mu\text{L}$  of acetal add Ethanol to make exactly 50 mL. To exactly 100  $\mu\text{L}$  of this solution add Ethanol to make exactly 10 mL, and use this solution as the standard solution (3). Separately, to exactly 100  $\mu\text{L}$  of benzene add Ethanol to make exactly 100 mL. To exactly 100  $\mu\text{L}$  of this solution add Ethanol to make exactly 50 mL, and use this solution as the standard solution (4). Perform the test with exactly 1  $\mu\text{L}$  each of Ethanol, the sample solution and standard solutions (1), (2), (3) and (4) as directed under Gas Chromatography <2.02> according to the following conditions, and determine the

peak areas of acetaldehyde,  $A_E$ , benzene,  $B_E$  and acetal,  $C_E$  obtained with Ethanol, and the peak area of methanol with the standard solution (1), the peak area of acetaldehyde,  $A_T$  with the standard solution (2), the peak area of acetal,  $C_T$  with the standard solution (3) and the peak area of benzene,  $B_T$ : the peak area of methanol is not more than 1/2 times the peak area of methanol with the standard solution (1). When calculate the amounts of the volatile impurities by the following equation, the total amount of acetaldehyde and acetal is not more than 10 vol ppm as acetaldehyde, and the amount of benzene is not more than 2 vol ppm. The total area of the peaks other than the peak mentioned above and the peak having the area not more than 3% that of 4-methylpentan-2-ol is not larger than the peak area of 4-methylpentan-2-ol.

Total amount (vol ppm) of acetaldehyde and acetal  
 $= \{(10 \times A_E)/(A_T - A_E)\} + \{(30 \times C_E)/(C_T - C_E)\}$

Amount (vol ppm) of benzene =  $2B_E/(B_T - B_E)$

If necessary, identify the peak of benzene by using a different stationary liquid phase and suitable chromatographic conditions.

**Operating conditions**—

Detector: A hydrogen flame-ionization detector

Column: A fused silica tube 0.32 mm in inside diameter and 30 m in length, coated with 6% cyanopropyl phenyl-94% dimethyl silicone polymer for gas chromatography in 1.8  $\mu\text{m}$  thickness.

Column temperature: Inject at a constant temperature of about 40°C, maintain the temperature for 12 minutes, then rise up to 240°C at the rate of 10°C per minute, and maintain at a constant temperature of about 240°C for 10 minutes.

Carrier gas: Helium

Flow rate: 35 cm/min.

Split ratio: 1: 20

**System suitability**—

System performance: When the procedure is run with 1  $\mu\text{L}$  of the standard solution (2) under the above operating conditions, acetaldehyde and methanol are eluted in this order with the resolution between these peaks being not less than 1.5.

(4) Other impurities (absorbance)—Perform the test with Ethanol as directed under Ultraviolet-visible Spectrophotometry <2.24>: the absorbances at 240 nm, between 250 nm and 260 nm and between 270 nm and 340 nm are not more than 0.40, 0.30, and 0.10, respectively. The absorption spectrum determined in a 5-cm cell using water as a blank shows a flat absorption curve between 235 nm and 340 nm.

(5) Residue on evaporation—Evaporate 100 mL of Ethanol, exactly measured, in a tared dish on a water bath, and dry for 1 hour at 105°C: the mass of the residue does not exceed 2.5 mg.


**Containers and storage** ♦Containers—Tight containers.♦

Storage—Without exposure to light.

## Anhydrous Ethanol

### Dehydrated Alcohol

無水エタノール



$\text{C}_2\text{H}_6\text{O}$ : 46.07

Ethanol [64-17-5]

This monograph is harmonized with the European Pharmacopoeia and the U.S. Pharmacopeia. The parts of the text that are not harmonized are marked with symbols (♦ ♦).

Anhydrous Ethanol contains not less than 99.5 vol% (by specific gravity) of  $\text{C}_2\text{H}_6\text{O}$  at 15°C.

♦**Description** Anhydrous Ethanol is a clear, colorless liquid.

It is miscible with water.

It is flammable and burns with a light blue flame on ignition.

It is volatile.

Boiling point: 78 – 79°C♦.

**Identification (1)** Determine the infrared absorption spectrum of Anhydrous Ethanol as directed in the solution method under Infrared Spectrophotometry <2.25>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wave numbers.

**Specific gravity** <2.56>  $d_{15}^{15}$ : 0.794 – 0.797

**Purity (1)** Clarity and color of solution—Anhydrous Ethanol is clear and colorless. To 1.0 mL of Anhydrous Ethanol add water to make 20 mL, and allow to stand for 5 minutes: the resulting liquid is clear. Control solution: water

(2) Acidity or alkalinity—To 20 mL of Anhydrous Ethanol add 20 mL of freshly boiled and cooled water and 0.1 mL of a solution obtained by addition of 7.0 mL of ethanol (95) and 2.0 mL of water to 1.0 mL of phenolphthalein TS: no color develops. Add 1.0 mL of 0.01 mol/L sodium hydroxide VS to this solution: a light red color develops.

(3) Volatile impurities—Pipet 500 mL of Anhydrous Ethanol, add 150  $\mu\text{L}$  of 4-methylpentan-2-ol, and use this solution as the sample solution. Separately, to 100  $\mu\text{L}$  of anhydrous ethanol add Anhydrous Ethanol to make exactly 50 mL. Pipet 5 mL of this solution, add Anhydrous Ethanol to make exactly 50 mL, and use this solution as the standard solution (1). Separately, to exactly 50  $\mu\text{L}$  each of anhydrous methanol and acetaldehyde add Anhydrous Ethanol to make exactly 50 mL. To exactly 100  $\mu\text{L}$  of this solution add Anhydrous Ethanol to make exactly 10 mL, and use this solution as the standard solution (2). Separately, to exactly 150  $\mu\text{L}$  of acetal add Anhydrous Ethanol to make exactly 50 mL. To exactly 100  $\mu\text{L}$  of this solution add Anhydrous Ethanol to make exactly 10 mL, and use this solution as the standard solution (3). Separately, to exactly 100  $\mu\text{L}$  of benzene add Anhydrous Ethanol to make exactly 100 mL. To exactly 100  $\mu\text{L}$  of this solution add Anhydrous Ethanol to make exactly 50 mL, and

use this solution as the standard solution (4). Perform the test with exactly 1  $\mu\text{L}$  each of Anhydrous Ethanol, the sample solution and standard solutions (1), (2), (3) and (4) as directed under Gas Chromatography <2.02> according to the following conditions, and determine the peak areas of acetaldehyde,  $A_E$ , benzene,  $B_E$  and acetal,  $C_E$  obtained with Anhydrous Ethanol, and the peak area of methanol with the standard solution (1), the peak area of acetaldehyde,  $A_T$  with the standard solution (2), the peak area of acetal,  $C_T$  with the standard solution (3) and the peak area of benzene,  $B_T$ : the peak area of methanol is not more than 1/2 times the peak area of methanol with the standard solution (1). When calculate the amounts of the volatile impurities by the following equation, the total amount of acetaldehyde and acetal is not more than 10 vol ppm as acetaldehyde, and the amount of benzene is not more than 2 vol ppm. The total area of the peaks other than the peak mentioned above and the peak having the area not more than 3% that of 4-methylpentan-2-ol is not larger than the peak area of 4-methylpentan-2-ol.

$$\text{Total amount (vol ppm) of acetaldehyde and acetal} \\ = \{(10 \times A_E)/(A_T - A_E)\} + \{(30 \times C_E)/(C_T - C_E)\}$$

$$\text{Amount (vol ppm) of benzene} = 2B_E/(B_T - B_E)$$

If necessary, identify the peak of benzene by using a different stationary liquid phase and suitable chromatographic conditions.

**Operating conditions—**

Detector: A hydrogen flame-ionization detector

Column: A fused silica tube 0.32 mm in inside diameter and 30 m in length, coated with 6% cyanopropyl phenyl-94% dimethyl silicone polymer for gas chromatography in 1.8  $\mu\text{m}$  thickness.

Column temperature: Inject at a constant temperature of about 40°C, maintain the temperature for 12 minutes, then rise up to 240°C at the rate of 10°C per minute, and maintain at a constant temperature of about 240°C for 10 minutes.

Carrier gas: Helium

Flow rate: 35 cm/min.

Split ratio: 1: 20

**System suitability—**

System performance: When the procedure is run with 1  $\mu\text{L}$  of the standard solution (2) under the above operating conditions, acetaldehyde and methanol are eluted in this order with the resolution between these peaks being not less than 1.5.

(4) Other impurities (absorbance)—Perform the test with Ethanol as directed under Ultraviolet-visible Spectrophotometry <2.24>: the absorbances at 240 nm, between 250 nm and 260 nm and between 270 nm and 340 nm are not more than 0.40, 0.30, and 0.10, respectively. The absorption spectrum determined in a 5-cm cell using water as a blank shows a flat absorption curve between 235 nm and 340 nm.

(5) Residue on evaporation—Evaporate 100 mL of Dehydrated Ethanol, exactly measured, in a tared dish on a water bath, and dry for 1 hour at 105°C: the mass of the residue does not exceed 2.5 mg.

**Containers and storage** ♦Containers—Tight containers.♦

Storage—Without exposure to light.

## Ethanol for Disinfection

### Alcohol for Disinfection

#### 消毒用エタノール

Ethanol for Disinfection contains not less than 76.9 vol% and not more than 81.4 vol% (by specific gravity) of ethanol (C<sub>2</sub>H<sub>6</sub>O: 46.07) at 15°C.

#### Method of preparation

|                 |                       |
|-----------------|-----------------------|
| Ethanol         | 830 mL                |
| Purified Water  | a sufficient quantity |
| To make 1000 mL |                       |

Prepare by mixing the above ingredients.

**Description** Ethanol for Disinfection is a colorless, clear liquid.

It is miscible with water.

It burns with a light blue flame on ignition.

It is volatile.

**Identification (1)** To 1 mL of Ethanol for Disinfection add 2 mL of iodine TS and 1 mL of sodium hydroxide TS, and mix: light yellow precipitates appear.

**(2)** To 1 mL of Ethanol for Disinfection add 1 mL of acetic acid (100) and 3 drops of sulfuric acid, and heat: the odor of ethyl acetate is produced.

**Specific gravity** <2.56>  $d_{4}^{15}$ : 0.860 – 0.873

**Purity** Proceed as directed in the Purity under Ethanol.


**Containers and storage** Containers—Tight containers.

Storage—Light-resistant.

## Ethenzamide

### Ethoxybenzamide

#### エテンザミド



C<sub>9</sub>H<sub>11</sub>NO<sub>2</sub>: 165.19

2-Ethoxybenzamide [938-73-8]

Ethenzamide, when dried, contains not less than 98.0% of C<sub>9</sub>H<sub>11</sub>NO<sub>2</sub>.

**Description** Ethenzamide occurs as white crystals or crystalline powder. It is odorless and tasteless.

It is soluble in ethanol (95) and in acetone, slightly soluble in diethyl ether, and practically insoluble in water.

Its saturated solution is neutral.

It begins to sublime slightly at about 105°C.

**Identification (1)** To 0.5 g of Ethenzamide add 5 mL of sodium hydroxide TS, and heat the mixture gently: the gas evolved turns moistened red litmus paper to blue.

**(2)** To 0.2 g of Ethenzamide add 10 mL of hydrobromic acid, and boil the mixture gently for 1 hour under a reflux condenser. Cool in ice water, collect the separated crystalline precipitate, wash with three 5-mL portions of ice water, and dry in a desiccator (in vacuum, silica gel) for 2 hours: the precipitate melts <2.60> between 158°C and 161°C.

**Melting point** <2.60> 131 – 134°C

**Purity (1) Chloride** <1.03>—Dissolve 0.5 g of Ethenzamide in 30 mL of acetone, add 6 mL of dilute nitric acid, and dilute with water to make 50 mL. Perform the test using this solution as the test solution. Prepare the control solution as follows: to 0.7 mL of 0.01 mol/L hydrochloric acid VS add 30 mL of acetone and 6 mL of dilute nitric acid, and dilute with water to make 50 mL (not more than 0.050%).

**(2) Sulfate** <1.14>—Dissolve 0.5 g of Ethenzamide in 30 mL of acetone, add 1 mL of dilute hydrochloric acid, and dilute with water to 50 mL. Perform the test using this solution as the test solution. Prepare the control solution as follows: to 0.50 mL of 0.005 mol/L sulfuric acid VS add 30 mL of acetone and 1 mL of dilute hydrochloric acid, and dilute with water to 50 mL (not more than 0.048%).

**(3) Heavy metals** <1.07>—Proceed with 2.0 g of Ethenzamide according to Method 2, and perform the test. Prepare the control solution with 2.0 mL of Standard Lead Solution (not more than 10 ppm).

**(4) Arsenic** <1.11>—To 0.40 g of Ethenzamide add 0.3 g of potassium nitrate and 0.5 g of anhydrous sodium carbonate, mix thoroughly, ignite the mixture gradually, and cool. Dissolve the residue in 10 mL of dilute sulfuric acid, and heat the solution until white fumes begin to evolve. After cooling, add water carefully to make 5 mL, use this solution as the test solution, and perform the test (not more than 5 ppm).

**(5) Salicylamide**—Dissolve 0.20 g of Ethenzamide in 15 mL of diluted ethanol (95) (2 in 3), and add 2 to 3 drops of dilute iron (III) chloride TS: no purple color develops.

**Loss on drying** <2.41> Not more than 1.0% (1 g, silica gel, 3 hours).

**Residue on ignition** <2.44> Not more than 0.1% (1 g).

**Assay** Weigh accurately about 20 mg each of Ethenzamide and Ethenzamide Reference Standard, previously dried, and dissolve each in 70 mL of ethanol (95) by warming, and after cooling, add ethanol (95) to make exactly 100 mL. Pipet 5 mL each of these solutions, add ethanol (95) to make exactly 50 mL, and use these solutions as the sample solution and standard solution, respectively. Determine the absorbances,  $A_T$  and  $A_S$ , of the sample solution and standard solution at 290 nm as directed under Ultraviolet-visible Spectrophotometry <2.24>, using ethanol (95) as the blank.

$$\begin{aligned} \text{Amount (mg) of C}_9\text{H}_{11}\text{NO}_2 \\ = W_S \times (A_T/A_S) \end{aligned}$$


$W_S$ : Amount (mg) of Ethenzamide Reference Standard

**Containers and storage** Containers—Well-closed containers.



## Ether

エーテル



$\text{C}_4\text{H}_{10}\text{O}$ : 74.12

Diethyl ether [60-29-7]

Ether contains not less than 96% and not more than 98% (by specific gravity) of  $\text{C}_4\text{H}_{10}\text{O}$ .

It contains a small quantity of ethanol and water. It cannot be used for anesthesia.

**Description** Ether is a colorless, clear, mobile liquid, having a characteristic odor.

It is miscible with ethanol (95).

It is soluble in water.

It is highly volatile and flammable.

It is slowly oxidized by the action of air and light, with the formation of peroxides.

Its vapor, when mixed with air and ignited, may explode violently.

Boiling point: 35 – 37°C

**Specific gravity** <2.56>  $d_{20}^{20}$ : 0.718 – 0.721

**Purity (1)** Foreign odor—Place 10 mL of Ether in an evaporating dish, and allow it to evaporate spontaneously to a volume of about 1 mL: no foreign odor is perceptible. Drop this residue onto a piece of clean, odorless filter paper to evaporate the ether: no foreign odor is perceptible.

**(2)** Acidity—Place 10 mL of diluted ethanol (95) (4 in 5) and 0.5 mL of phenolphthalein TS in a 50-mL glass-stoppered flask, and add 0.2 mol/L sodium hydroxide dropwise to produce a red color which persists after shaking for 30 seconds. Add 25 mL of Ether, stopper the flask, shake gently, and add 0.40 mL of 0.02 mol/L sodium hydroxide VS with shaking: a red color develops.

**(3)** Aldehyde—Place 10 mL of Ether in a Nessler tube, add 1 mL of potassium hydroxide TS, and allow the mixture to stand for 2 hours, protecting from light, with occasional shaking: no color is produced in the ether layer and the aqueous layer.

**(4)** Peroxide—Place 10 mL of Ether in a Nessler tube, add 1 mL of a freshly prepared solution of potassium iodide (1 in 10), shake for 1 minute, then add 1 mL of starch TS, and shake well: no color is produced in the ether layer and in the aqueous layer.


**(5)** Residue on evaporation—Evaporate 140 mL of Ether, and dry the residue at 105°C for 1 hour: the mass of the residue does not more than 1.0 mg.

**Containers and storage** Containers—Tight containers.

Storage—Without fill up, light-resistant, remote from fire, and not exceeding 25°C.

## Anesthetic Ether

麻醉用エーテル



$\text{C}_4\text{H}_{10}\text{O}$ : 74.12

Diethyl ether [60-29-7]

Anesthetic Ether contains not less than 96% and not more than 98% (by specific gravity) of  $\text{C}_4\text{H}_{10}\text{O}$ .

It contains small quantities of ethanol and water. Suitable stabilizers may be added.

It is not to be used for anesthesia if it has been removed from the original container for more than 24 hours.

**Description** Anesthetic Ether occurs as a colorless, clear, mobile liquid, having a characteristic odor.

It is miscible with ethanol (95).

It is soluble in water.

It is highly volatile and flammable.

It is slowly oxidized by the action of air and light, with the formation of peroxides.

Its vapor, when mixed with air and ignited, may explode violently.

Boiling point: 35 – 37°C

**Specific gravity** <2.56>  $d_{20}^{20}$ : 0.718 – 0.721

**Purity (1)** Foreign odor—Place 10 mL of Anesthetic Ether in an evaporating dish, and allow it to evaporate spontaneously to a volume of about 1 mL: no foreign odor is perceptible. Drop this residue onto a piece of clean, odorless filter paper to evaporate the ether: no foreign odor is perceptible.

**(2)** Acidity—Place 10 mL of diluted ethanol (95) (4 in 5) and 0.5 mL of phenolphthalein TS in a 50-mL glass-stoppered flask, and add 0.2 mol/L sodium hydroxide dropwise to produce a red color which persists after shaking for 30 seconds. Add 25 mL of Anesthetic Ether, stopper the flask, shake gently, and add 0.40 mL of 0.02 mol/L sodium hydroxide VS with shaking: a red color develops.

**(3)** Aldehyde—To 100 mL of water in a 200-mL glass-stoppered flask add 10 mL of Anesthetic Ether and 1 mL of a solution of sodium hydrogen sulfite (1 in 1000), stopper tightly, shake vigorously for 10 seconds, and allow the mixture to stand in a cool place for 30 minutes, protected from light. Add 2 mL of starch TS, and add dropwise 0.01 mol/L iodine VS until a pale blue color develops. Shake with about 2 g of sodium hydrogen carbonate to decolorize the solution, and add 1 mL of diluted 0.01 mol/L iodine VS (9 in 40): a blue color develops. Keep the temperature of the solution below 18°C during the procedure.

**(4)** Peroxide—Place 10 mL of Anesthetic Ether in a Nessler tube, add 1 mL of a freshly prepared solution of potassium iodide (1 in 10), shake occasionally for 1 hour, protecting from light, then add 1 mL of starch TS, and shake well: no color is produced and in the aqueous layer and in the ether layer.

**(5)** Residue on evaporation—Evaporate 50 mL of Anesthetic Ether, and dry the residue at 105°C for 1 hour:


the mass of the residue is not more than 1.0 mg.

**Containers and storage** Containers—Tight containers.

Storage—Without fill up, light-resistant, remote from fire, and not exceeding 25°C.

## Ethinylestradiol

エチニルエストラジオール



$C_{20}H_{24}O_2$ : 296.40

17 $\alpha$ -Ethinylestra-1,3,5(10)-triene-3,17 $\beta$ -diol [57-63-6]

Ethinylestradiol, when dried, contains not less than 98.0% of  $C_{20}H_{24}O_2$ .

**Description** Ethinylestradiol occurs as white to pale yellow crystals or crystalline powder. It is odorless.

It is freely soluble in pyridine and in tetrahydrofuran, soluble in ethanol (95) and in diethyl ether, and practically insoluble in water.

It dissolves in sodium hydroxide TS.

**Identification (1)** Dissolve 2 mg of Ethinylestradiol in 1 mL of a mixture of ethanol (95) and sulfuric acid (1:1); a purplish red color develops with a yellow-green fluorescence. Add carefully 2 mL of water to this solution: the color of the solution changes to red-purple.

**(2)** Transfer 0.02 g of Ethinylestradiol to a glass-stoppered test tube, dissolve in 10 mL of a solution of potassium hydroxide (1 in 20), add 0.1 g of benzoyl chloride, and shake. Collect the resulting precipitate, recrystallize from methanol, and dry in a desiccator (in vacuum, phosphorus (V) oxide): the precipitate melts <2.60> between 200°C and 202°C.

**Optical rotation** <2.49>  $[\alpha]_D^{20}$ : -26 - -31° (after drying, 0.1 g, pyridine, 25 mL, 200 mm).

**Melting point** <2.60> 180 - 186°C or 142 - 146°C

**Purity** Estrone—Dissolve 5 mg of Ethinylestradiol in 0.5 mL of ethanol (95), and add 0.05 g of 1,3-dinitrobenzene. Add 0.5 mL of freshly prepared dilute potassium hydroxide-ethanol TS, allow to stand in a dark place for 1 hour, and add 10 mL of ethanol (95): the solution has no more color than the following control solution.

Control solution: Proceed in the same manner as mentioned above, omitting Ethinylestradiol.

**Loss on drying** <2.41> Not more than 0.5% (0.5 g, in vacuum, phosphorus (V) oxide, 4 hours).

**Residue on ignition** <2.44> Not more than 0.1% (0.5 g).

**Assay** Weigh accurately about 0.2 g of Ethinylestradiol, previously dried, and dissolve in 40 mL of tetrahydrofuran. Add 10 mL of a solution of silver nitrate (1 in 20), and titrate <2.50> with 0.1 mol/L sodium hydroxide VS (potentiometric titration).

Each mL of 0.1 mol/L sodium hydroxide VS

= 29.64 mg of  $C_{20}H_{24}O_2$

**Containers and storage** Containers—Tight containers.

Storage—Light-resistant.

## Ethinylestradiol Tablets

エチニルエストラジオール錠

Ethinylestradiol Tablets contain not less than 90% and not more than 110% of the labeled amount of ethinylestradiol ( $C_{20}H_{24}O_2$ : 296.40).

**Method of preparation** Prepare as directed under Tablets, with Ethinylestradiol.

**Identification (1)** Evaporate to dryness 5 mL of the sample solution obtained in Assay, and add 2 mL of a mixture of sulfuric acid and ethanol (95) (2:1) to the residue: a light red color with a yellow fluorescence develops. To the solution add carefully 4 mL of water: the color of the solution changes to red-purple.

**(2)** Evaporate to dryness 10 mL of the sample solution obtained in Assay, add 0.2 mL of acetic acid (31) and 2 mL of phosphoric acid to the residue, and heat on a water bath for 5 minutes: a red color with a yellow-green fluorescence develops.

**Uniformity of dosage units** <6.02> Perform the test according to the following method: it meets the requirement of the Content uniformity test.

Place 1 tablet of Ethinylestradiol Tablets in a separator, add 10 mL of 2nd fluid for disintegration test, and shake until the tablet is disintegrated. Add 10 mL of dilute sulfuric acid and 20 mL of chloroform, shake vigorously for 5 minutes, and filter the chloroform layer into a conical flask through filter paper on which 5 g of anhydrous sodium sulfate is placed. Extract the aqueous layer with two 20-mL portions of chloroform, proceed with the extracts in the same manner as before, and combine the filtrates with the previous one. Evaporate gently the combined filtrate on a water bath with the aid of a current of nitrogen, dissolve the residue in exactly 100 mL of methanol, and centrifuge, if necessary. Pipet  $x$  mL of the supernatant liquid, add methanol to make exactly  $V$  mL of a solution containing about 0.04  $\mu$ g of ethinylestradiol ( $C_{20}H_{24}O_2$ ) per mL, and use this solution as the sample solution. Separately, weigh accurately about 10 mg of Ethinylestradiol Reference Standard, previously dried in a desiccator (in vacuum, phosphorus (V) oxide) for 4 hours, dissolve in methanol, dilute to a volume containing about 0.04  $\mu$ g of ethinylestradiol ( $C_{20}H_{24}O_2$ ) per mL, and use this solution as the standard solution. Pipet 4 mL each of sulfuric acid-methanol TS into three glass-stoppered test tubes, T, S and B, cool in ice, to each tube add exactly 1 mL each of the sample solution, the standard solution and methanol, shake immediately, and allow to stand in a water bath at 30°C for 40 minutes, then allow to stand in a water bath at 20°C for 5 minutes. Perform the test with these solutions as directed under Fluorometry <2.22>. Determine the fluorescence intensities,  $F_T$ ,  $F_S$  and  $F_B$ , of these solutions using the fluorophotometer, at about 460 nm of the excitation and at about 493 nm of the fluorescence.

Amount (mg) of ethinylestradiol ( $C_{20}H_{24}O_2$ )

$$= W_S \times \{(F_T - F_B)/(F_S - F_B)\} \times (V/2500) \times (1/x)$$

$W_S$ : Amount (mg) of Ethinylestradiol Reference Standard

**Assay** (i) Chromatographic tube: Pack a pledget of glass wool in the bottom of a tube 25 mm in inside diameter and 300 mm in length, and place 5 g of anhydrous sodium sulfate on the glass wool.

(ii) Chromatographic column: Place 5 g of siliceous earth for chromatography in a 200-mL beaker, soak well in 4 mL of 1 mol/L hydrochloric acid TS, and mix uniformly. Put the siliceous earth into the chromatographic tube in small portions to make 60 to 80 mm in height in proper hardness with a tamping rod.

(iii) Standard solution: Weigh accurately about 10 mg of Ethinylestradiol Reference Standard, previously dried in a desiccator (in vacuum, phosphorus (V) oxide) for 4 hours, and dissolve in chloroform to make exactly 100 mL. Pipet 5 mL of this solution, and add chloroform to make exactly 100 mL.

(iv) Sample: Weigh accurately not less than 20 Ethinylestradiol Tablets, and powder. Weigh accurately a portion of the powder, equivalent to about 0.5 mg of ethinylestradiol ( $C_{20}H_{24}O_2$ ), place in a 50-mL beaker, add 2 mL of water, shake well, add 3 mL of chloroform, and shake well again. Add 4 g of siliceous earth for chromatography, mix well until the contents do not stick to the inner wall of the beaker, and use the substance as the sample.

(v) Procedure: To the chromatographic column add the sample with a funnel, and pack in proper hardness. Mix well the sample sticking to the beaker with 0.5 g of siliceous earth for chromatography, and place in the chromatographic tube. Wipe off the sample solution sticking to the beaker and the tamping rod with glass wool, and place it in the chromatographic tube. Push down the sample, and press lightly on the chromatographic column to make the height of the column 110 mm to 130 mm. Take 70 mL of chloroform, rinse the inner wall of the chromatographic tube with a portion of the chloroform, and transfer the remaining portion to the chromatographic tube. Collect the effluent solution at a flow rate not more than 0.8 mL per minute. After completing the elution, rinse the lower end of the chromatographic tube with a small quantity of chloroform, add chloroform to make exactly 100 mL, and use this solution as the sample solution. Transfer 6 mL each of the sample solution and standard solution to each separator, and add 20 mL each of isooctane. Add exactly 10 mL of a mixture of sulfuric acid and methanol (7:3), shake vigorously for 5 minutes, allow to stand in a dark place for 15 minutes, and centrifuge. Perform the test with the resulting color solutions as directed under Ultraviolet-visible Spectrophotometry <2.24>, using a solution, prepared with 6 mL of chloroform in the same manner, as the blank. Determine the absorbances,  $A_T$  and  $A_S$ , of the subsequent solutions obtained from the sample solution and standard solution at 540 nm, respectively.


$$\begin{aligned} &\text{Amount (mg) of ethinylestradiol } (C_{20}H_{24}O_2) \\ &= W_S \times (A_T/A_S) \times (1/20) \end{aligned}$$

$W_S$ : Amount (mg) of Ethinylestradiol Reference Standard

**Containers and storage** Containers—Well-closed containers.

## Ethionamide

エチオナミド



$C_8H_{10}N_2S$ : 166.24

2-Ethylpyridine-4-carbothioamide [536-33-4]

Ethionamide, when dried, contains not less than 98.5% and not more than 101.0% of  $C_8H_{10}N_2S$ .

**Description** Ethionamide occurs as yellow crystals or crystalline powder, having a characteristic odor.

It is soluble in methanol and in acetic acid (100), sparingly soluble in ethanol (99.5) and in acetone, and practically insoluble in water.

**Identification** (1) Determine the absorption spectrum of a solution of Ethionamide in methanol (3 in 160,000) as directed under Ultraviolet-visible Spectrophotometry <2.24>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wavelengths.

(2) Determine the infrared absorption spectrum of Ethionamide as directed in the potassium bromide disk method under Infrared Spectrophotometry <2.25>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wave numbers.

**Melting point** <2.60> 161 – 165°C

**Purity** (1) Acidity—Dissolve 3.0 g of Ethionamide in 30 mL of methanol by warming, add 90 mL of water, allow to stand in ice water for 1 hour, and filter. To 80 mL of the filtrate add 0.8 mL of cresol red TS and 0.20 mL of 0.1 mol/L sodium hydroxide VS: a red color develops.

(2) Heavy metals <1.07>—Proceed with 1.0 g of Ethionamide according to Method 2, and perform the test. Prepare the control solution with 2.0 mL of Standard Lead Solution (not more than 20 ppm).

(3) Arsenic <1.11>—Prepare the test solution with 1.0 g of Ethionamide according to Method 3. Add 10 mL of a solution of magnesium nitrate hexahydrate in ethanol (95) (1 in 50), then add 1.5 mL of hydrogen peroxide (30), and fire to burn (not more than 2 ppm).

(4) Related substances—Conduct this procedure without exposure to light, using light-resistant vessels. Dissolve 0.20 g of Ethionamide in 10 mL of acetone, and use this solution as the sample solution. Pipet 0.5 mL of the sample solution, add acetone to make exactly 100 mL, and use this solution as the standard solution (1). Separately, pipet exactly 0.2 mL of the sample solution, add acetone to make exactly 100 mL, and use this solution as the standard solution (2). Perform the test with these solutions as directed under Thin-layer Chromatography <2.03>. Spot 10  $\mu$ L each of the sample solution and standard solutions (1) and (2) on a plate of silica gel with fluorescent indicator for thin-layer chromatography, de-

velop with a mixture of ethyl acetate, hexane and methanol (6:2:1) to a distance of about 15 cm, and air-dry the plate. Examine under ultraviolet light (main wavelength: 254 nm): the spot other than the principal spot obtained with the sample solution is not more intense than the spot with the standard solution (1), and number of the spot other than the principal spot obtained with the sample solution which is more intense than the spot with the standard solution (2) is not more than one.

**Loss on drying** <2.41> Not more than 0.5% (1 g, 105°C, 3 hours).

**Residue on ignition** <2.44> Not more than 0.1% (1 g).


**Assay** Weigh accurately about 0.3 g of Ethionamide, previously dried, dissolve in 50 mL of acetic acid (100), and titrate <2.50> with 0.1 mol/L perchloric acid VS until the color of the solution changes from orange-red to dark orange-brown (indicator: 2 mL of *p*-naphtholbenzein TS). Perform a blank determination in the same manner, and make any necessary correction.

Each mL of 0.1 mol/L perchloric acid VS  
= 16.62 mg of C<sub>8</sub>H<sub>10</sub>N<sub>2</sub>S

**Containers and storage** Containers—Well-closed containers.

## Ethosuximide

エトスクシミド



C<sub>7</sub>H<sub>11</sub>NO<sub>2</sub>: 141.17  
(2*RS*)-2-Ethyl-2-methylsuccinimide [77-67-8]

Ethosuximide contains not less than 98.5% of C<sub>7</sub>H<sub>11</sub>NO<sub>2</sub> calculated on the anhydrous basis.

**Description** Ethosuximide occurs as a white, paraffin-like solid or powder. It is odorless or has a slight, characteristic odor.

It is very soluble in methanol, in ethanol (95), in diethyl ether, and in *N,N*-dimethylformamide, and freely soluble in water.

Melting point: about 48°C

**Identification** (1) To 0.2 g of Ethosuximide add 10 mL of sodium hydroxide TS, and boil: the gas evolved turns a moistened red litmus paper blue.

(2) Dissolve 0.05 g of Ethosuximide in 1 mL of ethanol (95), add 3 drops of a solution of copper (II) acetate monohydrate (1 in 100), warm slightly, and add 1 to 2 drops of sodium hydroxide TS: a purple color is produced.

(3) Determine the absorption spectrum of a solution of Ethosuximide in ethanol (95) (1 in 2000) as directed under Ultraviolet-visible Spectrophotometry <2.24>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wavelengths.

**Purity** (1) Clarity and color of solution—Dissolve 1.0 g of

Ethosuximide in 10 mL of water: the solution is clear and colorless.

(2) Chloride <1.03>—With 1.0 g of Ethosuximide, perform the test. Prepare the control solution with 0.30 mL of 0.01 mol/L hydrochloric acid VS (not more than 0.011%).

(3) Heavy metals <1.07>—Proceed with 1.0 g of Ethosuximide according to Method 1, and perform the test. Prepare the control solution with 2.0 mL of Standard Lead Solution (not more than 20 ppm).

(4) Arsenic <1.11>—Prepare the test solution with 1.0 g of Ethosuximide, according to Method 1, and perform the test (not more than 2 ppm).

(5) Acid anhydride—Dissolve 0.50 g of Ethosuximide in 1 mL of ethanol (95), add 1 mL of hydroxylammonium chloride-iron (III) chloride TS, and allow to stand for 5 minutes. Add 3 mL of water, mix, and allow to stand for 5 minutes: the red to red-purple color of this solution is not more intense than that of the following control solution.

Control solution: Dissolve 0.070 g of succinic anhydride in ethanol (95) to make exactly 100 mL. To 1.0 mL of this solution add 1 mL of hydroxylammonium chloride-iron (III) chloride TS, and proceed in the same manner.

(6) Cyanide—Dissolve 1.0 g of Ethosuximide in 10 mL of ethanol (95), and add 3 drops of iron (II) sulfate TS, 1 mL of sodium hydroxide TS and 2 to 3 drops of iron (III) chloride TS. Warm gently, and acidify with dilute sulfuric acid: not a blue precipitate and a blue color are produced within 15 minutes.

**Water** <2.48> Not more than 0.5% (2 g, direct titration).

**Residue on ignition** <2.44> Not more than 0.1% (1 g).

**Assay** Weigh accurately about 0.2 g of Ethosuximide, dissolve in 20 mL of *N,N*-dimethylformamide, and titrate <2.50> with 0.1 mol/L tetramethylammonium hydroxide VS (potentiometric titration). Perform a blank determination.

Each mL of 0.1 mol/L tetramethylammonium hydroxide VS  
= 14.12 mg of C<sub>7</sub>H<sub>11</sub>NO<sub>2</sub>


**Containers and storage** Containers—Tight containers.

## Ethyl Aminobenzoate

Anesthamine

Benzocaine

アミノ安息香酸エチル



C<sub>9</sub>H<sub>11</sub>NO<sub>2</sub>: 165.19  
Ethyl 4-aminobenzoate [94-09-7]

Ethyl Aminobenzoate, when dried, contains not less than 99.0% of C<sub>9</sub>H<sub>11</sub>NO<sub>2</sub>.

**Description** Ethyl Aminobenzoate occurs as white crystals or crystalline powder. It is odorless. It has a slightly bitter

taste, numbing the tongue.

It is freely soluble in ethanol (95) and in diethyl ether, and very slightly soluble in water.

It dissolves in dilute hydrochloric acid.

**Identification (1)** Dissolve 0.01 g of Ethyl Aminobenzoate in 1 mL of dilute hydrochloric acid and 4 mL of water. This solution responds to the Qualitative Tests <1.09> for primary aromatic amines.

(2) Dissolve 0.1 g of Ethyl Aminobenzoate in 5 mL of water with the aid of dilute hydrochloric acid added dropwise, and add iodine TS dropwise: a brown precipitate is produced.

(3) Warm 0.05 g of Ethyl Aminobenzoate with 2 drops of acetic acid (31) and 5 drops of sulfuric acid: the odor of ethyl acetate is perceptible.

**Melting point** <2.60> 89 – 91°C

**Purity (1)** Acidity—Dissolve 1.0 g of Ethyl Aminobenzoate in 10 mL of neutralized ethanol, and add 10 mL of water, 2 drops of phenolphthalein TS and 0.50 mL of 0.01 mol/L sodium hydroxide VS: a red color is produced.

(2) Chloride—Dissolve 0.20 g of Ethyl Aminobenzoate in 5 mL of ethanol (95), add 2 to 3 drops each of dilute nitric acid and of silver nitrate TS: no change occurs immediately.

(3) Heavy metals <1.07>—Dissolve 2.0 g of Ethyl Aminobenzoate in 20 mL of ethanol (95), add 2 mL of dilute acetic acid and ethanol (95) to make 50 mL, and perform the test using this solution as the test solution. Prepare the control solution as follows: to 2.0 mL of Standard Lead Solution add 2 mL of dilute acetic acid and sufficient ethanol (95) to make 50 mL (not more than 10 ppm).

(4) Readily carbonizable substances <1.15>—Perform the test with 0.5 g of Ethyl Aminobenzoate: the solution has no more color than Matching Fluid A.

**Loss on drying** <2.41> Not more than 1.0% (1 g, silica gel, 3 hours).

**Residue on ignition** <2.44> Not more than 0.1% (1 g).

**Assay** Weigh accurately about 0.25 g of Ethyl Aminobenzoate, previously dried, dissolve in 10 mL of hydrochloric acid and 70 mL of water, add 10 mL of a solution of potassium bromide (3 in 10), and cool to a temperature below 15°C. Then titrate <2.50> with 0.1 mol/L sodium nitrite VS by the potentiometric titration or the amperometric titration.


Each mL of 0.1 mol/L sodium nitrite VS  
= 16.52 mg of C<sub>9</sub>H<sub>11</sub>NO<sub>2</sub>

**Containers and storage** Containers—Well-closed containers.

## Ethyl L-Cysteine Hydrochloride

### Ethyl Cysteine Hydrochloride

L-エチルシステイン塩酸塩



C<sub>5</sub>H<sub>11</sub>NO<sub>2</sub>S.HCl: 185.67

Ethyl (2*R*)-2-amino-3-sulfanylpropanoate  
monohydrochloride  
[868-59-7]

Ethyl L-Cysteine Hydrochloride, when dried, contains not less than 98.5% of C<sub>5</sub>H<sub>11</sub>NO<sub>2</sub>S.HCl.

**Description** Ethyl L-Cysteine Hydrochloride occurs as white crystals or crystalline powder. It has a characteristic odor, and has a bitter taste at first with a burning aftertaste.

It is very soluble in water, and freely soluble in ethanol (95).

Melting point: about 126°C (with decomposition).

**Identification (1)** Determine the infrared absorption spectrum of Ethyl L-Cysteine Hydrochloride as directed in the potassium bromide disk method under Infrared Spectrophotometry <2.25>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wave numbers.

(2) A solution of Ethyl L-Cysteine Hydrochloride (1 in 20) responds to the Qualitative Tests <1.09> (1) for chloride.

**Optical rotation** <2.49> [ $\alpha$ ]<sub>D</sub><sup>20</sup>: –10.0 – –13.0° (after drying, 2.0 g, 1 mol/L hydrochloric acid TS, 25 mL, 100 mm).

**Purity (1)** Sulfate <1.14>—Perform the test with 0.6 g of Ethyl L-Cysteine Hydrochloride. Prepare the control solution with 0.35 mL of 0.005 mol/L sulfuric acid (not more than 0.028%).

(2) Heavy metals <1.07>—Proceed with 1.0 g of Ethyl L-Cysteine Hydrochloride according to Method 1, and perform the test. Prepare the control solution with 1.0 mL of Standard Lead Solution (not more than 10 ppm).

(3) Related substances—Conduct this procedure rapidly. Dissolve 0.05 g each of Ethyl L-Cysteine Hydrochloride and *N*-ethylmaleimide in 5 mL of mobile phase, allow to stand for 30 minutes, and use this solution as the sample solution. Pipet 3 mL of the sample solution, add the mobile phase to make exactly 200 mL, and use this solution as the standard solution. Perform the test with exactly 2  $\mu$ L each of the sample solution and standard solution as directed under Liquid Chromatography <2.01> according to the following conditions. Determine each peak area of these solutions by the automatic integration method: a peak area from the sample solution with the ratio of the retention time to ethyl L-cysteine-*N*-ethylmaleimide complex from the standard solution being about 0.7 is not larger than the peak area of ethyl L-cysteine-*N*-ethylmaleimide complex from the standard solution. Each area of all peaks other than the peaks of ethyl L-cysteine-*N*-ethylmaleimide complex and *N*-ethylmaleimide from the sample solution is not larger than 1/3 of the peak area of ethyl L-cysteine-*N*-ethylmaleimide complex from the standard solution.

**Operating conditions—**

**Detector:** An ultraviolet absorption photometer (wavelength: 250 nm).

**Column:** A stainless steel column about 6 mm in inside diameter and about 15 cm in length, packed with octadecylsilanized silica gel for liquid chromatography (5  $\mu$ m in particle diameter).

**Column temperature:** A constant temperature of about 25°C.

**Mobile phase:** A mixture of 0.02 mol/L monobasic potas-

sium phosphate TS and acetonitrile (2:1).

**Flow rate:** Adjust the flow rate so that the retention time of ethyl L-cysteine-*N*-ethylmaleimide complex is about 4 minutes.

**Selection of column:** Dissolve 0.05 g of Ethyl L-Cysteine Hydrochloride, 0.01 g of L-cysteine hydrochloride and 0.05 g of *N*-ethylmaleimide in 25 mL of the mobile phase, and allow to stand for 30 minutes. Proceed with 2  $\mu$ L of this solution under the above conditions, and calculate the resolution. Use a column giving elution of L-cysteine-*N*-ethylmaleimide complex, ethyl L-cysteine-*N*-ethylmaleimide complex and *N*-ethylmaleimide in this order, complete resolution of each component, and the resolution of the peaks of L-cysteine-*N*-ethylmaleimide complex and ethyl L-cysteine-*N*-ethylmaleimide complex being not less than 3.

**Detection sensitivity:** Adjust the detection sensitivity so that the peak height of ethyl L-cysteine-*N*-ethylmaleimide complex obtained from 2  $\mu$ L of the standard solution is between 10 mm and 20 mm.

**Time span of measurement:** About 3 times as long as the retention time of ethyl L-cysteine-*N*-ethylmaleimide complex.

**Loss on drying** <2.41> Not more than 0.5% (1 g, in vacuum, phosphorus oxide (V), 5 hours).

**Residue on ignition** <2.44> Not more than 0.1% (1 g).


**Assay** Weigh accurately about 0.25 g of Ethyl L-Cysteine Hydrochloride, previously dried, transfer into a glass-stoppered flask, and dissolve in 10 mL of water previously freshly boiled and cooled to a temperature not exceeding 5°C in a stream of nitrogen. Add exactly 20 mL of 0.05 mol/L iodine VS, previously cooled to a temperature not exceeding 5°C, and allow to stand for 30 seconds, then titrate <2.50> with 0.1 mol/L sodium thiosulfate VS, on cooling below 5°C (indicator: 1 mL of starch TS). Perform a blank determination, and make any necessary correction.

Each mL of 0.05 mol/L iodine VS  
= 18.57 mg of C<sub>22</sub>H<sub>34</sub>NO<sub>2</sub>SHCl

**Containers and storage** Containers—Tight containers.

## Ethyl Icosapentate

イコサペント酸エチルエステル



C<sub>22</sub>H<sub>34</sub>O<sub>2</sub>: 330.50

Ethyl (5Z,8Z,11Z,14Z,17Z)-icosa-5,8,11,14,17-pentaenoate  
[86227-47-6]

Ethyl Icosapentate contains not less than 96.5% and not more than 101.0% of C<sub>22</sub>H<sub>34</sub>O<sub>2</sub>.

It may contain a suitable antioxidant.

**Description** Icosapentate is a colorless or pale yellow, clear liquid. It has a faint, characteristic odor.

It is miscible with ethanol (99.5), with acetic acid (100) and with hexane. It is practically insoluble in water and in ethylene glycol.

**Identification** (1) To 20 mg of Ethyl Icosapentate add 3

mL of a solution of potassium hydroxide in ethylene glycol (21 in 100), stopper tightly while passing a current of nitrogen, and heat at 180°C for 15 minutes. After cooling, add methanol to make 100 mL. To 4 mL of this solution add methanol to make 100 mL, and use this solution as the sample solution. Determine the absorption spectrum of the sample solution as directed under Ultraviolet-visible Spectrophotometry <2.24>, using a solution, prepared in the same manner as the sample solution with 3 mL of the solution of potassium hydroxide in ethylene glycol (21 in 100), as a control, and compare the spectrum with the Reference Spectrum or the spectrum of a solution of Ethyl Icosapentate Reference Standard prepared in the same manner as the sample solution: both spectra exhibit similar intensities of absorption at the same wavelengths.

(2) Determine the infrared absorption spectrum of Ethyl Icosapentate as directed in the liquid film method under Infrared Spectrophotometry <2.25>, and compare the spectrum with the Reference Spectrum or the spectrum of Ethyl Icosapentate Reference Standard: both spectra exhibit similar intensities of absorption at the same wave numbers.

**Refractive index** <2.45>  $n_D^{20}$ : 1.481 – 1.491

**Specific gravity** <2.56>  $d_{20}^{20}$ : 0.905 – 0.915

**Acid value** <1.13> Not more than 0.5

**Saponification value** <1.13> 165 – 175

**Iodine value** <1.13> 365 – 395. Perform the test with 20 mg of Ethyl Icosapentate.

**Purity** (1) Heavy metals <1.07>—Mix 1.0 g of Ethyl Icosapentate with ethanol (99.5), and add 2 mL of dilute acetic acid and ethanol (99.5) to make 50 mL. Perform the test with this solution as the test solution. Control solution: To 1.0 mL of Standard Lead Solution add 2 mL of dilute acetic acid and ethanol (99.5) to make 50 mL (not more than 10 ppm).

(2) Arsenic <1.11>—Prepare the test solution with 1.0 g of Ethyl Icosapentate according to Method 3, and perform the test (not more than 2 ppm).

(3) Related substances—To 0.40 g of Ethyl Icosapentate add hexane to make 50 mL, and use this solution as the sample solution. Perform the test with 1.5  $\mu$ L of the sample solution as directed under Gas Chromatography <2.02> according to the following conditions. Determine each peak area by the automatic integration method, and calculate the amounts of these peaks by the area percentage method: the area of the peak, having the relative retention time of about 0.53 with respect to ethyl icosapentate, is not more than 0.5%, the areas of the peaks other than the principal peak and other than the peak mentioned above are not more than 1.0%, and the total amount of these peaks other than the principal peak is not more than 3.5%.

(the rest is omitted)

(4) Peroxide—Weigh accurately about 1 g of Ethyl Icosapentate, put in a 200-mL glass-stoppered conical flask, add 25 mL of a mixture of acetic acid (100) and chloroform (3:2), and dissolve by gentle shaking. Add 1 mL of saturated potassium iodide solution, immediately stopper tightly, shake gently, and allow to stand in a dark place for 10 minutes. Then add 30 mL of water, shake vigorously for 5 to 10 seconds, and titrate <2.50> with 0.01 mol/L sodium thiosulfate VS until the blue color of the solution disappears

after addition of 1 mL of starch TS. Calculate the amount of peroxide by the following equation: not more than 2 meq/kg.

$$\text{Amount (meq/kg) of peroxide} = (V/W) \times 10$$

$V$ : Volume (mL) of 0.01 mol/L sodium thiosulfate VS consumed

$W$ : Amount (g) of the sample

**Residue on ignition** <2.44> Not more than 0.1% (1 g).

**Assay** Weigh accurately about 0.4 g of Ethyl Icosapentate, and add hexane to make exactly 50 mL. Pipet 2 mL of this solution, add exactly 2 mL of the internal standard solution, and use this solution as the sample solution. Separately, weigh accurately about 80 mg of Ethyl Icosapentate Reference Standard, and add hexane to make exactly 10 mL. Pipet 2 mL of this solution, add exactly 2 mL of the internal standard solution, and use this solution as the standard solution. Perform the test with 3  $\mu$ L each of the sample solution and standard solution as directed under Gas Chromatography <2.02> according to the following conditions, and determine the ratios,  $Q_T$  and  $Q_S$ , of the peak area of ethyl icosapentate to that of the internal standard.

$$\begin{aligned} &\text{Amount (mg) of ethyl icosapentate (C}_{22}\text{H}_{34}\text{O}_2) \\ &= W_S \times (Q_T/Q_S) \times 5 \end{aligned}$$

$W_S$ : Amount (mg) of Ethyl Icosapentate Reference Standard

**Internal standard solution**—A solution of methyl docosanate in hexane (1 in 125).

**Operating conditions**—

Detector: A hydrogen-flame ionization detector

Column: A glass column in 4 mm inside diameter and 1.8 m in length, packed with siliceous earth for gas chromatography (175 to 246  $\mu$ m in particle diameter), coated with diethylene glycol succinate polyester for gas chromatography in the ratio of 25%.

Column temperature: A constant temperature of about 190°C

Carrier gas: Nitrogen

Flow rate: Adjust the flow rate so that the retention time of ethyl icosapentate is about 30 minutes.

**System suitability**—

System performance: When the procedure is run with 3  $\mu$ L of the standard solution under the above operating conditions, the internal standard and ethyl icosapentate are eluted in this order with the resolution between these peaks being not less than 3.


System repeatability: When the test is repeated 6 times with 3  $\mu$ L of the standard solution under the above operating conditions, the relative standard deviation of the ratio of the peak area of ethyl icosapentate to that of the internal standard is not more than 1.0%.

**Containers and storage** Containers—Tight containers.

Storage—Being fully filled, or replacing the air with Nitrogen.

## Ethyl Parahydroxybenzoate

パラオキシ安息香酸エチル



$\text{C}_9\text{H}_{10}\text{O}_3$ : 166.17

Ethyl 4-hydroxybenzoate [120-47-8]

This monograph is harmonized with the European Pharmacopoeia and the U.S. Pharmacopeia. The parts of the text that are not harmonized are marked with symbols (♦ ♦).

Ethyl Parahydroxybenzoate contains not less than 98.0% and not more than 102.0% of  $\text{C}_9\text{H}_{10}\text{O}_3$ .

♦**Description** Ethyl Parahydroxybenzoate occurs as colorless crystals or a white, crystalline powder.

It is freely soluble in ethanol (95) and in acetone, and very slightly soluble in water. ♦

**Identification** (1) The melting point <2.60> of Ethyl Parahydroxybenzoate is between 115°C and 118°C.

♦(2) Determine the infrared absorption spectrum of Ethyl Parahydroxybenzoate as directed in the potassium bromide disk method under Infrared Spectrophotometry <2.25>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wave numbers. ♦

**Purity** (1) Clarity and color of solution—Dissolve 1.0 g of Ethyl Parahydroxybenzoate in 10 mL of ethanol (95): the solution is clear and not more intensely colored than the following control solution.

Control solution: To 5.0 mL of cobalt (II) chloride colorimetric stock solution, 12.0 mL of iron (III) chloride colorimetric stock solution and 2.0 mL of copper (II) sulfate colorimetric stock solution add water to make 1000 mL.

(2) Acidity—Dissolve 0.20 g of Ethyl Parahydroxybenzoate in 5 mL of ethanol (95), add 5 mL of freshly boiled and cooled water and 0.1 mL of bromocresol green-sodium hydroxide-ethanol TS, then add 0.1 mL of 0.1 mol/L sodium hydroxide VS: the solution shows a blue color.

♦(3) Heavy metals <1.07>—Dissolve 1.0 g of Ethyl Parahydroxybenzoate in 25 mL of acetone, add 2 mL of dilute acetic acid and water to make 50 mL, and perform the test using this solution as the test solution. Prepare the control solution as follows: to 2.0 mL of Standard Lead Solution add 25 mL of acetone, 2 mL of dilute acetic acid, and water to make 50 mL (not more than 20 ppm). ♦

(4) Related substances—Dissolve 0.10 g of Ethyl Parahydroxybenzoate in 10 mL of acetone, and use this solution as the sample solution. Pipet 0.5 mL of the sample solution, add acetone to make exactly 100 mL, and use this solution as the standard solution. Perform the test with these solutions as directed under Thin-layer Chromatography <2.03>. Spot 2  $\mu$ L each of the sample solution and standard solution on a plate of silica gel with fluorescent indicator for thin-layer chromatography. Develop the plate with a mixture of methanol, water and acetic acid (100) (70:30:1) to a distance

of about 15 cm, and air-dry the plate. Examine under ultraviolet light (main wavelength: 254 nm): the spot other than the principal spot is not more intense than the spot obtained with the standard solution.

**Residue on ignition** <2.44> Not more than 0.1% (1 g).


**Assay** Weigh accurately about 1.0 g of Ethyl Parahydroxybenzoate, add exactly 20 mL of 1 mol/L sodium hydroxide VS, heat at about 70°C for 1 hour, and immediately cool in ice. Titrate <2.50> the excess sodium hydroxide with 0.5 mol/L sulfuric acid VS up to the second equivalent point. (potentiometric titration). Perform a blank determination.

Each mL of 1 mol/L sodium hydroxide VS  
= 166.2 mg of  $C_9H_{10}O_3$

♦**Containers and storage** Containers—Well-closed containers. ♦

## Ethylenediamine

エチレンジアミン



$C_2H_8N_2$ : 60.10

Ethane-1,2-diamine [107-15-3]

Ethylenediamine contains not less than 97.0% of  $C_2H_8N_2$ .

**Description** Ethylenediamine is a clear, colorless to pale yellow liquid. It has an ammonia-like odor.

It is miscible with water, with ethanol (95) and with diethyl ether.

It has a caustic nature and an irritating property.

It is gradually affected by air.

Specific gravity  $d_{20}^{20}$ : about 0.898

**Identification (1)** A solution of Ethylenediamine (1 in 500) is alkaline.

(2) To 2 mL of copper (II) sulfate TS add 2 drops of Ethylenediamine: a blue-purple color develops.

(3) To 0.04 g of Ethylenediamine add 6 drops of benzoyl chloride and 2 mL of a solution of sodium hydroxide (1 in 10), warm for 2 to 3 minutes with occasional shaking, collect the white precipitate formed, and wash with water. Dissolve the precipitate in 8 mL of ethanol (95) by warming, promptly add 8 mL of water, cool, filter the crystals, wash with water, and dry at 105°C for 1 hour: it melts <2.60> between 247°C and 251°C.

**Purity (1)** Heavy metals <1.07>—Place 1.0 g of Ethylenediamine in a porcelain crucible, evaporate to dryness on a water bath, cover loosely, ignite at a low temperature until charred, proceed according to Method 2, and perform the test. Prepare the control solution with 2.0 mL of Standard Lead Solution (not more than 20 ppm).

(2) Residue on evaporation—Pipet 5 mL of Ethylenediamine, heat on a water bath to dryness, and dry to constant mass at 105°C: the mass of the residue does not exceed 3.0 mg.

**Distilling range** <2.57> 114 – 119°C, not less than 95 vol%.

**Assay** Weigh accurately about 0.7 g of Ethylenediamine in a glass-stoppered conical flask, add 50 mL of water, and titrate <2.50> with 1 mol/L hydrochloric acid VS (indicator: 3 drops of bromophenol blue TS).


Each mL of 1 mol/L hydrochloric acid VS  
= 30.05 mg of  $C_2H_8N_2$

**Containers and storage** Containers—Tight containers.  
Storage—Light-resistant, and almost well-filled.

## Ethylmorphine Hydrochloride Hydrate

Dionin

エチルモルヒネ塩酸塩水和物



$C_{19}H_{23}NO_3 \cdot HCl \cdot 2H_2O$ : 385.88

(5R,6S)-4,5-Epoxy-3-ethoxy-17-methyl-7,8-didehydromorphinan-6-ol monohydrochloride dihydrate [125-30-4, anhydride]

Ethylmorphine Hydrochloride Hydrate contains not less than 98.0% of ethylmorphine hydrochloride ( $C_{19}H_{23}NO_3 \cdot HCl$ : 349.86), calculated on the anhydrous basis.

**Description** Ethylmorphine Hydrochloride Hydrate occurs as white to pale yellow crystals or crystalline powder.

It is very soluble in methanol and in acetic acid (100), freely soluble in water, soluble in ethanol (95), sparingly soluble in acetic anhydride, and practically insoluble in diethyl ether.

It is affected by light.

Melting point: about 123°C (with decomposition).

**Identification (1)** Determine the absorption spectrum of a solution of Ethylmorphine Hydrochloride Hydrate (1 in 10,000) as directed under Ultraviolet-visible Spectrophotometry <2.24>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wavelengths.

(2) Determine the infrared absorption spectrum of Ethylmorphine Hydrochloride Hydrate as directed in the potassium bromide disk method under Infrared Spectrophotometry <2.25>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wave numbers.

(3) A solution of Ethylmorphine Hydrochloride Hydrate (1 in 50) responds to the Qualitative Tests <1.09> (2) for chloride.

**Optical rotation** <2.49>  $[\alpha]_D^{20}$ :  $-103 - -106^\circ$  (0.4 g calculated on the anhydrous basis, water, 20 mL, 100 mm).

**pH** <2.54> Dissolve 0.10 g of Ethylmorphine Hydrochloride



Hydrate in 10 mL of water: the pH of this solution is between 4.0 and 6.0.

**Purity** Related substances—Dissolve 0.20 g of Ethylmorphine Hydrochloride Hydrate in 10 mL of diluted ethanol (95) (1 in 2), and use this solution as the sample solution. Pipet 0.5 mL of the sample solution, add diluted ethanol (95) (1 in 2) to make exactly 100 mL, and use this solution as the standard solution. Perform the test with these solutions as directed under Thin-layer Chromatography <2.03>. Spot 10  $\mu$ L each of the sample solution and standard solution on a plate of silica gel with fluorescent indicator for thin-layer chromatography. Develop the plate with a mixture of ethanol (99.5), toluene, acetone and ammonia solution (28) (14:14:7:1) to a distance of about 10 cm, and air-dry the plate. Examine under ultraviolet light (main wavelength: 254 nm): the spots other than the principal spot from the sample solution are not more intense than the spot from the standard solution.

**Water** <2.48> 8.0 – 10.0% (0.25 g, direct titration).

**Residue on ignition** <2.44> Not more than 0.1% (0.5 g).


**Assay** Weigh accurately about 0.5 g of Ethylmorphine Hydrochloride Hydrate, and dissolve in 50 mL of a mixture of acetic anhydride and acetic acid (100) (7:3), and titrate <2.50> with 0.1 mol/L perchloric acid VS (potentiometric titration). Perform a blank determination, and make any necessary correction.

Each mL of 0.1 mol/L perchloric acid VS  
= 34.99 mg of  $C_{19}H_{23}NO_3 \cdot HCl$

**Containers and storage** Containers—Tight containers.  
Storage—Light-resistant.

## Etidronate Disodium

エチドロン酸二ナトリウム



$C_2H_6Na_2O_7P_2$ : 249.99

Disodium dihydrogen (1-hydroxyethylidene)diphosphonate  
[7414-83-7]

Etidronate Disodium, when dried, contains not less than 98.0% and not more than 101.0% of  $C_2H_6Na_2O_7P_2$ .

**Description** Etidronate Disodium occurs as a white powder. It is freely soluble in water, and practically insoluble in ethanol (99.5).

The pH of a solution prepared by dissolving 0.10 g of Etidronate Disodium in 10 mL of water is between 4.4 and 5.4.

It is hygroscopic.

**Identification** (1) To 5 mL of a solution of Etidronate Disodium (1 in 100) add 1 mL of copper (II) sulfate TS, and mix for 10 minutes: a blue precipitate is formed.

(2) Determine the infrared absorption spectrum of Etidronate Disodium, previously dried, as directed in the potassium bromide disk method under Infrared Spec-

trophotometry <2.25>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wave numbers.

(3) A solution of Etidronate Disodium (1 in 100) responds to the Qualitative Tests <1.09> for sodium salt.

**Purity** (1) Heavy metals <1.07>—Proceed with 1.0 g of Etidronate Disodium according to Method 4, and perform the test using the supernatant liquid obtained by centrifuging after addition of 2 mL of dilute acetic acid. Prepare the control solution with 2.0 mL of Standard Lead Solution (not more than 20 ppm).

(2) Arsenic <1.11>—Prepare the test solution with 1.0 g of Etidronate Disodium according to Method 1, and perform the test (not more than 2 ppm).

(3) Phosphite—Weigh accurately about 3.5 g of Etidronate Disodium, dissolve in 100 mL of 0.1 mol/L sodium dihydrogen phosphate TS adjusted the pH to 8.0 with sodium hydroxide TS, add exactly 20 mL of 0.05 mol/L iodine VS, and immediately stopper tightly. Allow to stand in a dark place for 30 minutes, add 1 mL of acetic acid (100), and titrate <2.50> the excess of iodine with 0.1 mol/L sodium thiosulfate VS (indicator: 1 mL of starch TS). Perform a blank determination in the same manner, and make any necessary correction. The amount of phosphite ( $NaH_2PO_3$ ) is not more than 1.0%.

Each mL of 0.05 mol/L iodine VS = 5.199 mg of  $NaH_2PO_3$

(4) Methanol—Weigh accurately about 0.5 g of Etidronate Disodium, dissolve in water to make exactly 5 mL, and use this solution as the sample solution. Separately, pipet 1 mL of methanol, and add water to make exactly 100 mL. Pipet 1 mL of this solution, add water to make exactly 100 mL, and use this solution as the standard solution. Perform the test with exactly 1  $\mu$ L each of the sample solution and standard solution as directed under Gas Chromatography <2.02> according to the following conditions, and determine the peak areas of methanol,  $A_T$  and  $A_S$ , and determine the amount of methanol ( $CH_4O$ ) by the following equation: not more than 0.1%.

Amount (%) of methanol ( $CH_4O$ )  
=  $(1/W) \times (A_T/A_S) \times (1/20) \times 0.79$

$W$ : Amount (g) of sample

0.79: Density (g/mL) of methanol

**Operating conditions**—

Detector: A hydrogen flame-ionization detector.

Column: A glass column 3 mm in inside diameter and 2 m in length, packed with porous copolymer beads for gas chromatography (180 – 250  $\mu$ m in particle diameter).

Column temperature: A constant temperature of about 130°C.

Carrier gas: Nitrogen

Flow rate: Adjust the flow rate so that the retention time of methanol is about 2 minutes.

**System suitability**—

System performance: To 1 mL of methanol and 1 mL of ethanol (99.5) add water to make 100 mL. To 1 mL of this solution add water to make 100 mL. When the procedure is run with 1  $\mu$ L of this solution under the above operating conditions, methanol and ethanol are eluted in this order with the resolution between these peaks being not less than 2.0.

System repeatability: When the test is repeated 6 times with

1  $\mu\text{L}$  of the standard solution under the above operating conditions, the relative standard deviation of the peak area of methanol is not more than 5.0%.

**Loss on drying** <2.41> Not more than 5.0% (0.5 g, 210°C, 2 hours).

**Assay** Weigh accurately about 0.5 g of Etidronate Disodium, previously dried, and dissolve in water to make exactly 50 mL. Transfer exactly 15 mL of this solution to a chromatographic column of 10 mm in internal diameter containing 5 mL of strongly acidic ion exchange resin for column chromatography (H type), allow to flow at a flow rate of about 1.5 mL per minute, and wash the column with two 25-mL portions of water. Combine the eluate and the washings, and titrate <2.50> with 0.1 mol/L sodium hydroxide VS (potentiometric titration). Perform a blank determination in the same manner, and make any necessary correction.

Each mL of 0.1 mol/L sodium hydroxide VS  
= 12.50 mg of  $\text{C}_2\text{H}_6\text{Na}_2\text{O}_7\text{P}_2$

**Containers and storage** Containers—Tight containers.

## Etidronate Disodium Tablets

エチドロン酸二ナトリウム錠

Etidronate Disodium Tablets contains not less than 93.0% and not more than 107.0% of the labeled amount of etidronate disodium ( $\text{C}_2\text{H}_6\text{Na}_2\text{O}_7\text{P}_2$ ; 249.99).

**Method of preparation** Prepare as directed under Tablets, with Etidronate Disodium.

**Identification** (1) Shake an amount of pulverized Etidronate Disodium Tablets, equivalent to 0.2 g of Etidronate Disodium according to the labeled amount, with 20 mL of water, and filter. Proceed with the filtrate as directed in the Identification (1) under Etidronate Disodium.

(2) Shake an amount of pulverized Etidronate Disodium Tablets, equivalent to 0.4 g of Etidronate Disodium according to the labeled amount, with 10 mL of water, and filter. Evaporate total amount of the filtrate to dryness under reduced pressure, shake the residue with 15 mL of ethanol (99.5), centrifuge, and dry the precipitate at 150°C for 4 hours. Determine the infrared absorption spectrum of the precipitate as directed in the potassium bromide disk method under Infrared Spectrophotometry <2.25>: it exhibits absorption at the wave numbers of about  $1170\text{ cm}^{-1}$ ,  $1056\text{ cm}^{-1}$ ,  $916\text{ cm}^{-1}$  and  $811\text{ cm}^{-1}$ .

**Uniformity of dosage unit** <6.02> It meets the requirement of the Mass variation test.

**Dissolution** <6.10> Perform the test according to the following method: it meets the requirement.

Perform the test with 1 tablet of Etidronate Disodium Tablets at 50 revolutions per minute according to the Paddle method, using 900 mL of water as the dissolution medium. Withdraw not less than 20 mL of the dissolution medium 60 minutes after start of the test, and filter through a membrane filter with pore size of not more than  $0.45\text{ }\mu\text{m}$ . Discard the first 10 mL of the filtrate, take exactly  $V\text{ mL}$  of the subse-

quent filtrate, add water to make exactly  $V'\text{ mL}$  so that each mL contains about 0.22 mg of etidronate disodium ( $\text{C}_2\text{H}_6\text{Na}_2\text{O}_7\text{P}_2$ ) according to the labeled amount, and use this solution as the sample solution. Separately, weigh accurately about 30 mg of etidronate disodium for assay, previously dried at 210°C for 2 hours, and dissolve in water to make exactly 100 mL. Dilute exactly a suitable amount of this solution with water to make solutions so that each mL contains about 0.12 mg, about 0.21 mg and about 0.24 mg of etidronate disodium ( $\text{C}_2\text{H}_6\text{Na}_2\text{O}_7\text{P}_2$ ), and use these solutions as the standard solutions. Pipet 2 mL each of the sample solution and standard solutions, add exactly 2 mL of a solution of copper (II) sulfate (7 in 10,000) and water to make exactly 10 mL. Determine the absorbances of these solutions at 233 nm as directed under Ultraviolet-visible Spectrophotometry <2.24> using a solution prepared by diluting exactly 2 mL of the solution of copper (II) sulfate (7 in 10,000) with water to make exactly 10 mL as the control. From the calibration curve obtained with the standard solutions determine the concentration of etidronate disodium,  $C_T$ , in the sample solution. The dissolution rate of Etidronate Disodium Tablets in 60 minutes is not less than 85%.

Dissolution rate (%) with respect to the labeled amount of etidronate disodium ( $\text{C}_2\text{H}_6\text{Na}_2\text{O}_7\text{P}_2$ )  
=  $C_T \times (V'/V) \times (1/C) \times 90$

$C_T$ : Concentration ( $\mu\text{g/mL}$ ) of etidronate disodium ( $\text{C}_2\text{H}_6\text{Na}_2\text{O}_7\text{P}_2$ ) in the sample solution


$C$ : Labeled amount (mg) of etidronate disodium ( $\text{C}_2\text{H}_6\text{Na}_2\text{O}_7\text{P}_2$ ) in 1 tablet

**Assay** Weigh accurately, and powder not less than 20 Etidronate Disodium Tablets. Weigh accurately a portion of the powder, equivalent to about 0.5 g of etidronate disodium ( $\text{C}_2\text{H}_6\text{Na}_2\text{O}_7\text{P}_2$ ), add 30 mL of water, shake vigorously for 10 minutes, add water to make exactly 50 mL, and filter. Proceed with the filtrate as directed in the Assay under Etidronate Disodium.

**Containers and storage** Containers—Tight containers.

## Etilefrine Hydrochloride

エチレフリン塩酸塩



$\text{C}_{10}\text{H}_{15}\text{NO}_2 \cdot \text{HCl}$ : 217.69  
(1*RS*)-2-Ethylamino-1-(3-hydroxyphenyl)ethanol  
monohydrochloride [943-17-9]

Etilefrine Hydrochloride, when dried, contains not less than 98.0% and not more than 101.0% of  $\text{C}_{10}\text{H}_{15}\text{NO}_2 \cdot \text{HCl}$ .

**Description** Etilefrine Hydrochloride occurs as white crystals or crystalline powder.

It is very soluble in water, freely soluble in ethanol (99.5), and sparingly soluble in acetic acid (100).

It is gradually colored to yellow-brown by light.

A solution of Etilefrine Hydrochloride (1 in 20) shows no optical rotation.

**Identification (1)** Dissolve 5 mg of Etilefrine Hydrochloride in 100 mL of diluted hydrochloric acid (1 in 1000). Determine the absorption spectrum of the solution as directed under Ultraviolet-visible Spectrophotometry <2.24>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wavelengths.

(2) Determine the infrared absorption spectrum of Etilefrine Hydrochloride as directed in the potassium chloride disk method under Infrared Spectrophotometry <2.25>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wave numbers.

(3) A solution of Etilefrine Hydrochloride (1 in 1000) responds to the Qualitative Tests <1.09> (2) for chloride.

**Melting point** <2.60> 118 – 122°C

**Purity (1)** Acidity or alkalinity—To 10 mL of a solution of Etilefrine Hydrochloride (1 in 50) add 0.1 mL of methyl red TS for acid or alkali test and 0.2 mL of 0.01 mol/L sodium hydroxide VS: a yellow color develops, and the necessary volume of 0.01 mol/L hydrochloric acid VS to change the color to red is not more than 0.4 mL.

(2) Clarity and color of solution—Dissolve 0.5 g of Etilefrine Hydrochloride in 10 mL of water: the solution is clear and colorless.

(3) Sulfate <1.14>—Perform the test with 0.85 g of Etilefrine Hydrochloride. Prepare the control solution with 0.35 mL of 0.005 mol/L sulfuric acid VS (not more than 0.020%).

(4) Heavy metals <1.07>—Dissolve 1.0 g of Etilefrine Hydrochloride in 30 mL of water and 2 mL of acetic acid (100), adjust with sodium hydroxide TS to a pH of 3.3, add water to make 50 mL, and perform the test. Prepare the control solution with 2.0 mL of Standard Lead Solution (not more than 20 ppm).

**Loss on drying** <2.41> Not more than 0.5% (1 g, 105°C, 4 hours).

**Residue on ignition** <2.44> Not more than 0.1% (1 g).

**Assay** Weigh accurately about 0.15 g of Etilefrine Hydrochloride, previously dried, dissolve in 20 mL of acetic acid (100), add 50 mL of acetic anhydride, and titrate <2.50> with 0.1 mol/L perchloric acid VS (potentiometric titration). Perform a blank determination, and make any necessary correction.

Each mL of 0.1 mol/L perchloric acid VS  
= 21.77 mg of  $C_{10}H_{15}NO_2 \cdot HCl$

**Containers and storage** Containers—Tight containers.  
Storage—Light-resistant.

## Etilefrine Hydrochloride Tablets

エチレフリン塩酸塩錠

Etilefrine Hydrochloride Tablets contain not less than 93.0% and not more than 107.0% of the labeled

amount of etilefrine hydrochloride ( $C_{10}H_{15}NO_2 \cdot HCl$ : 217.69).

**Method of preparation** Prepare as directed under Tablets, with Etilefrine Hydrochloride.

**Identification** To a quantity of powdered Etilefrine Hydrochloride Tablets, equivalent to 5 mg of Etilefrine Hydrochloride according to the labeled amount, add 60 mL of diluted hydrochloric acid (1 in 1000), shake well, add 40 mL of diluted hydrochloric acid (1 in 1000), and filter. Determine the absorption spectrum of the filtrate as directed under Ultraviolet-visible Spectrophotometry <2.24>, using diluted hydrochloric acid (1 in 1000) as the blank: it exhibits a maximum between 271 nm and 275 nm.

**Uniformity of dosage units** <6.02> Perform the test according to the following method: it meets the requirements of the Content uniformity test.

To 1 tablet of Etilefrine Hydrochloride Tablets add 60 mL of diluted hydrochloric acid (1 in 1000), and proceed as directed in the Assay.

Amount (mg) of etilefrine hydrochloride ( $C_{10}H_{15}NO_2 \cdot HCl$ )  
=  $W_s \times (A_T/A_S) \times (1/10)$

$W_s$ : Amount (mg) of etilefrine hydrochloride for assay

**Assay** Weigh accurately the mass of not less than 20 Etilefrine Hydrochloride Tablets, and powder. Weigh accurately a portion of the powder, equivalent to about 5 mg of etilefrine hydrochloride ( $C_{10}H_{15}NO_2 \cdot HCl$ ), add 60 mL of diluted hydrochloric acid (1 in 1000), shake for 10 minutes, add diluted hydrochloric acid (1 in 1000) to make exactly 100 mL, and filter. Discard the first 20 mL of the filtrate, and use the subsequent filtrate as the sample solution. Separately, weigh accurately about 50 mg of etilefrine hydrochloride for assay, previously dried at 105°C for 4 hours, and dissolve in diluted hydrochloric acid (1 in 1000) to make exactly 100 mL. Pipet 10 mL of this solution, add diluted hydrochloric acid (1 in 1000) to make exactly 100 mL, and use this solution as the standard solution. Perform the test with exactly 20  $\mu$ L each of the sample solution and standard solution as directed under Liquid Chromatography <2.01> according to the following conditions, and determine the peak areas,  $A_T$  and  $A_S$ , of etilefrine.

Amount (mg) of etilefrine hydrochloride ( $C_{10}H_{15}NO_2 \cdot HCl$ )  
=  $W_s \times (A_T/A_S) \times (1/10)$

$W_s$ : Amount (mg) of etilefrine hydrochloride for assay

**Operating conditions—**

Detector: An ultraviolet absorption photometer (wavelength: 220 nm).

Column: A stainless steel column 4.6 mm in inside diameter and 25 cm in length, packed with octylsilanized silica gel for liquid chromatography (5  $\mu$ m in particle diameter).

Column temperature: A constant temperature of about 40°C.

Mobile phase: Dissolve 5 g of sodium lauryl sulfate in 940 mL of water and 500 mL of acetonitrile, and adjust the pH to 2.3 with phosphoric acid.

Flow rate: Adjust the flow rate so that the retention time of etilefrine is about 6 minutes.

**System suitability—**

System performance: Dissolve 4 mg of bamethan sulfate

and 4 mg of etilefrine hydrochloride in the mobile phase to make 50 mL. When the procedure is run with 20  $\mu$ L of this solution under the above operating conditions, etilefrine and bamethan are eluted in this order with the resolution between these peaks being not less than 5.


**System repeatability:** When the test is repeated 6 times with 20  $\mu$ L of the standard solution under the above operating conditions, the relative standard deviation of the peak area of etilefrine is not more than 1.0%.

**Containers and storage** Containers—Tight containers.

Storage—Light-resistant.

## Etizolam

エチゾラム



$C_{17}H_{15}ClN_4S$ : 342.85

4-(2-Chlorophenyl)-2-ethyl-9-methyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepine  
[40054-69-1]

Etizolam contains not less than 98.5% and not more than 101.0% of etizolam ( $C_{17}H_{15}ClN_4S$ ).

**Description** Etizolam occurs as a white to pale yellowish white crystalline powder.

It is soluble in ethanol (99.5), sparingly soluble in acetonitrile and in acetic anhydride, and practically insoluble in water.

**Identification** (1) Determine the absorption spectrum of a solution of Etizolam in ethanol (99.5) (1 in 100,000) as directed under Ultraviolet-visible Spectrophotometry <2.24>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wavelengths.

(2) Determine the infrared absorption spectrum of Etizolam as directed in the potassium bromide disk method under Infrared Spectrophotometry <2.25>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wave numbers.

**Melting point** <2.60> 146 – 149°C

**Purity** (1) Heavy metals <1.07>—Proceed with 2.0 g of Etizolam according to Method 2, and perform the test. Prepare the control solution with 2.0 mL of Standard Lead Solution (not more than 10 ppm).

(2) Related substances—Dissolve 20 mg of Etizolam in 50 mL of acetonitrile, and use this solution as the sample solution. Pipet 1 mL of the sample solution, and add acetonitrile to make exactly 20 mL. Pipet 1 mL of this solution, add acetonitrile to make exactly 50 mL, and use this solution as the standard solution. Perform the test with exactly 10  $\mu$ L each of the sample solution and standard solution as directed

under Liquid Chromatography <2.01> according to the following conditions, and determine each peak area by the automatic integration method: the area of the peak other than etizolam obtained from the sample solution is not more than the peak area of etizolam from the standard solution.

**Operating conditions—**

**Detector:** An ultraviolet absorption photometer (wavelength: 240 nm).

**Column:** A stainless steel column 4.6 mm in inside diameter and 15 cm in length, packed with octadecylsilanized silica gel for liquid chromatography (5  $\mu$ m in particle diameter).

**Column temperature:** A constant temperature of about 35°C.

**Mobile phase:** Dissolve 1.36 g of potassium dihydrogen phosphate in water to make 1000 mL, and adjust the pH to 3.5 with diluted phosphoric acid (1 in 10). To 550 mL of this solution add 450 mL of acetonitrile.

**Flow rate:** Adjust the flow rate so that the retention time of etizolam is about 6 minutes.

**Time span of measurement:** About 5 times as long as the retention time of etizolam beginning after the solvent peak.

**System suitability—**

**Test for required detectability:** Measure exactly 2 mL of the standard solution, and add acetonitrile to make exactly 20 mL. Confirm that the peak area of etizolam obtained from 10  $\mu$ L of this solution is equivalent to 8 to 12% of that from 10  $\mu$ L of the standard solution.

**System performance:** Dissolve 0.02 g each of Etizolam and ethyl parahydroxybenzoate in the mobile phase to make 50 mL. To 1 mL of this solution add the mobile phase to make 50 mL. When the procedure is run with 10  $\mu$ L of this solution under the above operating conditions, ethyl parahydroxybenzoate and etizolam are eluted in this order with the resolution between these peaks being not less than 3.

**System repeatability:** When the test is repeated 6 times with 10  $\mu$ L of the standard solution under the above operating conditions, the relative standard deviation of the peak area of etizolam is not more than 2%.

**Loss on drying** <2.41> Not more than 0.5% (1 g, 105°C, 3 hours).

**Residue on ignition** <2.44> Not more than 0.1% (1 g).

**Assay** Weigh accurately about 0.3 g of Etizolam, previously dried, dissolve in 70 mL of a mixture of acetic anhydride and acetic acid (100) (7:3), and titrate <2.50> with 0.1 mol/L perchloric acid VS (potentiometric titration). The end point is the second equivalent point. Perform a blank determination, and make any necessary correction.


Each mL of 0.1 mol/L perchloric acid VS  
= 17.14 mg of  $C_{17}H_{15}ClN_4S$

**Containers and storage** Containers—Tight containers.

Storage—Light-resistant.

## Etodolac

エトドラク



$C_{17}H_{21}NO_3$ : 287.35

2-[(1*R,S*)-1,8-Diethyl-1,3,4,9-tetrahydropyrano[3,4-*b*]indol-1-yl]acetic acid

[41340-25-4]

Etodolac, when dried, contains not less than 98.5% and not more than 101.0% of  $C_{17}H_{21}NO_3$ .

**Description** Etodolac occurs as white to pale yellow, crystals or crystalline powder.

It is freely soluble in methanol and in ethanol (99.5), and practically insoluble in water.

A solution of Etodolac in methanol (1 in 50) shows no optical rotation.

Melting point: about 147°C (with decomposition).

**Identification (1)** Determine the absorption spectrum of a solution of Etodolac in ethanol (99.5) (3 in 200,000) as directed under Ultraviolet-visible Spectrophotometry <2.24>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wavelengths.

(2) Determine the infrared absorption spectrum of Etodolac as directed in the potassium bromide disk method under Infrared Spectrophotometry <2.25>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wave numbers.

**Purity (1)** Heavy metals <1.07>—Proceed with 2.0 g of Etodolac according to Method 2, and perform the test. Prepare the control solution with 2.0 mL of Standard Lead Solution (not more than 10 ppm).

(2) Related substances—Dissolve 0.5 g of Etodolac in 10 mL of methanol, and use this solution as the sample solution. Pipet 2 mL of the sample solution, and add methanol to make exactly 100 mL. Pipet 5 mL of this solution, add methanol to make exactly 20 mL, and use this solution as the standard solution (1). Pipet 4 mL of the standard solution (1), add methanol to make exactly 10 mL, and use this solution as the standard solution (2). Perform the test with these solutions as directed under Thin-layer Chromatography <2.03>. Previously develop a plate of silica gel with fluorescent indicator for thin-layer chromatography in a developing container containing 2 cm depth of a solution of L-ascorbic acid in a mixture of methanol and water (4:1) (1 in 200 mL) to the distance of 3 cm, and air-dry for 30 minutes. Spot 10  $\mu$ L each of the sample solution and standard solutions (1) and (2) on the plate 2.5 cm away from the bottom of the plate, then immediately develop with a mixture of toluene, ethanol (95) and acetic acid (100) (140:60:1) to a distance of about 15 cm, and air-dry the plate. Examine under ultraviolet light (main wavelength: 254 nm): any spot other than the principal spot

obtained with the sample solution is not more intense than the spot with the standard solution (1), and the number of spots which are more intense than the spot with the standard solution (2) is not more than 2.

**Loss on drying** <2.41> Not more than 0.5% (1 g, in vacuum, 60°C, 4 hours).

**Residue on ignition** <2.44> Not more than 0.1% (1 g).

**Assay** Weigh accurately about 0.3 g of Etodolac, previously dried, dissolve in 50 mL of ethanol (99.5), and titrate <2.50> with 0.1 mol/L sodium hydroxide VS (potentiometric titration). Perform a blank determination in the same manner, and make any necessary correction.


Each mL of 0.1 mol/L sodium hydroxide VS

= 28.74 mg of  $C_{17}H_{21}NO_3$

**Containers and storage** Containers—Tight containers. Storage—Light-resistant.

## Etoposide

エトポシド



$C_{29}H_{32}O_{13}$ : 588.56

(5*R*,5*aR*,8*aR*,9*S*)-9-{[4,6-*O*-(1*R*)-Ethylidene- $\beta$ -D-glucopyranosyl]oxy}-5-(4-hydroxy-3,5-dimethoxyphenyl)-5,8,8*a*,9-tetrahydrofuro[3',4':6,7]naphtho[2,3-*d*]-1,3-dioxol-6(5*aH*)-one [33419-42-0]

Etoposide contains not less than 98.0% and not more than 102.0% of  $C_{29}H_{32}O_{13}$ , calculated on the anhydrous basis.

**Description** Etoposide occurs as white crystals or crystalline powder.

It is sparingly soluble in methanol, slightly soluble in ethanol (99.5), and very slightly soluble in water.

Melting point: about 260°C (with decomposition).

**Identification (1)** Determine the absorption spectrum of a solution of Etoposide in methanol (1 in 10,000) as directed under Ultraviolet-visible Spectrophotometry <2.24>, and compare the spectrum with the Reference Spectrum or the spectrum of a solution of Etoposide Reference Standard prepared in the same manner as the sample solution: both spectra exhibit similar intensities of absorption at the same wavelengths.

(2) Determine the infrared absorption spectrum of Etoposide as directed in the potassium bromide disk method

under Infrared Spectrophotometry <2.25>, and compare the spectrum with the Reference Spectrum or the spectrum of Etoposide Reference Standard: both spectra exhibit similar intensities of absorption at the same wave numbers.

**Optical rotation** <2.49>  $[\alpha]_D^{20}$ :  $-100$  –  $-105^\circ$  (0.1 g calculated on the anhydrous basis, methanol, 20 mL, 100 mm).

**Purity** (1) Heavy metals <1.07>—Proceed with 2.0 g of Etoposide according to Method 2, and perform the test. Prepare the control solution with 2.0 mL of Standard Lead Solution (not more than 10 ppm).

(2) Related substances—Dissolve 50 mg of Etoposide in 10 mL of methanol, add the mobile phase to make 50 mL, and use this solution as the sample solution. Pipet 2 mL of the sample solution, add the mobile phase to make exactly 200 mL, and use this solution as the standard solution. Perform the test with exactly 50  $\mu$ L each of the sample solution and standard solution as directed under Liquid Chromatography <2.01> according to the following conditions, and determine each peak area by the automatic integration method: the area of the peak other than etoposide is not larger than 1/5 times the peak area of etoposide with the standard solution, and the total area of the peaks other than the peak of etoposide with the sample solution is not larger than 1/2 times the peak area of etoposide with the standard solution.

**Operating conditions—**

Detector, column, column temperature, mobile phase, and flow rate: Proceed as directed in the operating conditions in the Assay.

Time span of measurement: About 3 times as long as the retention time of etoposide beginning after the solvent peak.

**System suitability—**

Test for required detectability: Measure exactly 1 mL of the standard solution, and add the mobile phase to make exactly 10 mL. Confirm that the peak area of etoposide obtained with 50  $\mu$ L of this solution is equivalent to 7 to 13% of that with 50  $\mu$ L of the standard solution.

System performance: Proceed as directed in the system suitability in the Assay.

System repeatability: When the test is repeated 6 times with 50  $\mu$ L of the standard solution under the above operating conditions, the relative standard deviation of the peak area of etoposide is not more than 2.0%.

**Water** <2.48> Not more than 4.0% (0.5 g, volumetric titration, direct titration).

**Residue on ignition** <2.44> Not more than 0.1% (1 g).

**Assay** Weigh accurately about 25 mg each of Etoposide and Etoposide Reference Standard (previously determined the water <2.48> in the same manner as Etoposide) dissolve separately in methanol to make exactly 25 mL. Pipet 10 mL each of these solutions, add exactly 5 mL of the internal standard solution and the mobile phase to make 50 mL, and use these solutions as the sample solution and standard solution. Perform the test with 50  $\mu$ L each of the sample solution and standard solution as directed under Liquid Chromatography <2.01> according to the following conditions, and determine the ratios,  $Q_T$  and  $Q_S$ , of the peak area of etoposide to that of the internal standard.

$$\text{Amount (mg) of } C_{29}H_{32}O_{13} = W_S \times (Q_T/Q_S)$$

$W_S$ : Amount (mg) of Etoposide Reference Standard,

calculated on the anhydrous basis

**Internal standard solution—**A solution of 2,6-dichlorophenol in methanol (3 in 2500).

**Operating conditions—**

Detector: An ultraviolet absorption photometer (wavelength: 290 nm).

Column: A stainless steel column 3.9 mm in inside diameter and 30 cm in length, packed with phenylsilanized silica gel for liquid chromatography (10  $\mu$ m in particle diameter).

Column temperature: A constant temperature of about 35°C.

Mobile phase: Dissolve 6.44 g of sodium sulfate decahydrate in diluted acetic acid (100) (1 in 100) to make 1000 mL, and add 250 mL of acetonitrile.

Flow rate: Adjust the flow rate so that the retention time of etoposide is about 20 minutes.

**System suitability—**

System performance: Dissolve 10 mg of Etoposide in 2 mL of methanol, add 8 mL of the mobile phase, and mix well. Add 0.1 mL of diluted acetic acid (100) (1 in 25) and 0.1 mL of phenolphthalein TS, and add sodium hydroxide TS until the color of the solution changes to faintly red. After allowing to stand for 15 minutes, add 0.1 mL of diluted acetic acid (100) (1 in 25). When the procedure is run with 10  $\mu$ L of this solution under the above operating conditions, the resolution between the peak of etoposide and the peak having the relative retention time of about 1.3 with respect to etoposide is not less than 3.

System repeatability: When the test is repeated 6 times with 50  $\mu$ L of the standard solution under the above operating conditions, the relative standard deviation of the ratio of the peak area of etoposide to that of the internal standard is not more than 1.0%.

**Containers and storage** Containers—Tight containers.

Storage—Light-resistant.

## Eucalyptus Oil

### *Oleum Eucalypti*

ユーカリ油

Eucalyptus Oil is the essential oil distilled with steam from the leaves of *Eucalyptus globulus* Labillardière or allied plants (*Myrtaceae*).

It contains not less than 70.0% of cineol ( $C_{10}H_{18}O$ : 154.25).

**Description** Eucalyptus Oil is a clear, colorless or pale yellow liquid. It has a characteristic, aromatic odor and a pungent taste.

It is neutral.

**Identification** Shake 1 mL of Eucalyptus Oil vigorously with 1 mL of phosphoric acid, and allow to stand: the solution congeals within 30 minutes.

**Refractive index** <2.45>  $n_D^{20}$ : 1.458 – 1.470

**Specific gravity** <1.13>  $d_{20}^{20}$ : 0.907 – 0.927

**Purity** (1) Clarity of solution—Mix 1.0 mL of Eucalyptus

Oil with 5 mL of diluted ethanol (7 in 10): the solution is clear.

(2) Heavy metals <1.07>—Proceed with 1.0 mL of Eucalyptus Oil according to Method 2, and perform the test. Prepare the control solution with 4.0 mL of Standard Lead Solution (not more than 40 ppm).

**Assay** Weigh accurately about 0.1 g of Eucalyptus Oil, and dissolve in hexane to make exactly 25 mL. Pipet 5 mL of this solution, add exactly 5 mL of the internal standard solution, then add hexane to make 100 mL, and use this solution as the sample solution. Separately, weigh accurately about 0.1 g of cineol for assay, proceed as directed in the sample solution, and use this solution as the standard solution. Perform the test with 2  $\mu$ L each of the sample solution and standard solution as directed under Gas Chromatography <2.02> according to the following conditions. Calculate the ratios,  $Q_T$  and  $Q_S$ , of the peak area of cineol to that of the internal standard of each solutions, respectively.

$$\begin{aligned} &\text{Amount (mg) of cineol (C}_{10}\text{H}_{18}\text{O)} \\ &= W_S \times (Q_T/Q_S) \end{aligned}$$

$W_S$ : Amount (mg) of cineol for assay

**Internal standard solution**—A solution of anisol in hexane (1 in 250).

**Operating conditions**—

Detector: A hydrogen flame-ionization detector.

Column: A glass column about 3 mm in inside diameter and about 5 m in length, having alkylene glycol phthalate ester for gas chromatography coated at the ratio of 10% on silanized siliceous earth for gas chromatography (150 to 180  $\mu$ m in particle diameter).

Column temperature: A constant temperature of about 120°C.

Carrier gas: Nitrogen.

Flow rate: Adjust the flow rate so that the retention time of cineol is about 11 minutes.


Selection of column: Dissolve 0.1 g each of cineol and limonene in 25 mL of hexane. To 1 mL of this solution add hexane to make 20 mL. Proceed with about 2  $\mu$ L of this solution under the above operating conditions, and calculate the resolution. Use a column giving elution of limonene and cineol in this order with the resolution between these peaks being not less than 1.5.

**Containers and storage** Containers—Tight containers.

Storage—Light-resistant.

## Famotidine

ファミモチジン



$\text{C}_8\text{H}_{15}\text{N}_7\text{O}_2\text{S}_3$ : 337.45

*N*-Aminosulfonyl-3-[(2-diaminomethyleneamino)-1,3-thiazol-4-yl]methylsulfanylpropanimidamide  
[76824-35-6]

Famotidine, when dried, contains not less than

98.5% of  $\text{C}_8\text{H}_{15}\text{N}_7\text{O}_2\text{S}_3$ .

**Description** Famotidine occurs as white to yellowish white crystals.

It is freely soluble in acetic acid (100), slightly soluble in ethanol (95), and very slightly soluble in water.

It dissolves in 0.5 mol/L hydrochloric acid TS.

It is gradually colored by light.

Melting point: about 164°C (with decomposition).

**Identification** (1) Determine the absorption spectrum of a solution of Famotidine in 0.05 mol/L potassium dihydrogenphosphate TS (1 in 50,000) as directed under Ultraviolet-visible Spectrophotometry <2.24>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wavelengths.

(2) Determine the infrared absorption spectrum of Famotidine, previously dried, as directed in the potassium bromide disk method under Infrared Spectrophotometry <2.25>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wave numbers.

**Purity** (1) Clarity and color of solution—Dissolve 0.5 g of Famotidine in 10 mL of 0.5 mol/L hydrochloric acid TS: the solution is clear and colorless to pale yellow.

(2) Heavy metals <1.07>—Proceed with 2.0 g of Famotidine according to Method 2, and perform the test. Prepare the control solution with 2.0 mL of Standard Lead Solution (not more than 10 ppm).

(3) Related substances—Dissolve 0.20 g of Famotidine in 10 mL of acetic acid (100), and use this solution as the sample solution. Pipet 1 mL of the sample solution, and add acetic acid (100) to make exactly 100 mL. Pipet 1 mL, 2 mL and 3 mL of this solution, add acetic acid (100) to make exactly 10 mL, respectively, and use these solutions as the standard solution (1), the standard solution (2) and the standard solution (3). Perform the test with these solutions as directed under Thin-layer Chromatography <2.03>. Spot 5  $\mu$ L each of the sample solution and standard solutions (1), (2) and (3) on a plate of silica gel (5 to 7  $\mu$ m) with fluorescent indicator for thin-layer chromatography, and dry in a stream of nitrogen.

Develop the plate with a mixture of ethyl acetate, methanol, toluene and ammonia solution (28) (40:25:20:2) to a distance of about 8 cm, and air-dry the plate. Examine under ultraviolet light (main wavelength: 254 nm): the spots other than the principal spot and other than the spot of the starting point from the sample solution are not more intense than the spot from the standard solution (3). Total intensity of the spots other than the principal spot and other than the spot of the starting point from the sample solution is not more than 0.5% calculated on the basis of intensities of the spots from the standard solution (1) and the standard solution (2) (each spot is equivalent to 0.1% and 0.2%, respectively).

**Loss on drying** <2.41> Not more than 0.5% (1 g, in vacuum, phosphorus (V) oxide, 80°C, 4 hours).

**Residue on ignition** <2.44> Not more than 0.1% (1 g).

**Assay** Weigh accurately about 0.3 g of Famotidine, previously dried, dissolve in 50 mL of acetic acid (100), and titrate <2.50> with 0.1 mol/L perchloric acid VS (potentiometric titration). Perform a blank determination, and make any necessary correction.

Each mL of 0.1 mol/L perchloric acid VS  
= 16.87 mg of  $C_8H_{15}N_7O_2S_3$

**Containers and storage** Containers—Tight containers.  
Storage—Light-resistant.

## Famotidine for Injection

注射用ファモチジン

Famotidine for Injection is a preparation for injection which is dissolved before use.

It contains not less than 94.0% and not more than 106.0% of the labeled amount of famotidine ( $C_8H_{15}N_7O_2S_3$ ; 337.45).

**Method of preparation** Prepare as directed under Injection, with Famotidine.

**Description** Famotidine for Injection occurs as white porous masses or powder.

**Identification** Dissolve an amount of Famotidine for Injection, equivalent to 0.01 g of Famotidine according to the labeled amount, in 50 mL of 0.05 mol/L potassium dihydrogenphosphate TS. To 5 mL of this solution add 0.05 mol/L potassium dihydrogenphosphate TS to make 50 mL, and determine the absorption spectrum of this solution as directed under Ultraviolet-visible Spectrophotometry <2.24>; it exhibits a maximum between 263 nm and 267 nm.

**pH** <2.54> Dissolve an amount of Famotidine for Injection, equivalent to 0.02 g of Famotidine according to the labeled amount, in 1 mL of water: the pH of this solution is between 4.9 and 5.5.

**Purity** (1) Clarity and color of solution—Dissolve an amount of Famotidine for Injection, equivalent to 0.02 g of Famotidine according to the labeled amount, in 1 mL of water: the solution is clear and colorless.

(2) Related substances—Take a number of Famotidine for Injection, equivalent to about 0.1 g of famotidine ( $C_8H_{15}N_7O_2S_3$ ), dissolve each content in water, wash the inside of the container with water, combine the solutions of the contents with the washings, add water to the combined solution to make exactly 100 mL, and use this solution as the sample solution. Pipet 1 mL of the sample solution, add water to make exactly 100 mL, and use this solution as the standard solution. Perform the test with exactly 5  $\mu$ L each of the sample solution and standard solution as directed under Liquid Chromatography <2.01> according to the following conditions, and determine each peak area of these solutions by the automatic integration method: the total area of the peaks other than peak of famotidine from the sample solution is not larger than peak area of famotidine from the standard solution.

**Operating conditions—**

Detector, column, column temperature, mobile phase, and flow rate: Proceed as directed in the operating conditions in the Assay.

Time span of measurement: About 2 times as long as the retention time of famotidine beginning after the solvent peak.

**System suitability—**

Test for required detection: To exactly 2 mL of the standard solution add the water to make exactly 20 mL. Confirm that the peak area of famotidine obtained from 5  $\mu$ L of this solution is equivalent to 8 to 12% of that of famotidine obtained from 5  $\mu$ L of the standard solution.

System performance: Proceed as directed in the system suitability in the Assay.

System repeatability: When the test is repeated 6 times with 5  $\mu$ L of the standard solution under the above operating conditions, the relative standard deviation of the peak area of famotidine is not more than 2.0%.

**Water** <2.48> Not more than 1.5% (0.1 g, coulometric titration).

**Bacterial endotoxins** <4.01> Not more than 15 EU/mg.

**Assay** Take a number of Famotidine for Injection, equivalent to about 0.1 g of famotidine ( $C_8H_{15}N_7O_2S_3$ ), dissolve each content in water, wash the inside of each container with water, combine the solutions of the contents with the washings, add water to the combined solution to make exactly 100 mL, and use this solution as the sample stock solution. Pipet 5 mL of this solution, add exactly 5 mL of the internal standard solution, add the mobile phase to make exactly 50 mL, and use this solution as the sample solution. Separately, weigh accurately about 50 mg of famotidine for assay, previously dried in vacuum with phosphorus (V) oxide at 80°C for 4 hours, dissolve in the mobile phase to make exactly 50 mL. Pipet 5 mL of this solution, add 5 mL of the internal standard solution, add the mobile phase to make exactly 50 mL, and use this solution as the standard solution. Perform the test with 5  $\mu$ L each of the sample solution and standard solution as directed under Liquid Chromatography <2.01> according to the following conditions, and calculate the ratios,  $Q_T$  and  $Q_S$ , of the peak area of famotidine to that of the internal standard.

$$\begin{aligned} &\text{Amount (mg) of famotidine (C}_8\text{H}_{15}\text{N}_7\text{O}_2\text{S}_3\text{)} \\ &= W_S \times (Q_T/Q_S) \times 2 \end{aligned}$$

$W_S$ : Amount (mg) of famotidine for assay

**Internal standard solution—**To 5 mL of a solution of methyl parahydroxybenzoate in methanol (1 in 500) add water to make 50 mL.

**Operating conditions—**

Detector: An ultraviolet absorption photometer (wavelength: 254 nm).

Column: A stainless steel column 4.6 mm in inside diameter and 15 cm in length, packed with octadecylsilanized silica gel for liquid chromatography (5  $\mu$ m in particle diameter).

Column temperature: A constant temperature of about 25°C.

Mobile phase: Dissolve 2 g of sodium 1-heptane sulfonate in 900 mL of water, adjust to pH 3.0 with acetic acid (100), and add water to make 1000 mL. To this solution add 240 mL of acetonitrile and 40 mL of methanol.

Flow rate: Adjust the flow rate so that the retention time of famotidine is about 6 minutes.

**System suitability—**

System performance: When the procedure is run with 5  $\mu$ L of the standard solution under the above operating condi-



tions, famotidine and the internal standard are eluted in this order with the resolution between these peaks being not less than 11.

**System repeatability:** When the test is repeated 6 times with 5  $\mu$ L of the standard solution under the above operating conditions, the relative standard deviation of the ratios of the peak area of famotidine to that of the internal standard is not more than 1.0%.

**Containers and storage** Containers—Hermetic containers.

## Famotidine Powder

ファミチジン散

Famotidine Powder contains not less than 94.0% and not more than 106.0% of the labeled amount of famotidine ( $C_8H_{15}N_7O_2S_3$ ; 337.45).

**Method of preparation** Prepare as directed under Powder, with Famotidine.

**Identification** Weigh a portion of Famotidine Powder, equivalent to 0.01 g of Famotidine according to the labeled amount, add 50 mL of 0.05 mol/L potassium dihydrogenphosphate TS, shake well, and centrifuge. To 5 mL of the supernatant liquid add 0.05 mol/L potassium dihydrogenphosphate TS to make 50 mL, and determine the absorption spectrum of this solution as directed under Ultraviolet-visible Spectrophotometry <2.24>: it exhibits a maximum between 263 nm and 267 nm.

**Dissolution** <6.10> Perform the test according to the following method: it meets the requirement.

Weigh accurately an amount of Famotidine Powder, equivalent to about 20 mg of famotidine ( $C_8H_{15}N_7O_2S_3$ ) according to the labeled amount, and perform the test at 50 revolutions per minute according to the Paddle method, using 900 mL of 0.05 mol/L acetic acid-sodium acetate buffer solution, pH 4.0 as the dissolution medium. Withdraw not less than 20 mL of the dissolution medium 15 minutes after starting the test, and filter through a membrane filter with pore size of not more than 0.5  $\mu$ m. Discard the first 10 mL of the filtrate, and use the subsequent filtrate as the sample solution. Separately, weigh accurately about 40 mg of famotidine for assay, previously dried in vacuum on phosphorus (V) oxide at 80°C for 4 hours, dissolve in 0.05 mol/L acetic acid-sodium acetate buffer solution, pH 4.0 to make exactly 100 mL. Pipet 5 mL of this solution, add the same buffer solution to make exactly 100 mL, and use this solution as the standard solution. Determine the absorbances,  $A_T$  and  $A_S$ , of the sample solution and standard solution at 266 nm as directed under Ultraviolet-visible Spectrophotometry <2.24>. The dissolution rate of a 20-mg/g powder and a 100-mg/g powder in 15 minutes of Famotidine Powder are not less than 80% and not less than 85%, respectively.

Dissolution rate (%) with respect to the labeled amount of famotidine ( $C_8H_{15}N_7O_2S_3$ )

$$= (W_S / W_T) \times (A_T / A_S) \times (1 / C) \times 45$$

$W_S$ : Amount (mg) of famotidine for assay

$W_T$ : Amount (mg) of the sample.

C: Labeled amount (mg) of famotidine ( $C_8H_{15}N_7O_2S_3$ ) in

1 g

**Assay** Weigh accurately a portion of Famotidine Powder, equivalent to about 20 mg of famotidine ( $C_8H_{15}N_7O_2S_3$ ), add 20 mL of water, and shake well. Add 20 mL of methanol, then shake well, add methanol to make exactly 50 mL, and centrifuge. Pipet 5 mL of the supernatant liquid, add exactly 2 mL of the internal standard solution, add the mobile phase to make 20 mL, and use this solution as the sample solution. Separately, weigh accurately about 0.1 g of famotidine for assay, previously dried in vacuum with phosphorus (V) oxide at 80°C for 4 hours, dissolve in methanol to make exactly 50 mL. Pipet 10 mL of this solution, and add methanol to make exactly 50 mL. Pipet 5 mL of this solution, add exactly 2 mL of the internal standard solution, add the mobile phase to make 20 mL, and use this solution as the standard solution. Perform the test with 5  $\mu$ L each of the sample solution and standard solution as directed under Liquid Chromatography <2.01> according to the following conditions, and calculate the ratios,  $Q_T$  and  $Q_S$ , of the peak area of famotidine to that of the internal standard.

$$\begin{aligned} &\text{Amount (mg) of famotidine (} C_8H_{15}N_7O_2S_3 \text{)} \\ &= W_S \times (Q_T / Q_S) \times (1 / 5) \end{aligned}$$

$W_S$ : Amount (mg) of famotidine for assay

**Internal standard solution**—To 5 mL of a solution of methyl parahydroxybenzoate in methanol (1 in 500) add water to make 50 mL.

**Operating conditions**—

**Detector:** An ultraviolet absorption photometer (wavelength: 254 nm).

**Column:** A stainless steel column 4.6 mm in inside diameter and 15 cm in length, packed with octadecylsilanized silica gel for liquid chromatography (5  $\mu$ m in particle diameter).

**Column temperature:** A constant temperature of about 25°C.

**Mobile phase:** Dissolve 2 g of sodium 1-heptane sulfonate in 900 mL of water, adjust to pH 3.0 with acetic acid (100), and add water to make 1000 mL. To this solution add 240 mL of acetonitrile and 40 mL of methanol.

**Flow rate:** Adjust the flow rate so that the retention time of famotidine is about 6 minutes.

**System suitability**—

**System performance:** When the procedure is run with 5  $\mu$ L of the standard solution under the above operating conditions, famotidine and the internal standard are eluted in this order with the resolution between these peaks being not less than 11.

**System repeatability:** When the test is repeated 6 times with 5  $\mu$ L of the standard solution under the above operating conditions, the relative standard deviation of the ratios of the peak area of famotidine to that of the internal standard is not more than 1.0%.

**Containers and storage** Containers—Tight containers.

## Famotidine Tablets

ファミチジン錠

Famotidine Tablets contain not less than 94.0% and

not more than 106.0% of the labeled amount of famotidine ( $C_8H_{15}N_7O_2S_3$ ; 337.45).

**Method of preparation** Prepare as directed under Tablets, with Famotidine.

**Identification** Weigh a portion of powdered Famotidine Tablets, equivalent to 0.01 g of Famotidine according to the labeled amount, add 50 mL of 0.05 mol/L potassium dihydrogenphosphate TS, shake well, and centrifuge. To 5 mL of the supernatant liquid add 0.05 mol/L potassium dihydrogenphosphate TS to make 50 mL, and determine the absorption spectrum of this solution as directed under Ultraviolet-visible Spectrophotometry <2.24>: it exhibits a maximum between 263 nm and 267 nm.

**Uniformity of dosage units** <6.02> Perform the test according to the following method: it meets the requirement of the Content uniformity test.

To 1 tablet of Famotidine Tablets add 2 mL of water, shake to disintegrate, then add a suitable amount of methanol, and shake well. Add methanol to make exactly  $V$  mL of a solution containing about 0.2 mg of famotidine ( $C_8H_{15}N_7O_2S_3$ ) per mL, and centrifuge. Pipet 10 mL of the supernatant liquid, add exactly 2 mL of the internal standard solution, add the mobile phase to make 20 mL, and use this solution as the sample solution. Separately, weigh accurately about 0.1 g of famotidine for assay, previously dried in vacuum with phosphorus (V) oxide at 80°C for 4 hours, dissolve in methanol to make exactly 100 mL. Pipet 10 mL of this solution, and add methanol to make exactly 50 mL. Pipet 10 mL of this solution, add exactly 2 mL of the internal standard solution, add the mobile phase to make 20 mL, and use this solution as the standard solution. Perform the test with 5  $\mu$ L each of the sample solution and standard solution as directed under Liquid Chromatography <2.01> according to the operating conditions described in the Assay, and calculate the ratios,  $Q_T$  and  $Q_S$ , of the peak area of famotidine to that of the internal standard.

$$\begin{aligned} &\text{Amount (mg) of famotidine (C}_8\text{H}_{15}\text{N}_7\text{O}_2\text{S}_3\text{)} \\ &= W_S \times (Q_T/Q_S) \times (V/500) \end{aligned}$$

$W_S$ : Amount (mg) of famotidine for assay

**Internal standard solution**—To 5 mL of a solution of methyl parahydroxybenzoate in methanol (1 in 500) add water to make 50 mL.

**Dissolution** Being specified separately.

**Assay** Take a number of Famotidine Tablets, equivalent to 0.2 g of famotidine ( $C_8H_{15}N_7O_2S_3$ ), add 50 mL of water, and disintegrate by shaking well. Add 100 mL of methanol, then shake well, add methanol to make exactly 200 mL, and centrifuge. Pipet 5 mL of the supernatant liquid, add exactly 5 mL of the internal standard solution, add the mobile phase to make 50 mL, and use this solution as the sample solution. Separately, weigh accurately about 0.1 g of famotidine for assay, previously dried in vacuum with phosphorus (V) oxide at 80°C for 4 hours, dissolve in methanol to make exactly 100 mL. Pipet 5 mL of this solution, add exactly 5 mL of the internal standard solution, add the mobile phase to make 50 mL, and use this solution as the standard solution. Perform the test with 5  $\mu$ L each of the sample solution and standard solution as directed under Liquid Chromatography <2.01> according to the following conditions, and calculate the ratios,

$Q_T$  and  $Q_S$ , of the peak area of famotidine to that of the internal standard.

$$\begin{aligned} &\text{Amount (mg) of famotidine (C}_8\text{H}_{15}\text{N}_7\text{O}_2\text{S}_3\text{)} \\ &= W_S \times (Q_T/Q_S) \times 2 \end{aligned}$$

$W_S$ : Amount (mg) of famotidine for assay

**Internal standard solution**—To 5 mL of a solution of methyl parahydroxybenzoate in methanol (1 in 500) add water to make 50 mL.

**Operating conditions**—

**Detector:** An ultraviolet absorption photometer (wavelength: 254 nm).

**Column:** A stainless steel column 4.6 mm in inside diameter and 15 cm in length, packed with octadecylsilanized silica gel for liquid chromatography (5  $\mu$ m in particle diameter).

**Column temperature:** A constant temperature of about 25°C.

**Mobile phase:** Dissolve 2 g of sodium 1-heptane sulfonate in 900 mL of water, adjust to pH 3.0 with acetic acid (100), and add water to make 1000 mL. To this solution add 240 mL of acetonitrile and 40 mL of methanol.

**Flow rate:** Adjust the flow rate so that the retention time of famotidine is about 6 minutes.

**System suitability**—


**System performance:** When the procedure is run with 5  $\mu$ L of the standard solution under the above operating conditions, famotidine and the internal standard are eluted in this order with the resolution between these peaks being not less than 11.

**System repeatability:** When the test is repeated 6 times with 5  $\mu$ L of the standard solution under the above operating conditions, the relative standard deviation of the ratios of the peak area of famotidine to that of the internal standard is not more than 1.0%.

**Containers and storage** Containers—Tight containers.

## Faropenem Sodium Hydrate

ファロペネムナトリウム水和物



$C_{12}H_{14}NNaO_5S \cdot \frac{1}{2}H_2O$ : 352.34

Monosodium (5*R*,6*S*)-6-[(1*R*)-1-hydroxyethyl]-7-oxo-3-[(2*R*)-tetrahydrofuran-2-yl]-4-thia-1-azabicyclo[3.2.0]hept-2-ene-2-carboxylate hemipentahydrate [122547-49-3, anhydride]

Faropenem Sodium Hydrate contains not less than 870  $\mu$ g (potency) and not more than 943  $\mu$ g (potency) per mg, calculated on the anhydrous basis. The potency of Faropenem Sodium Hydrate is expressed as mass (potency) of faropenem ( $C_{12}H_{15}NO_5S$ ; 285.32).

**Description** Faropenem Sodium Hydrate occurs as white to light yellow, crystals or crystalline powder.

It is freely soluble in water and in methanol, slightly solu-

ble in ethanol (95), and practically insoluble in diethyl ether.

**Identification (1)** Dissolve 5 mg of Faropenem Sodium Hydrate in 1 mL of hydroxylammonium chloride-ethanol TS, allow to stand for 3 minutes, add 1 mL of acidic ammonium iron (III) sulfate TS, and shake: a red-brown to brown color develops.

(2) Determine the absorption spectra of solutions of Faropenem Sodium Hydrate and Faropenem Sodium Reference Standard (1 in 20,000) as directed under Ultraviolet-visible Spectrophotometry <2.24>, and compare the spectra: both spectra exhibit similar intensities of absorption at the same wavelengths.

(3) Determine the infrared absorption spectra of Faropenem Sodium Hydrate and Faropenem Sodium Reference Standard as directed in the potassium bromide disk method under Infrared Spectrophotometry <2.25>, and compare the spectra: both spectra exhibit similar intensities of absorption at the same wave numbers.

**Optical rotation** <2.49>  $[\alpha]_D^{20}$ : +145 – +150° (0.5 g calculated as the anhydrous basis, water, 50 mL, 100 mm).

**Purity (1)** Clarity and color of solution—Being specified separately.

(2) Heavy metals <1.07>—Proceed with 2.0 g of Faropenem Sodium Hydrate according to Method 4, and perform the test. Prepare the control solution with 2.0 mL of Standard Lead Solution (not more than 10 ppm).

(3) Related substances—Being specified separately.

**Water** <2.48> Not less than 12.6% and not more than 13.1% (20 mg, coulometric titration).

**Assay** Weigh accurately an amount of Faropenem Sodium Hydrate and Faropenem Sodium Reference Standard, equivalent to about 25 mg (potency), each of these, add exactly 10 mL each of the internal standard solution, add water to make 50 mL, and use these solutions as the sample solution and the standard solution, respectively. Perform the test with 20  $\mu$ L of the sample solution and standard solution as directed under Liquid Chromatography <2.01> according to the following conditions, and calculate the ratios,  $Q_T$  and  $Q_S$ , of the peak area of faropenem to that of the internal standard.

$$\begin{aligned} &\text{Amount } [\mu\text{g (potency)}] \text{ of faropenem (C}_{12}\text{H}_{15}\text{NO}_5\text{S)} \\ &= W_S \times (Q_T/Q_S) \times 1000 \end{aligned}$$

$W_S$ : amount (mg) of Faropenem Sodium Reference Standard

**Internal standard solution**—Dissolve 0.5 g of *m*-hydroxyacetophenone in 20 mL of acetonitrile, and add water to make 200 mL.

**Operating conditions**—

**Detector:** An ultraviolet absorption photometer (wavelength: 305 nm).

**Column:** A stainless steel column 4.6 mm in inside diameter and 25 cm in length, packed with octadecylsilanized silica gel for liquid chromatography (5  $\mu$ m in particle diameter).

**Column temperature:** A constant temperature of about 40°C.

**Mobile phase:** Dissolve 4.8 g of potassium dihydrogenphosphate, 5.4 g of disodium hydrogen phosphate dodecahydrate and 1.0 g of tetra *n*-butyl ammonium bromide in water

to make 1000 mL. To 870 mL of this solution add 130 mL of acetonitrile.

**Flow rate:** Adjust the flow rate so that the retention time of faropenem is about 11 minutes.

**System suitability**—

**System performance:** When the procedure is run with 20  $\mu$ L of the standard solution under the above operating conditions, the internal standard and faropenem are eluted in this order with the resolution between these peaks being not less than 1.5.

**System repeatability:** When the test is repeated 6 times with 20  $\mu$ L of the standard solution under the above operating conditions, the relative standard deviation of the ratios of the peak area of faropenem to that of the internal standard is not more than 1.0%.

**Containers and storage** Containers—Tight containers.

## Faropenem Sodium for Syrup

シロップ用ファロペネムナトリウム

Faropenem Sodium for Syrup is a preparation for syrup, which is dissolved before use. It contains not less than 93.0% and not more than 106.0% of the labeled amount of faropenem (C<sub>12</sub>H<sub>15</sub>NO<sub>5</sub>S: 285.32).

**Method of preparation** Prepare as directed under Syrups, with Faropenem Sodium Hydrate.

**Identification** Dissolve an amount of pulverized Faropenem Sodium for Syrup, equivalent to 25 mg (potency) of Faropenem Sodium Hydrate according to the labeled amount, in 50 mL of water. To 5 mL of this solution add water to make 50 mL, centrifuge, if necessary, and determine the absorption spectrum of the solution so obtained as directed under Ultraviolet-visible Spectrophotometry <2.24>: it exhibits maxima between 254 nm and 258 nm, and between 304 nm and 308 nm.

**Water** <2.48> Not less than 1.5% and not more than 2.1% (80 mg, coulometric titration).

**Uniformity of dosage units** <6.02> Faropenem Sodium for Syrup in single-unit container meets the requirement of the Mass variation test.

**Assay** Powder, if necessary, and weigh accurately an amount of Faropenem Sodium for Syrup, equivalent to about 25 mg (potency) of faropenem (C<sub>12</sub>H<sub>15</sub>NO<sub>5</sub>S), add exactly 10 mL of the internal standard solution and a suitable amount of water, shake well, and add water to make 50 mL. Filter, discard the first 10 mL of the filtrate, and use the subsequent filtrate as the sample solution. Separately, weigh accurately about 25 mg (potency) of Faropenem Sodium Reference Standard, add exactly 10 mL of the internal standard solution and water to make 50 mL, and use this solution as the standard solution. Proceed as directed in the Assay under Faropenem Sodium Hydrate.

$$\begin{aligned} &\text{Amount [mg (potency)] of faropenem (C}_{12}\text{H}_{15}\text{NO}_5\text{S)} \\ &= W_S \times (Q_T/Q_S) \end{aligned}$$

$W_S$ : Amount [mg (potency)] of Faropenem Sodium Reference Standard

**Internal standard solution**—Dissolve 0.5 g of *m*-hydroxycetophenone in 20 mL of acetonitrile, and add water to make 200 mL.

**Containers and storage** Containers—Tight containers.  
Storage—Light-resistant.

## Faropenem Sodium Tablets

ファロペネムナトリウム錠

Faropenem Sodium Tablets contain not less than 94.0% and not more than 106.0% of the labeled amount of faropenem ( $C_{12}H_{15}NO_5S$ ; 285.32).

**Method of preparation** Prepare as directed under Tablets, with Faropenem Sodium Hydrate.

**Identification** To pulverized Faropenem Sodium Tablets, equivalent to 70 mg (potency) of Faropenem Sodium Hydrate according to the labeled amount, add water to make 100 mL. To 5 mL of this solution add water to make 100 mL, filter, if necessary, and determine the absorption spectrum of this solution as directed under Ultraviolet-visible Spectrophotometry <2.24>: it exhibits maxima between 254 nm and 258 nm and between 304 nm and 308 nm.

**Uniformity of dosage units** <6.02> Perform the test according to the following method: it meets the requirement of the Content uniformity test.

To 1 tablet of Faropenem Sodium Tablets add 180 mL of water, shake vigorously until the tablets are disintegrated, and add water to make exactly *V* mL so that each mL contains about 1 mg (potency) of Faropenem Sodium Hydrate. Pipet 5 mL of this solution, add water to make exactly 100 mL, and filter. Discard the first 10 mL of the filtrate, and use the subsequent filtrate as the sample solution. Separately, weigh accurately about 25 mg (potency) of Faropenem Sodium Reference Standard, and dissolve in water to make exactly 50 mL. Pipet 10 mL of this solution, add water to make exactly 100 mL, and use this solution as the standard solution. Determine the absorbances,  $A_{T275}$ ,  $A_{T305}$ ,  $A_{T354}$ ,  $A_{S275}$ ,  $A_{S305}$  and  $A_{S354}$ , of the sample solution and standard solution at 275 nm, 305 nm and 354 nm as directed under Ultraviolet-visible Spectrophotometry <2.24>, and calculate  $A_T$  and  $A_S$ , using the following equations.

$$A_T = A_{T305} - (49 \times A_{T275} + 30 \times A_{T354})/79$$

$$A_S = A_{S305} - (49 \times A_{S275} + 30 \times A_{S354})/79$$

$$\text{Amount [mg (potency)] of faropenem (C}_{12}\text{H}_{15}\text{NO}_5\text{S)} \\ = W_S \times (A_T/A_S) \times (V/25)$$

$W_S$ : Amount [mg (potency)] of Faropenem Sodium Reference Standard

**Assay** Weigh accurately the mass of not less than 5 Faropenem Sodium Tablets, and powder. Weigh accurately a portion of the powder, equivalent to about 25 mg (potency) of faropenem ( $C_{12}H_{15}NO_5S$ ), add exactly 10 mL of the internal standard solution, shake well, and add water to make exactly 50 mL. Filter, discard the first 10 mL of the filtrate, and use the subsequent filtrate as the sample solution. Separately, weigh accurately about 25 mg (potency) of Faropenem Sodium

Hydrate Reference Standard, add exactly 10 mL of the internal standard solution and water to make exactly 50 mL, and use this solution as the standard solution. Proceed as directed in the Assay under Faropenem Sodium Hydrate.

$$\text{Amount [mg (potency)] of faropenem (C}_{12}\text{H}_{15}\text{NO}_5\text{S)} \\ = W_S \times (Q_T/Q_S)$$


$W_S$ : Amount [mg (potency)] of Faropenem Sodium Hydrate Reference Standard

**Internal standard solution**—Dissolve 0.5 g of *m*-hydroxycetophenone in 20 mL of acetonitrile, and add water to make 200 mL.

**Containers and storage** Containers—Tight containers.

## Fenbufen

フェンブフェン



$C_{16}H_{14}O_3$ ; 254.28

4-(Biphenyl-4-yl)-4-oxobutanoic acid [36330-85-5]

Fenbufen, when dried, contains not less than 98.0% of  $C_{16}H_{14}O_3$ .

**Description** Fenbufen occurs as a white crystalline powder. It has a bitter taste.

It is sparingly soluble in acetone, slightly soluble in methanol, in ethanol (95) and in diethyl ether, and practically insoluble in water.

Melting point: about 188°C (with decomposition).

**Identification** (1) Determine the absorption spectrum of a solution of Fenbufen in ethanol (95) (1 in 200,000) as directed under Ultraviolet-visible Spectrophotometry <2.24>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wavelengths.

(2) Determine the infrared absorption spectrum of Fenbufen, previously dried, as directed in the potassium bromide disk method under Infrared Spectrophotometry <2.25>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wave numbers.

**Purity** (1) Heavy metals <1.07>—Take 2.0 g of Fenbufen, add 2 mL of sulfuric acid, and carbonize by gentle heating, proceed according to Method 2, and perform the test. Prepare the control solution with 2.0 mL of Standard Lead Solution (not more than 10 ppm).

(2) Arsenic <1.11>—Prepare the test solution with 1.0 g of Fenbufen according to Method 3, and perform the test (not more than 2 ppm).

(3) Related substances—Dissolve 0.1 g of Fenbufen in 20 mL of acetone, and use this solution as the sample solution. Pipet 1 mL of the sample solution, add acetone to make exactly 100 mL, and use this solution as the standard solution.

Perform the test with these solutions as directed under Thin-layer Chromatography <2.03>. Spot 10  $\mu\text{L}$  each of the sample solution and standard solution on a plate of silica gel with fluorescent indicator for thin-layer chromatography. Develop the plate with a mixture of dichloromethane, methanol and water (80:20:3) to a distance of about 10 cm, and air-dry the plate. Examine under ultraviolet light (main wavelength: 254 nm); the spots other than the principal spot from the sample solution are not more intense than the spot from the standard solution.

**Loss on drying** <2.41> Not more than 0.3% (1 g, 105°C, 3 hours).

**Residue on ignition** <2.44> Not more than 0.1% (1 g).


**Assay** Weigh accurately about 0.2 g of Fenbuten, previously dried, dissolve in 100 mL of ethanol (99.5), and titrate <2.50> with 0.1 mol/L potassium hydroxide-ethanol VS (potentiometric titration). Perform a blank determination, and make any necessary correction.

Each mL of 0.1 mol/L potassium hydroxide-ethanol VS  
= 25.43 mg of  $\text{C}_{16}\text{H}_{14}\text{O}_3$

**Containers and storage** Containers—Tight containers.

## Fentanyl Citrate

フェンタニルクエン酸塩



$\text{C}_{22}\text{H}_{28}\text{N}_2\text{O} \cdot \text{C}_6\text{H}_8\text{O}_7$ ; 528.59

*N*-(1-Phenethylpiperidin-4-yl)-*N*-phenylpropanamide  
monocitrate [990-73-8]

Fentanyl Citrate contains not less than 98.0% of  $\text{C}_{22}\text{H}_{28}\text{N}_2\text{O} \cdot \text{C}_6\text{H}_8\text{O}_7$ , calculated on the dried basis.

**Description** Fentanyl Citrate occurs as white crystals or crystalline powder.

It is freely soluble in methanol and in acetic acid (100), sparingly soluble in water and in ethanol (95), and very slightly soluble in diethyl ether.

**Identification** (1) Dissolve 0.05 g of Fentanyl Citrate in 10 mL of 0.1 mol/L hydrochloric acid TS and ethanol (95) to make 100 mL. Determine the absorption spectrum of the solution as directed under Ultraviolet-visible Spectrophotometry <2.24>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wavelengths.

(2) Determine the infrared absorption spectrum of Fentanyl Citrate, previously dried, as directed in the potassium bromide disk method under Infrared Spectrophotometry <2.25>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wave numbers.

(3) A solution of Fentanyl Citrate (1 in 100) responds to

the Qualitative Tests <1.09> (1) for citrate.

**pH** <2.54> Dissolve 0.10 g of Fentanyl Citrate in 10 mL of water: the pH of this solution is between 3.0 and 5.0.

**Melting point** <2.60> 150 – 154°C

**Purity** (1) Heavy metals <1.07>—Proceed with 0.5 g of Fentanyl Citrate according to Method 2, and perform the test. Prepare the control solution with 1.0 mL of Standard Lead Solution (not more than 20 ppm).

(2) Related substances—Dissolve 0.10 g of Fentanyl Citrate in 5 mL of methanol, and use this solution as the sample solution. Pipet 1 mL of the sample solution, add methanol to make exactly 100 mL, and use this solution as the standard solution. Perform the test with these solutions as directed under Thin-layer Chromatography <2.03>. Spot 5  $\mu\text{L}$  each of the sample solution and standard solution on a plate of silica gel for thin-layer chromatography. Develop the plate with a mixture of 1-butanol, water and acetic acid (100) (3:1:1) to a distance of about 10 cm, and air-dry the plate. Spray evenly Dragendorff's TS for spraying on the plate: the spots other than the principal spot from the sample solution are not more intense than the spot from the standard solution.

**Loss on drying** <2.41> Not more than 0.5% (0.2 g, in vacuum, silica gel, 60°C, 2 hours).

**Residue on ignition** <2.44> Not more than 0.2% (0.5 g).

**Assay** Weigh accurately about 0.075 g of Fentanyl Citrate, dissolve in 50 mL of acetic acid (100), and titrate <2.50> with 0.02 mol/L perchloric acid VS (potentiometric titration). Perform a blank determination, and make any necessary correction.

Each mL of 0.02 mol/L perchloric acid VS  
= 10.57 mg of  $\text{C}_{22}\text{H}_{28}\text{N}_2 \cdot \text{C}_6\text{H}_8\text{O}_7$

**Containers and storage** Containers—Tight containers.  
Storage—Light-resistant.

## Ferrous Sulfate Hydrate

硫酸鉄水和物

$\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$ ; 278.01

Ferrous Sulfate Hydrate contains not less than 98.0% and not more than 104.0% of  $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$ .

**Description** Ferrous Sulfate Hydrate occurs as pale green crystals or crystalline powder. It is odorless, and has an astringent taste.

It is freely soluble in water, and practically insoluble in ethanol (95) and in diethyl ether.

It is efflorescent in dry air, and its surface becomes yellowish brown in moist air.

**Identification** A solution of Ferrous Sulfate Hydrate (1 in 10) responds to the Qualitative Tests <1.09> for ferrous salt and for sulfate.

**Purity** (1) Clarity of solution—Dissolve 1.0 g of Ferrous Sulfate Hydrate in 20 mL of water and 1 mL of dilute sulfuric acid: the solution is clear.

(2) **Acidity**—To 5.0 g of powdered Ferrous Sulfate Hydrate add 50 mL of ethanol (95), shake well for 2 minutes, and filter the mixture. To 25 mL of the filtrate add 50 mL of water, 3 drops of bromothymol blue TS and 0.5 mL of dilute sodium hydroxide TS: a blue color develops.

(3) **Heavy metals** <1.07>—Take 1.0 g of Ferrous Sulfate Hydrate in a porcelain dish, add 3 mL of aqua regia, and dissolve. Then evaporate on a water bath to dryness. To the residue add 5 mL of 6 mol/L hydrochloric acid TS, and dissolve. Transfer this solution to a separator. Wash the porcelain dish with two 5-mL portions of 6 mol/L hydrochloric acid TS, and combine the washings and the solution in the separator. Pour two 40-mL portions and one 20-mL portion of diethyl ether in the separator, shaking each time to mix. Allow to stand, and discard each separated diethyl ether layer. To the aqueous layer add 0.05 g of hydroxylammonium chloride, dissolve, and heat on a water bath for 10 minutes. Cool, adjust the solution to a pH of 3 to 4 by dropping strong ammonia solution, add water to make 50 mL, and perform the test using this solution as the test solution. Prepare the control solution as follows: take 2.5 mL of Standard Lead Solution in a porcelain dish, add 3 mL of aqua regia, and proceed as directed for the preparation of the test solution (not more than 25 ppm).

(4) **Arsenic** <1.11>—Prepare the test solution with 1.0 g of Ferrous Sulfate Hydrate according to Method 1, and perform the test (not more than 2 ppm).


**Assay** Dissolve about 0.7 g of Ferrous Sulfate Hydrate, accurately weighed, in a mixture of 20 mL of water and 20 mL of dilute sulfuric acid, add 2 mL of phosphoric acid, and immediately titrate <2.50> with 0.02 mol/L potassium permanganate VS.

Each mL of 0.02 mol/L potassium permanganate VS  
= 27.80 mg of  $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$

**Containers and storage** Containers—Tight containers.

## Flavin Adenine Dinucleotide Sodium

フラビンアデニンジヌクレオチドナトリウム



$\text{C}_{27}\text{H}_{31}\text{N}_9\text{Na}_2\text{O}_{15}\text{P}_2$ : 829.51

Disodium adenosine 5'-[(2*R*,3*S*,4*S*)-5-(7,8-dimethyl-2,4-dioxo-3,4-dihydrobenzo[*g*]pteridin-10(2*H*)-yl)-2,3,4-trihydroxypentyl diphosphate] [84366-81-4]

Flavin Adenine Dinucleotide Sodium contains not less than 93.0% of  $\text{C}_{27}\text{H}_{31}\text{N}_9\text{Na}_2\text{O}_{15}\text{P}_2$ , calculated on

the anhydrous basis.

**Description** Flavin Adenine Dinucleotide Sodium occurs as an orange-yellow to light yellow-brown powder. It is odorless or has a slight, characteristic odor, and has a slightly bitter taste.

It is freely soluble in water, and practically insoluble, in methanol, in ethanol (95), in ethyleneglycol and in diethyl ether.

It is hygroscopic.

It is decomposed by light.

**Identification** (1) A solution of Flavin Adenine Dinucleotide Sodium (1 in 100,000) is light yellow-green in color, and shows a strong yellow-green fluorescence. To 5 mL of the solution add 0.02 g of hydrosulfite sodium: the color and the fluorescence of the solution disappear, and gradually reappear when the solution is shaken in air. Add dilute hydrochloric acid or sodium hydroxide TS dropwise: the fluorescence of the solution disappears.

(2) Determine the infrared absorption spectrum of Flavin Adenine Dinucleotide Sodium as directed in the potassium bromide disk method under Infrared Spectrophotometry <2.25>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wave numbers.

(3) To 0.1 g of Flavin Adenine Dinucleotide Sodium add 10 mL of nitric acid, evaporate on a water bath to dryness, and ignite. To the residue add 10 mL of diluted nitric acid (1 in 50), boil for 5 minutes, and after cooling, neutralize with ammonia TS, then filter the solution if necessary: the solution responds to the Qualitative Tests <1.09> for sodium salt and the Qualitative Tests <1.09> (1) and (3) for phosphate.

**Optical rotation** <2.49>  $[\alpha]_{\text{D}}^{20}$ :  $-21.0 - -25.5^\circ$  (0.3 g, calculated on the anhydrous basis, water, 20 mL, 100 mm).

**pH** <2.54> Dissolve 1.0 g of Flavin Adenine Dinucleotide Sodium in 100 mL of water: the pH of this solution is between 5.5 and 6.5.

**Purity** (1) Clarity and color of solution—Dissolve 0.20 g of Flavin Adenine Dinucleotide Sodium in 10 mL of water: the solution is clear and orange-yellow in color.

(2) Free phosphoric acid—Weigh accurately about 0.02 g of Flavin Adenine Dinucleotide Sodium, dissolve in 10 mL of water, and use this solution as the sample solution. Separately, measure exactly 2 mL of Standard Phosphoric Acid Solution, add 10 mL of water, and use this solution as the standard solution. To each of the sample solution and standard solution add 2 mL of diluted perchloric acid (100 in 117), then add 1 mL of hexaammonium heptamolybdate TS and 2 mL of 2,4-diaminophenol hydrochloride TS, respectively, shake, add water to make exactly 25 mL, and allow to stand at  $20 \pm 1^\circ\text{C}$  for 30 minutes. Perform the test with these solutions as directed under Ultraviolet-visible Spectrophotometry <2.24>, using a solution prepared in the same manner with 2 mL of water, as the blank, and determine the absorbances,  $A_T$  and  $A_S$ , of the subsequent solutions of the sample solution and standard solution at 730 nm, respectively: the amount of free phosphoric acid is less than 0.25%.

Amount (%) of free phosphoric acid ( $\text{H}_3\text{PO}_4$ )  
=  $(A_T/A_S) \times (1/W) \times 5.16$

*W*: Amount (mg) of flavin adenine dinucleotide sodium,

calculated on the anhydrous basis.

(3) Heavy metals <1.07>—Proceed with 1.0 g of Flavin Adenine Dinucleotide Sodium according to Method 2, and perform the test. Prepare the control solution with 2.0 mL of Standard Lead Solution (not more than 20 ppm).

(4) Arsenic <1.11>—Prepare the test solution with 2.0 g of Flavin Adenine Dinucleotide Sodium according to Method 3, and perform the test (not more than 1 ppm).

(5) Related substances—Dissolve 0.10 g of Flavin Adenine Dinucleotide Sodium in 200 mL of the mobile phase, and use this solution as the sample solution. Perform the test with 20  $\mu$ L of the sample solution as directed under Liquid Chromatography <2.01> according to the following conditions. Determine the peak area,  $A$ , of flavin adenine dinucleotide and the total area,  $S$ , of peaks other than the peak of flavin adenine dinucleotide by the automatic integration method:  $S/(A + S)$  is not more than 0.10.

*Operating conditions—*

Column, column temperature, mobile phase, flow rate, and time span of measurement: Proceed as directed in the operating conditions in the Procedure (ii) under Assay (1).

Detector: An ultraviolet absorption photometer (wavelength: 260 nm).

*System suitability—*

Test for required detection: To exactly 2 mL of the sample solution add the mobile phase to make exactly 20 mL, and use this solution as the solution for system suitability test. Confirm that the peak area of flavin adenine dinucleotide obtained from 20  $\mu$ L of this solution is equivalent to 8 to 12% of that of flavin adenine dinucleotide obtained from 20  $\mu$ L of the sample solution.

System performance: Proceed as directed in the system suitability in the Procedure (ii) under Assay (1).

System repeatability: When the test is repeated 6 times with 20  $\mu$ L of the solution for system suitability test under the above operating conditions, the relative standard deviation of the peak area of flavin adenine dinucleotide is not more than 1.0%.

**Water <2.48>** Take 50 mL of a mixture of methanol for Karl Fischer method and ethyleneglycol for Karl Fischer method (1:1) into a dry titration flask, and titrate with Karl Fischer TS until end point. Weigh accurately about 0.1 g of Flavin Adenine Dinucleotide Sodium, transfer quickly to the titration flask, add an excess and constant volume of Karl Fischer TS, dissolve by stirring for 10 minutes, and perform the test: the water content is not more than 10.0%.

**Assay (1) Procedure (i) Total flavin content—**Conduct this procedure without exposure to daylight, using light-resistant vessels. Weigh accurately about 0.1 g of Flavin Adenine Dinucleotide Sodium, and dissolve in water to make exactly 200 mL. Pipet 5 mL of this solution, add 5 mL of zinc chloride TS, and heat in a water bath for 30 minutes. After cooling, add water to make exactly 100 mL, and use this solution as the sample solution. Separately, weigh accurately about 50 mg of Riboflavin Reference Standard, previously dried at 105°C for 2 hours, dissolve in 200 mL of diluted acetic acid (100) (1 in 100) by warming, cool, add water to make exactly 500 mL. Pipet 10 mL of this solution, add water to make exactly 100 mL, and use this solution as the standard solution. Determine the absorbances,  $A_T$  and  $A_S$ , of the sample solution and standard solution at 450 nm as

directed under Ultraviolet-visible Spectrophotometry <2.24>, using water as the blank.

$$\begin{aligned} &\text{Total amount (mg) of flavin} \\ &= W_S \times (A_T/A_S) \times (4/5) \end{aligned}$$

$W_S$ : Amount (mg) of Riboflavin Reference Standard

(ii) Peak area ratio of flavin adenine dinucleotide—Dissolve 0.1 g of Flavin Adenine Dinucleotide Sodium in 200 mL of water, and use this solution as the sample solution. Perform the test with 5  $\mu$ L of this solution as directed under the Liquid Chromatography <2.01> according to the following conditions. Determine the peak area,  $A$  of flavin adenine dinucleotide, and the total area,  $S$ , of the peaks other than flavin adenine dinucleotide by the automatic integration method.

$$\begin{aligned} &\text{Peak area ratio of flavin adenine dinucleotide} \\ &= 1.08A/(1.08A + S) \end{aligned}$$

*Operating conditions—*

Detector: A visible spectrophotometer (wavelength: 450 nm).

Column: A stainless steel column 4 mm in inside diameter and 15 cm in length, packed with octadecylsilanized silica gel for liquid chromatography (5  $\mu$ m in particle diameter).

Column temperature: A constant temperature of about 35°C.

Mobile phase: A mixture of a solution of potassium dihydrogen phosphate (1 in 500) and methanol (4:1).

Flow rate: Adjust the flow rate so that the retention time of flavin adenine dinucleotide is about 10 minutes.

Time span of measurement: About 4.5 times as long as the retention time of flavin adenine dinucleotide.

*System suitability—*

Test for required detection: To exactly 2 mL of the sample solution add water to make exactly 20 mL, and use this solution as the solution for system suitability test. Pipet 2 mL of the solution, and add water to make exactly 20 mL. Confirm that the peak area of flavin adenine dinucleotide obtained from 5  $\mu$ L of this solution is equivalent to 8 to 12% of that of flavin adenine dinucleotide obtained from 5  $\mu$ L of the solution for system suitability test.

System performance: Dissolve 20 mg each of Flavin Adenine Dinucleotide Sodium and riboflavin sodium phosphate in 100 mL of water. When the procedure is run with 5  $\mu$ L of this solution under the above operating conditions, flavin adenine dinucleotide and riboflavin phosphate are eluted in this order with the resolution between these peaks being not less than 2.0.

System repeatability: When the test is repeated 6 times with 5  $\mu$ L of the solution for system suitability test under the above operating conditions, the relative standard deviation of the peak area of flavin adenine dinucleotide is not more than 1.0%.

(2) Calculation

$$\begin{aligned} &\text{Amount (mg) of } C_{27}H_{31}N_9Na_2O_{15}P_2 \\ &= f_T \times f_R \times 2.2040 \end{aligned}$$


$f_T$ : Total amount (mg) of flavin in Flavin Adenine Dinucleotide Sodium obtained from the procedure (i).

$f_R$ : Peak area ratio of flavin adenine dinucleotide in Flavin Adenine Dinucleotide Sodium obtained from the procedure (ii).

**Containers and storage** Containers—Tight containers.  
Storage—Light-resistant.

## Flavoxate Hydrochloride

フラボキサート塩酸塩



$C_{24}H_{25}NO_4 \cdot HCl$ : 427.92

2-(Piperidin-1-yl)ethyl 3-methyl-4-oxo-2-phenyl-4H-chromene-8-carboxylate monohydrochloride  
[3717-88-2]

Flavoxate Hydrochloride, when dried, contains not less than 99.0% of  $C_{24}H_{25}NO_4 \cdot HCl$ .

**Description** Flavoxate Hydrochloride occurs as white crystals or crystalline powder.

It is sparingly soluble in acetic acid (100) and in chloroform, slightly soluble in water and in ethanol (95), and practically insoluble in acetonitrile and in diethyl ether.

**Identification (1)** Determine the absorption spectrum of a solution of Flavoxate Hydrochloride in 0.01 mol/L hydrochloric acid TS (1 in 50,000) as directed under Ultraviolet-visible Spectrophotometry <2.24>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wavelengths.

**(2)** Determine the infrared absorption spectrum of Flavoxate Hydrochloride, previously dried, as directed in the potassium bromide disk method under Infrared Spectrophotometry <2.25>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wave numbers.

**(3)** A solution of Flavoxate Hydrochloride (1 in 100) responds to the Qualitative Tests <1.09> for chloride.

**Purity (1)** Heavy metals <1.07>—Proceed with 2.0 g of Flavoxate Hydrochloride according to Method 2, and perform the test. Prepare the control solution with 2.0 mL of Standard Lead Solution (not more than 10 ppm).

**(2)** Arsenic <1.11>—Prepare the test solution with 2.0 g of Flavoxate Hydrochloride according to Method 4, and perform the test (not more than 1 ppm).

**(3)** Related substances—Dissolve 80 mg of Flavoxate Hydrochloride in 10 mL of chloroform, and use this solution as the sample solution. Pipet 1 mL of the sample solution, add chloroform to make exactly 20 mL, then pipet 1 mL of this solution, add chloroform to make exactly 20 mL, and use this solution as the standard solution. Perform the test with these solutions as directed under Thin-layer Chromatography <2.03>. Spot 5  $\mu$ L each of the sample solution and standard solution on a plate of silica gel with fluorescent indicator for thin-layer chromatography. Develop the plate with a mixture of 1-butanol, water and acetic acid (100) (3:1:1) to a distance of about 12 cm, and air-dry the plate. Examine under ultraviolet light (main wavelength: 254 nm): the spots other than the principal spot from the sample solu-

tion are not more intense than the spot from the standard solution.

**Loss on drying** <2.41> Not more than 1.0% (1 g, reduced pressure, silica gel, 2 hours).

**Residue on ignition** <2.44> Not more than 0.1% (1 g).


**Assay** Weigh accurately about 0.6 g of Flavoxate Hydrochloride, previously dried, add 10 mL of acetic acid (100) and 40 mL of acetonitrile to dissolve, add 50 mL of acetic anhydride, and titrate <2.50> with 0.1 mol/L perchloric acid VS (potentiometric titration). Perform a blank determination, and make any necessary correction.

Each mL of 0.1 mol/L perchloric acid VS  
= 42.79 mg of  $C_{24}H_{25}NO_4 \cdot HCl$

**Containers and storage** Containers—Tight containers.

## Flomoxef Sodium

フロモキセフナトリウム



$C_{15}H_{17}F_2N_6NaO_7S_2$ : 518.45

Monosodium (6R,7R)-7-  
{[(difluoromethylsulfanyl)acetyl]amino}-  
3-[1-(2-hydroxyethyl)-1H-tetrazol-5-ylsulfanylmethyl]-  
7-methoxy-8-oxo-5-oxa-1-azabicyclo[4.2.0]oct-2-ene-  
2-carboxylate [92823-03-5]

Flomoxef Sodium contains not less than 870  $\mu$ g (potency) and not more than 985  $\mu$ g (potency) per mg, calculated on the anhydrous basis. The potency of Flomoxef Sodium is expressed as mass (potency) of flomoxef ( $C_{15}H_{18}F_2N_6O_7S_2$ : 496.47).

**Description** Flomoxef Sodium occurs as white to light yellowish white, powder or masses.

It is very soluble in water, freely soluble in methanol, and sparingly soluble in ethanol (99.5).

**Identification (1)** Decompose 0.01 g of Flomoxef Sodium as directed under Oxygen Flask Combustion Method <1.06>, using a mixture of 0.5 mL of 0.01 mol/L sodium hydroxide TS and 20 mL of water as the absorbing liquid. To 2 mL of the test solution so obtained add 1.5 mL of a mixture of alizarin complexone TS, acetic acid-potassium acetate buffer solution, pH 4.3 and cerium (III) nitrate TS (1:1:1): blue-purple color develops.

**(2)** Determine the absorption spectrum of a solution of Flomoxef Sodium (3 in 100,000) as directed under Ultraviolet-visible Spectrophotometry <2.24>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wavelengths.

**(3)** Determine the infrared absorption spectrum of Flomoxef Sodium as directed in the potassium bromide disk



method under Infrared Spectrophotometry <2.25>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wave numbers.

(4) Determine the spectrum of a solution of Flomoxef Sodium in heavy water for nuclear magnetic resonance spectroscopy (1 in 10) as directed under Nuclear Magnetic Resonance Spectroscopy <2.21> (<sup>1</sup>H), using sodium 3-trimethylsilylpropanesulfonate for nuclear magnetic resonance spectroscopy as an internal reference compound: it exhibits a single signal A at around δ3.5 ppm, a single signal or a sharp multiple signal B at around δ3.7 ppm, and a single signal C at around δ5.2 ppm. The ratio of the integrated intensity of these signals, A:B:C, is about 3:2:1.

(5) Flomoxef Sodium responds to the Qualitative Tests <1.09> (1) for sodium salt.

**Optical rotation** <2.49>  $[\alpha]_D^{20}$ :  $-8 - -13^\circ$  (1 g calculated on the anhydrous basis, a mixture of water and ethanol (99.5) (4:1), 50 mL, 100 mm).

**pH** <2.54> The pH of a solution obtained by dissolving 0.5 g of Flomoxef Sodium in 5 mL of water is between 4.0 and 5.5.

**Purity** (1) Clarity and color of solution—Dissolve 1.0 g of Flomoxef Sodium in 10 mL of water: the solution is clear and pale yellow.

(2) Heavy metals <1.07>—Proceed with 1.0 g of Flomoxef Sodium in a quartz crucible according to Method 2, and perform the test. Prepare the control solution with 2.0 mL of Standard Lead Solution (not more than 20 ppm).

(3) Arsenic <1.11>—To 1.0 g of Flomoxef Sodium 5 mL of sulfuric acid and 5 mL of nitric acid, heat carefully until the solution changes to colorless to light yellow with occasional addition of 2 mL of nitric acid. After cooling, add 10 mL of ammonium oxalate TS, heat until white fumes evolve, and concentrate to 2 to 3 mL. After cooling, add water to make 10 mL, and perform the test using this solution as the test solution: the color is not darker than that of the control solution.

Control solution: Proceed to prepare a solution in the same manner as the test solution without Flomoxef Sodium, and transfer 10 mL of the solution so obtained to the generator bottle, add exactly 2 mL of Standard Arsenic Solution, and proceed in the same manner as the test solution (not more than 2 ppm).

(4) 1-(2-Hydroxyethyl)-1*H*-tetrazol-5-thiol—Weigh accurately about 20 mg of 1-(2-hydroxyethyl)-1*H*-tetrazol-5-thiol, and dissolve in water to make exactly 100 mL. Pipet 5 mL of this solution, add exactly 25 mL of the internal standard solution and water to make 50 mL, and use this solution as the standard solution. Perform the test with 5 μL each of the sample solution obtained in the Assay and standard solution as directed under Liquid Chromatography <2.01> according to the following conditions, and determine the ratios,  $Q_T$  and  $Q_S$ , of the peak area of 1-(2-hydroxyethyl)-1*H*-tetrazol-5-thiol to that of the internal standard: the amount of 1-(2-hydroxyethyl)-1*H*-tetrazol-5-thiol is not more than 1.0% of the amount of Flomoxef Sodium calculated on the anhydrous basis.

Amount (mg) of 1-(2-hydroxyethyl)-1*H*-tetrazol-5-thiol ( $C_3H_6N_4OS$ )

$$= W_S \times (Q_T/Q_S) \times (1/10)$$

$W_S$ : Amount (mg) of 1-(2-hydroxyethyl)-1*H*-tetrazol-5-

thiol

**Internal standard solution**—A solution of *m*-cresol (3 in 1000)

**Operating conditions**—

Proceed as directed in the operating conditions in the Assay.

**System suitability**—

Proceed as directed in the system suitability in the Assay.

**Water** <2.48> Not more than 1.5% (0.5 g, volumetric titration, back titration).

**Assay** Weigh accurately an amount of Flomoxef Sodium and Flomoxef Triethylammonium Reference Standard, equivalent to about 50 mg (potency), and dissolve each in exactly 50 mL of the internal standard solution, add water to make 100 mL, and use these solutions as the sample solution and standard solution. Perform the test with 5 μL each of the sample solution and standard solution as directed under Liquid Chromatography <2.01> according to the following conditions, and determine the ratios,  $Q_T$  and  $Q_S$ , of the peak area of flomoxef to that of the internal standard.

$$\begin{aligned} &\text{Amount } [\mu\text{g (potency)}] \text{ of flomoxef } (C_{15}H_{18}F_2N_6O_7S_2) \\ &= W_S \times (Q_T/Q_S) \times 1000 \end{aligned}$$

$W_S$ : Amount [mg (potency)] of Flomoxef Triethylammonium Reference Standard

**Internal standard solution**—A solution of *m*-cresol (3 in 1000).

**Operating conditions**—

Detector: An ultraviolet absorption photometer (wavelength: 246 nm).

Column: A stainless steel column 4 mm in inside diameter and 20 cm in length, packed with octadecylsilanized silica gel for liquid chromatography (5–10 μm in particle diameter).

Column temperature: A constant temperature of about 25°C.

Mobile phase: Dissolve 6.94 g of potassium dihydrogen phosphate, 3.22 g of disodium hydrogen phosphate dodecahydrate and 1.60 g of tetra-*n*-butylammonium bromide in water to make 1000 mL. To 750 mL of this solution add 250 mL of methanol.

Flow rate: Adjust the flow rate so that the retention time of flomoxef is about 9 minutes.

**System suitability**—

System performance: When the procedure is run with 5 μL of the standard solution under the above operating conditions, flomoxef and the internal standard are eluted in this order with the resolution between these peaks being not less than 10.

System repeatability: When the test is repeated 3 times with 5 μL of the standard solution under the above operating conditions, the relative standard deviation of the ratios of the peak area of flomoxef to that of the internal standard is not more than 1.0%.

**Containers and storage** Containers—Tight containers.

Storage—At 5°C or below.

## Flomoxef Sodium for Injection

注射用フロモキシセフナトリウム

Flomoxef Sodium for Injection is a preparation for injection which is dissolved before use.

It contains not less than 90.0% and not more than 110.0% of the labeled amount of flomoxef ( $C_{15}H_{18}F_2N_6O_7S_2$ ; 496.47).

**Method of preparation** Prepare as directed under Injections, with Flomoxef Sodium.

**Description** Flomoxef Sodium for Injection occurs as white to light yellowish white, friable masses or powder.

**Identification** Proceed as directed in the Identification (3) under Flomoxef Sodium.

**pH** <2.54> The pH of a solution obtained by dissolving an amount of Flomoxef Sodium for Injection, equivalent to 0.5 g (potency) of Flomoxef Sodium according to the labeled amount, in 5 mL of water is between 4.0 and 5.5.

**Purity** (1) Clarity and color of solution—Dissolve an amount of Flomoxef Sodium for Injection, equivalent to 1.0 g (potency) of Flomoxef Sodium according to the labeled amount, in 10 mL of water: the solution is clear and colorless or pale yellow.

(2) 1-(2-Hydroxyethyl)-1*H*-tetrazol-5-thiol—Use the sample solution obtained in the Assay as the sample solution. Weigh accurately about 20 mg of 1-(2-hydroxyethyl)-1*H*-tetrazol-5-thiol, and dissolve in water to make exactly 100 mL. Pipet 5 mL of this solution, add exactly 25 mL of the internal standard solution and water to make 50 mL, and use this solution as the standard solution. Perform the test with 5  $\mu$ L each of the sample solution and standard solution as directed under Liquid Chromatography <2.01> according to the following conditions, and determine the ratios,  $Q_T$  and  $Q_S$ , of the peak area of 1-(2-hydroxyethyl)-1*H*-tetrazol-5-thiol to that of the internal standard. Calculate the amount of 1-(2-hydroxyethyl)-1*H*-tetrazol-5-thiol per 1 g (potency) of Flomoxef Sodium for Injection by the following formula: not more than 10 mg.

$$\begin{aligned} &\text{Amount (mg) of 1-(2-hydroxyethyl)-1H-tetrazol-5-thiol} \\ & \quad (C_3H_6N_4OS) \\ &= W_S \times (Q_T/Q_S) \times (1/10) \end{aligned}$$

$W_S$ : Amount (mg) of 1-(2-hydroxyethyl)-1*H*-tetrazol-5-thiol

**Operating conditions—**

Proceed as directed in the Assay.

**System suitability—**

Test for required detectability: Pipet 1 mL of the standard solution, and add water to make exactly 20 mL. Confirm that the peak area of 1-(2-hydroxyethyl)-1*H*-tetrazol-5-thiol obtained from 5  $\mu$ L of this solution is equivalent to 3.5 – 6.5% of that obtained from 5  $\mu$ L of the standard solution.

**System performance:** When the procedure is run with 5  $\mu$ L of the standard solution under the above operating conditions, 1-(2-hydroxyethyl)-1*H*-tetrazol-5-thiol and the internal standard are eluted in this order with the resolution between these peaks being not less than 20.

**System repeatability:** When the test is repeated 3 times with 5  $\mu$ L of the standard solution under the above operating conditions, the relative standard deviation of the ratio of the peak area of 1-(2-hydroxyethyl)-1*H*-tetrazol-5-thiol to that of the internal standard is not more than 1.0%.

**Water** <2.48> Not more than 1.5% (0.5 g, volumetric titration, back titration).

**Bacterial endotoxins** <4.01> Less than 0.025 EU/mg (potency).

**Uniformity of dosage units** <6.02> It meets the requirement of the Mass variation test.

**Foreign insoluble matter** <6.06> Perform the test according to Method 2: it meets the requirement.

**Insoluble particulate matter** <6.07> Perform the test according to Method 1: it meets the requirement.

**Sterility** <4.06> Perform the test according to the Membrane filtration method: it meets the requirement.

**Assay** Weigh accurately the mass of the contents of not less than 10 Flomoxef Sodium for Injection, and calculate the average mass of the content. Spread out thinly about 1 g of the content in a petri dish, allow the dish to stand in a desiccator containing a saturated solution of magnesium bromide without light exposure to equilibrate the sample to constant water content. Determine the water content, separately, with about 0.1 g of the sample according to the method described in Water. Weigh accurately an amount of the sample, equivalent to about 50 mg (potency) of Flomoxef Sodium according to the labeled amount, add exactly 50 mL of the internal standard solution to dissolve, add water to make 100 mL, and use this solution as the sample solution. Separately weigh accurately about 50 mg (potency) of Flomoxef Triethylammonium Reference Standard, add exactly 50 mL of the internal standard solution to dissolve, add water to make 100 mL, and use this solution as the standard solution. Proceed as directed in the Assay under Flomoxef Sodium.

$$\begin{aligned} &\text{Amount } [\mu\text{g (potency)}] \text{ of flomoxef } (C_{15}H_{18}F_2N_6O_7S_2) \\ &= W_S \times (Q_T/Q_S) \times 1000 \end{aligned}$$


$W_S$ : Amount [mg (potency)] of Flomoxef Triethylammonium Reference Standard

**Internal standard solution—**A solution of *m*-cresol (3 in 1000).

**Containers and storage** Containers—Hermetic containers. Polyethylene or polypropylene containers for aqueous injection may be used.

## Flopropione

フロプロピオン



$C_9H_{10}O_4$ : 182.17

1-(2,4,6-Trihydroxyphenyl)propan-1-one [2295-58-1]

Flopropione contains not less than 98.0% of  $C_9H_{10}O_4$ , calculated on the anhydrous basis.

**Description** Flopropione occurs as a white to pale yellow-brown, crystalline powder. It is odorless, and has a bitter taste.

It is very soluble in *N,N*-dimethylformamide, freely soluble in methanol, in ethanol (99.5) and in diethyl ether, and practically insoluble in water.

**Identification (1)** To 1 mL of a solution of Flopropione in ethanol (99.5) (1 in 200) add 4 mL of water and 1 mL of iron (III) nitrate TS: a red-purple color develops.

**(2)** Determine the absorption spectrum of a solution of Flopropione in ethanol (99.5) (1 in 200,000) as directed under Ultraviolet-visible Spectrophotometry <2.24>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wavelengths.

**Melting point** <2.60> 177 – 181°C

**Purity (1)** Clarity and color of solution—Dissolve 1.0 g of Flopropione in 10 mL of ethanol (99.5): the solution is clear, and has no more color than Matching Fluid H.

**(2)** Heavy metals <1.07>—Proceed with 1.0 g of Flopropione according to Method 4, and perform the test. Prepare the control solution with 2.0 mL of Standard Lead Solution (not more than 20 ppm).

**(3)** Arsenic <1.11>—Prepare the test solution with 1.0 g of Flopropione according to Method 3, and perform the test (not more than 2 ppm).

**(4)** Related substances—Dissolve 0.10 g of Flopropione in 10 mL of ethanol (99.5), and use this solution as the sample solution. Pipet 1 mL of the sample solution, add ethanol (99.5) to make exactly 200 mL, and use this solution as the standard solution. Perform the test with these solutions as directed under Thin-layer Chromatography <2.03>. Spot 10  $\mu$ L each of the sample solution and standard solution on a plate of silica gel for thin-layer chromatography. Develop the plate with a mixture of hexane, ethanol (99.5) and acetic acid (100) (40:20:1) to a distance of about 10 cm, and air-dry the plate. Spray evenly *p*-nitrobenzenediazonium TS for spraying on the plate, and dry in cold wind for about 5 minutes: the spots other than the principal spot from the sample solution are not more intense than the spot from the standard solution.

**Water** <2.48> Not more than 4.0% (0.5 g, direct titration).

**Residue on ignition** <2.44> Not more than 0.1% (1 g).

**Assay** Weigh accurately about 0.3 g of Flopropione, dis-

solve in 30 mL of *N,N*-dimethylformamide, and titrate <2.50> with 0.1 mol/L tetramethylammonium hydroxide VS (potentiometric titration). Perform a blank determination, and make any necessary correction.

Each mL of 0.1 mol/L tetramethylammonium hydroxide VS  
= 18.22 mg of  $C_9H_{10}O_4$

**Containers and storage** Containers—Tight containers.  
Storage—Light-resistant.

## Flopropione Capsules

フロプロピオンカプセル

Flopropione Capsules contain not less than 93.0% and not more than 107.0% of the labeled amount of flopropione ( $C_9H_{10}O_4$ : 182.17).

**Method of preparation** Prepare as directed under the Capsules, with Flopropione.

**Identification (1)** Powder the contents of Flopropione Capsules. To a portion of the powder, equivalent to 60 mg of Flopropione according to the labeled amount, add 40 mL of water, shake well, and filter. To 5 mL of the filtrate add 1 mL of iron (III) nitrate TS: a red-purple color appears.

**(2)** Powder the contents of Flopropione Capsules. To a portion of the powder, equivalent to 90 mg of Flopropione according to the labeled amount, add 100 mL of ethanol (99.5), shake well, and filter. To 5 mL of the filtrate add ethanol (99.5) to make 50 mL. To 5 mL of this solution add ethanol (99.5) to make 100 mL, and use this solution as the sample solution. Determine the absorption spectrum of the sample solution as directed under Ultraviolet-visible Spectrophotometry <2.24>: it exhibits a maximum between 283 nm and 287 nm.

**Uniformity of dosage units** <6.02> Perform the test according to the following method: it meets the requirement of the Content uniformity test.

To 1 capsule of Flopropione Capsules add 43 mL of a mixture of water and phosphoric acid (86:1), and disintegrate the capsule in a water bath at 50°C. After cooling, add a suitable amount of acetonitrile to make exactly *V* mL of a solution containing about 0.4 mg of flopropione ( $C_9H_{10}O_4$ ) per mL. Stir the solution for 10 minutes, centrifuge a part of the solution at 3000 rpm for 5 minutes, and use the supernatant liquid as the sample solution. Proceed as directed in the Assay.

Amount (mg) of flopropione ( $C_9H_{10}O_4$ )  
=  $W_S \times (A_T/A_S) \times (V/100)$

$W_S$ : Amount (mg) of flopropione for assay, calculated on the anhydrous basis

**Assay** Take out the contents of not less than 20 Flopropione Capsules, weigh accurately the mass of the contents, and power. Weigh accurately a part of the powder, equivalent to about 40 mg of flopropione ( $C_9H_{10}O_4$ ), and add the mobile phase to make exactly 100 mL. Stir the solution for 10 minutes, centrifuge a part of this solution for 5 minutes at 3000 rpm, and use the supernatant liquid as the sample solution. Separately, weigh accurately about 40 mg of

flopropione for assay (previously determine the water <2.48> in the same manner as Flopropione), add 70 mL of the mobile phase, and dissolve by exposure for 10 minutes to ultrasonic vibration. Add the mobile phase to make exactly 100 mL, and use this solution as the standard solution. Perform the test with exactly 5  $\mu$ L each of the sample solution and standard solution as directed under Liquid chromatography <2.01> according to the following conditions, and determine the peak areas,  $A_T$  and  $A_S$ , of flopropione.

Amount (mg) of flopropione ( $C_9H_{10}O_4$ ) =  $W_S \times (A_T/A_S)$

$W_S$ : Amount (mg) of flopropione for assay, calculated on the anhydrous basis

#### Operating conditions—

Detector: An ultraviolet absorption photometer (wavelength: 267 nm).

Column: A stainless steel column 4.6 mm in inside diameter and 15 cm in length, packed with octadecylsilanized silica gel for liquid chromatography (5  $\mu$ m in particle diameter).

Column temperature: A constant temperature of about 35°C.

Mobile phase: A mixture of acetonitrile, water and phosphoric acid (114:86:1)

Flow rate: Adjust the flow rate so that the retention time of flopropione is about 3 minutes.

#### System suitability—


System performance: Dissolve 50 mg of flopropione in 50 mL of the mobile phase. To 20 mL of the solution add 25 mL of a solution prepared by dissolving 25 mg of ethyl parahydroxybenzoate in 30 mL of acetonitrile and add water to make 50 mL, and then add the mobile phase to make 50 mL. When the procedure is run with 5  $\mu$ L of this solution under the above operating conditions, Flopropione and ethyl parahydroxybenzoate are eluted in this order with the resolution between these peaks being not less than 2.0.

System repeatability: When the test is repeated 6 times with 5  $\mu$ L of the standard solution under the above operating conditions, the relative standard deviation of the peak area of flopropione is not more than 1.0%.

**Containers and storage** Containers—Tight containers.

## Flucytosine

フルシトシン



$C_4H_4FN_3O$ : 129.09

5-Fluorocytosine [2022-85-7]

Flucytosine, when dried, contains not less than 98.5% of  $C_4H_4FN_3O$ , and not less than 14.0% and not more than 15.5% of fluorine (F: 19.00).

**Description** Flucytosine occurs as a white, crystalline powder.

It is odorless.

It is sparingly soluble in water, slightly soluble in methanol, in ethanol (95), in acetic anhydride and in acetic acid (100), and practically insoluble in diethyl ether.

It dissolves in 0.1 mol/L hydrochloric acid TS.

The pH of a solution of Flucytosine (1 in 100) is between 5.5 and 7.5.

It is slightly hygroscopic.

Melting point: about 295°C (with decomposition).

**Identification (1)** Add 0.2 mL of bromine TS to 5 mL of a solution of Flucytosine (1 in 500): a yellow-brown color of bromine TS is immediately discharged. Further add 2 mL of barium hydroxide TS: a purple precipitate is formed.

(2) Proceed with 0.1 g of Flucytosine as directed under Oxygen Flask Combustion Method <1.06>, using a mixture of 0.5 mL of 0.01 mol/L sodium hydroxide TS and 20 mL of water as the absorbing liquid. The solution responds to the Qualitative Tests <1.09> (2) for fluoride.

(3) Determine the absorption spectrum of a solution of Flucytosine in 0.1 mol/L hydrochloric acid TS (1 in 125,000) as directed under Ultraviolet-visible Spectrophotometry <2.24>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wavelengths.

**Purity (1)** Clarity and color of solution—Dissolve 1.0 g of Flucytosine in 100 mL of water: the solution is clear and colorless.

(2) Chloride <1.03>—Dissolve 1.0 g of Flucytosine in 80 mL of water by heating on a water bath. After cooling, to 40 mL of this solution add 6 mL of dilute nitric acid and water to make 50 mL. Perform the test using this solution as the test solution. Prepare the control solution with 0.20 mL of 0.01 mol/L hydrochloric acid VS (not more than 0.014%).

(3) Fluoride—Dissolve 0.10 g of Flucytosine in 10.0 mL of diluted 0.01 mol/L sodium hydroxide TS (1 in 20). Transfer 5.0 mL of this solution to a 20-mL volumetric flask, add 10 mL of a mixture of alizarin complexone TS, acetic acid-potassium acetate buffer solution, pH 4.3, and cerrous nitrate TS (1:1:1), and add water to make 20 mL. Allow the mixture to stand for 1 hour, and use this solution as the sample solution. Separately, transfer 4.0 mL of Standard Fluorine Solution to a 20-mL volumetric flask, add 5.0 mL of diluted 0.01 mol/L sodium hydroxide TS (1 in 20), add 10 mL of a mixture of alizarin complexone TS, acetic acid-potassium acetate buffer solution, pH 4.3, and cerrous nitrate TS (1:1:1). Proceed in the same manner as directed in the preparation of the sample solution, and use this solution as the standard solution. Transfer 5.0 mL of diluted 0.01 mol/L sodium hydroxide TS (1 in 20) to a 20-mL volumetric flask, proceed in the same manner as directed in the preparation of the standard solution, and use this solution as the blank solution. Determine the absorbances,  $A_T$  and  $A_S$ , of the sample solution and standard solution at 600 nm, using the blank solution as the control as directed under Ultraviolet-visible Spectrophotometry <2.24>:  $A_T$  is not larger than  $A_S$  (not more than 0.048%).

(4) Heavy metals <1.07>—Proceed with 1.0 g of Flucytosine according to Method 2, and perform the test. Prepare the control solution with 2.0 mL of Standard Lead Solution (not more than 20 ppm).

(5) Arsenic <1.11>—Prepare the test solution with 1.0 g of Flucytosine according to Method 2, and perform the test (not more than 2 ppm).

(6) Related substances—Dissolve 50 mg of Flucytosine in 5 mL of diluted methanol (1 in 2), and use this solution as the sample solution. Measure accurately 1 mL of this solution, add diluted methanol (1 in 2) to make exactly 25 mL. Measure accurately 1 mL of this solution, add diluted methanol (1 in 2) to make exactly 20 mL, and use this solution as the standard solution. Perform the test with these solutions as directed under Thin-layer Chromatography <2.03>. Spot 20  $\mu$ L each of the sample solution and standard solution on a plate of silica gel with fluorescent indicator for thin-layer chromatography. Develop the chromatogram with a mixture of ethyl acetate, methanol and water (5:3:2) to a distance of about 12 cm, air-dry the plate, and observe the spots under ultraviolet light (main wavelength: 254 nm): the spots other than the principal spot from the sample solution are not more intense than the spot from the standard solution.

**Loss on drying** <2.41> Not more than 1.0% (1 g, 105°C, 4 hours).

**Residue on ignition** <2.44> Not more than 0.1% (1 g).

**Assay** (1) Flucytosine—Weigh accurately about 0.2 g of Flucytosine, previously dried, dissolve in 40 mL of acetic acid (100), add 100 mL of acetic anhydride, and titrate <2.50> with 0.1 mol/L perchloric acid VS (potentiometric titration). Perform a blank determination.

Each mL of 0.1 mol/L perchloric acid VS  
= 12.91 mg of  $C_4H_4FN_3O$


(2) Fluorine—Weigh accurately about 0.01 g of Flucytosine, previously dried, and proceed as directed in the determination of fluorine under Oxygen Flask Combustion Method <1.06>, using a mixture of 0.5 mL of 0.01 mol/L sodium hydroxide VS and 20 mL of water as the absorbing liquid.

**Containers and storage** Containers—Tight containers.

Storage—Light-resistant.

## Fludiazepam

フルジアゼパム



$C_{16}H_{12}ClFN_2O$ : 302.73

7-Chloro-5-(2-fluorophenyl)-1-methyl-1,3-dihydro-2H-1,4-benzodiazepin-2-one [3900-31-0]

Fludiazepam, when dried, contains not less than 99.0% of  $C_{16}H_{12}ClFN_2O$ .

**Description** Fludiazepam occurs as white to light yellow crystals or crystalline powder.

It is very soluble in chloroform, freely soluble in methanol, in ethanol (95), in acetic acid (100) and in diethyl ether, and practically insoluble in water.

**Identification** (1) Prepare the test solution with 0.01 g of

Fludiazepam as directed under Oxygen Flask Combustion Method <1.06>, using a mixture of 0.5 mL of 0.01 mol/L sodium hydroxide TS and 20 mL of water as the absorbing liquid: the test solution responds to the Qualitative Tests <1.09> (2) for fluoride.

(2) Determine the absorption spectrum of a solution of Fludiazepam in methanol (1 in 200,000) as directed under Ultraviolet-visible Spectrophotometry <2.24>, and compare the spectrum with the Reference Spectrum 1: both spectra exhibit similar intensities of absorption at the same wavelengths. Separately, determine the absorption spectrum of a solution of Fludiazepam in methanol (1 in 20,000) as directed under Ultraviolet-visible Spectrophotometry <2.24>, and compare the spectrum with the Reference Spectrum 2: both spectra exhibit similar intensities of absorption at the same wavelengths.

(3) Determine the infrared absorption spectrum of Fludiazepam, previously dried, as directed in the potassium bromide disk method under Infrared Spectrophotometry <2.25>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wave numbers.

(4) Perform the test with Fludiazepam as directed under Flame Coloration Test <1.04> (2): a green color appears.

**Melting point** <2.60> 91 – 94°C

**Purity** (1) Chloride <1.03>—Dissolve 1.0 g of Fludiazepam in 50 mL of diethyl ether, add 50 mL of water, and shake. Separate the water layer, wash it with two 20-mL portions of diethyl ether, and filter the water layer. To 20 mL of the filtrate add 6 mL of dilute nitric acid and water to make 50 mL. Perform the test using this solution as the test solution. Prepare the control solution with 0.40 mL of 0.01 mol/L hydrochloric acid VS (not more than 0.036%).

(2) Heavy metals <1.07>—Proceed with 2.0 g of Fludiazepam according to Method 2, and perform the test. Prepare the control solution with 2.0 mL of Standard Lead Solution (not more than 10 ppm).

(3) Related substances—Dissolve 0.10 g of Fludiazepam in 20 mL of chloroform, and use this solution as the sample solution. Pipet 1 mL of the sample solution, and add chloroform to make exactly 50 mL. Pipet 2 mL of this solution, add chloroform to make exactly 20 mL, and use this solution as the standard solution. Perform the test with these solutions as directed under Thin-layer Chromatography <2.03>. Spot 20  $\mu$ L each of the sample solution and standard solution on a plate of silica gel with fluorescent indicator for thin-layer chromatography. Develop the plate with a mixture of chloroform and ethyl acetate (10:7) to a distance of about 12 cm, and air-dry the plate. Examine under ultraviolet light (main wavelength: 254 nm): the spots other than the principal spot from the sample solution are not more intense than the spot from the standard solution.

**Loss on drying** <2.41> Not more than 0.30% (1 g, in vacuum, 60°C, 3 hours).

**Residue on ignition** <2.44> Not more than 0.1% (1 g, platinum crucible).

**Assay** Weigh accurately about 0.5 g of Fludiazepam, previously dried, dissolve in 50 mL of acetic acid (100), and titrate <2.50> with 0.1 mol/L perchloric acid VS (potentiometric titration). Perform a blank determination, and make any


necessary correction.

Each mL of 0.1 mol/L perchloric acid VS  
= 30.27 mg of  $C_{16}H_{12}ClFN_2O$

**Containers and storage** Containers—Tight containers.

## Flunitrazepam

フルニトラゼパム



$C_{16}H_{12}FN_3O_3$ : 313.28  
5-(2-Fluorophenyl)-1-methyl-7-nitro-1,3-dihydro-2H-1,4-benzodiazepin-2-one [1622-62-4]

Flunitrazepam, when dried, contains not less than 99.0% of  $C_{16}H_{12}FN_3O_3$ .

**Description** Flunitrazepam occurs as a white to pale yellow crystalline powder.

It is freely soluble in acetic acid (100), soluble in acetic anhydride and in acetone, slightly soluble in ethanol (99.5) and in diethyl ether, and practically insoluble in water.

**Identification (1)** Determine the absorption spectrum of a solution of Flunitrazepam in ethanol (99.5) (1 in 100,000) as directed under Ultraviolet-visible Spectrophotometry <2.24>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wavelengths.

(2) Determine the infrared absorption spectrum of Flunitrazepam, previously dried, as directed in the potassium bromide disk method under Infrared Spectrophotometry <2.25>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wave numbers.

**Melting point** <2.60> 168 – 172°C

**Purity (1)** Chloride <1.03>—To 1.0 g of Flunitrazepam add 50 mL of water, allow to stand for 1 hour with occasional stirring, and filter. To 20 mL of the filtrate add 6 mL of dilute nitric acid and water to make 50 mL, and perform the test with this solution. Prepare the control solution with 0.25 mL of 0.01 mol/L hydrochloric acid VS (not more than 0.022%).

(2) Heavy metals <1.07>—Proceed with 2.0 g of Flunitrazepam according to Method 4 using a platinum crucible, and perform the test. Prepare the control solution with 2.0 mL of Standard Lead Solution (not more than 10 ppm).

(3) Related substances—Dissolve 50 mg of Flunitrazepam in 10 mL of acetone, and use this solution as the sample solution. Pipet 2 mL of the sample solution, and add acetone to make exactly 20 mL. Pipet 1 mL of this solution, add acetone to make exactly 25 mL, and use this solution as the standard solution. Perform the test with these solutions as directed under Thin-layer Chromatography <2.03>. Spot 10

$\mu$ L each of the sample solution and standard solution on a plate of silica gel with fluorescent indicator for thin-layer chromatography. Develop the plate with a mixture of 1,2-dichloroethane, diethyl ether and ammonia solution (28) (200:100:3) to a distance of about 12 cm, and air-dry the plate. Examine under ultraviolet light (main wavelength: 254 nm): number of the spots other than the principal spot from the sample solution is not more than 2, and they are not more intense than the spot from the standard solution.

**Loss on drying** <2.41> Not more than 0.5% (1 g, 105°C, 4 hours).

**Residue on ignition** <2.44> Not more than 0.1% (1 g, platinum crucible).


**Assay** Weigh accurately about 0.5 g of Flunitrazepam, previously dried, dissolve in 20 mL of acetic acid (100), add 50 mL of acetic anhydride, and titrate <2.50> with 0.1 mol/L perchloric acid VS (potentiometric titration). Perform a blank determination, and make any necessary correction.

Each mL of 0.1 mol/L perchloric acid VS  
= 31.33 mg of  $C_{16}H_{12}FN_3O_3$

**Containers and storage** Containers—Tight containers.  
Storage—Light-resistant.

## Fluocinolone Acetonide

フルオシノロンアセトニド



$C_{24}H_{30}F_2O_6$ : 452.49  
6 $\alpha$ ,9-Difluoro-11 $\beta$ ,21-dihydroxy-16 $\alpha$ ,17-(1-methylethylidenedioxy)pregna-1,4-diene-3,20-dione [67-73-2]

Fluocinolone Acetonide, when dried, contains not less than 97.0% and not more than 102.0% of  $C_{24}H_{30}F_2O_6$ .

**Description** Fluocinolone Acetonide occurs as white crystals or crystalline powder. It is odorless.

It is freely soluble in acetic acid (100) and in acetone, soluble in ethanol (95) and ethanol (99.5), sparingly soluble in methanol and in chloroform, slightly soluble in acetonitrile, very slightly soluble in diethyl ether, and practically insoluble in water.

Melting point: 266 – 274°C (with decomposition).

**Identification (1)** To 2 mg of Fluocinolone Acetonide add 2 mL of sulfuric acid: a yellow color is produced.

(2) Dissolve 0.01 g of Fluocinolone Acetonide in 1 mL of methanol, add 1 mL of Fehling's TS, and heat: a red precipitate is produced.

(3) Proceed as directed under Oxygen Flask Combustion Method <1.06> with 0.01 g of Fluocinolone Acetonide, using

a mixture of 0.5 mL of 0.01 mol/L sodium hydroxide TS and 20 mL of water as the absorbing liquid. When the process is completed, shake well, and force the combustion gas into the absorbing liquid: this liquid responds to the Qualitative Tests <1.09> for fluoride.

(4) Determine the infrared absorption spectrum of Fluocinolone Acetonide, previously dried, as directed in the potassium bromide disk method under Infrared Spectrophotometry <2.25>, and compare the spectrum with the Reference Spectrum or the spectrum of previously dried Fluocinolone Acetonide Reference Standard: both spectra exhibit similar intensities of absorption at the same wave numbers. If any difference appears between the spectra, dissolve Fluocinolone Acetonide and Fluocinolone Acetonide Reference Standard in acetone, respectively, then evaporate the acetone to dryness, and repeat the test on the residues.

**Optical rotation** <2.49>  $[\alpha]_D^{20}$ : +98 – +108° (after drying, 0.1 g, methanol, 10 mL, 100 mm).

**Purity** Related substances—Dissolve 15 mg of Fluocinolone Acetonide in 25 mL of the mobile phase, and use this solution as the sample solution. Pipet 2 mL of the sample solution, add the mobile phase to make exactly 100 mL, and use this solution as the standard solution. Perform the test with exactly 20  $\mu$ L each of the sample solution and standard solution as directed under Liquid Chromatography <2.01> according to the following conditions. Determine each peak area of each solution by the automatic integration method: the total area of the peaks other than the peak of fluocinolone acetonide from the sample solution is not larger than the peak area of fluocinolone acetonide from the standard solution.

**Operating conditions—**

Detector: An ultraviolet absorption photometer (wavelength: 254 nm).

Column: A stainless steel column 4.6 mm in inside diameter and 25 cm in length, packed with silica gel (5  $\mu$ m in particle diameter).

Column temperature: A constant temperature of about 30°C.

Mobile phase: A mixture of water-saturated chloroform, methanol and acetic acid (100) (200:3:2).

Flow rate: Adjust the flow rate so that the retention time of fluocinolone acetonide is about 12 minutes.

Time span of measurement: About twice as long as the retention time of fluocinolone acetonide beginning after the solvent peak.

**System suitability—**

Test for required detection: To exactly 5 mL of the standard solution add the mobile phase to make exactly 100 mL. Confirm that the peak area of fluocinolone acetonide obtained from 20  $\mu$ L of this solution is equivalent to 4 to 6% of that of fluocinolone acetonide obtained from 20  $\mu$ L of the standard solution.

System performance: Dissolve 15 mg each of Fluocinolone Acetonide and triamcinolone acetonide in 25 mL of the mobile phase. To 5 mL of this solution add the mobile phase to make 20 mL. When the procedure is run with 20  $\mu$ L of this solution under the above operating conditions, triamcinolone acetonide and fluocinolone acetonide are eluted in this order with the resolution between these peaks being not less than 1.9.

System repeatability: When the test is repeated 6 times with

20  $\mu$ L of the standard solution under the above operating conditions, the relative standard deviation of the peak areas of fluocinolone acetonide is not more than 1.0%.

**Loss on drying** <2.41> Not more than 1.0% (0.2 g, in vacuum, 105°C, 3 hours).

**Residue on ignition** <2.44> Not more than 0.1% (0.2 g, platinum crucible).

**Assay** Dissolve about 20 mg each of Fluocinolone Acetonide and Fluocinolone Acetonide Reference Standard, previously dried and accurately weighed, in 40 mL each of methanol, add exactly 10 mL each of the internal standard solution, then add water to make 100 mL, and use these solutions as the sample solution and standard solution. Perform the test with 20  $\mu$ L each of these solutions as directed under Liquid Chromatography <2.01> according to the following conditions, and calculate the ratios,  $Q_T$  and  $Q_S$ , of the peak area of fluocinolone acetonide to that of the internal standard, respectively.

$$\begin{aligned} &\text{Amount (mg) of fluocinolone acetonide (C}_{24}\text{H}_{30}\text{F}_2\text{O}_6) \\ &= W_S \times (Q_T/Q_S) \end{aligned}$$

$W_S$ : Amount (mg) of Fluocinolone Acetonide Reference Standard

**Internal standard solution—**A solution of ethyl parahydroxybenzoate (1 in 2500).

**Operating conditions—**

Detector: An ultraviolet absorption photometer (wavelength: 254 nm).

Column: A stainless steel column 4.6 mm in inside diameter and 25 cm in length, packed with octadecylsilanized silica gel (5  $\mu$ m in particle diameter).

Column temperature: A constant temperature of about 40°C.

Mobile phase: A mixture of water and acetonitrile (7:3).

Flow rate: Adjust the flow rate so that the retention time of fluocinolone acetonide is about 20 minutes.

**System suitability—**

System performance: Dissolve 5 mg each of isopropyl parahydroxybenzoate and propyl parahydroxybenzoate in 50 mL of acetonitrile, and add water to make 100 mL. When the procedure is run with 20  $\mu$ L of this solution under the above operating conditions, isopropyl parahydroxybenzoate and propyl parahydroxybenzoate are eluted in this order with the resolution between these peaks being not less than 1.9.


System repeatability: When the test is repeated 6 times with 20  $\mu$ L of the standard solution under the above operating conditions, the relative standard deviation of the ratios of the peak area of fluocinolone acetonide to that of the internal standard is not more than 1.0%.

**Containers and storage** Containers—Tight containers.

Storage—Light-resistant.

## Fluocinonide

フルオシノニド



$C_{26}H_{32}F_2O_7$ : 494.52

6 $\alpha$ ,9-Difluoro-11 $\beta$ ,21-dihydroxy-16 $\alpha$ ,17-(1-methylethylidenedioxy)pregna-1,4-diene-3,20-dione 21-acetate [356-12-7]

Fluocinonide, when dried, contains not less than 97.0% and not more than 103.0% of  $C_{26}H_{32}F_2O_7$ .

**Description** Fluocinonide occurs as white crystals or crystalline powder.

It is sparingly soluble in chloroform, slightly soluble in acetonitrile, in methanol, in ethanol (95) and in ethyl acetate, very slightly soluble in diethyl ether, and practically insoluble in water.

**Identification** (1) To 0.01 g of Fluocinonide add 4 mL of water and 1 mL of Fehling's TS, and heat: a red precipitate is formed.

(2) Prepare the test solution with 0.01 g of Fluocinonide as directed under Oxygen Flask Combustion Method <1.06>, using a mixture of 0.5 mL of 0.01 mol/L sodium hydroxide TS and 20 mL of water as an absorbing liquid: the test solution responds to the Qualitative Tests <1.09> for fluoride.

(3) Determine the absorption spectrum of a solution of Fluocinonide in methanol (1 in 100,000) as directed under Ultraviolet-visible Spectrophotometry <2.24>, and compare the spectrum with the Reference Spectrum or the spectrum of a solution of Fluocinonide Reference Standard prepared in the same manner as the sample solution: both spectra exhibit similar intensities of absorption at the same wavelengths.

(4) Determine the infrared absorption spectra of Fluocinonide and Fluocinonide Reference Standard, previously dried, as directed in the potassium bromide disk method under Infrared Spectrophotometry <2.25>, and compare both spectra: both the sample and the Reference Standard exhibit similar intensities of absorption at the same wave numbers. If any difference appears in the absorption spectra, dissolve the sample and the Reference Standard in ethyl acetate, respectively, evaporate the ethyl acetate, and perform the test with the residue in the same manner.

**Optical rotation** <2.49>  $[\alpha]_D^{20}$ : +81 – +89° (after drying, 0.2 g, chloroform, 20 mL, 100 mm).

**Purity** Related substances—Dissolve 10 mg of Fluocinonide in 2 mL of chloroform, and use this solution as the sample solution. Pipet 1 mL of the sample solution, add chloroform to make exactly 100, and use this solution as the standard solution. Perform the test with these solutions as

directed under Thin-layer Chromatography <2.03>. Spot 10  $\mu$ L each of the sample solution and standard solution on a plate of silica gel for thin-layer chromatography. Develop the plate with a mixture of chloroform and methanol (97:3) to a distance of about 12 cm, and air-dry the plate. Spray evenly alkaline blue tetrazolium TS on the plate: the spots other than the principal spot from the sample solution are not more intense than the spot from the standard solution.

**Loss on drying** <2.41> Not more than 1.0% (0.5 g, 105°C, 3 hours).

**Residue on ignition** <2.44> Not more than 0.1% (0.5 g, platinum crucible).

**Assay** Weigh accurately about 20 mg of Fluocinonide and Fluocinonide Reference Standard, previously dried, dissolve each in 50 mL of acetonitrile, to each add exactly 8 mL of the internal standard solution and water to make 100 mL, and use these solutions as the sample solution and standard solution, respectively. Perform the test with 20  $\mu$ L each of the sample solution and standard solution as directed under Liquid Chromatography <2.01> according to the following conditions, and calculate the ratios,  $Q_T$  and  $Q_S$ , of the peak area of fluocinonide to that of the internal standard, respectively.

$$\begin{aligned} \text{Amount (mg) of fluocinonide (C}_{26}\text{H}_{32}\text{F}_2\text{O}_7) \\ = W_S \times (Q_T/Q_S) \end{aligned}$$

$W_S$ : Amount (mg) of Fluocinonide Reference Standard

**Internal standard solution**—A solution of propyl benzoate in acetonitrile (1 in 100).

**Operating conditions**—

**Detector**: An ultraviolet absorption photometer (wavelength: 254 nm).

**Column**: A stainless steel column 4.6 mm in inside diameter and 25 cm in length, packed with octadecylsilanized silica gel for liquid chromatography (5  $\mu$ m in particle diameter).

**Column temperature**: A constant temperature of about 40°C.

**Mobile phase**: A mixture of acetonitrile and water (1:1).

**Flow rate**: Adjust the flow rate so that the retention time of fluocinonide is about 8 minutes.

**System suitability**—

**System performance**: When the procedure is run with 20  $\mu$ L of the standard solution under the above operating conditions, fluocinonide and the internal standard are eluted in this order with the resolution between these peaks being not less than 6.


**System repeatability**: When the test is repeated 6 times with 20  $\mu$ L of the standard solution under the above operating conditions, the relative standard deviation of the ratios of the peak area of fluocinonide to that of the internal standard is not more than 1.0%.

**Containers and storage** Containers—Well-closed containers.



## Fluorescein Sodium

フルオレセインナトリウム



$C_{20}H_{10}Na_2O_5$ : 376.27

Disodium 2-(6-oxido-3-oxo-3*H*-xanthen-9-yl)benzoate  
[518-47-8]

Fluorescein Sodium contains not less than 98.5% of  $C_{20}H_{10}Na_2O_5$ , calculated on the dried basis.

**Description** Fluorescein Sodium occurs as an orange powder. It is odorless, and tasteless.

It is freely soluble in water, in methanol and in ethanol (95), and practically insoluble in diethyl ether.

It is hygroscopic.

**Identification (1)** To a solution of Fluorescein Sodium (1 in 100) having a strong green fluorescence, add a large quantity of water: the fluorescence remains. Acidify the solution with hydrochloric acid: the fluorescence disappears. Then render the solution alkaline with sodium hydroxide TS: the fluorescence reappears.

**(2)** Place 1 drop of a solution of Fluorescein Sodium (1 in 2000) on a piece of filter paper: a yellow spot develops. Expose the spot, while moist, to the vapor of bromine for 1 minute and then to ammonia vapor: the yellow color of the spot changes to red.

**(3)** Char 0.5 g of Fluorescein Sodium by ignition, cool, mix the residue with 20 mL of water, and filter: the filtrate responds to the Qualitative Tests <1.09> for sodium salt.

**Purity (1)** Clarity and color of solution—Dissolve 1 g of Fluorescein Sodium in 10 mL of water: the solution is clear, and shows a red color.

**(2)** Chloride <1.03>—Dissolve 0.15 g of Fluorescein Sodium in 20 mL of water, add 6 mL of dilute nitric acid and water to make 30 mL, and filter. To 20 mL of the filtrate add 2 mL of dilute nitric acid and water to make 50 mL. Perform the test using this solution as the test solution. Prepare the control solution with 1.0 mL of 0.01 mol/L hydrochloric acid VS (not more than 0.355%).

**(3)** Sulfate <1.14>—Dissolve 0.20 g of Fluorescein Sodium in 30 mL of water, add 2.5 mL of dilute hydrochloric acid and water to make 40 mL, and filter. To 20 mL of the filtrate add water to make 50 mL. Perform the test using this solution as the test solution. Prepare the control solution with 1.0 mL of 0.005 mol/L sulfuric acid VS (not more than 0.480%).

**(4)** Zinc—Dissolve 0.10 g of Fluorescein Sodium in 10 mL of water, add 2 mL of hydrochloric acid, and filter. To the filtrate add 0.1 mL of potassium hexacyanoferrate (II) TS: no turbidity is produced immediately.

**(5)** Related substances—Dissolve 0.20 g of Fluorescein Sodium in 10 mL of methanol, and use this solution as the sample solution. Perform the test with this solution as directed under Thin-layer Chromatography <2.03>. Spot 5  $\mu$ L of

the sample solution on a plate of silica gel for thin-layer chromatography. Develop the plate with a mixture of chloroform, methanol and ammonia solution (28) (30:15:1) to a distance of about 10 cm, and air-dry the plate: any colored spot other than the principal spot does not appear.

**Loss on drying** <2.41> Not more than 10.0% (1 g, 105°C, constant mass).

**Assay** Transfer about 0.5 g of Fluorescein Sodium, accurately weighed, to a separator. Dissolve in 20 mL of water, add 5 mL of dilute hydrochloric acid, and extract the solution with four 20-mL portions of a mixture of 2-methyl-1-propanol and chloroform (1:1). Wash each extract successively with the same 10 mL of water. Evaporate the combined extracts on a water bath with the aid of a current of air. Dissolve the residue in 10 mL of ethanol (99.5), evaporate the solution on a water bath to dryness, dry the residue at 105°C for 1 hour, and weigh as fluorescein ( $C_{20}H_{12}O_5$ : 332.31).


Amount (mg) of  $C_{20}H_{10}Na_2O_5$

= amount (mg) of fluorescein ( $C_{20}H_{12}O_5$ )  $\times$  1.1323

**Containers and storage** Containers—Tight containers.

## Fluorometholone

フルオロメトロン



$C_{22}H_{29}FO_4$ : 376.46

9-Fluoro-11 $\beta$ ,17-dihydroxy-6 $\alpha$ -methylpregna-1,4-diene-3,20-dione [426-13-1]

Fluorometholone, when dried, contains not less than 97.0% and not more than 103.0% of  $C_{22}H_{29}FO_4$ .

**Description** Fluorometholone occurs as a white to light yellowish white, odorless, crystalline powder.

It is freely soluble in pyridine, slightly soluble in methanol, in ethanol (99.5) and in tetrahydrofuran, and practically insoluble in water and in diethyl ether.

**Identification (1)** Proceed with 7 mg of Fluorometholone as directed under Oxygen Flask Combustion Method <1.06>, using a mixture of 0.5 mL of 0.01 mol/L sodium hydroxide TS and 20 mL of water as an absorbing liquid: the liquid responds to the Qualitative Tests <1.09> (2) for fluoride.

**(2)** Determine the absorption spectrum of a solution of Fluorometholone in methanol (1 in 100,000) as directed under Ultraviolet-visible Spectrophotometry <2.24>, and compare the spectrum with the Reference Spectrum or the spectrum of a solution of Fluorometholone Reference Standard prepared in the same manner as the sample solution: both spectra exhibit similar intensities of absorption at the same wavelengths.

**(3)** Determine the infrared absorption spectrum of Fluorometholone, previously dried, as directed in the potassi-

um bromide disk method under Infrared Spectrophotometry <2.25>, and compare the spectrum with the Reference Spectrum or the spectrum of previously dried Fluorometholone Reference Standard: both spectra exhibit similar intensities of absorption at the same wave numbers.

**Optical rotation** <2.49>  $[\alpha]_D^{20}$ : +52 – +60° (after drying, 0.1 g, pyridine, 10 mL, 100 mm).

**Purity (1)** Heavy metals <1.07>—Proceed with 1.0 g of Fluorometholone according to Method 3, and perform the test. Prepare the control solution with 2.0 mL of Standard Lead Solution (not more than 20 ppm).

(2) Related substances—Dissolve 20 mg of Fluorometholone in 10 mL of tetrahydrofuran, and use this solution as the sample solution. Pipet 1 mL of the sample solution, add tetrahydrofuran to make exactly 100 mL, and use this solution as the standard solution. Perform the test with these solutions as directed under Thin-layer chromatography <2.03>. Spot 25  $\mu$ L each of the sample solution and standard solution on a plate of silica gel with fluorescent indicator for thin-layer chromatography. Develop the plate with a mixture of dichloromethane, acetone and methanol (45:5:1) to a distance of about 12 cm, and air-dry the plate. Examine under ultraviolet light (main wavelength: 254 nm): the spots other than the principal spot from the sample solution are not more intense than the spot from the standard solution.

**Loss on drying** <2.41> Not more than 1.0% (0.2 g, in vacuum, phosphorus (V) oxide, 60°C, 3 hours).

**Residue on ignition** <2.44> Not more than 0.2 % (0.2 g, platinum crucible).

**Assay** Weigh accurately about 0.1 g each of Fluorometholone and Fluorometholone Reference Standard, previously dried, and dissolve each in methanol to make exactly 100 mL. Pipet 5 mL each of these solutions, and add diluted methanol (7 in 10) to make exactly 50 mL. Pipet 10 mL each of these solutions, add exactly 10 mL of the internal standard solution and diluted methanol (7 in 10) to make 100 mL, and use these solutions as the sample solution and standard solution, respectively. Perform the test with 20  $\mu$ L each of the sample solution and standard solution as directed under Liquid Chromatography <2.01> according to the following conditions and determine the ratios,  $Q_T$  and  $Q_S$ , of the peak area of fluorometholone to that of the internal standard.

$$\begin{aligned} &\text{Amount (mg) of fluorometholone (C}_{22}\text{H}_{29}\text{FO}_4) \\ &= W_S \times (Q_T/Q_S) \end{aligned}$$

$W_S$ : Amount (mg) of Fluorometholone Reference Standard

**Internal standard solution**—A solution of butyl parahydroxybenzoate in methanol (1 in 10,000).

**Operating conditions**—

**Detector:** An ultraviolet absorption photometer (wavelength: 254 nm).

**Column:** A stainless steel column about 4 mm in inside diameter and 25 to 30 cm in length, packed with octadecylsilanized silica gel for liquid chromatography (5  $\mu$ m in particle diameter).

**Column temperature:** A constant temperature of about 35°C.

**Mobile phase:** Diluted methanol (7 in 10).

**Flow rate:** Adjust the flow rate so that the retention time of

fluorometholone is about 8 minutes.


**Selection of column:** Proceed with 20  $\mu$ L of the standard solution under the above operating conditions, and calculate the resolution. Use a column giving elution of fluorometholone and the internal standard in this order with the resolution between these peaks being not less than 4.

**Containers and storage** Containers—Well-closed containers.

Storage—Light-resistant.

## Fluorouracil

フルオロウラシル



$\text{C}_4\text{H}_3\text{FN}_2\text{O}_2$ : 130.08

5-Fluorouracil [51-21-8]

Fluorouracil, when dried, contains not less than 98.5% of  $\text{C}_4\text{H}_3\text{FN}_2\text{O}_2$ , and not less than 13.1% and not more than 16.1% of fluorine (F: 19.00).

**Description** Fluorouracil occurs as white crystals or crystalline powder. It is odorless.

It is freely soluble in *N,N*-dimethylformamide, sparingly soluble in water, slightly soluble in ethanol (95), and practically insoluble in diethyl ether.

Melting point: about 282°C (with decomposition).

**Identification (1)** Add 0.2 mL of bromine TS to 5 mL of a solution of Fluorouracil (1 in 500): the color of bromine TS is discharged. Further add 2 mL of barium hydroxide TS: a purple precipitate is formed.

(2) Determine the absorption spectrum of a solution of Fluorouracil in 0.1 mol/L hydrochloric acid TS (1 in 100,000) as directed under Ultraviolet-visible Spectrophotometry <2.24>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wavelengths.

(3) Proceed with 0.01 g of Fluorouracil as directed under Oxygen Flask Combustion Method <1.06>, using a mixture of 0.5 mL of 0.01 mol/L sodium hydroxide TS and 20 mL of water as the absorbing liquid. After combustion is completed, shake well to absorb the gas evolved: the solution responds to the Qualitative Tests <1.09> for fluoride.

**Purity (1)** Clarity and color of solution—Add 20 mL of water to 0.20 g of Fluorouracil, and dissolve by warming: the solution is clear and colorless.

(2) Fluoride—Dissolve 0.10 g of Fluorouracil in 10.0 mL of diluted 0.01 mol/L sodium hydroxide TS (1 in 20). Transfer 5.0 mL of this solution to a 20-mL volumetric flask, add 10 mL of a mixture of alizarin complexone TS, acetic acid-potassium acetate buffer solution, pH 4.3, and cerium (III) nitrate TS (1:1:1), and add water to make 20 mL. Allow to stand for 1 hour, and use this solution as the sample solution. Separately, transfer 1.0 mL of Standard Fluorine Solution to a 20-mL volumetric flask, add 5.0 mL of diluted 0.01 mol/L sodium hydroxide TS (1 in 20), and add 10 mL of a mixture

of alizarin complexone TS, acetic acid-potassium acetate buffer solution, pH 4.3, and cerium (III) nitrate TS (1:1:1). Proceed in the same manner as directed for the preparation of the sample solution, and use this solution as the standard solution. Perform the test as directed under Ultraviolet-visible Spectrophotometry <2.24>, using a solution, prepared with 5.0 mL of diluted 0.01 mol/L sodium hydroxide TS (1 in 20) in the same manner, as the blank: the absorbance of the sample solution at 600 nm is not larger than that of the standard solution (not more than 0.012%).

(3) Heavy metals <1.07>—Proceed with 1.0 g of Fluorouracil according to Method 2, and perform the test. Prepare the control solution with 2.0 mL of Standard Lead Solution (not more than 20 ppm).

(4) Arsenic <1.11>—To 1.0 g of Fluorouracil in a crucible add 10 mL of a solution of magnesium nitrate hexahydrate in ethanol (95) (1 in 10), ignite the ethanol to burn, and incinerate by strong heating at 750°C to 850°C. If a carbonized substance remains in this method, moisten with a small amount of nitric acid, and incinerate by strong heating. Cool, add 10 mL of dilute hydrochloric acid to the residue, dissolve it by warming on a water bath, use this solution as the test solution, and perform the test (not more than 2 ppm).

(5) Related substances—Dissolve 0.10 g of Fluorouracil in 10 mL of water, and use this solution as the sample solution. Measure exactly 1 mL of this solution, add water to make exactly 200 mL, and use this solution as the standard solution. Perform the test with these solutions as directed under Thin-layer Chromatography <2.03>. Spot 10  $\mu$ L each of the sample solution and standard solution on a plate of silica gel with fluorescent indicator for thin-layer chromatography. Develop the plate with a mixture of ethyl acetate, acetone and water (7:4:1) to a distance of about 12 cm, air-dry the plate, and examine under ultraviolet light (main wavelength: 254 nm): the spots other than the principal spot from the sample solution are not more intense than the spot from the standard solution.

**Loss on drying** <2.41> Not more than 0.5% (1 g, in vacuum, 80°C, 4 hours).

**Residue on ignition** <2.44> Not more than 0.1% (1 g).

**Assay** (1) Fluorouracil—Weigh accurately about 0.2 g of Fluorouracil, previously dried, dissolve in 20 mL of *N,N*-dimethylformamide, and titrate <2.50> with 0.1 mol/L tetramethylammonium hydroxide VS until the color of the solution changes from yellow through blue-green to blue (indicator: 3 drops of thymol blue-dimethylformamide TS). Perform a blank determination.

Each mL of 0.1 mol/L tetramethylammonium hydroxide VS


= 13.01 mg of  $C_4H_3FN_2O_2$

(2) Fluorine—Weigh accurately about 4 mg of Fluorouracil, previously dried, and proceed as directed in the determination of fluorine under Oxygen Flask Combustion Method <1.06>, using a mixture of 0.5 mL of 0.01 mol/L sodium hydroxide TS and 20 mL of water as the absorbing liquid.

**Containers and storage** Containers—Tight containers.

## Fluoxymesterone

フルオキシメステロン



$C_{20}H_{29}FO_3$ ; 336.44

9-Fluoro-11 $\beta$ ,17 $\beta$ -dihydroxy-17-methylandro-4-en-3-one  
[76-43-7]

Fluoxymesterone, when dried, contains not less than 97.0% and not more than 102.0% of  $C_{20}H_{29}FO_3$ .

**Description** Fluoxymesterone occurs as white crystals or crystalline powder. It is odorless.

It is sparingly soluble in methanol, slightly soluble in ethanol (95) and in chloroform, very slightly soluble in diethyl ether, and practically insoluble in water.

**Identification** (1) Dissolve 5 mg of Fluoxymesterone in 2 mL of sulfuric acid: a yellow color develops.

(2) Prepare the test solution with 0.01 g of Fluoxymesterone as directed under Oxygen Flask Combustion Method <1.06>, using a mixture of 0.5 mL of 0.01 mol/L sodium hydroxide TS and 20 mL of water as an absorbing liquid: the test solution responds to the Qualitative Tests <1.09> (2) for fluoride.

(3) Determine the absorption spectrum of a solution of Fluoxymesterone in ethanol (95) (1 in 100,000) as directed under Ultraviolet-visible Spectrophotometry <2.24>, and compare the spectrum with the Reference Spectrum or the spectrum of a solution of Fluoxymesterone Reference Standard prepared in the same manner as the sample solution: both spectra exhibit similar intensities of absorption at the same wavelengths.

(4) Determine the infrared absorption spectrum of Fluoxymesterone, previously dried, as directed in the potassium bromide disk method under Infrared Spectrophotometry <2.25>, and compare the spectrum with the Reference Spectrum or the spectrum of previously dried Fluoxymesterone Reference Standard: both spectra exhibit similar intensities of absorption at the same wave numbers. If any difference appears between the spectra, dissolve Fluoxymesterone and Fluoxymesterone Reference Standard in ethanol (99.5), respectively, then evaporate the ethanol to dryness, and repeat the test on the residues.

**Optical rotation** <2.49>  $[\alpha]_D^{20}$ : +104 – +112° (after drying, 0.1 g, ethanol (95), 10 mL, 100 mm).

**Purity** (1) Heavy metals <1.07>—Proceed with 0.5 g of Fluoxymesterone according to Method 2, and perform the test. Prepare the control solution with 1.5 mL of Standard Lead Solution (not more than 30 ppm).

(2) Related substances—Dissolve 0.03 g of Fluoxymesterone in 10 mL of methanol, and use this solution as the sample solution. Pipet 1 mL of the sample solution, add methanol to make exactly 100 mL, and use this solution as the standard solution. Perform the test with these solutions as directed

under Thin-layer Chromatography <2.03>. Spot 10  $\mu\text{L}$  each of the sample solution and standard solution on a plate of silica gel with fluorescent indicator for thin-layer chromatography. Develop the plate with a mixture of toluene, ethanol (95) and ethyl acetate (3:1:1) to a distance of about 12 cm, and air-dry the plate. Examine under ultraviolet light (main wavelength: 254 nm): the spots other than the principal spot from the sample solution are not more intense than the spot from the standard solution.

**Loss on drying** <2.41> Not more than 1.0% (1 g, 105°C, 3 hours).

**Residue on ignition** <2.44> Not more than 0.2% (0.5 g, platinum crucible).

**Assay** Weigh accurately about 25 mg each of Fluoxymesterone and Fluoxymesterone Reference Standard, previously dried, dissolve each in the internal standard solution to make exactly 100 mL, and use these solutions as the sample solution and standard solution, respectively. Perform the test with 10  $\mu\text{L}$  each of the sample solution and standard solution as directed under Liquid Chromatography <2.01> according to the following conditions, and calculate the ratios,  $Q_T$  and  $Q_S$ , of the peak area of fluoxymesterone to that of the internal standard, respectively.

$$\begin{aligned} &\text{Amount (mg) of } \text{C}_{20}\text{H}_{29}\text{FO}_3 \\ &= W_S \times (Q_T/Q_S) \end{aligned}$$

$W_S$ : Amount (mg) of Fluoxymesterone Reference Standard

**Internal standard solution**—A solution of methylprednisolone in a mixture of chloroform and methanol (19:1) (1 in 5000).

**Operating conditions**—

**Detector**: An ultraviolet absorption photometer (wavelength: 254 nm).

**Column**: A stainless steel column 4.6 mm in inside diameter and 30 cm in length, packed with silica gel for liquid chromatography (5  $\mu\text{m}$  in particle diameter).

**Column temperature**: A constant temperature of about 25°C.

**Mobile phase**: A mixture of *n*-butyl chloride, water-saturated *n*-butyl chloride, tetrahydrofuran, methanol and acetic acid (100) (95:95:14:7:6).

**Flow rate**: Adjust the flow rate so that the retention time of fluoxymesterone is about 9 minutes.

**System suitability**—

**System performance**: When the procedure is run with 10  $\mu\text{L}$  of the standard solution under the above operating conditions, fluoxymesterone and the internal standard are eluted in this order with the resolution between these peaks being not less than 6.


**System repeatability**: When the test is repeated 6 times with 10  $\mu\text{L}$  of the standard solution under the above operating conditions, the relative standard deviation of the ratios of the peak area of fluoxymesterone to that of the internal standard is not more than 1.5%.

**Containers and storage** Containers—Well-closed containers.

Storage—Light-resistant.

## Fluphenazine Enanthate

フルフェナジンエナント酸エステル



$\text{C}_{29}\text{H}_{38}\text{F}_3\text{N}_3\text{O}_2\text{S}$ : 549.69

2-(4-{3-[2-(Trifluoromethyl)-10*H*-phenothiazin-10-yl]propyl} piperazin-1-yl)ethyl heptanoate [2746-81-8]

Fluphenazine Enanthate, when dried, contains not less than 98.5% of  $\text{C}_{29}\text{H}_{38}\text{F}_3\text{N}_3\text{O}_2\text{S}$ .

**Description** Fluphenazine Enanthate is a light yellow to yellowish orange viscous liquid. It is generally clear, and can be opaque by producing crystals.

It is freely soluble in methanol and in diethyl ether, soluble in acetic acid (100) and in ethanol (95), and practically insoluble in water.

**Identification** (1) Prepare the test solution with 0.01 g of Fluphenazine Enanthate as directed under Oxygen Flask Combustion Method <1.06>, using a mixture of 0.5 mL of 0.01 mol/L sodium hydroxide TS and 20 mL of water as the absorbing liquid: the test solution responds to the Qualitative Tests <1.09> for fluoride.

(2) Dissolve 2 mg of Fluphenazine Enanthate in 200 mL of a solution of hydrochloric acid in methanol (17 in 2000). Determine the absorption spectrum of the solution as directed under Ultraviolet-visible Spectrophotometry <2.24>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wavelengths.

(3) Determine the infrared absorption spectrum of Fluphenazine Enanthate as directed in the liquid film method under Infrared Spectrophotometry <2.25>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wave numbers.

**Purity** (1) Heavy metals <1.07>—Proceed with 1.0 g of Fluphenazine Enanthate according to Method 2, and perform the test. Prepare the control solution with 3.0 mL of Standard Lead Solution (not more than 30 ppm).

(2) Related substances—Dissolve 0.25 g of Fluphenazine Enanthate in 10 mL of methanol, and use this solution as the sample solution. Pipet 1 mL of the sample solution, add methanol to make exactly 100 mL, and use this solution as the standard solution. Perform the test with these solutions as directed under Thin-layer Chromatography <2.03>. Spot 20  $\mu\text{L}$  each of the sample solution and standard solution on a plate of silica gel with fluorescent indicator for thin-layer chromatography. Develop the plate with a mixture of acetone, hexane and ammonia solution (28) (16:6:1) to a distance of about 15 cm, and air-dry the plate. Examine under ultraviolet light (main wavelength: 254 nm): the spots other than the principal spot from the sample solution are not more intense than the spot from the standard solution. Then spray evenly diluted sulfuric acid (1 in 2) on the plate: the spots

other than the principal spot from the sample solution are not more intense than the spot from the standard solution.

**Loss on drying** <2.41> Not more than 1.0% (1 g, in vacuum, 60°C, 3 hours).

**Residue on ignition** <2.44> Not more than 0.2% (1 g).


**Assay** Weigh accurately about 0.5 g of Flurazepam Enanthate, previously dried, dissolve in 50 mL of acetic acid (100), and titrate <2.50> with 0.1 mol/L perchloric acid VS (indicator: 2 drops of crystal violet TS). Perform a blank determination, and make any necessary correction.

Each mL of 0.1 mol/L perchloric acid VS  
= 27.49 mg of  $C_{29}H_{38}F_3N_3O_2S$

**Containers and storage** Containers—Tight containers.  
Storage—Light-resistant.

## Flurazepam

フルラゼパム



$C_{21}H_{23}ClFN_3O$ : 387.88

7-Chloro-1-[2-(diethylamino)ethyl]-5-(2-fluorophenyl)-1,3-dihydro-2H-1,4-benzodiazepin-2-one [17617-23-1]

Flurazepam, when dried, contains not less than 99.0% of  $C_{21}H_{23}ClFN_3O$ .

**Description** Flurazepam occurs as white to light yellow crystals or crystalline powder.

It is very soluble in chloroform, freely soluble in methanol, in ethanol (95), in acetic anhydride and in diethyl ether, and practically insoluble in water.

**Identification** (1) Dissolve 0.01 g of Flurazepam in 3 mL of sulfuric acid: the solution shows a greenish yellow fluorescence under ultraviolet light (main wavelength: 365 nm).

(2) Dissolve 0.01 g of Flurazepam in 3 mL of citric acid-acetic acid TS, and heat in a water bath for 4 minutes: a dark red color develops.

(3) Prepare the test solution with 0.01 g of Flurazepam as directed under Oxygen Flask Combustion Method <1.06>, using a mixture of 0.5 mL of 0.01 mol/L sodium hydroxide TS and 20 mL of water as an absorbing liquid: the test solution responds to the Qualitative Tests <1.09> (2) for fluoride.

(4) Determine the absorption spectrum of a solution of Flurazepam in methanol (1 in 100,000) as directed under Ultraviolet-visible Spectrophotometry <2.24>, and compare the spectrum with the Reference Spectrum 1: both spectra exhibit similar intensities of absorption at the same wavelengths. Separately, determine the absorption spectrum of a solution of Flurazepam in methanol (1 in 10,000) as directed under Ultraviolet-visible Spectrophotometry <2.24>,

and compare the spectrum with the Reference Spectrum 2: both spectra exhibit similar intensities of absorption at the same wavelengths.

(5) Perform the test with Flurazepam as directed under Flame Coloration Test <1.04> (2): a green color appears.

**Melting point** <2.60> 79 – 83°C

**Purity** (1) Clarity and color of solution—Dissolve 1.0 g of Flurazepam in 10 mL of ethanol (95): the solution is clear and colorless to light yellow.

(2) Chloride <1.03>—Dissolve 1.0 g of Flurazepam in 50 mL of diethyl ether, add 46 mL of water and 4 mL of sodium carbonate TS, shake, separate the water layer, wash with two 20-mL portions of diethyl ether, and filter the water layer. Neutralize 20 mL of the filtrate with dilute nitric acid, and add 6 mL of dilute nitric acid and water to make 50 mL. Perform the test using this solution as the test solution. Prepare the control solution with 0.40 mL of 0.01 mol/L hydrochloric acid VS (not more than 0.036%).

(3) Sulfate <1.14>—Neutralize 20 mL of the filtrate obtained in (2) with dilute hydrochloric acid, and add 1 mL of dilute hydrochloric acid and water to make 50 mL. Perform the test using this solution as the test solution. Prepare the control solution with 0.40 mL of 0.005 mol/L sulfuric acid VS (not more than 0.048%).

(4) Heavy metals <1.07>—Proceed with 2.0 g of Flurazepam according to Method 2, and perform the test. Prepare the control solution with 2.0 mL of Standard Lead Solution (not more than 10 ppm).

(5) Arsenic <1.11>—Prepare the test solution with 1.0 g of Flurazepam according to Method 3, and perform the test (not more than 2 ppm).

(6) Related substances—Dissolve 0.20 g of Flurazepam in 20 mL of chloroform, and use this solution as the sample solution. Pipet 1 mL of the sample solution, and add chloroform to make exactly 20 mL. Pipet 3 mL of this solution, add chloroform to make exactly 50 mL, and use this solution as the standard solution. Perform the test with these solutions as directed under Thin-layer Chromatography <2.03>. Spot 10  $\mu$ L each of the sample solution and standard solution on a plate of silica gel with fluorescent indicator for thin-layer chromatography. Develop the plate with a mixture of cyclohexane, acetone and ammonia solution (28) (60:40:1) to a distance of about 12 cm, and air-dry the plate. Examine under ultraviolet light (main wavelength: 254 nm): the spots other than the principal spot from the sample solution are not more intense than the spot from the standard solution.

**Loss on drying** <2.41> Not more than 0.20% (1 g, in vacuum, 60°C, 2 hours).

**Residue on ignition** <2.44> Not more than 0.1% (1 g, platinum crucible).

**Assay** Weigh accurately about 0.3 g of Flurazepam, previously dried, dissolve in 50 mL of acetic anhydride, and titrate <2.50> with 0.1 mol/L perchloric acid VS to the second equivalence point (potentiometric titration). Perform a blank determination, and make any necessary correction.

Each mL of 0.1 mol/L perchloric acid VS  
= 19.39 mg of  $C_{21}H_{23}ClFN_3O$

**Containers and storage** Containers—Well-closed containers.

Storage—Light-resistant.

## Flurazepam Capsules

フルラゼパムカプセル

Flurazepam Capsules contain not less than 93% and not more than 107% of the labeled amount of flurazepam ( $C_{21}H_{23}ClFN_3O$ : 387.88).

**Method of preparation** Prepare as directed under Capsules, with Flurazepam.

**Identification (1)** Powder the contents of Flurazepam Capsules. To a quantity of the powder, equivalent to 0.1 g of Flurazepam according to the labeled amount, add 100 mL of 0.1 mol/L hydrochloric acid TS, stir, and filter. To 40 mL of the filtrate add 80 mL of a solution of sodium hydroxide (1 in 250) and 100 mL of hexane, extract by shaking well, and use the hexane layer as the sample solution. Evaporate 25 mL of the sample solution on a water bath to dryness. Dissolve the residue in 3 mL of sulfuric acid: the solution shows a greenish yellow fluorescence under ultraviolet light.

(2) Evaporate 25 mL of the sample solution obtained in (1) on a water bath to dryness. Dissolve the residue in 3 mL of citric acid-acetic acid TS, and heat in a water bath for 4 minutes: a dark red color develops.

(3) Determine the absorption spectrum of the sample solution obtained in the Assay as directed under Ultraviolet-visible Spectrophotometry <2.24>: it exhibits a maximum between 315 nm and 319 nm, and a minimum between 297 nm and 301 nm.

**Assay** Weigh accurately the contents of not less than 20 Flurazepam Capsules, and powder the combined contents. Weigh accurately a portion of the powder, equivalent to about 0.05 g of flurazepam ( $C_{21}H_{23}ClFN_3O$ ), add 30 mL of methanol, stir well for 10 minutes, and add methanol to make exactly 50 mL. Filter this solution, discard the first 20 mL of the filtrate, pipet 6 mL of the subsequent filtrate, add methanol to make exactly 50 mL, and use this solution as the sample solution. Separately, weigh accurately about 0.05 g of flurazepam for assay, previously dried in vacuum at 60°C for 2 hours, and dissolve in methanol to make exactly 50 mL. Pipet 6 mL of this solution, add methanol to make exactly 50 mL, and use this solution as the standard solution. Determine the absorbances,  $A_T$  and  $A_S$ , of the sample solution and standard solution at 317 nm as directed under Ultraviolet-visible Spectrophotometry <2.24>.


$$\begin{aligned} &\text{Amount (mg) of flurazepam (C}_{21}\text{H}_{23}\text{ClFN}_3\text{O)} \\ &= W_S \times (A_T/A_S) \end{aligned}$$

$W_S$ : Amount (mg) of flurazepam for assay

**Containers and storage** Containers—Tight containers.

## Flurazepam Hydrochloride

フルラゼパム塩酸塩



$C_{21}H_{23}ClFN_3O \cdot HCl$ : 424.34

7-Chloro-1-[2-(diethylamino)ethyl]-5-(2-fluorophenyl)-1,3-dihydro-2H-1,4-benzodiazepin-2-one monohydrochloride [36105-20-1]

Flurazepam Hydrochloride, when dried, contains not less than 99.0% of  $C_{21}H_{23}ClFN_3O \cdot HCl$ .

**Description** Flurazepam Hydrochloride occurs as white to yellowish white crystals or crystalline powder.

It is freely soluble in water, in ethanol (95), in ethanol (99.5) and in acetic acid (100).

Melting point: about 197°C (with decomposition).

**Identification (1)** Determine the absorption spectrum of a solution of Flurazepam Hydrochloride in sulfuric acid-ethanol TS (1 in 100,000) as directed under Ultraviolet-visible Spectrophotometry <2.24>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wavelengths.

(2) Determine the infrared absorption spectrum of Flurazepam Hydrochloride, previously dried, as directed in the potassium chloride disk method under Infrared Spectrophotometry <2.25>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wave numbers.

(3) A solution of Flurazepam Hydrochloride (1 in 20) responds to the Qualitative Tests <1.09> for chloride.

**pH** <2.54> Dissolve 1.0 g of Flurazepam Hydrochloride in 20 mL of water: the pH of this solution is between 5.0 and 6.0.

**Purity (1)** Clarity and color of solution—Dissolve 1.0 g of Flurazepam Hydrochloride in 10 mL of water: the solution is clear and colorless to pale yellow.

(2) Sulfate <1.14>—Perform the test with 1.5 g of Flurazepam Hydrochloride. Prepare the control solution with 0.35 mL of 0.005 mol/L sulfuric acid VS (not more than 0.011%).

(3) Heavy metals <1.07>—Proceed with 1.0 g of Flurazepam Hydrochloride in a platinum crucible according to Method 2, and perform the test. Prepare the control solution with 2.0 mL of Standard Lead Solution (not more than 20 ppm).

(4) Related substances—Dissolve 0.05 g of Flurazepam Hydrochloride in 5 mL of ethanol (95), and use this solution as the sample solution. Pipet 1 mL of the sample solution, add ethanol (95) to make exactly 50 mL. Pipet 1 mL of this solution, add ethanol (95) to make exactly 10 mL, and use

this solution as the standard solution. Perform the test with these solutions as directed under Thin-layer Chromatography <2.03>. Spot 20  $\mu$ L each of the sample solution and standard solution on a plate of silica gel with fluorescent indicator for thin-layer chromatography. Place the plate in a chamber filled with ammonia vapor, allow to stand for about 15 minutes, and immediately develop the plate with a mixture of diethyl ether and diethylamine (39:1) to a distance of about 12 cm, and air-dry the plate. Examine under ultraviolet light (main wavelength: 254 nm): not more than 3 spots other than the principal spot and the spot on the starting point from the sample solution appear, and are not more intense than the spot from the standard solution.

**Loss on drying** <2.41> Not more than 0.5% (1 g, 105°C, 4 hours).

**Residue on ignition** <2.44> Not more than 0.1% (1 g).


**Assay** Weigh accurately about 0.3 g of Flurbiprofen Hydrochloride, previously dried, dissolve in 10 mL of acetic acid (100), add 40 mL of acetic anhydride, and titrate <2.50> with 0.1 mol/L perchloric acid VS (potentiometric titration). Perform a blank determination, and make any necessary correction.

Each mL of 0.1 mol/L perchloric acid VS  
= 21.22 mg of  $C_{21}H_{23}ClFN_3O \cdot HCl$

**Containers and storage** Containers—Tight containers.

## Flurbiprofen

フルルビプロフェン



$C_{15}H_{13}FO_2$ : 244.26  
(2*RS*)-2-(2-Fluorobiphenyl-4-yl)propanoic acid  
[5104-49-4]

Flurbiprofen, when dried, contains not less than 98.0% of  $C_{15}H_{13}FO_2$ .

**Description** Flurbiprofen occurs as a white, crystalline powder. It has a slightly irritating odor.

It is freely soluble in methanol, in ethanol (95), in acetone and in diethyl ether, soluble in acetonitrile, and practically insoluble in water.

A solution of Flurbiprofen in ethanol (95) (1 in 50) shows no optical rotation.

**Identification** (1) Determine the absorption spectrum of a solution of Flurbiprofen in methanol (1 in 200,000) as directed under Ultraviolet-visible Spectrophotometry <2.24>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wavelengths.

(2) Determine the infrared absorption spectrum of Flurbiprofen, previously dried, as directed in the potassium bromide disk method under Infrared Spectrophotometry

<2.25>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wave numbers.

**Melting point** <2.60> 114 – 117°C

**Purity** (1) Chloride <1.03>—Dissolve 0.6 g of Flurbiprofen in 40 mL of acetone, and add 6 mL of dilute nitric acid and water to make 50 mL. Perform the test using this solution as the test solution. Prepare the control solution as follows: to 0.25 mL of 0.01 mol/L hydrochloric acid VS add 40 mL of acetone, 6 mL of dilute nitric acid and water to make 50 mL (not more than 0.015%).

(2) Heavy metals <1.07>—Dissolve 2.0 g of Flurbiprofen in 30 mL of acetone, and add 2 mL of dilute acetic acid and water to make 50 mL. Perform the test using this solution as the test solution. Prepare the control solution as follows: to 2.0 mL of Standard Lead Solution add 30 mL of acetone, 2 mL of dilute acetic acid and water to make 50 mL (not more than 10 ppm).

(3) Related substances—Dissolve 20 mg of Flurbiprofen in 10 mL of a mixture of water and acetonitrile (11:9), and use this solution as the sample solution. Pipet 1 mL of the sample solution, and add a mixture of water and acetonitrile (11:9) to make exactly 200 mL, and use this solution as the standard solution. Perform the test with exactly 20  $\mu$ L each of the sample solution and standard solution as directed under Liquid Chromatography <2.01> according to the following conditions. Determine each peak area of both solutions by the automatic integration method: each area of the peaks other than the peak of flurbiprofen from the sample solution is not larger than the peak area of flurbiprofen from the standard solution, and the total area of these peaks is not larger than twice the peak area of flurbiprofen from the standard solution.

**Operating conditions**—

**Detector:** An ultraviolet absorption photometer (wavelength: 254 nm).

**Column:** A stainless steel column 4.6 mm in inside diameter and 15 cm in length, packed with octadecylsilanized silica gel for liquid chromatography (5  $\mu$ m in particle diameter).

**Column temperature:** A constant temperature of about 30°C.

**Mobile phase:** A mixture of water, acetonitrile and acetic acid (100) (12:7:1).

**Flow rate:** Adjust the flow rate so that the retention time of flurbiprofen is about 20 minutes.

**Time span of measurement:** About twice as long as the retention time of flurbiprofen beginning after the solvent peak.

**System suitability**—

**Test for required detection:** To exactly 5 mL of the standard solution add a mixture of water and acetonitrile (11:9) to make exactly 25 mL. Confirm that the peak area of flurbiprofen obtained from 20  $\mu$ L of this solution is equivalent to 16 to 24% of that of flurbiprofen obtained from 20  $\mu$ L of the standard solution.

**System performance:** Dissolve 0.04 g of flurbiprofen and 0.02 g of butyl parahydroxybenzoate in 100 mL of a mixture of water and acetonitrile (11:9). To 5 mL of this solution add a mixture of water and acetonitrile (11:9) to make 50 mL. When the procedure is run with 20  $\mu$ L of this solution under the above operating conditions, butyl parahydroxybenzoate

and flurbiprofen are eluted in this order with the resolution between these peaks being not less than 12.

**System repeatability:** When the test is repeated 6 times with 20  $\mu$ L of the standard solution under the above operating conditions, the relative standard deviation of the peak area of flurbiprofen is not more than 2.0%.

**Loss on drying** <2.41> Not more than 0.10% (1 g, in vacuum at a pressure not exceeding 0.67 kPa, silica gel, 4 hours).

**Residue on ignition** <2.44> Not more than 0.1% (1 g, platinum crucible).


**Assay** Weigh accurately about 0.6 g of Flurbiprofen, previously dried, dissolve in 50 mL of ethanol (95), and titrate <2.50> with 0.1 mol/L sodium hydroxide VS (indicator: 3 drops of phenolphthalein TS). Perform a blank determination, and make any necessary correction.

Each mL of 0.1 mol/L sodium hydroxide VS  
= 24.43 mg of  $C_{15}H_{13}FO_2$

**Containers and storage** Containers—Well-closed containers.

## Folic Acid

葉酸



$C_{19}H_{19}N_7O_6$ ; 441.40

*N*-{4-[(2-Amino-4-hydroxypteridin-6-ylmethyl)amino]benzoyl}-L-glutamic acid [59-30-3]

Folic Acid contains not less than 98.0% and not more than 102.0% of  $C_{19}H_{19}N_7O_6$ , calculated on the anhydrous basis.

**Description** Folic Acid occurs as a yellow to orange-yellow, crystalline powder. It is odorless.

It is practically insoluble in water, in methanol, in ethanol (95), in pyridine and in diethyl ether.

It dissolves in hydrochloric acid, in sulfuric acid, in dilute sodium hydroxide TS and in a solution of sodium carbonate decahydrate (1 in 100), and these solutions are yellow in color.

It is slowly affected by light.

**Identification** (1) Dissolve 1.5 mg of Folic Acid in dilute sodium hydroxide TS to make 100 mL. Determine the absorption spectrum of the solution as directed under Ultraviolet-visible Spectrophotometry <2.24>, and compare the spectrum with the Reference Spectrum or the spectrum of a solution of Folic Acid Reference Standard prepared in the same manner as the sample solution: both spectra exhibit similar intensities of absorption at the same wavelengths.

(2) To 10 mL of the solution obtained in (1) add 1 drop of potassium permanganate TS, and mix well until the color changes to blue, and immediately observe under ultraviolet light (main wavelength: 365 nm): a blue fluorescence is

produced.

**Purity** (1) Clarity and color of solution—Dissolve 0.10 g of Folic Acid in 10 mL of dilute sodium hydroxide TS: the solution is clear and yellow in color.

(2) Free amines—Pipet 30 mL of the sample solution obtained in the Assay, add 20 mL of dilute hydrochloric acid and water to make exactly 100 mL, and use this solution as the sample solution. Weigh accurately about 50 mg of *p*-Aminobenzoylglutamic Acid Reference Standard, previously dried in a desiccator (in vacuum, silica gel) for 4 hours, dissolve in diluted ethanol (95) (2 in 5) to make exactly 100 mL. Pipet 3 mL of this solution, add water to make exactly 1000 mL, and use this solution as the standard solution. Pipet 4 mL each of the sample solution and the standard solution, proceed as directed in the Assay, and perform the test as directed under Ultraviolet-visible Spectrophotometry <2.24>. Determine the absorbances,  $A_T$  and  $A_S$ , of subsequent solutions of the sample solution and the standard solution at 550 nm: the content of free amines is not more than 1.0%.

Content (%) of free amines =  $(A_T/A_S) \times (W'/W)$

$W$ : Amount (mg) of Folic Acid, calculated on the anhydrous basis

$W'$ : Amount (mg) of *p*-Aminobenzoylglutamic Acid Reference Standard

**Water** <2.48> Not more than 8.5% (10 mg, coulometric titration).

**Residue on ignition** <2.44> Not more than 0.5% (1 g).

**Assay** Weigh accurately about 50 mg each of Folic Acid and Folic Acid Reference Standard. To each add 50 mL of dilute sodium hydroxide TS, mix well to dissolve, add dilute sodium hydroxide TS to make exactly 100 mL, and use these solutions as the sample solution and standard solution. To 30 mL each of these solutions, accurately measured, add 20 mL of dilute hydrochloric acid and water to make exactly 100 mL. To 60 mL each of these solutions add 0.5 g of zinc powder, and allow to stand with frequent shaking for 20 minutes. Filter each mixture through a dry filter paper, and discard the first 10 mL of the filtrate. Pipet 10 mL each of the subsequent filtrate, and add water to make exactly 100 mL. To 4 mL each of solutions, accurately measured, add 1 mL of water, 1 mL of dilute hydrochloric acid and 1 mL of a solution of sodium nitrite (1 in 1000), mix well, and allow to stand for 2 minutes. To each solution add 1 mL of a solution of ammonium amidosulfate (1 in 200), mix thoroughly, and allow to stand for 2 minutes. To each of these solutions, add 1 mL of a solution of *N*-(1-naphthyl)-*N'*-diethylethylenediamine oxalate (1 in 1000), shake, allow to stand for 10 minutes, and add water to make exactly 20 mL. Separately, to 30 mL of the sample solution, accurately measured, add 20 mL of dilute hydrochloric acid and water to make exactly 100 mL. Pipet 10 mL of this solution, add 18 mL of dilute hydrochloric acid and water to make exactly 100 mL. Pipet 4 mL of this solution, and prepare the blank solution in the same manner as the sample solution. Perform the test with these solutions as directed under Ultraviolet-visible Spectrophotometry <2.24>, using a solution prepared with 4 mL of water in the same manner as a blank. Determine the absorbances,  $A_T$ ,  $A_S$  and  $A_C$ , of the subsequent solution of the sample solution, the standard solution and the blank solution at 550 nm.

Amount (mg) of  $C_{19}H_{19}N_7O_6$



$$= W_S \times \{(A_T - A_C)/A_S\}$$

$W_S$ : Amount (mg) of Folic Acid Reference Standard, calculated on the anhydrous basis

**Containers and storage** Containers—Tight containers.

Storage—Light-resistant.

## Folic Acid Injection

葉酸注射液

Folic Acid Injection is an aqueous solution for injection.

It contains not less than 95% and not more than 115% of the labeled amount of folic acid ( $C_{19}H_{19}N_7O_6$ : 441.40).

**Method of preparation** Dissolve Folic Acid in water with the aid of Sodium Hydroxide or Sodium Carbonate, and prepare as directed under Injections.

**Description** Folic Acid Injection is a yellow to orange-yellow, clear liquid.

pH: 8.0 – 11.0

**Identification (1)** To a volume of Folic Acid Injection, equivalent to 1.5 mg of Folic Acid according to the labeled amount, add dilute sodium hydroxide TS to make 100 mL. Proceed as directed in the Identification (2) under Folic Acid, using this solution as the sample solution.

(2) Determine the absorption spectrum of the sample solution obtained in (1) as directed under Ultraviolet-visible Spectrophotometry <2.24>: it exhibits maxima between 255 nm and 257 nm, between 281 nm and 285 nm and between 361 nm and 369 nm. Separately, determine the maximal absorbances of the sample solution,  $A_1$  and  $A_2$ , between 255 nm and 257 nm and between 361 nm and 369 nm, respectively: the ratio of  $A_1/A_2$  is between 2.80 and 3.00.

(3) Folic Acid Injection responds to the Qualitative Tests <1.09> (1) for sodium salt.

**Extractable volume** <6.05> It meets the requirement.

**Assay** To an exactly measured volume of Folic Acid Injection, equivalent to about 50 mg of folic acid ( $C_{19}H_{19}N_7O_6$ ) add dilute sodium hydroxide TS to make exactly 100 mL, and use this solution as the sample solution. Separately, weigh accurately about 50 mg of Folic Acid Reference Standard, dissolve in dilute sodium hydroxide TS to make exactly 100 mL, and use this solution as the standard solution. Proceed with 30 mL each of the sample solution and standard solution, exactly measured, as directed in the Assay under Folic Acid.

$$\begin{aligned} &\text{Amount (mg) of folic acid } (C_{19}H_{19}N_7O_6) \\ &= W_S \times \{(A_T - A_C)/A_S\} \end{aligned}$$

$W_S$ : Amount (mg) of Folic Acid Reference Standard, calculated on the anhydrous basis

**Containers and storage** Containers—Hermetic containers, and colored containers may be used.

Storage—Light-resistant.

## Folic Acid Tablets

葉酸錠

Folic Acid Tablets contain not less than 90% and not more than 115% of the labeled amount of folic acid ( $C_{19}H_{19}N_7O_6$ : 441.40).

**Method of preparation** Prepare as directed under Tablets, with Folic Acid.

**Identification (1)** Take a quantity of powdered Folic Acid Tablets, equivalent to 1.5 mg of Folic Acid according to the labeled amount, add 100 mL of dilute sodium hydroxide TS, shake, and filter. Discard the first 10 mL of the filtrate, use the subsequent filtrate as the sample solution, and proceed as directed in the Identification (2) under Folic Acid.

(2) Determine the absorption spectrum of the filtrate obtained in (1) as directed under Ultraviolet-visible Spectrophotometry <2.24>: it exhibits maxima between 255 nm and 257 nm, between 281 nm and 285 nm and between 361 nm and 369 nm. Separately, determine the maximal absorbances of the filtrate,  $A_1$  and  $A_2$ , between 255 nm and 257 nm and between 361 nm and 369 nm, respectively: the ratio of  $A_1/A_2$  is between 2.80 and 3.00.

**Assay** Weigh accurately and powder not less than 20 Folic Acid Tablets. Weigh accurately a portion of the powder, equivalent to about 50 mg of folic acid ( $C_{19}H_{19}N_7O_6$ ). Add 50 mL of dilute sodium hydroxide TS, shake frequently, then filter into a 100-mL volumetric flask, and wash with dilute sodium hydroxide TS. To the combined filtrate and washings add dilute sodium hydroxide TS to make exactly 100 mL, and use this solution as the sample solution. Separately, weigh accurately about 50 mg of Folic Acid Reference Standard, dissolve in dilute sodium hydroxide TS to make exactly 100 mL, and use this solution as the standard solution. Take 30 mL each of the sample solution and standard solution, exactly measured, and proceed as directed in the Assay under Folic Acid.

$$\begin{aligned} &\text{Amount (mg) of folic acid } (C_{19}H_{19}N_7O_6) \\ &= W_S \times \{(A_T - A_C)/A_S\} \end{aligned}$$

$W_S$ : Amount (mg) of Folic Acid Reference Standard, calculated on the anhydrous basis

**Containers and storage** Containers—Well-closed containers.

Storage—Light-resistant.

## Formalin

ホルマリン

Formalin contains not less than 35.0% and not more than 38.0% of formaldehyde ( $CH_2O$ : 30.03.)

It contains 5% to 13% of methanol to prevent polymerization.

**Description** Formalin is a clear, colorless liquid. Its vapor is

irritating to the mucous membrane.

It is miscible with water and with ethanol (95).

When stored for a long time, especially in a cold place, it may become cloudy.

**Identification (1)** Dilute 2 mL of Formalin with 10 mL of water in a test tube, and add 1 mL of silver nitrate-ammonia TS: a gray precipitate is produced, or a silver mirror is formed on the wall of the test tube.

**(2)** To 5 mL of sulfuric acid in which 0.1 g of salicylic acid has been dissolved add 2 drops of Formalin, and warm the solution: a persistent, dark red color develops.

**Purity** Acidity—Dilute 20 mL of Formalin with 20 mL of water, and add 5.0 mL of 0.1 mol/L sodium hydroxide VS and 2 drops of bromothymol blue TS: a blue color develops.

**Residue on ignition** <2.44> Not more than 0.06 w/v% (5 mL, after evaporation).

**Assay** Weigh accurately a weighing bottle containing 5 mL of water, add about 1 g of Formalin, and weigh accurately again. Add water to make exactly 100 mL. Pipet 10 mL of this solution, add exactly 50 mL of 0.05 mol/L iodine VS and 20 mL of potassium hydroxide TS, and allow to stand for 15 minutes at an ordinary temperature. To this mixture add 15 mL of dilute sulfuric acid, and titrate <2.50> the excess iodine with 0.1 mol/L sodium thiosulfate VS (indicator: 1 mL of starch TS). Perform a blank determination.

Each mL of 0.05 mol/L iodine VS = 1.501 mg of CH<sub>2</sub>O

**Containers and storage** Containers—Tight containers.

Storage—Light-resistant.

## Formalin Water

ホルマリン水

Formalin Water contains not less than 0.9 w/v% and not more than 1.1 w/v% of formaldehyde (CH<sub>2</sub>O: 30.03).

### Method of preparation

|                         |                       |
|-------------------------|-----------------------|
| Formalin                | 30 mL                 |
| Water or Purified Water | a sufficient quantity |
| To make 1000 mL         |                       |

Prepare by mixing the above ingredients.

**Description** Formalin Water is a clear, colorless liquid. It has a slight odor of formaldehyde.

It is almost neutral.


**Assay** Transfer 20 mL of Formalin Water, measured exactly, to a 100-mL volumetric flask containing 2.5 mL of 1 mol/L sodium hydroxide VS, and add water to make 100 mL. Pipet 10 mL of this solution, and proceed as directed in the Assay under Formalin.

Each mL of 0.05 mol/L iodine VS = 1.501 mg of CH<sub>2</sub>O

**Containers and storage** Containers—Tight containers.

## Formoterol Fumarate Hydrate

ホルモテロールフマル酸塩水和物



(C<sub>19</sub>H<sub>24</sub>N<sub>2</sub>O<sub>4</sub>)<sub>2</sub>·C<sub>4</sub>H<sub>4</sub>O<sub>4</sub>·2H<sub>2</sub>O: 840.91

*N*-(2-Hydroxy-5-[(1*RS*)-1-hydroxy-

2-[(1*RS*)-2-(4-methoxyphenyl)-

1-methylethylamino]ethyl] phenyl)formamide

hemifumarate monohydrate [43229-80-7, anhydride]

Formoterol Fumarate Hydrate contains not less than 98.5% of formoterol fumarate [(C<sub>19</sub>H<sub>24</sub>N<sub>2</sub>O<sub>4</sub>)<sub>2</sub>·C<sub>4</sub>H<sub>4</sub>O<sub>4</sub>: 804.88], calculated on the anhydrous basis.

**Description** Formoterol Fumarate Hydrate occurs as a white to yellowish white, crystalline powder.

It is freely soluble in acetic acid (100), soluble in methanol, very slightly soluble in water and in ethanol (95), and practically insoluble in diethyl ether.

A solution of Formoterol Fumarate in methanol (1 in 100) shows no optical rotation.

Melting point: about 138°C (with decomposition).

**Identification (1)** Dissolve 0.5 g of Formoterol Fumarate Hydrate in 20 mL of 0.5 mol/L sulfuric acid TS, and extract with three 25-mL portions of diethyl ether. Wash the combined diethyl ether extracts with 10 mL of 0.5 mol/L sulfuric acid TS, and evaporate the ether layer under reduced pressure, and dry the residue at 105°C for 3 hours: the residue melts <2.60> at about 290°C (with decomposition, in a sealed tube).

**(2)** Determine the absorption spectrum of a solution of Formoterol Fumarate Hydrate in methanol (1 in 40,000) as directed under Ultraviolet-visible Spectrophotometry <2.24>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wavelengths.

**(3)** Determine the infrared absorption spectrum of Formoterol Fumarate Hydrate as directed in the potassium bromide disk method under Infrared Spectrophotometry <2.25>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wave numbers.

**Purity (1)** Heavy metals <1.07>—Proceed with 1.0 g of Formoterol Fumarate Hydrate according to Method 2, and perform the test. Prepare the control solution with 2.0 mL of Standard Lead Solution (not more than 20 ppm).

**(2)** Related Substances—Dissolve 0.20 g of Formoterol Fumarate Hydrate in 10 mL of methanol, and use this solution as the sample solution. Pipet 1 mL of the sample solution, add methanol to make exactly 200 mL, and use this solution as the standard solution. Perform the test with these solutions as directed under Thin-layer Chromatography <2.03>. Spot 5 μL each of the sample solution and standard

solution on a plate of silica gel for thin-layer chromatography. Develop the plate with a mixture of chloroform, 1,4-dioxane, ethanol (99.5) and ammonia solution (28) (20:20:10:3) to a distance of about 12 cm, and air-dry the plate. Allow the plate to stand for 5 minutes in iodine vapor: the spots other than the principal spot from the sample solution are not more intense than the spot from the standard solution.

**Water** <2.48> 4.0 – 5.0% (0.5 g, direct titration).

**Residue on ignition** <2.44> Not more than 0.1% (1 g).

**Assay** Weigh accurately about 0.7 g of Formoterol Fumarate Hydrate, dissolve in 50 mL of acetic acid (100), and titrate <2.50> with 0.1 mol/L perchloric acid VS (potentiometric titration). Perform a blank determination, and make any necessary correction.


Each mL of 0.1 mol/L perchloric acid VS  
= 40.24 mg of  $(C_{19}H_{24}N_2O_4)_2 \cdot C_4H_4O_4$

**Containers and storage** Containers—Tight containers.

## Fosfestrol

### Diethylstilbestrol Diphosphate

ホスフェストロール



$C_{18}H_{22}O_8P_2$ : 428.31

(*E*)-4,4'-(Hex-3-ene-3,4-diyl)bis(phenyl dihydrogen phosphate) [522-40-7]

Fosfestrol, when dried, contains not less than 98.5% of  $C_{18}H_{22}O_8P_2$ .

**Description** Fosfestrol occurs as a white, crystalline powder. It is odorless.

It is freely soluble in ethanol (95), soluble in formamide, slightly soluble in water, and practically insoluble in acetonitrile and in diethyl ether.

It dissolves in sodium hydroxide TS.

Melting point: about 234°C (with decomposition).

**Identification** (1) Dissolve 15 mg of Fosfestrol in 1 mL of sulfuric acid: a yellow to orange color develops. To this solution add 10 mL of water: the color of the solution disappears.

(2) Determine the infrared absorption spectrum of Fosfestrol as directed in the potassium bromide disk method under Infrared Spectrophotometry <2.25>, and compare the spectrum with the Reference Spectrum or with the spectrum of Fosfestrol Reference Standard: both spectra exhibit similar intensities of absorption at the same wave numbers.

(3) Place 0.4 g of Fosfestrol in a crucible, wet by adding 0.1 mL of sulfuric acid, and heat to carbonize. Add 10 mL of water to the residue, stir well, and filter. Add 0.1 mL of nitric acid to the filtrate, and heat in a water bath for 15 minutes: this solution responds to the Qualitative Tests <1.09> for phosphate.

**pH** <2.54> Dissolve 0.10 g of Fosfestrol in 30 mL of water: the pH of this solution is between 1.0 and 2.5.

**Purity** (1) Clarity and color of solution—Dissolve 1.0 g of Fosfestrol in 15 mL of sodium hydroxide TS: the solution is clear and colorless.

(2) Chloride <1.03>—Dissolve 0.10 g of Fosfestrol in 30 mL of ethanol (95), add 6 mL of dilute nitric acid and water to make 50 mL. Perform the test using this solution as the test solution. Prepare the control solution as follows: to 0.70 mL of 0.01 mol/L hydrochloric acid VS add 6 mL of dilute nitric acid, 30 mL of ethanol (95) and water to make 50 mL (not more than 0.248%).

(3) Heavy metals <1.07>—Proceed with 1.0 g of Fosfestrol according to Method 4, and perform the test. Use a solution of magnesium nitrate hexahydrate in ethanol (95) (1 in 5). Prepare the control solution with 2.0 mL of Standard Lead Solution (not more than 20 ppm).

(4) Arsenic <1.11>—Prepare the test solution with 1.0 g of Fosfestrol according to Method 3, and perform the test. Use a solution of magnesium nitrate hexahydrate in ethanol (95) (1 in 5) (not more than 2 ppm).

(5) Free phosphoric acid—Weigh accurately about 0.4 g of Fosfestrol, dissolve in a mixture of water and formamide (1:1) to make exactly 200 mL, and use this solution as the sample solution. Separately, weigh accurately 0.112 g of monobasic potassium phosphate, previously dried in a desiccator (silica gel) to constant mass, dissolve in 10 mL of diluted sulfuric acid (1 in 10) and water to make exactly 1000 mL. Measure exactly 10 mL of this solution, add 100 mL of formamide and water to make exactly 200 mL, and use this solution as the standard solution. Measure exactly 10 mL each of the sample solution and standard solution, and place in a 25-mL volumetric flask, respectively. To each of these solutions add 2.5 mL of hexaammonium heptamolybdate-sulfuric acid TS and 1 mL of 1-amino-2-naphthol-4-sulfonic acid TS, shake, add water to make 25 mL, and allow to stand at  $20 \pm 1^\circ\text{C}$  for 30 minutes. Perform the test with these solutions, using a solution obtained in the same manner with 10 mL of a mixture of water and formamide (1:1) as the blank, as directed under Ultraviolet-visible Spectrophotometry <2.24>. Determine the absorbances,  $A_T$  and  $A_S$ , of the solutions obtained from the sample solution and standard solution at 740 nm: the amount of free phosphoric acid is not more than 0.2%.

Amount (%) of free phosphoric acid ( $H_3PO_4$ )  
=  $(A_T/A_S) \times (1/W) \times 80.65$

$W$ : Amount (mg) of Fosfestrol

(6) Related substances—Dissolve 20 mg of Fosfestrol in 100 mL of the mobile phase, and use this solution as the sample solution. Measure exactly 5 mL of this solution, and add the mobile phase to make exactly 50 mL. Pipet 3 mL of this solution, add the mobile phase to make exactly 20 mL, and use this solution as the standard solution. Perform the test with exactly 10  $\mu\text{L}$  each of the sample solution and standard solution as directed under Liquid Chromatography <2.01> according to the following conditions. Determine each peak area of these solutions by the automatic integration method: the total area of the peaks other than the peak of fosfestrol from the sample solution is not larger than the peak area of fosfestrol from the standard solution.

**Operating conditions—**

**Detector:** An ultraviolet absorption photometer (wavelength: 240 nm).

**Column:** A stainless steel column about 4 mm in inside diameter and about 15 cm in length, packed with octadecylsilanized silica gel for liquid chromatography (5  $\mu$ m in particle diameter).

**Column temperature:** A constant temperature of about 25°C.

**Mobile phase:** A mixture of a solution of potassium dihydrogen phosphate (1 in 500), acetonitrile and tetrabutylammonium hydroxide TS (70:30:1).

**Flow rate:** Adjust the flow rate so that the retention time of fosfestrol is about 8 minutes.

**Selection of column:** Dissolve 0.02 g of Fosfestrol and 8 mg of methyl parahydroxybenzoate in 100 mL of the mobile phase. Proceed with 10  $\mu$ L of this solution under the above operating conditions, and calculate the resolution. Use a column giving elution of methyl parahydroxybenzoate and fosfestrol in this order with the resolution between these peaks being not less than 3.

**Detection sensitivity:** Adjust the detection sensitivity so that the peak height of fosfestrol obtained from 10  $\mu$ L of the standard solution is between 5 mm and 15 mm.

**Time span of measurement:** Three times as long as the retention time of fosfestrol.

**Loss on drying** <2.41> Not more than 1.0% (1 g, 105°C, 4 hours).

**Assay** Weigh accurately about 0.2 g of Fosfestrol, previously dried, dissolve in 60 mL of water, and titrate <2.50> with 0.1 mol/L sodium hydroxide VS (potentiometric titration). The end point is the second equivalent point. Perform a blank determination, and make any necessary correction.

$$\begin{aligned} \text{Each mL of 0.1 mol/L sodium hydroxide VS} \\ = 10.71 \text{ mg of } C_{18}H_{22}O_8P_2 \end{aligned}$$

**Containers and storage** Containers—Tight containers.

## Fosfestrol Tablets

### Diethylstilbestrol Diphosphate Tablets

ホスフェストロール錠

Fosfestrol Tablets contain not less than 93% and not more than 107% of the labeled amount of fosfestrol ( $C_{18}H_{22}O_8P_2$ ; 428.31).

**Method of preparation** Prepare as directed under Tablets, with Fosfestrol.

**Identification (1)** To a quantity of powdered Fosfestrol Tablets, equivalent to 0.5 g of Fosfestrol according to the labeled amount, add 50 mL of 0.1 mol/L hydrochloric acid TS, shake well, and filter. To the filtrate add 100 mL of diethyl ether, extract, and evaporate carefully the diethyl ether extract on a water bath to dryness. Proceed with 0.015 g of the residue as directed in the Identification (1) under Fosfestrol.

**(2)** Dry 0.01 g of the residue obtained in (1) at 105°C for 4 hours, and determine the infrared absorption spectrum as

directed in the potassium bromide disk method under Infrared Spectrometry <2.25>: it exhibits absorption at the wave numbers of about 2970  $\text{cm}^{-1}$ , 1605  $\text{cm}^{-1}$ , 1505  $\text{cm}^{-1}$ , 1207  $\text{cm}^{-1}$  and 1006  $\text{cm}^{-1}$ .

**Dissolution** <6.10> Perform the test according to the following method: it meets the requirement.

Perform the test with 1 tablet of Fosfestrol Tablets at 50 revolutions per minute according to the Paddle method, using 900 mL of water. Take 20 mL or more of the dissolved solution 20 minutes after starting the test, and filter through a membrane filter with pore size of not more than 0.8  $\mu$ m. Discard the first 10 mL of the filtrate, pipet 2 mL of the subsequent, add a solution of sodium hydroxide (1 in 250) to make exactly 20 mL, and use this solution as the sample solution. Separately, weigh accurately about 50 mg of Fosfestrol Reference Standard, previously dried at 105°C for 4 hours, and dissolve in a solution of sodium hydroxide (1 in 250) to make exactly 100 mL. Pipet 2 mL of this solution, add a solution of sodium hydroxide (1 in 250) to make exactly 100 mL, and use this solution as the standard solution. Determine the absorbances,  $A_T$  and  $A_S$ , of the sample solution and standard solution at 242 nm as directed under Ultraviolet-visible Spectrophotometry <2.24>.

The dissolution rate of Fosfestrol Tablets in 20 minutes is not less than 80%.

Dissolution rate (%) with respect to the labeled amount of fosfestrol ( $C_{18}H_{22}O_8P_2$ )

$$= W_S \times (A_T/A_S) \times (1/C) \times 180$$

$W_S$ : Amount (mg) of Fosfestrol Reference Standard

$C$ : Labeled amount (mg) of fosfestrol ( $C_{18}H_{22}O_8P_2$ ) in 1 tablet

**Assay** Weigh accurately not less than 20 Fosfestrol Tablets, and powder. Weigh accurately a quantity of the powder, equivalent to about 1 g of fosfestrol ( $C_{18}H_{22}O_8P_2$ ) according to the labeled amount, add 100 mL of a solution of sodium hydroxide (1 in 125), shake well, add water to make exactly 500 mL. Filter this solution, discard the first 30 mL of the filtrate, pipet the subsequent 2 mL of the filtrate, add 30 mL of a solution of sodium hydroxide (1 in 125) and water to make exactly 250 mL, and use this solution as the sample solution. Separately, weigh accurately about 80 mg of Fosfestrol Reference Standard, previously dried at 105°C for 4 hours, and dissolve in a solution of sodium hydroxide (1 in 125) to make exactly 50 mL. Pipet 1 mL of this solution, add 10 mL of a solution of sodium hydroxide (1 in 125) and water to make exactly 100 mL, and use this solution as the standard solution. Determine the absorbances,  $A_T$  and  $A_S$ , of the sample solution and standard solution at 242 nm as directed under Ultraviolet-visible Spectrophotometry <2.24>.


$$\begin{aligned} \text{Amount (mg) of fosfestrol } (C_{18}H_{22}O_8P_2) \\ = W_S \times (A_T/A_S) \times (25/2) \end{aligned}$$

$W_S$ : Amount (mg) of Fosfestrol Reference Standard

**Containers and storage** Containers—Tight containers.

## Fosfomycin Calcium Hydrate

ホスホマイシンカルシウム水和物



$C_3H_5CaO_4P \cdot H_2O$ : 194.14

Monocalcium (2*R*,3*S*)-3-methyloxiran-2-ylphosphonate monohydrate [26016-98-8]

Fosfomycin Calcium Hydrate is the calcium salt of a substance having antibacterial activity produced by the growth of *Streptomyces fradiae* or by the chemical synthesis.

It contains not less than 725  $\mu$ g (potency) and not more than 805  $\mu$ g (potency) per mg, calculated on the anhydrous basis. The potency of Fosfomycin Calcium Hydrate is expressed as mass (potency) of fosfomycin ( $C_3H_7O_4P$ : 138.06).

**Description** Fosfomycin Calcium Hydrate occurs as a white crystalline powder.

It is slightly soluble in water, and practically insoluble in methanol and in ethanol (99.5).

**Identification (1)** Determine the infrared absorption spectrum of Fosfomycin Calcium Hydrate as directed in the potassium bromide disk method under Infrared Spectrophotometry <2.25>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wave numbers.

(2) Determine the spectrum of a solution of Fosfomycin Calcium Hydrate in heavy water for nuclear magnetic resonance spectroscopy (1 in 300), using sodium 3-trimethylsilylpropanesulfonate for nuclear magnetic resonance spectroscopy as an internal reference compound, as directed under Nuclear Magnetic Resonance Spectroscopy <2.21> ( $^1H$ ): it exhibits a double signal at around  $\delta$  1.5 ppm, a double signal at around  $\delta$  2.9 ppm, a multiple signal at around  $\delta$  3.3 ppm, and no signal at around  $\delta$  1.4 ppm.

(3) A solution of Fosfomycin Calcium Hydrate (1 in 500) responds to the Qualitative Tests <1.09> (3) for calcium salt.

**Optical rotation** <2.49>  $[\alpha]_D^{20}$ :  $-2.5$  –  $-5.4^\circ$  (0.5 g calculated on the anhydrous bases, 0.4 mol/L disodium dihydrogen ethylenediamine tetraacetate TS, pH 8.5, 10 mL, 100 mm).

**Phosphorus Content** Weigh accurately about 0.1 g of Fosfomycin Calcium Hydrate, add 40 mL of sodium periodate (107 in 10,000) and 2 mL of perchloric acid, and heat in a water bath for 1 hour. After cooling, add water to make exactly 200 mL. Pipet 10 mL of this solution, and add 1 mL of potassium iodide TS. To this solution add sodium thiosulfate TS until the solution is colorless, add water to make exactly 100 mL, and use this solution as the sample stock solution. Separately, weigh accurately about 70 mg of potassium dihydrogenphosphate, proceed with this solution in the same manner as directed for the preparation of the sample stock solution, and use the solution so obtained as the standard stock solution. Proceed and prepare a solution in the same manner for the preparation of the sample stock solution without using Fosfomycin Calcium, and use the solution so

obtained as the blank stock solution. Pipet 5 mL each of the sample stock solution, the standard stock solution, and the blank stock solution, add 2.5 mL of ammonium molybdate-sulfuric acid TS and 1 mL of 1-amino-2-naphthol-4-sulfonic acid TS, mix, and add water to make exactly 25 mL, and use these solutions as the sample solution, the standard solution, and the blank solution, respectively. After allowing these solutions to stand for 30 minutes at  $20 \pm 1^\circ C$ , perform the test with these solutions as directed under Ultraviolet-visible Spectrophotometry <2.24>, using water as a blank, and determine the absorbances at 740 nm,  $A_T$ ,  $A_S$  and  $A_B$ , of the sample solution, the standard solution and the blank solution: the content of phosphorus is 15.2 – 16.7%.

Amount (mg) of phosphorus (P)

$$= W_S \times \{(A_T - A_B)/(A_S - A_B)\} \times 0.22760$$

$W_S$ : Amount (mg) of potassium dihydrogenphosphate

**Calcium Content** Weigh accurately about 0.2 g of Fosfomycin Calcium Hydrate, add 4 mL of 1 mol/L Hydrochloric acid TS, and shake well until the sample is completely dissolved. To this solution add 100 mL of water, 9 mL of sodium hydroxide TS and 0.1 g of methylthymol blue-sodium chloride indicator, and titrate <2.50> with 0.05 mol/L disodium dihydrogen ethylenediamine tetraacetate VS until the color of the solution changes from clear blue to gray or gray-purple: calcium content is 19.6 – 21.7%. Perform a blank determination in the same manner, and make any necessary correction.

Each mL of 0.05 mol/L disodium dihydrogen ethylenediamine tetraacetate VS = 2.004 mg of Ca

**Purity (1)** Heavy metals <1.07>—To 1.0 g of Fosfomycin Calcium Hydrate add 40 mL of 0.25 mol/L acetic acid TS and water to make 50 mL. Proceed with this solution according to Method 1, and perform the test. Prepare the control solution with 2.0 mL of Standard Lead Solution (not more than 20 ppm).

(2) Arsenic <1.11>—Prepare the test solution with 1.0 g of Fosfomycin Calcium Hydrate according to Method 3, and perform the test (not more than 2 ppm).

**Water** <2.48> Not more than 12.0% (0.1 g, volumetric titration, direct titration. Use a mixture of formamide for water determination and methanol for water determination (2:1) instead of methanol for water determination).

**Assay** Perform the test according to the Cylinder-plate method as directed under Microbial Assay for Antibiotics <4.02> according to the following conditions.

(i) Test organism—*Proteus* sp. (MB838)

(ii) Culture medium—Dissolve 5.0 g of peptone, 3.0 g of meat extract, 2.0 g of yeast extract, and 15 g of agar in 1000 mL of water, sterilize, and use as the agar media for base layer and seed layer with the pH of between 6.5 and 6.6 after sterilization.

(iii) Seeded agar layer—Incubate the test organism on the slant of the agar medium for transferring test organisms at  $37^\circ C$  for 40 – 48 hours. Subcultures at least three times. Inoculate the grown organisms onto the surface of 300 mL of the agar medium for transferring test organisms in a Roux bottle, incubate at  $37^\circ C$  for 40 – 48 hours, and suspend the grown organisms in about 30 mL of water. To the suspension add water, and use this as the stock suspension of test organ-

ism. The amount of the water to be added is adjust so that the percent transmission at 560 nm of the suspension diluted ten times with water is 17%. Keep the stock suspension at 10°C or below and use within 7 days. Add 1.0–2.0 mL of the stock suspension of test organism to 100 mL of the agar medium for seed layer previously kept at 48°C, mix thoroughly, and use this as the deeded agar layer.


(iv) Standard solutions—Weigh accurately an amount of Fosfomycin Phenethylammonium Reference Standard equivalent to about 20 mg (potency), dissolve in 0.05 mol/L Tris buffer solution, pH 7.0 to make exactly 50 mL, and use this solution as the standard stock solution. Keep the standard stock solution at 5°C or below and use within 7 days. Take exactly a suitable amount of the standard stock solution before use, add 0.05 mol/L Tris buffer solution, pH 7.0 to make solutions so that each mL contains 10 µg (potency) and 5 µg (potency), and use these solutions as the high concentration standard solution and low concentration standard solution, respectively.

(v) Sample solutions—Weigh accurately an amount of Fosfomycin Calcium Hydrate equivalent to about 20 mg (potency), and dissolve in 0.05 mol/L Tris buffer solution, pH 7.0 to make exactly 50 mL. To exactly a suitable amount of this solution add 0.05 mol/L Tris buffer solution, pH 7.0 to make solutions so that each mL contains 10 µg (potency) and 5 µg (potency), and use these solutions as the high concentration sample solution and low concentration sample solution, respectively.

**Containers and storage** Containers—Tight containers.

## Fosfomycin Sodium

ホスホマイシンナトリウム



$C_3H_5Na_2O_4P$ : 182.02

Disodium (2*R*,3*S*)-3-methyloxiran-2-ylphosphonate  
[26016-99-9]

Fosfomycin Sodium is the sodium salt of a substance having antibacterial activity produced by the growth of *Streptomyces fradiae* or by the chemical synthesis.

It contains not less than 725 µg (potency) and not more than 770 µg (potency) per mg, calculated on the anhydrous basis. The potency of Fosfomycin Sodium is expressed as mass (potency) of fosfomycin ( $C_3H_7O_4P$ : 138.06).

**Description** Fosfomycin Sodium occurs as a white crystalline powder.

It is very soluble in water, sparingly soluble in methanol, and practically insoluble in ethanol (99.5).

**Identification** (1) Determine the infrared absorption spectrum of Fosfomycin Sodium as directed in the potassium bromide disk method under Infrared Spectrophotometry <2.25>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at

the same wave numbers.

(2) Determine the spectrum of a solution of Fosfomycin Sodium in heavy water for nuclear magnetic resonance spectroscopy (1 in 300), using sodium 3-trimethylsilylpropanesulfonate for nuclear magnetic resonance spectroscopy as an internal reference compound, as directed under Nuclear Magnetic Resonance Spectroscopy <2.21> ( $^1H$ ): it exhibits a double signal at around  $\delta$  1.5 ppm, a double signal at around  $\delta$  2.8 ppm, a multiple signal at around  $\delta$  3.3 ppm, and no signal at around  $\delta$  1.3 ppm.

(3) A solution of Fosfomycin Sodium (1 in 500) responds to the Quantitative Tests <1.09> (1) for sodium salt.

**Optical rotation** <2.49>  $[\alpha]_D^{20}$ :  $-3.5 - -5.5^\circ$  (0.5 g calculated on the anhydrous bases, water, 10 mL, 100 mm).

**pH** <2.54> Dissolve 0.70 g of Fosfomycin Sodium in 10 mL of water: the pH of the solution is between 8.5 and 10.5.

**Phosphorus Content** Weigh accurately about 0.1 g of Fosfomycin Sodium, add 40 mL of a solution of sodium periodate (107 in 10,000) and 2 mL of perchloric acid, and heat in a water bath for 1 hour. After cooling, add water to make exactly 200 mL. Pipet 10 mL of this solution, and add 1 mL of potassium iodide TS. To this solution add sodium thiosulfate TS until the solution is colorless, add water to make exactly 100 mL, and use this solution as the sample stock solution. Separately, weigh accurately about 70 mg of potassium dihydrogenphosphate, proceed with this solution in the same manner as directed for the preparation of the sample stock solution, and use the solution so obtained as the standard stock solution. Proceed and prepare a solution in the same manner for the preparation of the sample stock solution without using Fosfomycin Sodium, and use the solution so obtained as the blank stock solution. Pipet 5 mL each of the sample stock solution, the standard stock solution, and the blank stock solution, add 2.5 mL of ammonium molybdate-sulfuric acid TS and 1 mL of 1-amino-2-naphthol-4-sulfonic acid TS, mix, and add water to make exactly 25 mL, and use these solutions as the sample solution, the standard solution, and the blank solution, respectively. After allowing these solutions to stand for 30 minutes at  $20 \pm 1^\circ C$ , perform the test with these solutions as directed under Ultraviolet-visible Spectrophotometry <2.24>, using water as a blank, and determine the absorbances at 740 nm,  $A_T$ ,  $A_S$  and  $A_B$ , of the sample solution, the standard solution and the blank solution: the content of phosphorus is 16.2–17.9%.

$$\begin{aligned} &\text{Amount (mg) of phosphorus (P)} \\ &= W \times \{(A_T - A_B)/(A_S - A_B)\} \times 0.22760 \end{aligned}$$

$W$ : Amount (mg) of potassium dihydrogenphosphate

**Purity** (1) Clarity and color of solution—Dissolve 1.0 g of Fosfomycin Sodium in 10 mL of water: the solution is clear and colorless.

(2) Heavy metals <1.07>—Proceed with 1.0 g of Fosfomycin Sodium according to Method 1, and perform the test. Prepare the control solution with 2.0 mL Standard Lead Solution (not more than 20 ppm).

(3) Arsenic <1.11>—Prepare the test solution with 1.0 g of Fosfomycin Sodium according to Method 3, and perform the test (not more than 2 ppm).

**Water** <2.48> Not more than 3.0% (0.2 g, volumetric titra-

tion, direct titration).

**Assay** Perform the test according to the Cylinder-plate method as directed under Microbial Assay for Antibiotics <4.02> according to the following conditions.

- (i) Test organism—*Proteus* sp. (MB838)
- (ii) Culture medium—Mix 5.0 g of peptone, 3.0 g of meat extract, 2.0 g of yeast extract, and 15 g of agar in 1000 mL of water, sterilize, and use as the agar media for base layer and seed layer with the pH of between 6.5 and 6.6 after sterilization.
- (iii) Seeded agar layer—Incubate the test organism on the slant of the agar medium for transferring test organisms at 37 °C for 40 – 48 hours. Subcultures at least three times. Inoculate the grown organisms onto the surface of 300 mL of the agar medium for transferring test organisms in a Roux bottle, incubate at 37 °C for 40 – 48 hours, and suspend the grown organisms in about 30 mL of water. To the suspension add water, and use this as the stock suspension of test organism. The amount of the water to be added is adjust so that the percent transmission at 560 nm of the suspension diluted ten times with water is 17%. Keep the stock suspension at 10 °C or below and use within 7 days. Add 1.0 – 2.0 mL of the stock suspension of test organism to 100 mL of the agar medium for seed layer previously kept at 48 °C, mix thoroughly, and use this as the deeded agar layer.

(iv) Standard solutions—Weigh accurately an amount of Fosfomycin Phenethylammonium Reference Standard equivalent to about 20 mg (potency), dissolve in 0.05 mol/L Tris buffer solution, pH 7.0 to make exactly 50 mL, and use this solution as the standard stock solution. Keep the standard stock solution at 5 °C or below and use within 7 days. Take exactly a suitable amount of the standard stock solution before use, add 0.05 mol/L Tris buffer solution, pH 7.0 to make solutions so that each mL contains 10 µg (potency) and 5 µg (potency), and use these solutions as the high concentration standard solution and low concentration standard solution, respectively.

(v) Sample solutions—Weigh accurately an amount of Fosfomycin Sodium equivalent to about 20 mg (potency), and dissolve in 0.05 mol/L Tris buffer solution, pH 7.0 to make exactly 50 mL. To exactly a suitable amount of this solution add 0.05 mol/L Tris buffer solution, pH 7.0 to make solutions so that each mL contains 10 µg (potency) and 5 µg (potency), and use these solutions as the high concentration sample solution and low concentration sample solution, respectively.

**Containers and storage** Containers—Hermetic containers.

## Fosfomycin Sodium for Injection

注射用ホスホマイシシナトリウム

Fosfomycin Sodium for Injection is a preparation for injection which is dissolved before use. It contains not less than 90.0% and not more than 110.0% of the labeled amount of fosfomycin ( $C_3H_7O_4P$ : 138.06).

**Method of preparation** Prepare as directed under Injections, with Fosfomycin Sodium.

**Description** Fosfomycin Sodium for Injection occurs as a

white crystalline powder.

**Identification (1)** Dissolve about 0.1 g of Fosfomycin Sodium for Injection in 3 mL of a solution of perchloric acid (1 in 4), add 1 mL of 0.1 mol/L sodium periodate solution, and heat in a water bath at 60 °C for 30 minutes. After cooling, add 50 mL of water, neutralize with saturated sodium hydrogen carbonate solution, and add 1 mL of potassium iodide TS; the solution does not reveal a red color, while the blank solution reveals a red color.

**(2)** To 2 mL of a solution of Fosfomycin Sodium for Injection (1 in 250) add 1 mL of perchloric acid and 2 mL of 0.1 mol/L sodium periodate solution, and heat in a water bath for 10 minutes. After cooling, add 1 mL of ammonium molybdate-sulfuric acid TS and 1 mL of 1-amino-2-naphthol-4-sulfonic acid TS, and allow to stand for 30 minutes: a blue color develops.

**(3)** Dissolve an amount of Fosfomycin Sodium for Injection, equivalent to 0.1 g (potency) of Fosfomycin Sodium, in 50 mL of water. Perform the test with this solution as directed in the Identification (3) under Fosfomycin Sodium.

**pH <2.54>** The pH of a solution prepared by dissolving an amount of Fosfomycin Sodium for Injection, equivalent to 1.0 g (potency) of Fosfomycin Sodium according to the labeled amount, in 20 mL of water is between 6.5 and 8.5.

**Purity** Clarity and color of solution—Dissolve an amount of Fosfomycin Sodium for Injection, equivalent to 1.0 g (potency) of Fosfomycin Sodium according to the labeled amount, in 10 mL of water: the solution is clear and colorless.

**Water <2.48>** Not more than 4.0% (25 mg, volumetric titration, direct titration).

**Bacterial endotoxins <4.01>** Less than 0.025 EU/mg (potency).

**Uniformity of dosage unit <6.02>** It meets the requirement of the Mass variation test.

**Foreign insoluble matter <6.06>** Perform the test according to the Method 2: it meets the requirement.

**Insoluble particulate matter <6.07>** Perform the test according to the Method 1: it meets the requirement.

**Sterility <4.06>** Perform the test according to the Membrane filtration method: it meets the requirement.

**Assay** Perform the test according to the Cylinder-plate method as directed under Microbial Assay for Antibiotics <4.02> according to the following conditions.

(i) Test organism, culture medium, seeded agar layer, and standard solutions—Proceed as directed in the Assay under Fosfomycin Sodium.


(ii) Sample solutions—Weigh accurately the mass of the contents of not less than 10 Fosfomycin Sodium for Injection. Weigh accurately an amount of the content, equivalent to about 20 mg (potency) of Fosfomycin Sodium according to the labeled amount, and dissolve in 0.05 mol/L tris buffer solution, pH 7.0 to make exactly 50 mL. Take exactly a suitable amount of this solution, add 0.05 mol/L tris buffer solution, pH 7.0 to make solutions so that each mL contains 10 µg (potency) and 5 µg (potency), and use these solutions as the high concentration sample solution and low concentration sample solution, respectively.

**Containers and storage** Containers—Hermetic containers. Polyethylene or polypropylene containers for aqueous injections may be used.

## Fradiomycin Sulfate

### Neomycin Sulfate

フラジオマイシン硫酸塩



$C_{23}H_{46}N_6O_{13} \cdot 3H_2SO_4$ : 908.88

Fradiomycin Sulfate B

2,6-Diamino-2,6-dideoxy- $\alpha$ -D-glucopyranosyl-(1 $\rightarrow$ 4)-[2,6-diamino-2,6-dideoxy- $\beta$ -L-idopyranosyl-(1 $\rightarrow$ 3)- $\beta$ -D-ribofuranosyl-(1 $\rightarrow$ 5)]-2-deoxy-D-streptamine trisulfate [119-04-0, Neomycin B]

Fradiomycin Sulfate C

2,6-Diamino-2,6-dideoxy- $\alpha$ -D-glucopyranosyl-(1 $\rightarrow$ 4)-[2,6-diamino-2,6-dideoxy- $\alpha$ -D-glucopyranosyl-(1 $\rightarrow$ 3)- $\beta$ -D-ribofuranosyl-(1 $\rightarrow$ 5)]-2-deoxy-D-streptamine trisulfate [66-86-4, Neomycin C] [1405-10-3, Neomycin Sulfate]

Fradiomycin Sulfate is the sulfate of a mixture of aminoglycoside substances having antibacterial activity produced by the growth of *Streptomyces fradiae*.

It, when dried, contains not less than 623  $\mu$ g (potency) and not more than 740  $\mu$ g (potency) per mg. The potency of Fradiomycin Sulfate is expressed as mass (potency) of fradiomycin ( $C_{23}H_{46}N_6O_{13}$ : 614.64).

**Description** Fradiomycin Sulfate occurs as a white to light yellow powder.

It is freely soluble in water, and practically insoluble in ethanol (95).

It is hygroscopic.

**Identification** (1) Dissolve 50 mg each of Fradiomycin Sulfate and Fradiomycin Sulfate Reference Standard in 1 mL of water, and use these solutions as the sample solution and the standard solution. Perform the test with these solutions as directed under Thin-layer Chromatography <2.03>. Spot 2  $\mu$ L each of the sample solution and standard solution on a plate of silica gel for thin-layer chromatography. Develop the

plate with a mixture of methanol, ammonia solution (28) and dichloromethane (3:2:1) to a distance of about 10 cm, and air-dry the plate. Spray evenly a solution of ninhydrin in acetone (1 in 50) on the plate, and heat at 110°C for 15 minutes: the  $R_f$  values of the principal spots from the sample solution and the standard solution are not different each other.

(2) A solution of Fradiomycin Sulfate (1 in 20) responds to the Qualitative Tests <1.09> (1) for sulfate.

**Optical rotation** <2.49>  $[\alpha]_D^{20}$ : +53.5 – +59.0° (1 g calculated on the dried basis, water, 10 mL, 100 mm).

**pH** <2.54> The pH of a solution obtained by dissolving 1.0 g of Fradiomycin Sulfate in 10 mL of water is between 5.0 and 7.5.

**Purity** (1) Heavy metals <1.07>—Proceed with 1.0 g of Fradiomycin Sulfate according to Method 2, and perform the test. Prepare the control solution with 2.0 mL of Standard Lead Solution (not more than 20 ppm).

(2) Arsenic <1.11>—Prepare the test solution with 1.0 g of Fradiomycin Sulfate according to Method 1, and perform the test (not more than 2 ppm).

(3) Related substances—Dissolve 0.63 g of Fradiomycin Sulfate in 5 mL of water, and use this solution as the sample solution. Pipet 1 mL of the sample solution, add water to make exactly 50 mL, and use this solution as the standard solution. Perform the test with these solutions as directed under Thin-layer Chromatography <2.03>. Spot 1  $\mu$ L each of the sample solution and standard solution on a plate of silica gel for thin-layer chromatography. Develop the plate with a mixture of methanol, ammonia solution (28) and dichloromethane (3:2:1) to a distance of about 10 cm, and air-dry the plate. Spray evenly a solution of ninhydrin in acetone (1 in 50) on the plate, and heat at 110°C for 15 minutes: the spot at around  $R_f$  0.4 from the sample solution is not more intense than the spot from the standard solution.

**Loss on drying** <2.41> Not more than 8.0% (0.2 g, in vacuum, 60°C, 3 hours).

**Residue on ignition** <2.44> Not more than 0.3% (1 g).

**Assay** Perform the test according to the Cylinder-plate method as directed under Microbial Assay for Antibiotics <4.02> according to the following conditions.

- (i) Test organism—*Staphylococcus aureus* ATCC 6538 P
- (ii) Agar medium for seed and base layer

|                 |         |
|-----------------|---------|
| Glucose         | 1.0 g   |
| Peptone         | 6.0 g   |
| Meat extract    | 1.5 g   |
| Yeast extract   | 3.0 g   |
| Sodium chloride | 2.5 g   |
| Agar            | 15.0 g  |
| Water           | 1000 mL |

Mix all the ingredients and sterilize. Adjust the pH after sterilization to 7.8 – 8.0 with sodium hydroxide TS.

(iii) Standard solutions—Weigh accurately an amount of Fradiomycin Sulfate Reference Standard, previously dried, equivalent to about 50 mg (potency), dissolve in 0.1 mol/L phosphate buffer solution for antibiotics, pH 8.0 to make exactly 50 mL, and use this solution as the standard stock solution. Keep the standard stock solution at 5°C or below and use within 14 days. Take exactly a suitable amount of the standard stock solution before use, add 0.1 mol/L phosphate buffer solution for antibiotics, pH 8.0 to make solutions so



that each mL contains 80  $\mu$ g (potency) and 20  $\mu$ g (potency), and use these solutions as the high concentration standard solution and low concentration standard solution, respectively.


(iv) Sample solutions—Weigh accurately an amount of Fradiomycin Sulfate, previously dried, equivalent to about 50 mg (potency), dissolve in 0.1 mol/L phosphate buffer solution for antibiotics, pH 8.0 to make exactly 50 mL. Take exactly a suitable amount of this solution, add 0.1 mol/L phosphate buffer solution for antibiotics, pH 8.0 to make solutions so that each mL contains 80  $\mu$ g (potency) and 20  $\mu$ g (potency), and use these solutions as the high concentration sample solution and low concentration sample solution, respectively.

**Containers and storage** Containers—Tight containers.

Storage—Light-resistant.

## Fructose

果糖



$C_6H_{12}O_6$ : 180.16

$\beta$ -D-Fructopyranose [57-48-7]

Fructose, when dried, contains not less than 98.0% of  $C_6H_{12}O_6$ .

**Description** Fructose occurs as colorless to white crystals or crystalline powder. It is odorless and has a sweet taste.

It is very soluble in water, sparingly soluble in ethanol (95) and practically insoluble in diethyl ether.

It is hygroscopic.

**Identification** (1) Add 2 to 3 drops of a solution of Fructose (1 in 20) to 5 mL of boiling Fehling's TS: a red precipitate is produced.

(2) Determine the infrared absorption spectrum of Fructose as directed in the paste method under Infrared Spectrophotometry <2.25>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wave numbers.

**pH** <2.54> Dissolve 4.0 g of Fructose in 20 mL of water: the pH of the solution is between 4.0 and 6.5.

**Purity** (1) Clarity and color of solution—Dissolve 25.0 g of Fructose in 50 mL of water: the solution is clear and has no more color than the following control solution.

Control solution: To a mixture of 1.0 mL of Cobaltous Chloride Stock CS, 3.0 mL of Ferric Chloride Stock CS and 2.0 mL of Cupric Sulfate Stock CS, and add water to make 10.0 mL. To 3.0 mL of the solution add water to make 50 mL.

(2) Acidity—Dissolve 5.0 g of Fructose in 50 mL of freshly boiled and cooled water, and add 3 drops of phenolphthalein TS and 0.60 mL of 0.01 mol/L sodium hydroxide VS: a red color develops.

(3) Chloride <1.03>—Perform the test with 2.0 g of Fructose. Prepare the control solution with 1.0 mL of 0.01 mol/L

hydrochloric acid VS (not more than 0.018%).

(4) Sulfate <1.14>—Perform the test with 2.0 g of Fructose. Prepare the control solution with 1.0 mL of 0.005 mol/L sulfuric acid VS (not more than 0.024%).

(5) Sulfite—Dissolve 0.5 g of Fructose in 5 mL of water, and add 0.25 mL of 0.02 mol/L iodine: the color of the solution is yellow.

(6) Heavy metals <1.07>—Proceed with 5.0 g of Fructose according to Method 2, and perform the test. Prepare the control solution with 2.0 mL of Standard Lead Solution (not more than 4 ppm).

(7) Calcium—Dissolve 0.5 g of Fructose in 5 mL of water, add 2 to 3 drops of ammonia TS and 1 mL of ammonium oxalate TS, and allow to stand for 1 minute: the solution is clear.

(8) Arsenic <1.11>—Dissolve 1.5 g of Fructose in 5 mL of water, heat with 5 mL of dilute sulfuric acid and 1 mL of bromine TS on a water bath for 5 minutes, concentrate to 5 mL, and cool. Perform the test with this solution as the test solution (not more than 1.3 ppm).

(9) 5-Hydroxymethylfurfurals—Dissolve 5.0 g of Fructose in 100 mL of water, and read the absorbance at 284 nm as directed under Ultraviolet-visible Spectrophotometry <2.24>: the absorbance is not more than 0.32.

**Loss on drying** <2.41> Not more than 0.5% (1 g, in vacuum, silica gel, 3 hours).

**Residue on ignition** <2.44> Not more than 0.1% (1 g).

**Assay** Weigh accurately about 4 g of Fructose, previously dried, dissolve in 0.2 mL of ammonia TS and 80 mL of water, and after standing for 30 minutes add water to make exactly 100 mL, and determine the optical rotation,  $\alpha_D$ , in a 100-mm cell at  $20 \pm 1^\circ\text{C}$  as directed under Optical Rotation Determination <2.49>.

$$\text{Amount (mg) of } C_6H_{12}O_6 = |\alpha_D| \times 1087.0$$

**Containers and storage** Containers—Tight containers.

## Fructose Injection

果糖注射液

Fructose Injection is an aqueous solution for injection.

It contains not less than 95% and not more than 105% of the labeled amount of fructose ( $C_6H_{12}O_6$ : 180.16).

**Method of preparation** Prepare as directed under Injections, with Fructose. No preservative is added.

**Description** Fructose Injection is a colorless to pale yellow, clear liquid. It has a sweet taste.

**Identification** (1) Take a volume of Fructose Injection, equivalent to 1 g of Fructose according to the labeled amount, dilute with water or concentrate on a water bath to 20 mL, if necessary, and use this solution as the sample solution. Add 2 to 3 drops of the sample solution to 5 mL of boiling Fehling's TS: a red precipitate is produced.

(2) To 10 mL of the sample solution obtained in (1) add 0.1 g of resorcinol and 1 mL of hydrochloric acid, and warm

in a water bath for 3 minutes: a red color develops.

**pH** <2.54> 3.0 – 6.5 In the case where the labeled concentration of the injection exceeds 5%, dilute to 5% with water before the test.

**Purity** (1) Heavy metals <1.07>—Take a volume of Fructose Injection, equivalent to 5.0 g of Fructose, according to the labeled amount, and evaporate on a water bath to dryness. With the residue, proceed according to Method 2, and perform the test. Prepare the control solution with 2.0 mL of Standard Lead Solution.

(2) Arsenic <1.11>—Take a volume of Fructose Injection, equivalent to 1.5 g of Fructose, according to the labeled amount, dilute with water or concentrate on a water bath to 5 mL, if necessary, and add 5 mL of dilute sulfuric acid and 1 mL of bromine TS. Proceed as directed in the purity (8) under Fructose.

**Residue on ignition** <2.44> Measure exactly a volume of Fructose Injection, equivalent to 2 g of Fructose according to the labeled amount, evaporate on a water bath to dryness, and perform the test: the residue weighs not more than 2 mg.

**Extractable volume** <6.05> It meets the requirement.

**Pyrogen** <4.04> Perform the test with Fructose Injection stored in a container in a volume exceeding 10 mL: it meets the requirement.


**Assay** Measure exactly a volume of Fructose Injection equivalent to about 4 g of fructose ( $C_6H_{12}O_6$ ), add 0.2 mL of ammonia TS, dilute with water to make exactly 100 mL, shake well, and after allowing to stand for 30 minutes, determine the optical rotation,  $\alpha_D$ , in a 100-mm cell at  $20 \pm 1^\circ C$  as directed under Optical Rotation Determination <2.49>.

Amount (mg) of fructose ( $C_6H_{12}O_6$ ) =  $|\alpha_D| \times 1087.0$

**Containers and storage** Containers—Hermetic containers. Plastic containers for aqueous injections may be used.

## Furosemide

フロセミド



$C_{12}H_{11}ClN_2O_5S$ : 330.74

4-Chloro-2-[(furan-2-ylmethyl)amino]-5-sulfamoylbenzoic acid [54-31-9]

Furosemide, when dried, contains not less than 98.0% and not more than 101.0% of  $C_{12}H_{11}ClN_2O_5S$ .

**Description** Furosemide occurs as white, crystals or crystalline powder.

It is freely soluble in *N,N*-dimethylformamide, soluble in methanol, sparingly soluble in ethanol (99.5), slightly soluble in acetonitrile and in acetic acid (100), and practically insoluble in water.

It dissolves in dilute sodium hydroxide TS.

It is gradually colored by light.

Melting point: about  $205^\circ C$  (with decomposition).

**Identification** (1) Dissolve 25 mg of Furosemide in 10 mL of methanol. To 1 mL of this solution add 10 mL of 2 mol/L hydrochloric acid TS. Heat the solution under a reflux condenser on a water bath for 15 minutes, cool, and add 18 mL of sodium hydroxide TS to make weakly acidic: the solution responds to the Qualitative Tests <1.09> for primary aromatic amines, producing a red to red-purple color.

(2) Determine the absorption spectrum of a solution of Furosemide in dilute sodium hydroxide TS (1 in 125,000) as directed under Ultraviolet-visible Spectrophotometry <2.24>, and compare the spectrum with the Reference Spectrum or the spectrum of a solution of Furosemide Reference Standard prepared in the same manner as the sample solution: both spectra exhibit similar intensities of absorption at the same wavelengths.

(3) Determine the infrared absorption spectrum of Furosemide as directed in the potassium bromide disk method under Infrared Spectrophotometry <2.25>, and compare the spectrum with the Reference Spectrum or the spectrum of Furosemide Reference Standard: both spectra exhibit similar intensities of absorption at the same wave numbers.

**Purity** (1) Clarity and color of solution—Dissolve 0.5 g of Furosemide in 10 mL of a solution of sodium hydroxide (1 in 50): the solution is clear and colorless.

(2) Chloride <1.03>—Dissolve 2.6 g of Furosemide in 90 mL of dilute sodium hydroxide TS, add 2 mL of nitric acid, and filter. To 25 mL of the filtrate add 6 mL of dilute nitric acid and water to make 50 mL, and perform the test using this solution as the test solution. Prepare the control solution as follows: To 0.40 mL of 0.01 mol/L hydrochloric acid VS add 6 mL of dilute nitric acid and water to make 50 mL (not more than 0.020%).

(3) Sulfate <1.14>—To 20 mL of the filtrate obtained in (2) add 1 mL of dilute hydrochloric acid and water to make 50 mL, and perform the test using this solution as the test solution. Prepare the control solution as follows: To 0.35 mL of 0.005 mol/L sulfuric acid VS add 1 mL of dilute hydrochloric acid and water to make 50 mL (not more than 0.030%).

(4) Heavy metals <1.07>—Proceed with 2.0 g of Furosemide according to Method 2, and perform the test. Prepare the control solution with 2.0 mL of Standard Lead Solution (not more than 10 ppm).

(5) Related substances—Dissolve 25 mg of Furosemide in 25 mL of the dissolving solution, and use this solution as the sample solution. Pipet 1 mL of the sample solution, add the dissolving solution to make exactly 200 mL, and use this solution as the standard solution. Perform the test with exactly 20  $\mu L$  each of the sample solution and standard solution as directed under Liquid Chromatography <2.01> according to the following conditions, and determine each peak area by the automatic integration method: the area of each peak appeared ahead of the peak of furosemide is not more than 2/5 times the peak area of furosemide from the standard solution, the area of each peak appeared behind the peak of furosemide is not more than 1/4 times the peak area of furosemide from the standard solution, and the total area of these peaks is not more than 2 times the peak area of furosemide from the standard solution.

**Dissolving solution**—To 22 mL of acetic acid (100) add a mixture of water and acetonitrile (1:1) to make 1000 mL.

**Operating conditions**—

**Detector:** An ultraviolet absorption photometer (wavelength: 272 nm).

**Column:** A stainless steel column 4.6 mm in inside diameter and 25 cm in length, packed with octadecylsilanized silica gel for liquid chromatography (5  $\mu$ m in particle diameter).

**Column temperature:** A constant temperature of about 25°C.

**Mobile phase:** A mixture of water, tetrahydrofuran and acetic acid (100) (70:30:1).

**Flow rate:** Adjust the flow rate so that the retention time of furosemide is about 18 minutes.

**Time span of measurement:** About 2.5 times as long as the retention time of furosemide beginning after the solvent peak.

**System suitability**—

**Test for required detectability:** Measure exactly 2 mL of the standard solution, and add the dissolving solution to make exactly 50 mL. Confirm that the peak area of furosemide obtained from 20  $\mu$ L of this solution is equivalent to 3.2 to 4.8% of that obtained from 20  $\mu$ L of the standard solution.

**System performance:** When the procedure is run with 20  $\mu$ L of the standard solution under the above operating conditions, the number of theoretical plates and the symmetry factor of the peak of furosemide is not less than 7000 and not more than 1.5, respectively.

**System repeatability:** When the test is repeated 6 times with 20  $\mu$ L of the standard solution under the above operating conditions, the relative standard deviation of the peak area of furosemide is not more than 2.0%.

**Loss on drying** <2.41> Not more than 0.5% (1 g, 105°C, 4 hours).

**Residue on ignition** <2.44> Not more than 0.1% (1 g).

**Assay** Weigh accurately about 0.5 g of Furosemide, previously dried, dissolve in 50 mL of *N,N*-dimethylformamide, and titrate <2.50> with 0.1 mol/L sodium hydroxide VS until the color of the solution changes from yellow to blue (indicator: 3 drops of bromothymol blue TS). Perform a blank determination with a mixture of 50 mL of *N,N*-dimethylformamide and 15 mL of water, and make any necessary correction

Each mL of 0.1 mol/L sodium hydroxide VS  
= 33.07 mg of  $C_{12}H_{11}ClN_2O_5S$

**Containers and storage** Containers—Tight containers.  
Storage—Light-resistant.

## Furosemide Tablets

フロセミド錠

Furosemide Tablets contain not less than 95.0% and not more than 105.0% of the labeled amount of furosemide ( $C_{12}H_{11}ClN_2O_5S$ : 330.74).

**Method of preparation** Prepare as directed under Tablets,

with Furosemide.

**Identification** (1) Shake well a quantity of powdered Furosemide Tablets, equivalent to 0.2 g of Furosemide according to the labeled amount, with 40 mL of acetone, and filter. To 0.5 mL of the filtrate add 10 mL of 2 mol/L hydrochloric acid TS, and heat under a reflux condenser on a water bath for 15 minutes. After cooling, add 18 mL of sodium hydroxide TS to make the solution slightly acetic: the solution responds to the Qualitative Tests <1.09> for primary aromatic amines, producing a red to red-purple color.

(2) Determine the absorption spectrum of the sample solution obtained in the Assay as directed under Ultraviolet-visible Spectrophotometry <2.24>: it exhibits maxima between 227 nm and 231 nm, between 269 nm and 273 nm, and between 330 nm and 336 nm.

**Purity** To a quantity of powdered Furosemide Tablets, equivalent to 40 mg of Furosemide according to the labeled amount, add about 30 mL of acetone, shake well, and add acetone to make exactly 50 mL. Centrifuge the solution, add 3.0 mL of water to 1.0 mL of the supernatant liquid, cool in ice, add 3.0 mL of dilute hydrochloric acid and 0.15 mL of sodium nitrite TS, shake, and allow to stand for 1 minute. Add 1.0 mL of ammonium amidosulfate TS, shake well, allow to stand for 3 minutes, add 1.0 mL of *N,N*-diethyl-*N'*-1-naphthylethylenediamine oxalate TS, shake well, and allow to stand for 5 minutes. Perform the test with this solution as directed under Ultraviolet-visible Spectrophotometry <2.24>, using a solution prepared in the same manner with 1.0 mL of acetone as the blank: the absorbance at 530 nm is not more than 0.10.

**Uniformity of dosage units** <6.02> Perform the test according to the following method: it meets the requirement of the Content uniformity test.

To 1 tablet of Furosemide Tablets add a suitable amount of 0.05 mol/L sodium hydroxide TS, shake to disintegrate, then add 0.05 mol/L sodium hydroxide TS to make exactly *V* mL so that each mL contains about 0.4 mg of furosemide ( $C_{12}H_{11}ClN_2O_5S$ ). Filter the solution, discard the first 10 mL of the filtrate, pipet the subsequent 2 mL of the filtrate, add 0.05 mol/L sodium hydroxide TS to make exactly 100 mL, and use this solution as the sample solution. Proceed as directed in the Assay.

$$\begin{aligned} &\text{Amount (mg) of furosemide } (C_{12}H_{11}ClN_2O_5S) \\ &= W_S \times (A_T/A_S) \times (V/50) \end{aligned}$$

$W_S$ : Amount (mg) of Furosemide Reference Standard

**Dissolution** <6.10> Perform the test according to the following method: it meets the requirement.

Perform the test with 1 tablet of Furosemide Tablets at 50 revolutions per minute according to the Paddle method, using 900 mL of 2nd fluid for dissolution test as the dissolution medium. Withdraw 20 mL or more of the dissolution medium 15 minutes after starting the test for a 20-mg tablet or 30 minutes after for a 40-mg tablet, and filter through a membrane filter with pore size of not more than 0.45  $\mu$ m. Discard the first 10 mL of the filtrate, pipet *V* mL of the subsequent filtrate, add 2nd fluid for dissolution test to make exactly *V'* mL so that each mL contains about 10  $\mu$ g of furosemide ( $C_{12}H_{11}ClN_2O_5S$ ) according to the labeled amount, and use this solution as the sample solution. Separately, weigh accurately about 20 mg of Furosemide Reference Standard, previously

dried at 105°C for 4 hours, and dissolve in 5 mL of methanol, and add 2nd fluid for dissolution test to make exactly 100 mL. Pipet 5 mL of this solution, add 2nd fluid for dissolution test to make exactly 100 mL, and use this solution as the standard solution. Determine the absorbances,  $A_T$  and  $A_S$ , of the sample solution and the standard solution at 277 nm as directed under Ultraviolet-visible Spectrophotometry <2.24>: the dissolution rates for a 20-mg tablet in 15 minutes and for a 40-mg tablet in 30 minutes are not less than 80%, respectively.

Dissolution rate (%) with respect to the labeled amount of furosemide ( $C_{12}H_{11}ClN_2O_5S$ )

$$= W_S \times (A_T/A_S) \times (V'/V) \times (1/C) \times 45$$

$W_S$ : Amount (mg) of Furosemide Reference Standard

$C$ : Labeled amount (mg) of furosemide ( $C_{12}H_{11}ClN_2O_5S$ ) in 1 tablet

**Assay** Weigh accurately the mass of not less than 20 Furosemide Tablets, and powder. Weigh accurately a portion of the powder, equivalent to about 40 mg of furosemide ( $C_{12}H_{11}ClN_2O_5S$ ), add about 70 mL of 0.05 mol/L sodium hydroxide TS, shake well, and add 0.05 mol/L sodium hydroxide TS to make exactly 100 mL. Filter, discard the first 10 mL of the filtrate, pipet 2 mL of the subsequent filtrate, add 0.05 mol/L sodium hydroxide TS to make exactly 100 mL, and use this solution as the sample solution. Separately, weigh accurately about 20 mg of Furosemide Reference Standard, previously dried at 105°C for 4 hours, and dissolve in 0.05 mol/L sodium hydroxide TS to make exactly 50 mL. Pipet 2 mL of this solution, add 0.05 mol/L sodium hydroxide TS to make exactly 100 mL, and use this solution as the standard solution. Determine the absorbances,  $A_T$  and  $A_S$ , of the sample solution and standard solution at 271 nm as directed under the Ultraviolet-visible Spectrophotometry <2.24>.

$$\begin{aligned} &\text{Amount (mg) of furosemide (} C_{12}H_{11}ClN_2O_5S \text{)} \\ &= W_S \times (A_T/A_S) \times 2 \end{aligned}$$


$W_S$ : Amount (mg) of Furosemide Reference Standard

**Containers and storage** Containers—Tight containers.

Storage—Light-resistant.

## Fursultiamine Hydrochloride

フルスルチアミン塩酸塩



$C_{17}H_{26}N_4O_3S_2 \cdot HCl$ : 435.00

*N*-(4-Amino-2-methylpyrimidin-5-ylmethyl)-*N*-(1*Z*)-4-hydroxy-1-methyl-2-[(2*RS*)-tetrahydrofuran-2-ylmethyl]disulfanylmethylbut-1-en-1-yl]formamide monohydrochloride [804-30-8, Fursultiamine]

Fursultiamine Hydrochloride contains not less than

98.5% of  $C_{17}H_{26}N_4O_3S_2 \cdot HCl$ , calculated on the dried basis.

**Description** Fursultiamine Hydrochloride occurs as white crystals or crystalline powder. It is odorless or has a characteristic odor, and has a bitter taste.

It is freely soluble in water, in methanol and in ethanol (95), and practically insoluble in diethyl ether.

**Identification (1)** Dissolve 5 mg of Fursultiamine Hydrochloride in 6 mL of 0.1 mol/L hydrochloric acid TS, add 0.1 g of zinc powder, allow to stand for several minutes, and filter. To 3 mL of the filtrate, add 3 mL of sodium hydroxide TS and 0.5 mL of potassium hexacyanoferrate (III) TS, then add 5 mL of 2-methyl-1-propanol, shake vigorously for 2 minutes, allow to stand to separate the 2-methyl-1-propanol layer, and examine under ultraviolet light (main wavelength: 365 nm): the 2-methyl-1-propanol layer shows a blue-purple fluorescence. The fluorescence disappears by acidifying, and appears again by alkalifying.

(2) Determine the infrared absorption spectrum of a solution of Fursultiamine Hydrochloride, previously dried in a desiccator (in vacuum, phosphorus (V) oxide) for 24 hours, as directed in the potassium bromide disk method under Infrared Spectrophotometry <2.25>, and compare the spectrum with the Reference Spectrum, or with the spectrum of Fursultiamine Hydrochloride Reference Standard, previously dried in a desiccator (in vacuum, phosphorus (V) oxide) for 24 hours: both spectra exhibit similar intensities of absorption at the same wave numbers. If any differences appear, dissolve the Fursultiamine Hydrochloride in water, evaporate the water, and dry the residue in a desiccator (in vacuum, phosphorus (V) oxide) for 24 hours, and repeat the test.

(3) A solution of Fursultiamine Hydrochloride (1 in 50) responds to the Qualitative Tests <1.09> (2) for chloride.

**Purity (1)** Clarity of solution—Dissolve 1.0 g of Fursultiamine Hydrochloride in 20 mL of water: the solution is clear and colorless.

(2) Sulfate <1.14>—Proceed with 1.5 g of Fursultiamine Hydrochloride, and perform the test. Prepare the control solution with 0.35 mL of 0.005 mol/L sulfuric acid VS (not more than 0.011%).

(3) Heavy metals <1.07>—Proceed with 1.0 g of Fursultiamine Hydrochloride according to Method 2, and perform the test. Prepare the control solution with 2.0 mL of Standard Lead Solution (not more than 20 ppm).

(4) Related substances—Dissolve 0.10 g of Fursultiamine Hydrochloride in 100 mL of the mobile phase, and use this solution as the sample solution. Pipet 1 mL of sample solution, add the mobile phase to make exactly 100 mL, and use this solution as the standard solution. Perform the test with exactly 10  $\mu$ L each of the sample solution and standard solution as directed under Liquid Chromatography <2.01> according to the following condition. Determine each peak area of each solution by the automatic integration method: the total area of the peaks other than the peak of fursultiamine from the sample solution is not larger than the peak area of fursultiamine from the standard solution.

**Operating conditions—**

Detector, column, column temperature, mobile phase, flow rate, and selection of column: Proceed as directed in the operating conditions in the Assay.

Detector sensitivity: Adjust the detection sensitivity so that

the peak height of fursultiamine from 10  $\mu$ L of the standard solution is between 20 mm and 30 mm.

Time span of measurement: About 3 times as long as the retention time of fursultiamine.

**Water** <2.48> Not more than 5.0% (0.3 g, direct titration).

**Residue on ignition** <2.44> Not more than 0.1% (1 g).

**Assay** Weigh accurately about 55 mg each of Fursultiamine Hydrochloride and Fursultiamine Hydrochloride Reference Standard (previously determined the water <2.48> in the same manner as Fursultiamine Hydrochloride) and dissolve each in 50 mL of water, and add exactly 10 mL each of the internal standard solution, then add water to make exactly 100 mL. To 8 mL each of the solution add water to make 50 mL, and use these solutions as the sample solution and standard solution, respectively. Perform the test with 10  $\mu$ L each of the sample solution and standard solution as directed under Liquid Chromatography <2.01> according to the following conditions, and calculate the ratios,  $Q_T$  and  $Q_S$ , of the peak area of fursultiamine to that of the internal standard, respectively.

$$\begin{aligned} &\text{Amount (mg) of } C_{17}H_{26}N_4O_3S_2 \cdot HCl \\ &= W_S \times (Q_T/Q_S) \end{aligned}$$

$W_S$ : Amount (mg) of Fursultiamine Hydrochloride Reference Standard, calculated on the anhydrous basis

**Internal standard solution**—A solution of isopropyl 4-aminobenzoate in ethanol (95) (3 in 400).

**Operating conditions**—

**Detector**: An ultraviolet absorption photometer (wavelength: 254 nm).

**Column**: A stainless steel column about 4 mm in inside diameter and about 15 cm in length, packed with octadecylsilanized silica gel for liquid chromatography (5  $\mu$ m in particle diameter).

**Column temperature**: A constant temperature of about 50°C.

**Mobile phase**: Dissolve 1.01 g of sodium 1-heptane sulfonate in 1000 mL of diluted acetic acid (100) (1 in 100). To 675 mL of this solution add 325 mL of a mixture of methanol and acetonitrile (3:2).


**Flow rate**: Adjust the flow rate so that the retention time of Fursultiamine is about 9 minutes.

**Selection of column**: Proceed with 10  $\mu$ L of the standard solution under the above operating conditions and calculate the resolution. Use a column giving elution of fursultiamine and the internal standard in this order with the resolution between these peaks being not less than 10.

**Containers and storage** Containers—Tight containers.

## Gabexate Mesilate

ガベキサートメシル酸塩



$C_{16}H_{23}N_3O_4 \cdot CH_4O_3S$ : 417.48

Ethyl 4-(6-guanidinohexanoyloxy)benzoate

monomethanesulfonate [56974-61-9]

Gabexate Mesilate, when dried, contains not less than 98.5% of  $C_{16}H_{23}N_3O_4 \cdot CH_4O_3S$ .

**Description** Gabexate Mesilate occurs as white crystals or crystalline powder.

It is very soluble in water, freely soluble in ethanol (95), and practically insoluble in diethyl ether.

**Identification** (1) To 4 mL of a solution of Gabexate Mesilate (1 in 2000) add 2 mL of 1-naphthol TS and 1 mL of diacetyl TS, and allow to stand for 10 minutes: a red color develops.

(2) Dissolve 1 g of Gabexate Mesilate in 5 mL of water, add 2 mL of sodium hydroxide TS, and heat in a water bath for 5 minutes. After cooling, add 2 mL of dilute nitric acid and 5 mL of ethanol (95), shake, add 5 drops of iron (III) chloride TS, and shake: a purple color develops.

(3) Determine the absorption spectrum of a solution of Gabexate Mesilate (1 in 100,000) as directed under Ultraviolet-visible Spectrophotometry <2.24>, and compare the spectrum with the Reference Spectrum or the spectrum of a solution of Gabexate Mesilate Reference Standard prepared in the same manner as the sample solution: both spectra exhibit similar intensities of absorption at the same wavelengths.

(4) To 0.1 g of Gabexate Mesilate add 0.2 g of sodium hydroxide, fuse by heating gently, and continue the heating for 20 to 30 seconds. After cooling, add 0.5 mL of water and 3 mL of dilute hydrochloric acid, and warm: the gas evolved changes a potassium iodate-starch paper to blue.

**pH** <2.54> Dissolve 1.0 g of Gabexate Mesilate in 10 mL of water: the pH of the solution is between 4.5 and 5.5.

**Melting point** <2.60> 90 – 93°C

**Purity** (1) Clarity and color of solution—Dissolve 1.0 g of Gabexate Mesilate in 10 mL of water: the solution is clear and colorless.

(2) Heavy metals <1.07>—Proceed with 2.0 g of Gabexate Mesilate according to Method 1, and perform the test. Prepare the control solution with 2.0 mL of Standard Lead Solution (not more than 10 ppm).

(3) Arsenic <1.11>—Dissolve 2.0 g of Gabexate Mesilate in 20 mL of 1 mol/L hydrochloric acid TS by heating in a water bath, and continue the heating for 20 minutes. After cooling, centrifuge, and use 10 mL of the supernatant liquid as the test solution. Perform the test (not more than 2 ppm).

(4) Ethyl parahydroxybenzoate—Weigh 50 mg of Gabexate Mesilate, previously dried, and dissolve in dilute ethanol to make exactly 100 mL. Pipet 5 mL of this solution, add exactly 5 mL of the internal standard solution, and use this solution as the sample solution. Separately, dissolve 5.0 mg of ethyl parahydroxybenzoate in dilute ethanol to make exactly 100 mL. Pipet 1 mL of this solution, and add dilute ethanol to make exactly 20 mL. To exactly 5 mL of this solution add exactly 5 mL of the internal standard solution, and use this solution as the standard solution. Perform the test with 3  $\mu$ L each of the sample solution and standard solution as directed under Liquid Chromatography <2.01> according to the following conditions, and calculate the ratios,  $Q_T$  and  $Q_S$ , of the peak area of ethyl parahydroxybenzoate to that of the internal standard:  $Q_T$  is not larger than  $Q_S$ .

the peak height of fursultiamine from 10  $\mu$ L of the standard solution is between 20 mm and 30 mm.

Time span of measurement: About 3 times as long as the retention time of fursultiamine.

**Water** <2.48> Not more than 5.0% (0.3 g, direct titration).

**Residue on ignition** <2.44> Not more than 0.1% (1 g).

**Assay** Weigh accurately about 55 mg each of Fursultiamine Hydrochloride and Fursultiamine Hydrochloride Reference Standard (previously determined the water <2.48> in the same manner as Fursultiamine Hydrochloride) and dissolve each in 50 mL of water, and add exactly 10 mL each of the internal standard solution, then add water to make exactly 100 mL. To 8 mL each of the solution add water to make 50 mL, and use these solutions as the sample solution and standard solution, respectively. Perform the test with 10  $\mu$ L each of the sample solution and standard solution as directed under Liquid Chromatography <2.01> according to the following conditions, and calculate the ratios,  $Q_T$  and  $Q_S$ , of the peak area of fursultiamine to that of the internal standard, respectively.

$$\begin{aligned} &\text{Amount (mg) of } C_{17}H_{26}N_4O_3S_2 \cdot HCl \\ &= W_S \times (Q_T/Q_S) \end{aligned}$$

$W_S$ : Amount (mg) of Fursultiamine Hydrochloride Reference Standard, calculated on the anhydrous basis

**Internal standard solution**—A solution of isopropyl 4-aminobenzoate in ethanol (95) (3 in 400).

**Operating conditions**—

**Detector**: An ultraviolet absorption photometer (wavelength: 254 nm).

**Column**: A stainless steel column about 4 mm in inside diameter and about 15 cm in length, packed with octadecylsilanized silica gel for liquid chromatography (5  $\mu$ m in particle diameter).

**Column temperature**: A constant temperature of about 50°C.

**Mobile phase**: Dissolve 1.01 g of sodium 1-heptane sulfonate in 1000 mL of diluted acetic acid (100) (1 in 100). To 675 mL of this solution add 325 mL of a mixture of methanol and acetonitrile (3:2).


**Flow rate**: Adjust the flow rate so that the retention time of Fursultiamine is about 9 minutes.

**Selection of column**: Proceed with 10  $\mu$ L of the standard solution under the above operating conditions and calculate the resolution. Use a column giving elution of fursultiamine and the internal standard in this order with the resolution between these peaks being not less than 10.

**Containers and storage** Containers—Tight containers.

## Gabexate Mesilate

ガベキサートメシル酸塩



$C_{16}H_{23}N_3O_4 \cdot CH_4O_3S$ : 417.48

Ethyl 4-(6-guanidinohexanoyloxy)benzoate

monomethanesulfonate [56974-61-9]

Gabexate Mesilate, when dried, contains not less than 98.5% of  $C_{16}H_{23}N_3O_4 \cdot CH_4O_3S$ .

**Description** Gabexate Mesilate occurs as white crystals or crystalline powder.

It is very soluble in water, freely soluble in ethanol (95), and practically insoluble in diethyl ether.

**Identification (1)** To 4 mL of a solution of Gabexate Mesilate (1 in 2000) add 2 mL of 1-naphthol TS and 1 mL of diacetyl TS, and allow to stand for 10 minutes: a red color develops.

**(2)** Dissolve 1 g of Gabexate Mesilate in 5 mL of water, add 2 mL of sodium hydroxide TS, and heat in a water bath for 5 minutes. After cooling, add 2 mL of dilute nitric acid and 5 mL of ethanol (95), shake, add 5 drops of iron (III) chloride TS, and shake: a purple color develops.

**(3)** Determine the absorption spectrum of a solution of Gabexate Mesilate (1 in 100,000) as directed under Ultraviolet-visible Spectrophotometry <2.24>, and compare the spectrum with the Reference Spectrum or the spectrum of a solution of Gabexate Mesilate Reference Standard prepared in the same manner as the sample solution: both spectra exhibit similar intensities of absorption at the same wavelengths.

**(4)** To 0.1 g of Gabexate Mesilate add 0.2 g of sodium hydroxide, fuse by heating gently, and continue the heating for 20 to 30 seconds. After cooling, add 0.5 mL of water and 3 mL of dilute hydrochloric acid, and warm: the gas evolved changes a potassium iodate-starch paper to blue.

**pH** <2.54> Dissolve 1.0 g of Gabexate Mesilate in 10 mL of water: the pH of the solution is between 4.5 and 5.5.

**Melting point** <2.60> 90 – 93°C

**Purity (1)** Clarity and color of solution—Dissolve 1.0 g of Gabexate Mesilate in 10 mL of water: the solution is clear and colorless.

**(2)** Heavy metals <1.07>—Proceed with 2.0 g of Gabexate Mesilate according to Method 1, and perform the test. Prepare the control solution with 2.0 mL of Standard Lead Solution (not more than 10 ppm).

**(3)** Arsenic <1.11>—Dissolve 2.0 g of Gabexate Mesilate in 20 mL of 1 mol/L hydrochloric acid TS by heating in a water bath, and continue the heating for 20 minutes. After cooling, centrifuge, and use 10 mL of the supernatant liquid as the test solution. Perform the test (not more than 2 ppm).

**(4)** Ethyl parahydroxybenzoate—Weigh 50 mg of Gabexate Mesilate, previously dried, and dissolve in dilute ethanol to make exactly 100 mL. Pipet 5 mL of this solution, add exactly 5 mL of the internal standard solution, and use this solution as the sample solution. Separately, dissolve 5.0 mg of ethyl parahydroxybenzoate in dilute ethanol to make exactly 100 mL. Pipet 1 mL of this solution, and add dilute ethanol to make exactly 20 mL. To exactly 5 mL of this solution add exactly 5 mL of the internal standard solution, and use this solution as the standard solution. Perform the test with 3  $\mu$ L each of the sample solution and standard solution as directed under Liquid Chromatography <2.01> according to the following conditions, and calculate the ratios,  $Q_T$  and  $Q_S$ , of the peak area of ethyl parahydroxybenzoate to that of the internal standard:  $Q_T$  is not larger than  $Q_S$ .

**Internal standard solution**—A solution of butyl parahydroxybenzoate in dilute ethanol (1 in 5000).

**Operating conditions**—

Proceed as directed in the operating conditions in the Assay.

**System suitability**—

Proceed as directed in the system suitability in the Assay.

(5) **Related substances**—Dissolve 0.20 g of Gabexate Mesilate in 5 mL of ethanol (95), and use this solution as the sample solution. Pipet 1 mL of the sample solution, add ethanol (95) to make exactly 100 mL, and use this solution as the standard solution. Perform the test with these solutions as directed under Thin-layer Chromatography <2.03>. Spot 5  $\mu$ L each of the sample solution and standard solution on a plate of silica gel for thin-layer chromatography. Develop the plate with a mixture of ethyl acetate, water and acetic acid (100) (3:1:1) to a distance of about 10 cm, and air-dry the plate until it has no acetic odor. Spray evenly a solution of 8-quinolinol in acetone (1 in 1000) on the plate, and after air-drying, spray evenly bromine-sodium hydroxide TS: the spots other than the principal spot from the sample solution are not more intense than the spot from the standard solution.

**Loss on drying** <2.41> Not more than 0.30% (1 g, in vacuum, silica gel, 4 hours).

**Residue on ignition** <2.44> Not more than 0.1% (1 g).

**Assay** Weigh accurately about 50 mg each of Gabexate Mesilate and Gabexate Mesilate Reference Standard, previously dried, and dissolve each in dilute ethanol to make exactly 100 mL. Pipet 5 mL each of these solutions, add exactly 5 mL each of the internal standard solution, and use these solutions as the sample solution and standard solution, respectively. Perform the test with 3  $\mu$ L each of the sample solution and standard solution as directed under Liquid Chromatography <2.01> according to the following conditions, and calculate the ratios,  $Q_T$  and  $Q_S$ , of the peak area of gabexate to that of the internal standard.

$$\begin{aligned} &\text{Amount (mg) of } C_{16}H_{23}N_3O_4 \cdot CH_4O_3S \\ &= W_S \times (Q_T/Q_S) \end{aligned}$$

$W_S$ : Amount (mg) of Gabexate Mesilate Reference Standard

**Internal standard solution**—A solution of butyl parahydroxybenzoate in dilute ethanol (1 in 5000).

**Operating conditions**—

**Detector**: An ultraviolet absorption photometer (wavelength: 245 nm).

**Column**: A stainless steel column 4.6 mm in inside diameter and 15 cm in length, packed with octadecylsilanized silica gel for liquid chromatography (5  $\mu$ m in particle diameter).

**Column temperature**: A constant temperature of about 25°C.

**Mobile phase**: A mixture of methanol, a solution of sodium lauryl sulfate (1 in 1000), a solution of sodium 1-heptane sulfonate (1 in 200) and acetic acid (100) (540:200:20:1).

**Flow rate**: Adjust the flow rate so that the retention time of gabexate is about 13 minutes.

**System suitability**—

**System performance**: When the procedure is run with 3  $\mu$ L of the standard solution under the above operating condi-

tions, the internal standard and gabexate are eluted in this order with the resolution between these peaks being not less than 5.

**System repeatability**: When the test is repeated 6 times with 3  $\mu$ L of the standard solution under the above operating conditions, the relative standard deviation of the ratios of the peak area of gabexate to that of the internal standard is not more than 1.0%.

**Containers and storage** Containers—Tight containers.

## $\beta$ -Galactosidase (*Aspergillus*)

$\beta$ -ガラクトシダーゼ (アスペルギルス)

[9031-11-2]

$\beta$ -Galactosidase (*Aspergillus*) contains an enzyme produced by *Aspergillus oryzae*. It is an enzyme drug having lactose decomposition activity.

It contains 8000 to 12000 units per g.

Usually, it is diluted with a mixture of Maltose Hydrate and Dextrin, Maltose Hydrate and D-Mannitol, or Maltose Hydrate, Dextrin and D-Mannitol.

**Description**  $\beta$ -Galactosidase (*Aspergillus*) occurs as a white to light yellow powder.

It is slightly soluble in water with a turbidity, and practically insoluble in ethanol (95) and in diethyl ether.

**Identification** (1) Dissolve 25 mg of  $\beta$ -Galactosidase (*Aspergillus*) in 100 mL of water, then to 1 mL of this solution add 9 mL of lactose substrate TS, and stand at 30°C for 10 minutes. To 1 mL of this solution add 6 mL of glucose detection TS, and stand at 30°C for 10 minutes: a red to red-purple color develops.

(2) Dissolve 0.1 g of  $\beta$ -Galactosidase (*Aspergillus*) in 100 mL of water, and filter the solution if necessary. Determine the absorption spectrum of the solution as directed under Ultraviolet-visible Spectrophotometry <2.24>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wavelengths.

**Purity** (1) **Odor**— $\beta$ -Galactosidase (*Aspergillus*) has no any rancid odor.

(2) **Heavy metals** <1.07>—Proceed with 1.0 g of  $\beta$ -Galactosidase (*Aspergillus*) according to Method 2, and perform the test. Prepare the control solution with 2.0 mL of Standard Lead Solution (not more than 20 ppm).

(3) **Arsenic** <1.11>—Prepare the test solution with 1.0 g of  $\beta$ -Galactosidase (*Aspergillus*) according to Method 3, and perform the test (not more than 2 ppm).

**Loss on drying** <2.41> Not more than 9.0% (0.5 g, in vacuum, 80°C, 4 hours).

**Residue on ignition** <2.44> Not more than 3% (0.5 g).

**Nitrogen content** Weigh accurately about 70 mg of  $\beta$ -Galactosidase (*Aspergillus*), and perform the test as directed under Nitrogen Determination <1.08>: the amount of nitrogen (N: 14.01) is between 0.5% and 5.0%, calculated on the dried basis.

**Assay** (i) Substrate solution: Dissolve 0.172 g of 2-nitrophenyl- $\beta$ -D-galactopyranoside in disodium hydrogen-phosphate-citric acid buffer solution, pH 4.5 to make 100 mL.

(ii) Procedure: Weigh accurately about 25 mg of  $\beta$ -Galactosidase (*Aspergillus*), dissolve in water to make exactly 100 mL, then pipet 2 mL of this solution, add water to make exactly 50 mL, and use this solution as the sample solution. Take exactly 3.5 mL of the substrate solution, stand at  $30 \pm 0.1^\circ\text{C}$  for 5 minutes, add exactly 0.5 mL of the sample solution, immediately mix, and stand at  $30 \pm 0.1^\circ\text{C}$  for exactly 10 minutes, then add exactly 1 mL of sodium carbonate TS and mix immediately. Perform the test as directed under Ultraviolet-visible Spectrophotometry <2.24>, and determine the absorbance,  $A_1$ , of this solution at 420 nm using water as the control. Separately, take exactly 3.5 mL of the substrate solution, add exactly 1 mL of sodium carbonate TS and mix, then add exactly 0.5 mL of the sample solution and mix. Determine the absorbance,  $A_2$ , of this solution in the same manner as above.

$$\begin{aligned} &\text{Units per g of } \beta\text{-Galactosidase (Aspergillus)} \\ &= (1/W) \times \{(A_1 - A_2)/0.917\} \times (1/0.5) \times (1/10) \end{aligned}$$

0.917: Absorbance of 1  $\mu\text{mol}$ /5 mL of *o*-nitrophenol

$W$ : Amount (g) of the sample in the sample solution per mL

Unit: One unit indicates an amount of the enzyme which decomposes 1  $\mu\text{mol}$  of 2-nitrophenyl- $\beta$ -D-galactopyranoside in 1 minute under the above conditions.

**Containers and storage** Containers—Tight containers.

Storage—In a cold place.

## $\beta$ -Galactosidase (Penicillium)

$\beta$ -ガラクトシダーゼ (ペニシリウム)

[9031-11-2]

$\beta$ -Galactosidase (*Penicillium*) contains an enzyme, having lactose decomposition activity, produced by *Penicillium multicolor*.

It contains not less than 8500 units and not more than 11,500 units in each g.

Usually, it is diluted with D-mannitol.

**Description**  $\beta$ -Galactosidase (*Penicillium*) occurs as a white to pale yellowish white, crystalline powder or powder.

It is soluble in water with a turbidity, and practically insoluble in ethanol (95).

It is hygroscopic.

**Identification** (1) Dissolve 0.05 g of  $\beta$ -Galactosidase (*Penicillium*) in 100 mL of water, then to 0.2 mL of this solution add 0.2 mL of lactose substrate TS, and allow to stand at  $30^\circ\text{C}$  for 10 minutes. To this solution add 3 mL of glucose detection TS, and allow to stand at  $30^\circ\text{C}$  for 10 minutes: a red to red-purple color develops.

(2) Dissolve 0.15 g of  $\beta$ -Galactosidase (*Penicillium*) in 100 mL of water, filter if necessary, and determine the absorption spectrum of this solution as directed under Ultraviolet-visible Spectrophotometry <2.24>: it exhibits a maxi-

mum between 278 nm and 282 nm.

**Purity** (1) Odor— $\beta$ -Galactosidase (*Penicillium*) has no any rancid odor.

(2) Heavy metals <1.07>—Proceed with 1.0 g of  $\beta$ -Galactosidase (*Penicillium*) according to Method 2, and perform the test. Prepare the control solution with 2.0 mL of Standard Lead Solution (not more than 20 ppm).

(3) Arsenic <1.11>—Prepare the test solution with 1.0 g of  $\beta$ -Galactosidase (*Penicillium*) according to Method 3, and perform the test (not more than 2 ppm).

(4) Nitrogen—Weigh accurately about 0.1 g of  $\beta$ -Galactosidase (*Penicillium*), and perform the test as directed under Nitrogen Determination <1.08>: not more than 3 mg of nitrogen (N: 14.01) is found for each labeled 1000 Units.

(5) Protein contaminants—Dissolve 0.15 g of  $\beta$ -Galactosidase (*Penicillium*) in 4 mL of water, and use this solution as the sample solution. Perform the test with 15  $\mu\text{L}$  of the sample solution as directed under Liquid Chromatography <2.01> according to the following conditions, and determine each peak area by the automatic integration method: the total area of the peaks other than the peak having retention time of about 19 minutes is not more than 75% of the total area of all peaks, and the areas of peaks other than the peaks having retention times of about 3, 16 and 19 minutes are not more than 15% of the total area of all peaks.

**Operating conditions**—

Detector: An ultraviolet absorption photometer (wavelength: 280 nm).

Column: A stainless steel column about 7.5 mm in inside diameter and about 75 mm in length, packed with strongly acidic ion-exchange resin for liquid chromatography of sulfopropyl group-binding hydrophilic polymer (10  $\mu\text{m}$  in particle diameter).

Column temperature: A constant temperature of about  $20^\circ\text{C}$ .

Mobile phase: A solution obtained by dissolving 2.83 g of sodium acetate in 1000 mL of water, and adjusting to pH 4.5 with acetic acid (100) (mobile phase A), and a solution obtained by dissolving 29.2 g of sodium chloride in 1000 mL of mobile phase A (mobile phase B).

Flow system: Adjust a linear concentration gradient from the mobile phase A to the mobile phase B immediately after injection of the sample so that the retention times of non-retaining protein and the enzyme protein are about 3 minutes and 19 minutes, respectively, when the flow runs 0.8 mL per minute, and then continue the running of the mobile phase B.

Selection of column: Dissolve 15 mg of  $\beta$ -lactoglobulin in 4.5 mL of water, add 0.5 mL of a solution of cytosine (1 in 5000), and use this solution as the column-selecting solution. Proceed with 15  $\mu\text{L}$  of the column-selecting solution under the above operating conditions, and calculate the resolution. Use a column giving elution of cytosine and  $\beta$ -lactoglobulin in this order with the resolution between these peaks being not less than 4.

Detection sensitivity: Adjust the detection sensitivity so that the peak height of  $\beta$ -lactoglobulin from 15  $\mu\text{L}$  of the column-selecting solution is between 5 cm and 14 cm.

Time span of measurement: About 1.4 times as long as the retention time of  $\beta$ -lactoglobulin.

**Loss on drying** <2.41> Not more than 5.0% (0.5 g, in vacuum, phosphorus (V) oxide, 4 hours).

**Residue on ignition** <2.44> Not more than 2% (1 g).



**Assay** (i) Substrate solution—Dissolve 0.603 g of 2-nitrophenyl- $\beta$ -D-galactopyranoside in disodium hydrogen phosphate-citric acid buffer solution, pH 4.5 to make 100 mL.

(ii) Procedure—Weigh accurately about 0.15 g of  $\beta$ -Galactosidase (Penicillium), dissolve in water with thorough shaking to make exactly 100 mL, and allow to stand at room temperature for an hour. Pipet 2 mL of this solution, add disodium hydrogen phosphate-citric acid buffer solution, pH 4.5 to make exactly 100 mL, and use this solution as the sample solution. Transfer exactly 0.5 mL of the sample solution to a test tube, stand at  $30 \pm 0.1^\circ\text{C}$  for 10 minutes, add exactly 0.5 mL of the substrate solution previously kept at  $30 \pm 0.1^\circ\text{C}$ , then mix immediately, and stand at  $30 \pm 0.1^\circ\text{C}$  for exactly 10 minutes. Then add exactly 1 mL of sodium carbonate TS, mix immediately to stop the reaction. To this solution add exactly 8 mL of water, mix, and use as the colored sample solution. Separately, pipet 0.5 mL of disodium hydrogen phosphate-citric acid buffer solution, pH 4.5, then proceed in the same manner as the sample solution, and use the solution so obtained as the colored blank solution. Perform the test with the colored sample solution and the colored blank solution as directed under Ultraviolet-visible Spectrophotometry <2.2.4>, using water as the blank, and determine the absorbances,  $A_T$  and  $A_B$ , at 420 nm.

$$\begin{aligned} &\text{Units per g of } \beta\text{-Galactosidase (Penicillium)} \\ &= (1/W) \times \{(A_T - A_B)/0.459\} \times (1/10) \end{aligned}$$

0.459: Absorbance of 1  $\mu\text{mol}/10\text{ mL}$  of *o*-nitrophenol

$W$ : Amount (g) of the sample in 0.5 mL of the sample solution

Unit: One unit indicates an amount of the enzyme which decomposes 1  $\mu\text{mol}$  of 2-nitrophenyl- $\beta$ -D-galactopyranoside in 1 minute under the above conditions.

**Containers and storage** Containers—Tight containers.

## Gallium ( $^{67}\text{Ga}$ ) Citrate Injection

クエン酸ガリウム ( $^{67}\text{Ga}$ ) 注射液

Gallium ( $^{67}\text{Ga}$ ) Citrate Injection is an aqueous solution for injection containing gallium-67 ( $^{67}\text{Ga}$ ) in the form of gallium citrate.

It conforms to the requirements of Gallium ( $^{67}\text{Ga}$ ) Citrate Injection in the Minimum Requirements for Radiopharmaceuticals.

Test for Extractable volume of Parenteral Preparations and Insoluble Particulate Matter Test for Injections are not applied to this injection.

**Description** Gallium ( $^{67}\text{Ga}$ ) Citrate Injection is a clear, colorless or light red liquid.

## Gas Gangrene Antitoxin, Equine

ガスエソウマ抗毒素

Gas Gangrene Antitoxin, Equine, is a liquid for in-

jection containing *Clostridium perfringens* (*C. welchii*) Type A antitoxin, *Clostridium septicum* (*Vibrio septique*) antitoxin and *Clostridium oedematiens* (*C. novyi*) antitoxin in immunoglobulin of horse origin.

It may contain also *Clostridium histolyticum* antitoxin.

It conforms to the requirements of Gas Gangrene Antitoxin, Equine, in the Minimum Requirements for Biological Products.

**Description** Gas Gangrene Antitoxin, Equine, is a colorless to light yellow-brown, clear liquid or a slightly whitish turbid liquid.

## Gelatin

ゼラチン

Gelatin is a product prepared from aqueous extract of raw collagen by heating. The raw collagen is obtained by acid or alkali treatment of the bone, skin, ligament or tendon of animals.

**Description** Gelatin occurs as colorless or white to light yellow-brown sheets, shreds, granules or powder. It is odorless and tasteless.

Gelatin is very soluble in hot water, and practically insoluble in ethanol (95) and in diethyl ether.

Gelatin does not dissolve in water, but slowly swells and softens when immersed in it, gradually absorbing water 5 to 10 times its own mass.

Gelatin derived from an acid-treated collagen exhibits an isoelectric point between pH 7.0 and 9.0, and Gelatin derived from an alkali-treated collagen exhibits an isoelectric point between pH 4.5 and 5.0.

**Identification** (1) To 5 mL of a solution of Gelatin (1 in 100) add chromium (VI) oxide TS or 2,4,6-trinitrophenol TS dropwise: a precipitate is formed.

(2) To 5 mL of a solution of Gelatin (1 in 5000) add tannic acid TS dropwise: the solution becomes turbid.

**Purity** (1) Foreign odor and water-insoluble substances—Dissolve 1.0 g of Gelatin in 40 mL of water by heating: the solution has no disagreeable odor. It is clear, or only slightly opalescent. The solution has no more color than Matching Fluid A.

(2) Sulfite—Take 20.0 g of Gelatin in a round-bottomed flask, dissolve in 150 mL of hot water, and add 3 to 5 drops of silicone resin, 5 mL of phosphoric acid and 1 g of sodium hydrogen carbonate. Attach a condenser, immediately distill the solution, immersing the end of the condenser into a receiver containing 50 mL of iodine TS, and continue the distillation until 50 mL of distillate is obtained. Acidify the distillate with 2 to 3 drops of hydrochloric acid, add 2 mL of barium chloride TS, and heat on a water bath until the color of iodine TS is discharged. Collect the precipitates, wash with water, and ignite: the mass of the residue is not more than 4.5 mg, but the mass of the residue obtained from Gelatin for use in the preparation of capsules and tablets is not more than 75 mg. Perform a blank determination, and make any necessary correction.

(3) Heavy metals <1.07>—Proceed with 0.5 g of Gelatin

according to Method 2, and perform the test. Prepare the control solution with 2.5 mL of Standard Lead Solution (not more than 50 ppm).

(4) **Arsenic <1.11>**—Take 15.0 g of Gelatin in a flask, add 60 mL of diluted hydrochloric acid (1 in 5), and heat until solution is effected. Add 15 mL of bromine TS, heat until the excess of bromine is expelled, neutralize with ammonia TS, add 1.5 g of disodium hydrogen phosphate dodecahydrate, and allow to cool. To this solution add 30 mL of magnesia TS, allow to stand for 1 hour, and collect the precipitates. Wash the precipitates with five 10-mL portions of diluted ammonia TS (1 in 4), and dissolve in diluted hydrochloric acid (1 in 4) to make exactly 50 mL. Perform the test with 5 mL of this solution: the solution has no more color than the following standard stain.

Standard stain: Proceed with 15 mL of Standard Arsenic Solution, instead of Gelatin, in the same manner (not more than 1 ppm).

(5) **Mercury**—Place 2.0 g of Gelatin in a decomposition flask, add 20 mL of diluted sulfuric acid (1 in 2) and 100 mL of a solution of potassium permanganate (3 in 50), heat gently under a reflux condenser, and boil for 2 hours. If the solution becomes clear during boiling, reduce the temperature of the solution to about 60°C, add further 5 mL of a solution of potassium permanganate (3 in 50), boil again, and repeat the above-mentioned procedure until the precipitate of manganese dioxide remains for about 20 minutes. Cool, add a solution of hydroxylammonium chloride (1 in 5) until the precipitate of manganese dioxide disappears, add water to make exactly 150 mL, and use the solution as the sample solution. Perform the test as directed under Atomic Absorption Spectrophotometry <2.23> (Cold vapor type) using the sample solution. Place the sample solution in a sample water bottle of the atomic absorption spectrophotometer, add 10 mL of tin (II) chloride-sulfuric acid TS, connect the bottle immediately to the atomic absorption spectrophotometer, and circulate air. Determine the absorbance  $A_T$  of the sample solution at 253.7 nm when the indication of the recorder has risen rapidly and become constant. On the other hand, place 2.0 mL of Standard Mercury Solution in a decomposition flask, add 20 mL of diluted sulfuric acid (1 in 2) and 100 mL of a solution of potassium permanganate (3 in 50), and proceed in the same manner as for the sample solution. Determine the absorbance  $A_S$  of the standard solution:  $A_T$  is not more than  $A_S$  (not more than 0.1 ppm).

**Loss on drying** Not more than 15.0%. Take about 1 g of Gelatin, accurately weighed, in a tared 200-mL beaker containing 10 g of sea sand (No. 1) previously dried at 110°C for 3 hours. Add 20 mL of water, allow to stand for 30 minutes with occasional shaking, evaporate to dryness on a water bath with occasional shaking, and dry the residue at 110°C for 3 hours.

**Residue on ignition <2.44>** Not more than 2.0% (0.5 g).

**Containers and storage** Containers—Tight containers.

## Purified Gelatin

精製ゼラチン

Purified Gelatin is a product prepared from aqueous

extract of raw collagen by heating. The raw collagen is obtained by acid or alkali treatment of the bone, skin, ligament, or tendon of animals.

**Description** Purified Gelatin occurs as colorless to light yellow sheets, shreds, pellets or powder. It is odorless and tasteless.

It is very soluble in hot water, and practically insoluble in ethanol (95) and in diethyl ether.

Purified Gelatin does not dissolve in water. It slowly swells and softens when immersed in water, and absorbs water 5 to 10 times its own mass.

Purified Gelatin derived from an acid-treated collagen has an isoelectric point at pH 7.0 to 9.0, and Purified Gelatin derived from an alkali-treated collagen has an isoelectric point at pH 4.5 to 5.0.

**Identification (1)** To 5 mL of a solution of Purified Gelatin (1 in 100) add chromium (VI) oxide TS or 2,4,6-trinitrophenol TS dropwise: a precipitate is formed.

(2) To 5 mL of a solution of Purified Gelatin (1 in 5000) add tannic acid TS dropwise: the solution becomes turbid.

**Purity (1)** Foreign odor and water-insoluble substances—Dissolve 1.0 g of Purified Gelatin in 40 mL of water by heating: the solution is clear, colorless and free from any disagreeable odor when the layer of the solution is 20 mm in depth.

(2) **Sulfite**—Take 20.0 g of Purified Gelatin in a round-bottomed flask, dissolve in 150 mL of hot water, and add 3 to 5 drops of silicone resin, 5 mL of phosphoric acid and 1 g of sodium hydrogen carbonate. Attach a condenser, immediately distill the solution, immersing the end of the condenser into a receiver containing 50 mL of iodine TS, and continue the distillation until 50 mL of distillate is obtained. Acidify the distillate by dropwise addition of hydrochloric acid, add 2 mL of barium chloride TS, and heat on a water bath until the color of iodine TS is discharged. Collect the precipitates, wash with water, and ignite: the mass of the residue is not more than 1.5 mg. Perform a blank determination, and make any necessary correction.

(3) **Heavy metals <1.07>**—Proceed with 1.0 g of Purified Gelatin according to Method 2, and perform the test. Prepare the control solution with 2.0 mL of Standard Lead Solution (not more than 20 ppm).

(4) **Arsenic <1.11>**—Place 15.0 g of Purified Gelatin in a flask, add 60 mL of diluted hydrochloric acid (1 in 5), and heat until solution is effected. Add 15 mL of bromine TS, heat until the excess of bromine is expelled, neutralize with ammonia TS, add 1.5 g of disodium hydrogen phosphate dodecahydrate, and allow to cool. To this solution add 30 mL of magnesia TS, allow to stand for 1 hour, and collect the precipitates. Wash the precipitates with five 10-mL portions of diluted ammonia TS (1 in 4), and dissolve in diluted hydrochloric acid (1 in 4) to make exactly 50 mL. Perform the test with 5 mL of this solution: the solution has no more color than the following standard solution.

Standard solution: Proceed with 15 mL of Standard Arsenic Solution, instead of Purified Gelatin, in the same manner (not more than 1 ppm).

(5) **Mercury**—Place 2.0 g of Purified Gelatin in a decomposition flask, add 20 mL of diluted sulfuric acid (1 in 2) and 100 mL of a solution of potassium permanganate (3 in 50), heat gently under a reflux condenser, and boil for 2 hours. If

the solution becomes clear during boiling, reduce the temperature of the solution to about 60°C, add further 5 mL of a solution of potassium permanganate (3 in 50), boil again, and repeat the above-mentioned procedure until the precipitate of manganese dioxide remains for about 20 minutes. Cool, add a solution of hydroxylammonium chloride (1 in 5) until the precipitate of manganese dioxide disappears, add water to make exactly 150 mL, and use the solution as the sample solution. Perform the test as directed under Atomic Absorption Spectrophotometry <2.23> (Cold vapor type) using the sample solution. Place the sample solution in a sample water bottle of the atomic absorption spectrophotometer, add 10 mL of tin (II) chloride-sulfuric acid TS, connect the bottle immediately to the atomic absorption spectrophotometer, and circulate air. Determine the absorbance  $A_T$  of the sample solution at 253.7 nm when the indication of the recorder has risen rapidly and become constant. On the other hand, place 2.0 mL of Standard Mercury Solution in a decomposition flask, add 20 mL of diluted sulfuric acid (1 in 2) and 100 mL of a solution of potassium permanganate (3 in 50), and proceed in the same manner as for the sample solution. Determine the absorbance  $A_S$  of the standard solution:  $A_T$  is not more than  $A_S$  (not more than 0.1 ppm).


**Loss on drying** Not more than 15.0%. Take about 1 g of Purified Gelatin, accurately weighed, in a tared 200-mL beaker containing 10 g of sea sand (No. 1), previously dried at 110°C for 3 hours. Add 20 mL of water, allow to stand for 30 minutes with occasional shaking, evaporate on a water bath to dryness with occasional shaking, and dry the residue at 110°C for 3 hours.

**Residue on ignition** <2.44> Not more than 2.0% (0.5 g).

**Containers and storage** Containers—Tight containers.

## Gentamicin Sulfate

ゲンタマイシン硫酸塩



Gentamicin Sulfate  $C_1$  :  $R^1 = CH_3$   $R^2 = NHCH_3$

Gentamicin Sulfate  $C_2$  :  $R^1 = CH_3$   $R^2 = NH_2$

Gentamicin Sulfate  $C_{1a}$  :  $R^1 = H$   $R^2 = NH_2$

Gentamicin Sulfate  $C_1$ :

(6R)-2-Amino-2,3,4,6-tetradeoxy-6-methylamino-6-methyl- $\alpha$ -D-erythro-hexopyranosyl-(1 $\rightarrow$ 4)-[3-deoxy-4-C-methyl-3-methylamino- $\beta$ -L-arabinopyranosyl-(1 $\rightarrow$ 6)]-2-deoxy-D-streptamine sulfate

Gentamicin Sulfate  $C_2$ :

(6R)-2,6-Diamino-2,3,4,6-tetradeoxy-6-methyl- $\alpha$ -D-erythro-hexopyranosyl-(1 $\rightarrow$ 4)-[3-deoxy-4-C-methyl-3-methylamino- $\beta$ -L-arabinopyranosyl-(1 $\rightarrow$ 6)]-2-deoxy-D-streptamine sulfate

Gentamicin Sulfate  $C_{1a}$ :

2,6-Diamino-2,3,4,6-tetradeoxy- $\alpha$ -D-erythro-hexopyranosyl-(1 $\rightarrow$ 4)-[3-deoxy-4-C-methyl-3-methylamino- $\beta$ -L-arabinopyranosyl-(1 $\rightarrow$ 6)]-2-deoxy-D-streptamine sulfate

[1405-41-0, Gentamicin Sulfate]

Gentamicin Sulfate is the sulfate of a mixture of aminoglycoside substances having antibacterial activity produced by the growth of *Micromonospora purpurea* or *Micromonospora echinospora*.

It contains not less than 590  $\mu$ g (potency) and not more than 775  $\mu$ g (potency) per mg, calculated on the dried basis. The potency of Gentamicin Sulfate is expressed as mass (potency) of gentamicin  $C_1$  ( $C_{21}H_{43}N_5O_7$ : 477.60).

**Description** Gentamicin Sulfate occurs as a white to light yellowish white powder.

It is very soluble in water, and practically insoluble in ethanol (99.5).

It is hygroscopic.

**Identification (1)** Dissolve 50 mg of Gentamicin Sulfate in 1 mL of water, and add 2 drops of a solution of 1-naphthol in ethanol (95) (1 in 500). Gently superimpose this solution on 1 mL of sulfuric acid: a blue-purple color develops at the zone of contact.

**(2)** Dissolve 50 mg each of Gentamicin Sulfate and Gentamicin Sulfate Reference Standard in 10 mL of water, and use these solutions as the sample solution and standard solution. Perform the test with these solutions as directed under Thin-layer Chromatography <2.03>. Spot 20  $\mu$ L of the sample solution and standard solution on a plate of silica gel for thin-layer chromatography. Separately, shake a mixture of chloroform, ammonia solution (28) and methanol (2:1:1) in a separator, and allow the mixture to stand for more than 1 hour. To 20 mL of the lower layer so obtained add 0.5 mL of methanol, and use this as the developing solvent. Develop the plate with the developing solvent to a distance of about 17 cm in a developing container with a cover, having an opening of about 20 mm<sup>2</sup>, and without putting a filter paper in the container, and air-dry the plate. Allow the plate to stand in iodine vapors: three principal spots obtained from the sample solution are the same with the corresponding spots from the standard solution in color tone and the  $R_f$  value, respectively.

**(3)** Dissolve 50 mg of Gentamicin Sulfate in 5 mL of water, and add 0.5 mL of barium chloride TS: a white precipitate is formed.

**Optical rotation** <2.49>  $[\alpha]_D^{25}$ : +107 – +121° (0.25 g calculated on the dried basis, water, 25 mL, 100 mm).

**pH** <2.54> The pH of a solution obtained by dissolving 0.20 g of Gentamicin Sulfate in 5 mL of water is between 3.5 and 5.5.

**Content ratio of the active principle** Dissolve 50 mg of Gentamicin Sulfate in water to make 10 mL, and use this solution as the sample solution. Perform the test with this solution as directed under Thin-layer Chromatography <2.03>. Spot 20

$\mu\text{L}$  of the sample solution on a plate of silica gel for thin-layer chromatography. Separately, shake a mixture of chloroform, ammonia solution (28) and methanol (2:1:1) in a separator, and allow the mixture to stand for more than 1 hour. To 20 mL of the lower layer so obtained add 0.5 mL of methanol, and use this as the developing solvent. Develop the plate with the developing solvent to a distance of about 17 cm in a developing container with a cover, having an opening of about 20 mm<sup>2</sup>, without putting a filter paper in the container, and air-dry the plate. Allow the plate to stand in iodine vapor. Determine the integral absorbances,  $A_a$ ,  $A_b$  and  $A_c$ , of the colored spots of gentamicin  $C_1$  ( $R_f$  value: about 0.3), gentamicin  $C_2$  ( $R_f$  value: about 0.2) and gentamicin  $C_{1a}$  ( $R_f$  value: about 0.1), respectively, by a densitometer (wavelength: 450 nm) while covering the plate with a glass plate, and calculate these amounts by the following formulae: gentamicin  $C_1$  is between 25% and 55%, gentamicin  $C_2$  is between 25% and 50%, and gentamicin  $C_{1a}$  is between 5% and 30%.

$$\begin{aligned} \text{Amount (\%)} \text{ of gentamicin } C_1 \\ = \{A_a / (A_a + 1.35A_b + A_c)\} \times 100 \end{aligned}$$

$$\begin{aligned} \text{Amount (\%)} \text{ of gentamicin } C_2 \\ = \{1.35A_b / (A_a + 1.35A_b + A_c)\} \times 100 \end{aligned}$$

$$\begin{aligned} \text{Amount (\%)} \text{ of gentamicin } C_{1a} \\ = \{A_c / (A_a + 1.35A_b + A_c)\} \times 100 \end{aligned}$$

**Purity (1)** Clarity and color of solution—Dissolve 1.0 g of Gentamicin Sulfate in 10 mL of water: the solution is clear and colorless to pale yellow.

**(2)** Heavy metals <1.07>—Proceed with 2.0 g of Gentamicin Sulfate according to Method 4, and perform the test. Prepare the control solution with 2.0 mL of Standard Lead Solution (not more than 10 ppm).

**(3)** Related substances—Dissolve 50 mg of Gentamicin Sulfate in water to make 10 mL, and use this solution as the sample solution. Pipet 1 mL of the sample solution, add water to make exactly 50 mL, and use this solution as the standard solution. Perform the test with these solutions as directed under Thin-layer Chromatography <2.03>. Spot 20  $\mu\text{L}$  of the sample solution and standard solution on a plate of silica gel for thin-layer chromatography. Separately, shake a mixture of chloroform, ammonia solution (28) and methanol (2:1:1) in a separator, and allow the mixture to stand for more than 1 hour. To 20 mL of the lower layer so obtained add 0.5 mL of methanol, and use this as the developing solvent. Develop the plate with the developing solvent to a distance of about 17 cm in a developing container with a cover, having an opening of about 20 mm<sup>2</sup>, without putting a filter paper in the container, and air-dry the plate. Allow the plate to stand in iodine vapor, and compare the colored spots while covering with a glass plate: the spots other than the spots of gentamicin  $C_1$  ( $R_f$  value: about 0.3), gentamicin  $C_2$  ( $R_f$  value: about 0.2) and gentamicin  $C_{1a}$  ( $R_f$  value: about 0.1) obtained from the sample solution are not more intense than the spot of gentamicin  $C_2$  from the standard solution.

**Loss on drying** <2.41> Not more than 18.0% (0.15 g, reduced pressure not exceeding 0.67 kPa, 110°C, 3 hours). Handle the sample avoiding absorption of moisture.

**Residue on ignition** <2.44> Not more than 1.0% (1 g).

**Assay** Perform the test according to the Cylinder-plate

method as directed under Microbial Assay for Antibiotics <4.02> according to the following conditions.

(i) Test organism—*Staphylococcus epidermidis* ATCC 12228

(ii) Agar media for seed and base layer—

|                 |         |
|-----------------|---------|
| Glucose         | 1.0 g   |
| Peptone         | 6.0 g   |
| Meat extract    | 1.5 g   |
| Yeast extract   | 3.0 g   |
| Sodium chloride | 10.0 g  |
| Agar            | 15.0 g  |
| Water           | 1000 mL |

Mix all the ingredients, and sterilize. Adjust the pH of the solution so that it will be 7.8 to 8.0 after sterilization.

(iii) Agar medium for transferring test organisms—Use the medium ii in 2) Medium for other organisms under (2) Agar media for transferring test organisms.


(iv) Standard solutions—Weigh accurately an amount of Gentamicin Sulfate Reference Standard, equivalent to about 25 mg (potency), dissolve in 0.1 mol/L phosphate buffer solution, pH 8.0 to make exactly 25 mL, and use this solution as the standard stock solution. Keep the standard stock solution at 15°C or lower, and use within 30 days. Take exactly a suitable amount of the standard stock solution before use, add 0.1 mol/L phosphate buffer solution, pH 8.0 to make solutions so that each mL contains 4  $\mu\text{g}$  (potency) and 1  $\mu\text{g}$  (potency), and use these solutions as the high concentration standard solution and low concentration standard solution, respectively.

(v) Sample solutions—Weigh accurately an amount of Gentamicin Sulfate, equivalent to about 25 mg (potency), and dissolve in 0.1 mol/L phosphate buffer solution, pH 8.0 to make exactly 25 mL. Take exactly a suitable amount of this solution, add 0.1 mol/L phosphate buffer solution, pH 8.0 to make solutions so that each mL contains 4  $\mu\text{g}$  (potency) and 1  $\mu\text{g}$  (potency), and use these solutions as the high concentration sample solution and low concentration sample solution, respectively.

**Containers and storage** Containers—Tight containers.

## Glibenclamide

グリベンクラミド



$\text{C}_{23}\text{H}_{28}\text{ClN}_3\text{O}_5\text{S}$ : 494.00

4-[2-(5-Chloro-2-methoxybenzoylamino)ethyl]-  
N-(cyclohexylcarbamoyl)benzenesulfonamide [10238-21-8]

Glibenclamide, when dried, contains not less than 98.5% of  $\text{C}_{23}\text{H}_{28}\text{ClN}_3\text{O}_5\text{S}$ .

**Description** Glibenclamide occurs as white to pale yellowish white crystals or crystalline powder.

It is freely soluble in dimethylformamide, sparingly soluble

in chloroform, slightly soluble in methanol and in ethanol (95), and practically insoluble in water.

**Identification (1)** Determine the absorption spectrum of a solution of Glibenclamide in methanol (1 in 10,000) as directed under Ultraviolet-visible Spectrophotometry <2.24>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wavelengths.

**(2)** Determine the infrared absorption spectrum of Glibenclamide, previously dried, as directed in the potassium bromide disk method under Infrared Spectrophotometry <2.25>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wave numbers.

**(3)** Perform the test with Glibenclamide as directed under Flame Coloration Test <1.04> (2): a green color appears.

**Melting point** <2.60> 169 – 174°C

**Purity (1)** Heavy metals <1.07>—Proceed with 1.0 g of Glibenclamide according to Method 2, and perform the test. Prepare the control solution with 2.0 mL of Standard Lead Solution (not more than 20 ppm).

**(2)** Related substances—Dissolve 0.20 g of Glibenclamide in 20 mL of chloroform, and use this solution as the sample solution. Pipet 1 mL of the sample solution, and add chloroform to make exactly 20 mL. Pipet 1 mL of this solution, add chloroform to make exactly 10 mL, and use this solution as the standard solution. Perform the test with these solutions as directed under Thin-layer Chromatography <2.03>. Spot 10  $\mu$ L each of the sample solution and standard solution on a plate of silica gel with fluorescent indicator for thin-layer chromatography. Develop the plate with a mixture of 1-propanol, chloroform and diluted ammonia TS (4 in 5) (11:7:2) to a distance of about 12 cm, and air-dry the plate. Examine under ultraviolet light (main wavelength: 254 nm): the spots other than the principal spot from the sample solution are not more intense than the spot from the standard solution.

**Loss on drying** <2.41> Not more than 0.5% (1 g, 105°C, 4 hours).


**Assay** Weigh accurately about 0.9 g of Glibenclamide, previously dried, dissolve in 50 mL of *N,N*-dimethylformamide, and titrate <2.50> with 0.1 mol/L sodium hydroxide VS (indicator: 3 drops of phenolphthalein TS). Perform a blank determination with a solution prepared by adding 18 mL of water to 50 mL of *N,N*-dimethylformamide, and make any necessary correction.

Each mL of 0.1 mol/L sodium hydroxide VS  
= 49.40 mg of  $C_{23}H_{28}ClN_3O_5S$

**Containers and storage** Containers—Tight containers.

## Glucose

ブドウ糖



$\alpha$ -D-Glucopyranose :  $R^1=H, R^2=OH$   
 $\beta$ -D-Glucopyranose :  $R^1=OH, R^2=H$

$C_6H_{12}O_6$ : 180.16

D-Glucopyranose [50-99-7]

Glucose is  $\alpha$ -D-glucopyranose,  $\beta$ -D-glucopyranose, or a mixture of them, and when dried, it contains not less than 99.5% of  $C_6H_{12}O_6$ .

**Description** Glucose occurs as white crystals or crystalline powder. It is odorless, and has a sweet taste.

It is freely soluble in water, slightly soluble in ethanol (95), and practically insoluble in diethyl ether.

**Identification** Add 2 to 3 drops of a solution of Glucose (1 in 20) to 5 mL of boiling Fehling's TS: a red precipitate is produced.

**Purity (1)** Clarity and color of solution—Add 25 g of Glucose to 30 mL of water in a Nessler tube, warm at 60°C in a water bath until solution is effected, cool, and add water to make 50 mL: the solution is clear and has no more color than the following control solution.

Control solution: To a mixture of 1.0 mL of Cobaltous Chloride Stock CS, 3.0 mL of Ferric Chloride stock CS, and 2.0 mL of Cupric Sulfate Stock CS, add water to make 10.0 mL. To 3.0 mL of this solution add water to make 50 mL.

**(2)** Acidity—Dissolve 5.0 g of Glucose in 50 mL of freshly boiled and cooled water, and add 3 drops of phenolphthalein TS and 0.60 mL of 0.01 mol/L sodium hydroxide VS: a red color develops.

**(3)** Chloride <1.03>—Perform the test with 2.0 g of Glucose. Prepare the control solution with 1.0 mL of 0.01 mol/L hydrochloric acid VS (not more than 0.018%).

**(4)** Sulfate <1.14>—Perform the test with 2.0 g of Glucose. Prepare the control solution with 1.0 mL of 0.005 mol/L sulfuric acid VS (not more than 0.024%).

**(5)** Heavy metals <1.07>—Proceed with 5.0 g of Glucose according to Method 2, and perform the test. Prepare the control solution with 2.0 mL of Standard Lead Solution (not more than 4 ppm).

**(6)** Arsenic <1.11>—Dissolve 1.5 g of Glucose in 5 mL of water, add 5 mL of dilute sulfuric acid and 1 mL of bromine TS, heat on a water bath for 5 minutes, and concentrate to 5 mL. After cooling, perform the test with this solution as the test solution (not more than 1.3 ppm).

**(7)** Dextrin—To 1.0 g of Glucose add 20 mL of ethanol (95), and boil under a reflux condenser: the solution is clear.

**(8)** Soluble starch and sulfite—Dissolve 1.0 g of Glucose in 10 mL of water, and add 1 drop of iodine TS: a yellow color develops.

**Loss on drying** <2.41> Not more than 1.0% (1 g, 105°C,

6 hours).

**Residue on ignition** <2.44> Not more than 0.1% (2 g).

**Assay** Weigh accurately about 10 g of Glucose, previously dried, dissolve in 0.2 mL of ammonia TS and water to make exactly 100 mL, allow to stand for 30 minutes, and determine the optical rotation,  $\alpha_D$ , of this solution at  $20 \pm 1^\circ\text{C}$  in a 100-mm cell as directed under Optical Rotation Determination <2.49>.

$$\text{Amount (mg) of } \text{C}_6\text{H}_{12}\text{O}_6 = \alpha_D \times 1895.4$$

**Containers and storage** Containers—Tight containers.

## Glucose Injection

ブドウ糖注射液

Glucose Injection is an aqueous solution for injection. It contains not less than 95% and not more than 105% of the labeled amount of glucose ( $\text{C}_6\text{H}_{12}\text{O}_6$ ; 180.16).

**Method of preparation** Prepare as directed under Injections, with Glucose. No preservative is added.

**Description** Glucose Injection is a clear, colorless liquid. It has a sweet taste. It occurs as a colorless to pale yellow, clear liquid when its labeled concentration exceeds 40%.

**Identification** Measure a volume of Glucose Injection, equivalent to 0.1 g of Glucose according to the labeled amount, and, if necessary, add water or evaporate on a water bath to a volume of 2 mL. Add 2 to 3 drops of the solution to 5 mL of boiling Fehling's TS: a red precipitate is produced.

**pH** <2.54> 3.5 – 6.5 In the case where the labeled concentration of the injection exceeds 5%, dilute to 5% with water before the test.

**Purity** 5-Hydroxymethylfurfural and related substances—Measure exactly a volume of Glucose Injection, equivalent to 2.5 g of Glucose according to the labeled amount, and add water to make exactly 100 mL. Determine the absorbance of this solution at 284 nm as directed under Ultraviolet-visible Spectrophotometry <2.24>: it is not more than 0.80.

**Bacterial endotoxins** <4.01> Less than 0.50 EU/mL.

**Extractable volume** <6.05> It meets the requirement.


**Assay** Measure accurately a volume of Glucose Injection, equivalent to about 4 g of glucose ( $\text{C}_6\text{H}_{12}\text{O}_6$ ), and add 0.2 mL of ammonia TS and water to make exactly 100 mL. Shake the solution well, allow to stand for 30 minutes, and determine the optical rotation,  $\alpha_D$ , at  $20 \pm 1^\circ\text{C}$  in a 100-mm cell as directed under Optical Rotation Determination <2.49>.

$$\text{Amount (mg) of glucose } (\text{C}_6\text{H}_{12}\text{O}_6) = \alpha_D \times 1895.4$$

**Containers and storage** Containers—Hermetic containers. Plastic containers for aqueous injections may be used.

## Glutathione

グルタチオン



$\text{C}_{10}\text{H}_{17}\text{N}_3\text{O}_6\text{S}$ ; 307.32

(2S)-2-Amino-4-[1-(carboxymethyl)carbamoyl-(2R)-2-sulfanylethylcarbamoyl]butanoic acid [70-18-8]

Glutathione, when dried, contains not less than 98.0% and not more than 101.0% of  $\text{C}_{10}\text{H}_{17}\text{N}_3\text{O}_6\text{S}$ .

**Description** Glutathione occurs as a white crystalline powder.

It is freely soluble in water, and practically insoluble in ethanol (99.5).

Melting point: about  $185^\circ\text{C}$  (with decomposition).

**Identification** Determine the infrared absorption spectrum of Glutathione, previously dried, as directed in the potassium bromide disk method under Infrared Spectrophotometry <2.25>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wave numbers.

**Optical rotation** <2.49>  $[\alpha]_D^{20}$ :  $-15.5 - -17.5^\circ$  (after drying, 2 g, water, 50 mL, 100 mm).

**Purity** (1) Clarity and color of solution—Dissolve 1.0 g of Glutathione in 10 mL of water: the solution is clear and colorless.

(2) Heavy metals <1.07>—Proceed with 2.0 g of Glutathione according to Method 2, and perform the test. Prepare the control solution with 2.0 mL of Standard Lead Solution (not more than 10 ppm).

(3) Arsenic <1.11>—Prepare the test solution with 1.0 g of Glutathione according to Method 1, and perform the test (not more than 2 ppm).

(4) Related substances—Dissolve 50 mg of Glutathione in 100 mL of the mobile phase, and use this solution as the sample solution. Pipet 2 mL of the sample solution, add the mobile phase to make exactly 100 mL, and use this solution as the standard solution. Perform the test with exactly  $10 \mu\text{L}$  each of the sample solution and standard solution as directed under Liquid Chromatography <2.01> according to the following conditions, and determine each peak area by the automatic integration method: the area of the peak having the relative retention time of about 4 with respect to glutathione is not more than 3/4 times the peak area of glutathione from the standard solution, and the total area of the peaks other than the peak of glutathione is not more than the peak area of glutathione from the standard solution.

**Operating conditions**—

**Detector:** An ultraviolet absorption photometer (wavelength: 210 nm).

**Column:** A stainless steel column 4.6 mm in inside diameter and 15 cm in length, packed with octadecylsilanized silica gel for liquid chromatography ( $5 \mu\text{m}$  in particle diameter).

Column temperature: A constant temperature of about 30°C.

Mobile phase: Dissolve 6.8 g of potassium dihydrogen phosphate and 2.02 g of sodium 1-heptane sulfonate in 1000 mL of water, and adjust the pH to 3.0 with phosphoric acid. To 970 mL of this solution add 30 mL of methanol.

Flow rate: Adjust the flow rate so that the retention time of glutathione is about 5 minutes.

Time span of measurement: About 6 times as long as the retention time of glutathione beginning after the solvent peak.

#### System suitability—

Test for required detectability: Pipet 10 mL of the standard solution, and add the mobile phase to make exactly 100 mL. Confirm that the peak area of glutathione obtained from 10  $\mu$ L of this solution is equivalent to 8 to 12% of that obtained from 10  $\mu$ L of the standard solution.

System performance: Dissolve 50 mg of glutathione, 10 mg of D-phenylglycine and 50 mg of ascorbic acid in 100 mL of water. When the procedure is run with 10  $\mu$ L of this solution under the above operating conditions, ascorbic acid, glutathione and D-phenylglycine are eluted in this order, and the resolutions between the peaks of ascorbic acid and glutathione and between the peaks of glutathione and D-phenylglycine are not less than 5, respectively.

System repeatability: When the test is repeated 6 times with 10  $\mu$ L of the standard solution under the above operating conditions, the relative standard deviation of the peak area of glutathione is not more than 1.5%.

**Loss on drying** <2.41> Not more than 0.5% (1 g, 105°C, 3 hours).

**Residue on ignition** <2.44> Not more than 0.1% (1 g).

**Assay** Weigh accurately about 0.5 g of Glutathione, previously dried, dissolve in 50 mL of a solution of metaphosphoric acid (1 in 50), and titrate <2.50> with 0.05 mol/L iodine VS (indicator: 1 mL of starch TS). Perform a blank determination in the same manner, and make any necessary correction.

Each mL of 0.05 mol/L iodine VS  
= 30.73 mg of  $C_{10}H_{17}N_3O_6S$

**Containers and storage** Containers—Tight containers.

## Glycerin

### Glycerol

グリセリン

$C_3H_8O_3$ : 92.09

Glycerin contains not less than 84.0% and not more than 87.0% of  $C_3H_8O_3$ .

**Description** Glycerin is a clear, colorless, viscous liquid. It has a sweet taste.

It is miscible with water and with ethanol (99.5).

It is hygroscopic.

**Identification** Determine the infrared absorption spectrum

of Glycerin as directed in the liquid film method under Infrared Spectrophotometry <2.25>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wave numbers.

**Refractive index** <2.45>  $n_D^{20}$ : 1.449 – 1.454

**Specific gravity** <2.56>  $d_{20}^{20}$ : 1.221 – 1.230

**Purity** (1) Color—Place 50 mL of Glycerin in a Nessler tube, and observe downward: the solution has no more color than the following control solution. Control solution: Place 0.40 mL of Ferric Chloride Colorimetric Stock Solution in a Nessler tube, and add water to make 50 mL.

(2) Acidity or alkalinity—To 2 mL of Glycerin add 8 mL of water and mix: the solution is neutral.

(3) Chloride <1.03>—Take 10.0 g of Glycerin, and perform the test: Prepare the control solution with 0.30 mL of 0.01 mol/L hydrochloric acid VS (not more than 0.001%).

(4) Sulfate <1.14>—Take 10.0 g of Glycerin, and perform the test. Prepare the control solution with 0.40 mL of 0.005 mol/L sulfuric acid VS (not more than 0.002%).

(5) Ammonium—To 5 mL of Glycerin add 5 mL of a solution of sodium hydroxide (1 in 10), and boil: the gas evolved does not change moistened red litmus paper to blue.

(6) Heavy metals <1.07>—Proceed with 5.0 g of Glycerin according to Method 1, and perform the test: Prepare the control solution with 2.5 mL of Standard Lead Solution (not more than 5 ppm).

(7) Calcium—To 5 mL of the solution obtained in (2) add 3 drops of ammonium oxalate TS: the solution remains unchanged.

(8) Arsenic <1.11>—Prepare the test solution with 1.0 g of Glycerin according to Method 1, and perform the test (not more than 2 ppm).

(9) Acrolein, glucose, and other reducing substances—To 1.0 g of Glycerin add 1 mL of ammonia TS, mix, and warm in a water bath at 60°C for 5 minutes: no yellow color is produced. Take the solution out of the water bath, add 3 drops of silver nitrate TS immediately, and allow to stand in a dark place for 5 minutes: the color of the solution does not change, and no turbidity is produced.

(10) Fatty acids and esters—Mix 50 g of Glycerin with 50 mL of freshly boiled and cooled water, add exactly 10 mL of 0.1 mol/L sodium hydroxide VS, boil the mixture for 15 minutes, cool, and titrate the excess sodium hydroxide with 0.1 mol/L hydrochloric acid VS: 0.1 mol/L sodium hydroxide VS consumed is not more than 3.0 mL (indicator: 3 drops of phenolphthalein TS). Perform a blank determination.

(11) Readily carbonizable substances—To 5 mL of Glycerin add carefully 5 mL of sulfuric acid for readily carbonizable substances, mix gently at a temperature between 18°C and 20°C, and allow to stand for 1 hour between 15°C and 25°C: the solution has not more color than Matching Fluid H.

**Water** <2.48> 13 – 17% (0.1 g, volumetric titration, direct titration).

**Residue on ignition** <2.44> Weigh accurately about 10 g of Glycerin in a tared crucible, heat to boiling, and fire to burn immediately. After cooling, moisten the residue with 1 to 2 drops of sulfuric acid, and ignite cautiously to constant mass: the mass of the residue is not more than 0.01%.

**Assay** Weigh accurately about 0.2 g of Glycerin, transfer into a glass-stoppered flask, add 50 mL of water, mix, add exactly 50 mL of sodium periodate TS, shake, and allow to stand in a dark place at a room temperature for about 30 minutes. Add 10 mL of a mixture of water and ethylene glycol (1:1), allow to stand for about 20 minutes, add 100 mL of water, and titrate <2.50> with 0.1 mol/L sodium hydroxide VS (indicator: 2 drops of phenolphthalein TS). Perform a blank determination, and make the necessary correction.


$$\begin{aligned} \text{Each mL of 0.1 mol/L sodium hydroxide VS} \\ = 9.209 \text{ mg of C}_3\text{H}_8\text{O}_3 \end{aligned}$$

**Containers and storage** Containers—Tight containers.

## Concentrated Glycerin

### Concentrated Glycerol

濃グリセリン



C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>: 92.09

Propane-1,2,3-triol [56-81-5]

Concentrated Glycerin contains not less than 98.0% and not more than 101.0% of glycerin (C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>), calculated of the anhydrous basis.

**Description** Concentrated Glycerin is a clear, colorless and viscous liquid. It has a sweet taste.

It is miscible with water and with ethanol (99.5).

It is hygroscopic.

**Identification** Determine the infrared absorption spectrum of Concentrated Glycerin as directed in the liquid film method under Infrared Spectrophotometry <2.25>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wave numbers.

**Refractive index** <2.45>  $n_D^{20}$ : Not less than 1.470.

**Specific gravity** <2.56>  $d_{20}^{20}$ : Not less than 1.258.

**Purity** (1) Color—Place 50 mL of Concentrated Glycerin in a Nessler tube, and observe downward: the solution has no more color than the following control solution.

Control solution: Pipet 0.40 mL of Ferric Chloride Colorimetric Stock Solution into a Nessler tube, and add water to make 50 mL.

(2) Acidity or alkalinity—To 2 mL of Concentrated Glycerin add 8 mL of water and mix: the solution is neutral.

(3) Chloride <1.03>—Take 10.0 g of Concentrated Glycerin, and perform the test. Prepare the control solution with 0.30 mL of 0.01 mol/L hydrochloric acid VS (not more than 0.001%).

(4) Sulfate <1.14>—Take 10.0 g of Concentrated Glycerin, and perform the test. Prepare the control solution with 0.40 mL of 0.005 mol/L sulfuric acid VS (not more than 0.002%).

(5) Ammonium—To 5 mL of Concentrated Glycerin add 5 mL of a solution of sodium hydroxide (1 in 10), and boil:

the gas evolved does not change moistened red litmus paper to blue.

(6) Heavy metals <1.07>—Proceed with 5.0 g of Concentrated Glycerin according to Method 1, and perform the test. Prepare the control solution with 2.5 mL of Standard Lead Solution (not more than 5 ppm).

(7) Calcium—To 5 mL of the solution obtained in (2) add 3 drops of ammonium oxalate TS: the solution remains unchanged.

(8) Arsenic <1.11>—Prepare the test solution with 1.0 g of Concentrated Glycerin according to Method 1, and perform the test (not more than 2 ppm).

(9) Acrolein, glucose, or other reducing substances—To 1.0 g of Concentrated Glycerin add 1 mL of ammonia TS, mix, and warm in a water bath at 60°C for 5 minutes: no yellow color is produced. Take the solution out of the water bath, add 3 drops of silver nitrate TS immediately, and allow to stand in a dark place for 5 minutes: the color of the solution does not change, and no turbidity is produced.

(10) Fatty acids and esters—Mix 50 g of Concentrated Glycerin with 50 mL of freshly boiled and cooled water, add 10 mL of 0.1 mol/L sodium hydroxide VS, accurately measured, boil the mixture for 15 minutes, cool, and titrate the excess sodium hydroxide with 0.1 mol/L hydrochloric acid VS: not more than 3.0 mL of 0.1 mol/L sodium hydroxide VS is consumed (indicator: 3 drops of phenolphthalein TS). Perform a blank determination.

(11) Readily carbonizable substances—To 5 mL of Concentrated Glycerin add carefully 5 mL of sulfuric acid for readily carbonizable substances, mix gently at a temperature between 18°C and 20°C, and allow to stand for 1 hour between 15°C and 25°C: the solution has no more color than Matching Fluid H.

**Water** <2.48> Not more than 2.0% (6 g, volumetric titration, direct titration).

**Residue on ignition** <2.44> Weigh accurately about 10 g of Concentrated Glycerin in a tared crucible, heat to boiling, and fire to burn immediately. Cool, moisten the residue with 1 to 2 drops of sulfuric acid, and ignite cautiously to constant mass: the mass of the residue is not more than 0.01%.

**Assay** Weigh accurately about 0.2 g of Concentrated Glycerin, transfer into a glass-stoppered flask, add 50 mL of water, mix, add exactly 50 mL of sodium periodate TS, shake, and allow to stand in a dark place at a room temperature for about 30 minutes. Add 10 mL of a mixture of water and ethylene glycol (1:1), allow to stand for about 20 minutes, add 100 mL of water, and titrate <2.50> with 0.1 mol/L sodium hydroxide VS (indicator: 2 drops of phenolphthalein TS). Perform a blank determination, and make the necessary correction.

$$\begin{aligned} \text{Each mL of 0.1 mol/L sodium hydroxide VS} \\ = 9.209 \text{ mg of C}_3\text{H}_8\text{O}_3 \end{aligned}$$

**Containers and storage** Containers—Tight containers.



## Glycerin and Potash Solution

グリセリンカリ液

### Method of preparation

|                         |                       |
|-------------------------|-----------------------|
| Potassium Hydroxide     | 3 g                   |
| Glycerin                | 200 mL                |
| Ethanol                 | 250 mL                |
| Aromatic substance      | a suitable quantity   |
| Water or Purified Water | a sufficient quantity |
| To make 1000 mL         |                       |

Dissolve Potassium Hydroxide in a portion of Water or Purified Water, add Glycerin, Ethanol, a suitable quantity of aromatic substance and another portion of Water or Purified Water to volume, and filter. Concentrated Glycerin may be used in place of Glycerin.

**Description** Glycerin and Potash Solution is a clear, colorless liquid, having an aromatic odor.

The pH of a solution of Glycerin and Potash Solution (1 in 5) is about 12.

Specific gravity  $d_{20}^{20}$ : about 1.02

**Identification (1)** A solution of Glycerin and Potash Solution (1 in 2) is alkaline (potassium hydroxide).

(2) Place 10 mL of a solution of Glycerin and Potash Solution (1 in 10) in a glass-stoppered test tube, add 2 mL of sodium hydroxide TS and 1 mL of copper (II) sulfate TS, and shake: a blue color is produced (glycerin).

(3) Glycerin and Potash Solution responds to the Qualitative Tests <1.09> for potassium salt.

**Containers and storage** Containers—Tight containers.

## Glyceryl Monostearate

モノステアリン酸グリセリン

Glyceryl Monostearate is a mixture of  $\alpha$ - and  $\beta$ -glyceryl monostearate and other fatty acid esters of glycerin.

**Description** Glyceryl Monostearate occurs as white to light yellow, waxy masses, thin flakes, or granules. It has a characteristic odor and taste.

It is very soluble in hot ethanol (95), soluble in chloroform, sparingly soluble in diethyl ether, and practically insoluble in water and in ethanol (95).

It is slowly affected by light.

**Identification (1)** Heat 0.2 g of Glyceryl Monostearate with 0.5 g of potassium hydrogen sulfate until thoroughly charred: the irritative odor of acrolein is perceptible.

(2) Dissolve 0.1 g of Glyceryl Monostearate in 2 mL of ethanol (95) by warming, heat with 5 mL of dilute sulfuric acid in a water bath for 30 minutes, and cool: a white to yellow solid is produced. This separated solid dissolves when shaken with 3 mL of diethyl ether.

**Melting point** <1.13> Not below 55°C.

**Acid value** <1.13> Not more than 15.

**Saponification value** <1.13> 157 – 170

**Iodine value** <1.13> Not more than 3.0. Use chloroform instead of cyclohexane.

**Purity** <1.13> Acidity or alkalinity—To 1.0 g of Glyceryl Monostearate add 20 mL of boiling water, and cool with swirling: the solution is neutral.

**Residue on ignition** <2.44> Not more than 0.1% (1 g).

**Containers and storage** Containers—Tight containers. Storage—Light-resistant.

## Glycine

### Aminoacetic Acid

グリシン



$\text{C}_2\text{H}_5\text{NO}_2$ : 75.07

Aminoacetic acid [56-40-6]

Glycine, when dried, contains not less than 98.5% of  $\text{C}_2\text{H}_5\text{NO}_2$ .

**Description** Glycine occurs as white crystals or crystalline powder. It is odorless. It has a sweet taste.

It is freely soluble in water and in formic acid, and practically insoluble in ethanol (95).

**Identification** Determine the infrared absorption spectrum of Glycine, previously dried, as directed in the potassium bromide disk method under Infrared Spectrophotometry <2.25>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wave numbers. If any difference appears between the spectra, dissolve Glycine in water, evaporate the water to dryness, and repeat the test with the residue.

**pH** <2.54> Dissolve 1.0 g of Glycine in 20 mL of water: the pH of the solution is between 5.6 and 6.6.

**Purity (1)** Clarity and color of solution—Dissolve 1.0 g of Glycine in 10 mL of water: the solution is clear and colorless.

(2) Chloride <1.03>—Perform the test with 0.5 g of Glycine. Prepare the control solution with 0.30 mL of 0.01 mol/L hydrochloric acid VS (not more than 0.021%).

(3) Sulfate <1.14>—Perform the test with 0.6 g of Glycine. Prepare the control solution with 0.35 mL of 0.005 mol/L sulfuric acid VS (not more than 0.028%).

(4) Ammonium<1.02>—Perform the test using 0.25 g of Glycine. Prepare the control solution with 5.0 mL of Standard Ammonium Solution (not more than 0.02%).

(5) Heavy metals <1.07>—Proceed with 1.0 g of Glycine according to Method 1, and perform the test. Prepare the control solution with 2.0 mL of Standard Lead Solution (not more than 20 ppm).

(6) Arsenic <1.11>—Prepare the test solution with 1.0 g of Glycine according to Method 1, and perform the test (not more than 2 ppm).

(7) Related substances—Dissolve 0.10 g of Glycine in 25

mL of water and use this solution as the sample solution. Pipet 1 mL of the sample solution, add water to make exactly 50 mL. Pipet 5 mL of this solution, add water to make exactly 20 mL, and use this solution as the standard solution. Perform the test with these solutions as directed under Thin-layer Chromatography <2.03>. Spot 5  $\mu$ L each of the sample solution and standard solution on a plate of silica gel for thin-layer chromatography. Develop the plate with a mixture of 1-butanol, water and acetic acid (100) (3:1:1) to a distance of about 10 cm, and dry the plate at 80°C for 30 minutes. Spray evenly a solution of ninhydrin in acetone (1 in 50), and heat at 80°C for 5 minutes: the spots other than the principal spot from the sample solution are not more intense than the spot from the standard solution.

**Loss on drying** <2.41> Not more than 0.30% (1 g, 105°C, 3 hours).

**Residue on ignition** <2.44> Not more than 0.1% (1 g).


**Assay** Weigh accurately about 80 mg of Glycine, previously dried, dissolve in 3 mL of formic acid, add 50 mL of acetic acid (100), and titrate <2.50> with 0.1 mol/L perchloric acid VS (potentiometric titration). Perform a blank determination, and make any necessary correction.

Each mL of 0.1 mol/L perchloric acid VS  
= 7.507 mg of  $C_{55}H_{75}N_{17}O_{13} \cdot 2C_2H_4O_2$

**Containers and storage** Containers—Well-closed containers.

## Gonadorelin Acetate

ゴナドレリン酢酸塩



$C_{55}H_{75}N_{17}O_{13} \cdot 2C_2H_4O_2$ : 1302.39

5-Oxo-L-prolyl-L-histidyl-L-tryptophyl-L-seryl-L-tyrosyl-glycyl-L-leucyl-L-arginyl-L-prolyl-glycinamide diacetate [34973-08-5]

Gonadorelin Acetate contains not less than 96.0% and not more than 102.0% of  $C_{55}H_{75}N_{17}O_{13} \cdot 2C_2H_4O_2$ , calculated on the anhydrous basis.

**Description** Gonadorelin Acetate occurs as a white to pale yellow powder. It is odorless or has a slight, acetic odor.

It is freely soluble in water, in methanol and in acetic acid (100), and sparingly soluble in ethanol (95).

It is hygroscopic.

**Identification** (1) Determine the absorption spectrum of a solution of Gonadorelin Acetate in methanol (1 in 10,000) as directed under Ultraviolet-visible Spectrophotometry <2.24>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wavelengths.

(2) Determine the infrared absorption spectrum of Gonadorelin Acetate as directed in the potassium bromide disk method under Infrared Spectrophotometry <2.25>, and compare the spectrum with the Reference Spectrum or the

spectrum of Gonadorelin Acetate Reference Standard: both spectra exhibit similar intensities of absorption at the same wave numbers.

(3) Dissolve 20 mg of Gonadorelin Acetate in 0.5 mL of ethanol (99.5), add 1 mL of sulfuric acid, and heat: the odor of ethyl acetate is perceptible.

**Optical rotation** <2.49>:  $[\alpha]_D^{20}$ :  $-53.0 - -57.0^\circ$  (0.10 g calculated on the anhydrous basis, diluted acetic acid (100) (1 in 100), 10 mL, 100 mm).

**pH** <2.54> Dissolve 0.10 g of Gonadorelin Acetate in 10 mL of water: the pH of this solution is between 4.8 and 5.8.

**Constituent amino acids** Put 10 mg of Gonadorelin Acetate in a test tube for hydrolysis, add 0.5 mL of hydrochloric acid and 0.5 mL of a solution of mercaptoacetic acid (2 in 25), seal the tube under reduced pressure, and heat at 110°C for 5 hours. After cooling, open the tube, transfer the hydrolyzate into a beaker, and evaporate to dryness on a water bath. Add exactly 100 mL of 0.02 mol/L hydrochloric acid TS to dissolve the residue, and use this solution as the sample solution. Separately, weigh exactly 0.105 g of L-serine, 0.147 g of L-glutamic acid, 0.115 g of L-proline, 75 mg of glycine, 0.131 g of L-leucine, 0.181 g of L-tyrosine, 0.210 g of L-histidine hydrochloride monohydrate, 0.204 g of L-tryptophan and 0.211 g of L-arginine hydrochloride, which are all previously dried at 105°C for 3 hours, add 50 mL of 1 mol/L hydrochloric acid TS to dissolve them, and add water to make exactly 1000 mL. Pipet 10 mL of this solution, add 0.02 mol/L hydrochloric acid TS to make exactly 200 mL, and use this solution as the standard solution. Perform the test with 50  $\mu$ L each of the sample solution and standard solution as directed under Liquid Chromatography <2.01> according to the following conditions: the peaks of nine constituent amino acids are observed on the chromatogram, and their respective molar ratios with respect to arginine are 0.7 – 1.0 for serine and tryptophan, 0.8 – 1.2 for proline, 0.9 – 1.1 for glutamic acid, leucine, tyrosine and histidine, respectively, and 1.8 – 2.2 for glycine.

**Operating conditions**—

**Detector:** A visible spectrophotometer (wavelength: 440 nm for proline and 570 nm for others).

**Column:** A stainless steel column 4 mm in inside diameter and 8 cm in length, packed with strongly acidic ion-exchange resin for liquid chromatography composed with a sulfonated polystyrene copolymer (5  $\mu$ m in particle diameter).

**Column temperature:** A constant temperature of about 57°C.

**Chemical reaction bath temperature:** A constant temperature of about 130°C.

**Mobile phase:** Prepare the mobile phases A, B, C and D according to the following table.

| Mobile phase                                     | A                   | B                   | C                   | D                   |
|--|---------------------|---------------------|---------------------|---------------------|
| Trisodium citrate dihydrate                      | 6.19 g              | 7.74 g              | 26.67 g             | —                   |
| Sodium hydroxide                                 | —                   | —                   | —                   | 8.00 g              |
| Sodium chloride                                  | 5.66 g              | 7.07 g              | 54.35 g             | —                   |
| Citric acid monohydrate                          | 19.80 g             | 22.00 g             | 6.10 g              | —                   |
| Ethanol (99.5)                                   | 130 mL              | 20 mL               | —                   | 100 mL              |
| Benzyl alcohol                                   | —                   | —                   | 5 mL                | —                   |
| Thiodiglycol                                     | 5 mL                | 5 mL                | —                   | —                   |
| Lauromacrogol solution in diethyl ether (1 in 4) | 4 mL                | 4 mL                | 4 mL                | 4 mL                |
| Caprylic acid                                    | 0.1 mL              | 0.1 mL              | 0.1 mL              | 0.1 mL              |
| Water  | a sufficient amount | a sufficient amount | a sufficient amount | a sufficient amount |
| Total volume                                     | 1000 mL             | 1000 mL             | 1000 mL             | 1000 mL             |

Flowing of mobile phase: Control the gradient by mixing the mobile phases A, B, C and D as directed in the following table.

| Time after injection of sample (min) | Mobile phase A (vol%) | Mobile phase B (vol%) | Mobile phase C (vol%) | Mobile phase D (vol%) |
|--------------------------------------|-----------------------|-----------------------|-----------------------|-----------------------|
| 0 ~ 9                                | 100                   | 0                     | 0                     | 0                     |
| 9 ~ 25                               | 0                     | 100                   | 0                     | 0                     |
| 25 ~ 61                              | 0                     | 100→0                 | 0→100                 | 0                     |
| 61 ~ 76                              | 0                     | 0                     | 100                   | 0                     |
| 76 ~ 96                              | 0                     | 0                     | 0                     | 100                   |

Reaction reagent: Dissolve 204 g of lithium acetate dihydrate in 336 mL of water, add 123 mL of acetic acid (100) and 401 mL of 1-methoxy-2-propanol, and use as Solution A. Separately, dissolve 39 g of ninhydrin and 81 mg of sodium borohydride in 979 mL of 1-methoxy-2-propanol, and use as Solution B. Mix the same volume of Solution A and Solution B before use.

Flow rate of mobile phase: 0.25 mL per minute.

Flow rate of reaction reagent: 0.3 mL per minute.

#### System suitability—

System performance: When the procedure is run with 50  $\mu$ L of the standard solution under the above operating conditions, serine, glutamic acid, proline, glycine, leucine, tyrosine, histidine, tryptophan and arginine are eluted in this order with enough separation between these peaks.

**Purity (1)** Clarity and color of solution—A solution obtained by dissolving 0.10 g of Gonadorelin Acetate in 10 mL of water is clear, and the absorbance of this solution at 350 nm determined as directed under Ultraviolet-visible Spectrophotometry <2.24> is not more than 0.10.

**(2)** Related substances—Dissolve 50 mg of Gonadorelin Acetate in 100 mL of the mobile phase A, and use this solution as the sample solution. Pipet 1 mL of the sample solution, add the mobile phase to make exactly 20 mL, and use this solution as the standard solution. Perform the test with 10  $\mu$ L each of the sample solution and standard solution as directed under Liquid Chromatography <2.01> according to the following conditions, and determine each peak area by the automatic integration method: the peak area other than gonadorelin from the sample solution is not more than 1/5 times the peak area of gonadorelin from the standard solution, and the total area of the peaks other than the peak of gonadorelin is not more than 3/5 of the peak area of

gonadorelin from the standard solution.

#### Operating conditions—

Detector, column, column temperature, mobile phase, and flow rate: Proceed as directed in the operating conditions in the Assay.

Time span of measurement: About 2.5 times as long as the retention time of gonadorelin beginning after the solvent peak.

#### System suitability—

Test for required detectability: To exactly 2 mL of the standard solution add the mobile phase to make exactly 100 mL. Confirm that the peak area of gonadorelin obtained from 10  $\mu$ L of this solution is equivalent to 1 to 3% of that obtained from 10  $\mu$ L of the standard solution.

System performance: Dissolve 4 mg of Gonadorelin Acetate in a suitable amount of the mobile phase, add 5 mL of a solution of phenacetin in acetonitrile (1 in 1000) and the mobile phase to make 50 mL. When the procedure is run with 10  $\mu$ L of this solution under the above operating conditions, gonadorelin and phenacetin are eluted in this order with the resolution between these peaks being not less than 3.

System repeatability: When the test is repeated 6 times with 10  $\mu$ L of the standard solution under the above operating conditions, the relative standard deviation of the peak area of gonadorelin is not more than 5%.

**Water** <2.48> Not more than 8.0% (0.15 g, volumetric titration, direct titration).

**Residue on ignition** <2.44> Not more than 0.2% (0.1 g).

**Assay** Weigh accurately about 40 mg of Gonadorelin Acetate and Gonadorelin Acetate Reference Standard (separately determine the water <2.48> in the same manner as Gonadorelin Acetate) and dissolve in diluted acetic acid (100) (1 in 1000) to make exactly 25 mL each. Pipet 5 mL each of these solutions, add exactly 5 mL of the internal standard solution and water to make 25 mL, and use these solutions as the sample solution and standard solution. Perform the test with 10  $\mu$ L each of the sample solution and standard solution as directed under Liquid Chromatography <2.01> according to the following conditions, and determine the ratios,  $Q_T$  and  $Q_S$ , of the peak area of gonadorelin to that of the internal standard.

Amount (mg) of  $C_{55}H_{75}N_{17}O_{13} \cdot 2C_2H_4O_2 = W_S \times (Q_T/Q_S)$

$W_S$ : Amount (mg) of Gonadorelin Acetate Reference Standard, calculated on the anhydrous basis

**Internal standard solution**—A solution of phenacetin in a mixture of water and acetonitrile (3:2) (1 in 1000).

#### Operating conditions—

Detector: An ultraviolet absorption photometer (wavelength: 220 nm).

Column: A stainless steel column 4.6 mm in inside diameter and 15 cm in length, packed with octadecylsilanized silica gel for liquid chromatography (5  $\mu$ m in particle diameter).

Column temperature: A constant temperature of about 40°C.

Mobile phase: A mixture of 0.05 mol/L potassium dihydrogen phosphate TS, pH 3.0 and acetonitrile (90:10).

Flow rate: Adjust the flow rate so that the retention time of gonadorelin is about 13 minutes.

#### System suitability—

**System performance:** When the procedure is run with 10  $\mu\text{L}$  of the standard solution under the above operating conditions, gonadorelin and the internal standard are eluted in this order with the resolution between these peaks being not less than 3.

**System repeatability:** When the test is repeated 6 times with 10  $\mu\text{L}$  of the standard solution under the above operating conditions, the relative standard deviation of the ratio of the peak area of gonadorelin to that of the internal standard is not more than 1.5%.

**Containers and storage** Containers—Tight containers.

Storage—Light-resistant.

## Human Chorionic Gonadotrophin

### Chorionic Gonadotrophin

ヒト絨毛性性腺刺激ホルモン

Human Chorionic Gonadotrophin is a dried preparation of gonad-stimulating hormone obtained from the urine of healthy pregnant women after the manufacturing process to remove or inactivate the virus.

It contains not less than 2500 human chorionic gonadotrophin Units per mg, and contains not less than 3000 chorionic gonadotrophin Units per mg protein.

It contains not less than 80% and not more than 125% of the labeled human chorionic gonadotrophin Units.

**Description** Human Chorionic Gonadotrophin occurs as a white to light yellow-brown powder.

It is freely soluble in water.

**Identification** Calculate  $b$  by the following equation, using  $Y_3$  and  $Y_4$  obtained in the Assay:  $b$  is not less than 120.

$$b = (E/I)$$

$$E = (Y_3 - Y_4)/f$$

$f$ : Number of test animals per group.

$$I = \log (T_H/T_L)$$

**Purity** (1) Clarity and color of solution—Dissolve 0.05 g of Human Chorionic Gonadotrophin in 5 mL of isotonic sodium chloride solution: the solution is clear and colorless or light yellow.

(2) Estrogen—Inject subcutaneously into each of three female albino rats or albino mice ovariectomized at least two weeks before the test, single dose of 100 units according to the labeled Units dissolved in 0.5 mL of isotonic sodium chloride solution. Take vaginal smear twice daily, on the third, fourth and fifth day. Place the smear thinly on a slide glass, dry, stain with Giemsa's TS, wash with water, and again dry: no estrus figure is shown microscopically <5.01>.

**Loss on drying** <2.41> Not more than 5.0% (0.1 g, in vacuum, phosphorus (V) oxide, 4 hours).

**Bacterial endotoxins** <4.01> Less than 0.03 EU/unit.

**Abnormal toxicity** Dilute Human Chorionic Gonadotrophin with isotonic sodium chloride solution so that each mL of the solution contains 120 Units, and use this solution as the sample solution. Inject 5.0 mL of the sample solution

into the peritoneal cavity of each of 2 or more of well-nourished, healthy guinea pigs weighing about 350 g, and observe the conditions of the animals for more than 7 days: all the animals exhibit no abnormalities.

**Specific activity** When calculate from the results obtained by the Assay and the following test, the specific activity is not less than 3000 human chorionic gonadotrophin Units per mg protein.

(i) Sample solution—To an exactly amount of Chorionic Gonadotrophin add water to make a solution so that each mL contains about 500 Units of human chorionic gonadotrophin according to the labeled amount.

(ii) Standard solution—Weigh accurately about 10 mg of bovine serum albumin, and dissolve in water to make exactly 20 mL. To a suitable volume of this solution add water to make four solutions containing exactly 300, 200, 100 and 50  $\mu\text{g}$  of the albumin per mL, respectively.

(iii) Procedure—Pipet 0.5 mL each of the sample solution and the standard solutions, put them in glass test tubes about 18 mm in inside diameter and about 130 mm in length, add exactly 5 mL of alkaline copper TS, mix, and allow the tubes to stand in a water bath at 30°C for 10 minutes. Then add exactly 0.5 mL of diluted Folin's TS (1 in 2), mix, and warm in a water bath at 30°C for 20 minutes. Determine the absorbances of these solutions at 750 nm as directed under Ultraviolet-visible Spectrophotometry <2.24> using a solution obtained in the same manner with 0.5 mL of water as the blank.

Plot the absorbances of the standard solutions on the vertical axis and their protein concentrations on the horizontal axis to prepare a calibration curve, and determine the protein content of the sample solution from its absorbance by using this curve. Then calculate the amount of the protein in the sample.

**Assay** (i) Test animals—Select healthy female albino rats weighing about 45 to 65 g.

(ii) Standard solution—Dissolve a quantity of Human Chorionic Gonadotrophin Reference Standard in bovine serum albumin-isotonic sodium chloride solution to prepare four kinds of solutions, having 7.5, 15, 30 and 60 Units per 2.5 mL, respectively. Inject these solutions into four groups consisting of five test animals each, and weigh their ovaries, as directed in procedure of (iv). Inject bovine serum albumin-isotonic sodium chloride solution to another group, and use this group as the control group. According to the result of this test, designate the concentration of the reference standard which will increase the masses of the ovaries about 2.5 times the mass of the ovaries of the control group as a low-dose concentration of the standard solution, and the concentration 1.5 to 2.0 times the low-dose concentration as a high-dose concentration. Dissolve a quantity of Human Chorionic Gonadotrophin Reference Standard, in bovine serum albumin-isotonic sodium chloride solution, and prepare a high-dose standard solution  $S_H$  and a low-dose standard solution  $S_L$  whose concentrations are equal to those determined by the above test.

(iii) Sample solution—According to the labeled units, weigh accurately a suitable quantity of Human Chorionic Gonadotrophin, dissolve in bovine serum albumin-isotonic sodium chloride solution, and prepare a high-dose sample solution  $T_H$  and a low-dose sample solution  $T_L$  having Units equal to the standard solutions in equal volumes, respectively.

(iv) Procedure—Divide the test animals at random into 4 groups, A, B, C and D, with not less than 10 animals and equal numbers in each group. Inject subcutaneously 0.5 mL of  $S_H$ ,  $S_L$ ,  $T_H$  and  $T_L$  in each group for 5 days. On the sixth day, excise the ovaries, remove the fat and other unwanted tissues attached to the ovaries, and remove the adhering water by lightly pressing between filter paper, and immediately weigh the ovaries.

(v) Calculation—Designate the mass of ovaries by  $S_H$ ,  $S_L$ ,  $T_H$  and  $T_L$  as  $y_1$ ,  $y_2$ ,  $y_3$  and  $y_4$ , respectively. Sum up  $y_1$ ,  $y_2$ ,  $y_3$  and  $y_4$  on each set to obtain  $Y_1$ ,  $Y_2$ ,  $Y_3$  and  $Y_4$ .

Units per mg of Human Chorionic Gonadotrophin  
= antilog  $M \times (\text{units per mL of } S_H) \times (b/a)$

$$M = IY_a/Y_b$$

$$I = \log (S_H/S_L) = \log (T_H/T_L)$$

$$Y_a = -Y_1 - Y_2 + Y_3 + Y_4$$

$$Y_b = Y_1 - Y_2 + Y_3 - Y_4$$

$a$ : Mass (mg) of sample.

$b$ : Total volume (mL) of the high dose of the test solution prepared by diluting with bovine serum albumin-isotonic sodium chloride solution.

$F'$  computed by the following equation should be smaller than  $F_1$  against  $n$  when  $s^2$  is calculated. And compute  $L$  ( $P = 0.95$ ) by the following equation:  $L$  should be not more than 0.3. If  $F'$  exceeds  $F_1$ , or if  $L$  exceeds 0.3, repeat the test increasing the number of the test animals or arranging the assay method in a better way until  $F'$  is smaller than  $F_1$  or  $L$  is not more than 0.3.

$$F' = (Y_1 - Y_2 - Y_3 + Y_4)^2 / 4fs^2$$

$f$ : Number of test animals per group.

$$s^2 = \{\Sigma y^2 - (Y/f)\} / n$$

$\Sigma y^2$ : The sum of the squares of each  $y_1$ ,  $y_2$ ,  $y_3$  and  $y_4$ .

$$Y = Y_1^2 + Y_2^2 + Y_3^2 + Y_4^2$$

$$n = 4(f-1)$$

$$L = 2\sqrt{(C-1)(CM^2 + I^2)}$$

$$C = Y_b^2 / (Y_b^2 - 4fs^2I^2)$$

$I^2$ : Value shown in the following table against  $n$  used to calculate  $s^2$ .

| $n$ | $I^2 = F_1$ | $n$ | $I^2 = F_1$ | $n$      | $I^2 = F_1$ |
|-----|-------------|-----|-------------|----------|-------------|
| 1   | 161.45      | 13  | 4.667       | 25       | 4.242       |
| 2   | 18.51       | 14  | 4.600       | 26       | 4.225       |
| 3   | 10.129      | 15  | 4.543       | 27       | 4.210       |
| 4   | 7.709       | 16  | 4.494       | 28       | 4.196       |
| 5   | 6.608       | 17  | 4.451       | 29       | 4.183       |
| 6   | 5.987       | 18  | 4.414       | 30       | 4.171       |
| 7   | 5.591       | 19  | 4.381       | 40       | 4.085       |
| 8   | 5.318       | 20  | 4.351       | 60       | 4.001       |
| 9   | 5.117       | 21  | 4.325       | 120      | 3.920       |
| 10  | 4.965       | 22  | 4.301       | $\infty$ | 3.841       |
| 11  | 4.844       | 23  | 4.279       |          |             |
| 12  | 4.747       | 24  | 4.260       |          |             |

**Containers and storage** Containers—Tight containers.

Storage—Light-resistant, and in a cold place.

## Human Chorionic Gonadotrophin for Injection

### Chorionic Gonadotrophin for Injection

注射用ヒト絨毛性性腺刺激ホルモン

Human Chorionic Gonadotrophin for Injection is a preparation for injection which is dissolved before use.

It contains not less than 80% and not more than 125% of the labeled human chorionic gonadotrophin Units.

**Method of preparation** Prepare as directed under Injections with Human Chorionic Gonadotrophin.

**Description** Human Chorionic Gonadotrophin for Injection occurs as a white to light yellow-brown powder or masses.

**Identification** Proceed as directed in the Identification under Human Chorionic Gonadotrophin.

**pH** <2.54> Prepare a solution so that each mL of isotonic sodium chloride solution contains 2 mg of Human Chorionic Gonadotrophin for Injection: the pH of this solution is between 5.0 and 7.0.

**Loss on drying** <2.41> Not more than 5.0% (0.1 g, in vacuum, phosphorus (V) oxide, 4 hours).

**Bacterial endotoxins** <4.01> Less than 0.03 EU/unit.

**Uniformity of dosage units** <6.02> When calculate the acceptance value using the mean of estimated contents of the units tested as M, it meets the requirements of the Mass variation test.

**Foreign insoluble matter** <6.06> Perform the test according to the Method 2: it meets the requirement.

**Insoluble particulate matter** <6.07> Perform the test according to the Method 1: it meets the requirement.

**Sterility** <4.06> Perform the test according to the Membrane filtration method: it meets the requirement.

**Assay** Proceed as directed in the Assay under Human Chorionic Gonadotrophin. The ratio of the assayed Units to the labeled Units should be calculated by the following equation.

$$\begin{aligned} &\text{The ratio of the assayed Units to the labeled Units} \\ &= \text{antilog } M \end{aligned}$$

**Containers and storage** Containers—Hermetic containers.  
Storage—Light-resistant, and in a cold place.

## Human Menopausal Gonadotrophin

ヒト下垂体性性腺刺激ホルモン

Human Menopausal Gonadotrophin is a dried preparation of gonad-stimulating hormone obtained

from the urine of postmenopausal healthy women, after processing for virus removal or inactivation. It has follicle-stimulating hormonal action and luteinizing hormonal action.

It contains not less than 40 follicle-stimulating hormone Units per mg.

**Description** Human Menopausal Gonadotrophin occurs as a white to pale yellow powder.

It is soluble in water.

**Purity** Interstitial cell-stimulating hormone—Perform the test according to the following method: the ratio of the unit of interstitial cell-stimulating hormone (luteinizing hormone) to that of follicle-stimulating hormone is not more than 1.

(i) Test animals—Select healthy male albino rats weighing about 45 to 65 g.

(ii) Standard solutions—Weigh accurately a suitable amount of Human Menopausal Gonadotrophin Reference Standard, and dissolve in bovine serum albumin-sodium chloride-phosphate buffer solution, pH 7.2, to prepare three kinds of solutions, containing 10, 20 and 40 interstitial cell-stimulating hormone (luteinizing hormone) units per 1.0 mL, respectively. Inject these solutions into three groups consisting of five test animals each, and weigh their seminal vesicles as directed in (iv). According to the result of the test, designate the concentration of the reference standard, which will make the mass of the seminal vesicle 20 to 35 mg, as the high-dose standard solution,  $S_H$ . Dilute the  $S_H$  to 1.5 to 2.0 times the initial volume with the bovine serum albumin-sodium chloride-phosphate buffer solution, pH 7.2, and designate this solution as the low-dose standard solution,  $S_L$ . Store these solutions at 2–8°C.

(iii) Sample solution—According to the labeled units, weigh accurately a suitable amount of Human Menopausal Gonadotrophin, and dissolve in the bovine serum albumin-sodium chloride-phosphate buffer solution, pH 7.2, to prepare the high-dose sample solution,  $T_H$  and the low-dose sample solution,  $T_L$ , so that their concentrations are similar to those of the corresponding standard solutions, respectively. Store these solutions at 2–8°C.

(iv) Procedure—Divide the test animals at random into 4 groups, A, B, C and D, with not less than 10 animals and equal numbers in each group. Inject subcutaneously once every day 0.2 mL each of  $S_H$ ,  $S_L$ ,  $T_H$  and  $T_L$  into the animals in each group. On the sixth day, excise the seminal vesicles, remove extraneous tissue, remove fluid adhering to the vesicles and the contents of the vesicles by lightly pressing between filter papers, and weigh the vesicles.

(v) Calculation—Proceed as directed in (v) in the Assay.

**Bacterial endotoxins** <4.01> Dissolve Human Menopausal Gonadotrophin in water for bacterial endotoxins test to prepare a solution containing 75 follicle-stimulating hormone Units per mL, and perform the test: less than 0.66 EU/ follicle-stimulating hormone Unit.

**Water** <2.48> Not more than 5.0% (0.2 g, volumetric titration, direct titration).

**Specific activity** Perform the test with Human Menopausal Gonadotrophin according to the following method, and calculate the specific activity using the amount (Unit) obtained in the Assay: it is not less than 50 follicle-stimulating hormone Units per 1 mg of protein.

(i) Sample solution—Weigh accurately about 10 mg of Human Menopausal Gonadotrophin, dissolve in water so that each mL contains exactly 200  $\mu$ g, and use this solution as the sample solution.

(ii) Standard solutions—Weigh accurately about 10 mg of bovine serum albumin, and dissolve in water to make exactly 20 mL. To this solution add water to make four solutions containing exactly 300  $\mu$ g, 200  $\mu$ g, 100  $\mu$ g and 50  $\mu$ g of the albumin per mL, respectively, and use these solutions as the standard solutions.

(iii) Procedure—To glass test tubes, about 18 mm in inside diameter and about 130 mm in height, add separately exactly 0.5 mL each of the sample solution and the standard solutions. To these tubes add exactly 5 mL of alkaline copper TS, warm in a water bath at 30°C for 10 minutes, then add exactly 0.5 mL of diluted Folin's TS (1 in 2), and warm in a water bath at 30°C for 20 minutes. Perform the test with these solutions as directed under Ultraviolet-visible Spectrometry <2.24>, and determine the absorbances at 750 nm, using a liquid obtained with 0.5 mL of water in the same manner as above as a blank. Prepare a calibration curve from the absorbances of the standard solutions, with absorbance on the vertical axis and concentration on the horizontal axis. Determine the amount of protein in the sample solution from the absorbance of the sample solution using the curve, and calculate the protein content of the sample.

#### Assay

(i) Test animals—Select healthy female albino rats weighing about 45 to 65 g.

(ii) Standard solutions—Weigh accurately a suitable amount of Human Menopausal Gonadotrophin Reference Standard, dissolve in human chorionic gonadotrophin TS to make three solutions which contain 0.75, 1.5 and 3.0 follicle-stimulating hormone Units per 1.0 mL, respectively. Inject these solutions into three groups consisting of five test animals each, and weigh their ovaries, as directed in (iv). According to the result of the test, designate the concentration of the reference standard, which will make the mass of the ovary about 120 to 160 mg, as the high-dose standard solution,  $S_H$ . Dilute the  $S_H$  to 1.5 to 2.0 times the initial volume with the human chorionic gonadotrophin TS, and designate the solution as the low-dose standard solution,  $S_L$ .

(iii) Sample solutions—According to the labeled units, weigh accurately a suitable amount of Human Menopausal Gonadotrophin, dissolve in human chorionic gonadotrophin TS, and prepare the high-dose sample solution,  $T_H$ , and the low-dose sample solution,  $T_L$ , which have similar numbers of units to those of corresponding standard solutions in equal volume, respectively.

(iv) Procedure—Divide the test animals at random into 4 groups, A, B, C and D, with not less than 10 animals and equal numbers in each group. Inject subcutaneously 0.2 mL each of  $S_H$ ,  $S_L$ ,  $T_H$  and  $T_L$  into the animals in each group, once in the afternoon on the first day, three times in the morning, noon and afternoon on the second day, and two times in the morning and afternoon on the third day. On the fifth day, excise the ovaries, remove the fat and extraneous tissue, remove fluid adhering to the ovaries by lightly pressing between filter papers, and immediately weigh the ovaries.

(v) Calculation—Designate the mass of ovaries by  $S_H$ ,  $S_L$ ,  $T_H$  and  $T_L$  as  $y_1$ ,  $y_2$ ,  $y_3$  and  $y_4$ , respectively. Sum up  $y_1$ ,  $y_2$ ,  $y_3$  and  $y_4$  on each set to obtain  $Y_1$ ,  $Y_2$ ,  $Y_3$  and  $Y_4$ .

Units per mg of Human Menopausal Gonadotrophin  
= antilog  $M \times (\text{units per mL of } S_H) \times (b/a)$

$M = IY_a/Y_b$

$I = \log (S_H/S_L) = \log (T_H/T_L)$

$Y_a = -Y_1 - Y_2 + Y_3 + Y_4$

$Y_b = Y_1 - Y_2 + Y_3 - Y_4$

$a$ : Mass (mg) of sample.

$b$ : Total volume (mL) of the high dose of the test solution prepared by diluting with bovine serum albumin-isotonic sodium chloride solution.

$F'$  computed by the following equation should be smaller than  $F_1$  against  $n$  when  $s^2$  is calculated. And compute  $L$  ( $P = 0.95$ ) by the following equation:  $L$  should be not more than 0.3. If  $F'$  exceeds  $F_1$ , or if  $L$  exceeds 0.3, repeat the test increasing the number of the test animals or arranging the assay method in a better way until  $F'$  is smaller than  $F_1$  or  $L$  is not more than 0.3.

$F' = (Y_1 - Y_2 - Y_3 - Y_4)^2 / (4fs^2)$

$f$ : Number of test animals per group.

$s^2 = \{ \sum y^2 - (Y/f) \} / n$

$\sum y^2$ : The sum of the squares of each  $y_1, y_2, y_3$  and  $y_4$ .

$Y = -Y_{12} - Y_{22} + Y_{32} + Y_{42}$

$n = 4(f - 1)$

$L = 2\sqrt{(C-1)(CM^2 + I^2)}$

$C = Y_b^2 / (Y_a^2 - 4fs^2t^2)$

$t^2$ : Value shown in the following table against  $n$  used to calculate  $s^2$ .

| $n$ | $t^2 = F_1$ | $n$ | $t^2 = F_1$ | $n$      | $t^2 = F_1$ |
|-----|-------------|-----|-------------|----------|-------------|
| 1   | 161.45      | 13  | 4.667       | 25       | 4.242       |
| 2   | 18.51       | 14  | 4.600       | 26       | 4.225       |
| 3   | 10.129      | 15  | 4.543       | 27       | 4.210       |
| 4   | 7.709       | 16  | 4.494       | 28       | 4.196       |
| 5   | 6.608       | 17  | 4.451       | 29       | 4.183       |
| 6   | 5.987       | 18  | 4.414       | 30       | 4.171       |
| 7   | 5.591       | 19  | 4.381       | 40       | 4.085       |
| 8   | 5.318       | 20  | 4.351       | 60       | 4.001       |
| 9   | 5.117       | 21  | 4.325       | 120      | 3.920       |
| 10  | 4.965       | 22  | 4.301       | $\infty$ | 3.841       |
| 11  | 4.844       | 23  | 4.279       |          |             |
| 12  | 4.747       | 24  | 4.260       |          |             |

**Containers and storage** Containers—Tight containers.

Storage—Light-resistant, and in a cold place.

## Serum Gonadotrophin

血清性腺刺激ホルモン

Serum Gonadotrophin is a dried preparation of gonad-stimulating hormone, obtained from pregnant mares' serum which has adequately inspected viruses, and subjected to a suitable process for removal or inactivation of viruses.

It contains not less than 2000 serum gonadotrophin Units per mg.

It contains not less than 80% and not more than 125% of the labeled serum gonadotrophin Units.

**Description** Serum Gonadotrophin occurs as a white powder.

It is freely soluble in water.

**Identification** Calculate  $b$  by the following equation, using  $Y_3$  and  $Y_4$  obtained in the Assay:  $b$  is not less than 120.

$$b = E/I$$

$$E = (Y_3 - Y_4)/f$$

$f$ : Number of test animals per group.

$I = \log (T_H/T_L)$

**Purity** Clarity and color of solution—Dissolve Serum Gonadotrophin in isotonic sodium chloride solution to prepare a solution containing 9000 units per mL according to the labeled Units: the solution is clear and colorless.

**Loss on drying** <2.41> Not more than 8.0% (0.1 g, in vacuum, phosphorus (V) oxide, 4 hours).

**Bacterial endotoxins** <4.01> Less than 0.1 EU/unit.

**Specific activity** When calculated from the results obtained by the Assay and the following test, Serum Gonadotrophin contains not less than 3000 serum gonadotrophin Units per mg of protein.

(i) Standard solutions—Dissolve about 3 mg of bovine serum albumin in water to make a solution containing 500  $\mu$ g of the albumin in each mL. To this solution add water to make four standard solutions so that each mL contains exactly 200  $\mu$ g, 150  $\mu$ g, 100  $\mu$ g and 50  $\mu$ g of the albumin, respectively.

(ii) Sample solution—Dissolve about 1 mg of Serum Gonadotrophin in water to make a solution containing exactly 180  $\mu$ g in each mL.

(iii) Sodium carbonate solution—Dissolve 2 g of sodium carbonate (standard reagent) in 0.1 mol/L sodium hydroxide TS to make 100 mL.

(iv) Sodium tartrate solution—Dissolve about 1 g of sodium tartrate dihydrate in water to make 100 mL.

(v) Copper (II) sulfate solution—Dissolve 0.5 g of copper (II) sulfate pentahydrate in the sodium tartrate solution to make 100 mL.

(vi) Alkaline copper solution—Mix 50 mL of the sodium carbonate solution and 1 mL of the copper (II) sulfate solution. Prepare before use. Use within the day of preparation.

(vii) Procedure—Pipet 0.5 mL each of the standard solutions and the sample solution in small test tubes, add 3 mL of the alkaline copper solution to them, and mix. Allow them to stand at the room temperature for not less than 10 minutes, add 0.3 mL of diluted Folin's TS (1 in 2), mix immediately, and allow to stand for not less than 30 minutes. Determine the absorbances of these solutions so obtained at 750 nm as directed under Ultraviolet-visible Spectrophotometry <2.24> using a solution, prepared in the same manner with 0.5 mL of water, as the blank. Plot the calibration curve from the absorbances obtained with the standard solutions, and determine the amount of protein in the sample solution from

this curve.

$$\text{Specific activity (unit/mg protein)} \\ = [(\text{units per mg, obtained in the Assay}) / \\ (\text{amount (\%) of protein in the sample})] \times 100$$

**Abnormal toxicity** Dissolve Serum Gonadotrophin in isotonic sodium chloride solution so that each 5 mL of the solution contains 4000 Units, and use this solution as the sample solution. Inject 5.0 mL of the sample solution into the peritoneal cavity of each of 2 or more of well-nourished, healthy guinea pigs weighing about 350 g, and inject 0.5 mL of the sample solution into the peritoneal cavity of each of 2 or more of well-nourished, healthy mice aged about 5 weeks. Observe the conditions of the animals for more than 7 days: all the animals exhibit no abnormalities.

#### Assay

(i) Test animals—Select healthy female albino rats weighing about 45 g.

(ii) Standard solution—Dissolve a quantity of Serum Gonadotrophin Reference Standard in bovine serum albumin-isotonic sodium chloride solution to prepare four kinds of solutions, containing 10, 20, 40 and 80 Units per 0.5 mL, respectively. Inject these solutions into four groups consisting of five test animals each, and weigh their ovaries, as directed in procedure of (iv). Inject bovine serum albumin-isotonic sodium chloride solution to another group, and use this group as the control group. According to the result of this test, designate the concentration of the reference standard which will increase the masses of the ovaries about 3 times the mass of the ovaries of the control group as a low-dose concentration of the standard solution, and the concentration 1.5 to 2.0 times the low-dose concentration as a high-dose concentration. Weigh accurately a suitable quantity of Serum Gonadotrophin Reference Standard, dissolve in bovine serum albumin-isotonic sodium chloride solution, and prepare a high-dose standard solution  $S_H$  and a low-dose standard solution  $S_L$  whose concentrations are equal to those determined by the above test.

(iii) Sample solution—According to the labeled units, weigh accurately a suitable quantity of Serum Gonadotrophin, dissolve in bovine serum albumin-isotonic sodium chloride solution, and prepare a high-dose sample solution  $T_H$  and a low-dose sample solution  $T_L$  having Units equal to the standard solutions in equal volumes, respectively.

(iv) Procedure—Divide the test animals at random into 4 groups, A, B, C and D, with not less than 10 animals and equal numbers in each group. Inject once subcutaneously 0.5 mL of  $S_H$ ,  $S_L$ ,  $T_H$  and  $T_L$  in each group. On the sixth day, excise the ovaries, remove the fat and other unwonted tissues attached to the ovaries, and remove the adhering water by lightly pressing between filter paper, and immediately weigh the ovaries.

(v) Calculation—Designate the mass of ovaries by  $S_H$ ,  $S_L$ ,  $T_H$  and  $T_L$  as  $y_1$ ,  $y_2$ ,  $y_3$  and  $y_4$ , respectively. Sum up  $y_1$ ,  $y_2$ ,  $y_3$  and  $y_4$  on each set to obtain  $Y_1$ ,  $Y_2$ ,  $Y_3$  and  $Y_4$ .

$$\text{Units per mg of Serum Gonadotrophin} \\ = \text{antilog } M \times (\text{units per mL of } S_H) \times (b/a)$$

$$M = IY_a/Y_b$$

$$I = \log (S_H/S_L) = \log (T_H/T_L)$$

$$Y_a = -Y_1 - Y_2 + Y_3 + Y_4$$

$$Y_b = Y_1 - Y_2 + Y_3 - Y_4$$

$a$ : Mass (mg) of sample.

$b$ : Total volume (mL) of the high dose of the test solution prepared by diluting with bovine serum albumin-isotonic sodium chloride solution.

$F'$  computed by the following equation should be smaller than  $F_1$  against  $n$  when  $s^2$  is calculated. And compute  $L$  ( $P = 0.95$ ) by the following equation:  $L$  should be not more than 0.3. If  $F'$  exceeds  $F_1$ , or if  $L$  exceeds 0.3, repeat the test increasing the number of the test animals or arranging the assay method in a better way until  $F'$  is smaller than  $F_1$  or  $L$  is not more than 0.3.

$$F' = (Y_1 - Y_2 - Y_3 + Y_4)^2 / 4fs^2$$

$f$ : Number of test animals per group.

$$s^2 = \{\Sigma y^2 - (Y/f)\} / n$$

$\Sigma y^2$ : The sum of the squares of each  $y_1$ ,  $y_2$ ,  $y_3$  and  $y_4$ .

$$Y = Y_1^2 + Y_2^2 + Y_3^2 + Y_4^2$$

$$n = 4(f - 1)$$

$$L = 2\sqrt{(C-1)(CM^2 + I^2)}$$

$$C = Y_b^2 / (Y_b^2 - 4fs^2I^2)$$

$I^2$ : Value shown in the following table against  $n$  used to calculate  $s^2$ .

| $n$ | $I^2 = F_1$ | $n$ | $I^2 = F_1$ | $n$      | $I^2 = F_1$ |
|-----|-------------|-----|-------------|----------|-------------|
| 1   | 161.45      | 13  | 4.667       | 25       | 4.242       |
| 2   | 18.51       | 14  | 4.600       | 26       | 4.225       |
| 3   | 10.129      | 15  | 4.543       | 27       | 4.210       |
| 4   | 7.709       | 16  | 4.494       | 28       | 4.196       |
| 5   | 6.608       | 17  | 4.451       | 29       | 4.183       |
| 6   | 5.987       | 18  | 4.414       | 30       | 4.171       |
| 7   | 5.591       | 19  | 4.381       | 40       | 4.085       |
| 8   | 5.318       | 20  | 4.351       | 60       | 4.001       |
| 9   | 5.117       | 21  | 4.325       | 120      | 3.920       |
| 10  | 4.965       | 22  | 4.301       | $\infty$ | 3.841       |
| 11  | 4.844       | 23  | 4.279       |          |             |
| 12  | 4.747       | 24  | 4.260       |          |             |

**Containers and storage** Containers—Tight containers.

Storage—Light-resistant, and in a cold place.

## Serum Gonadotrophin for Injection

注射用血清性腺刺激ホルモン

Serum Gonadotrophin for Injection is a preparation for injection which is dissolved before use.

It contains not less than 80% and not more than 125% of the labeled Serum Gonadotrophin Units.

**Method of preparation** Prepare as directed under Injections with Serum Gonadotrophin.

**Description** Serum Gonadotrophin for Injection occurs as white powder or masses.

**Identification** Proceed as directed in the Identification under Serum Gonadotrophin.

**pH <2.54>** Dissolve 30 mg of Serum Gonadotrophin for In-



jection in 20 mL of isotonic sodium chloride solution: the pH of this solution is between 5.0 and 7.0.

**Loss on drying** <2.41> Not more than 5.0% (0.1 g, in vacuum, phosphorus (V) oxide, 4 hours).

**Bacterial endotoxins** <4.01> Less than 0.1 EU/unit.

**Assay** Proceed as directed in the Assay under Serum Gonadotrophin. The ratio of the Units assayed to the labeled Units should be calculated by the following equation.

$$\frac{\text{The ratio of the assayed Units to the labeled Units}}{= \text{antilog } M}$$

**Containers and storage** Containers—Hermetic containers.  
Storage—Light-resistant, and in a cold place.

## Gramicidin

グラミシジン

[1405-97-6]

Gramicidin is a mixture of peptide substances having antibacterial activity produced by the growth of *Bacillus brevis* Dubos.

It contains not less than 900 µg (potency) per mg, calculated on the dried basis. The potency of Gramicidin is expressed as mass (potency) of gramicidin.

**Description** Gramicidin occurs as a white to light yellowish white crystalline powder.

It is freely soluble in methanol, soluble in ethanol (99.5), and practically insoluble in water.

**Identification** (1) To 10 mg of Gramicidin add 2 mL of 6 mol/L hydrochloric acid TS, and heat in a water bath for 30 minutes with occasional stirring. After cooling, neutralize with 6 mol/L sodium hydroxide TS, add 1 mL of ninhydrin TS and 0.5 mL of pyridine, and heat for 2 minutes: a blue-purple to red-purple color develops.

(2) Determine the absorption spectrum of a solution of Gramicidin in ethanol (95) (1 in 20,000), as directed under Ultraviolet-visible Spectrophotometry <2.24>, and compare the spectrum with the Reference Spectrum or the spectrum of a solution of Gramicidin Reference Standard prepared in the same manner as the sample solution: both spectra exhibit similar intensities of absorption at the same wavelengths.

**Loss on drying** <2.41> Not more than 3.0% (0.1 g, in vacuum, 60°C, 3 hours).

**Residue on ignition** <2.44> Not more than 1.0% (1 g).

**Assay** Perform the test according to the Turbidimetric method as directed under Microbial Assay for Antibiotics <4.02> according to the following conditions.

- (i) Test organism—*Enterococcus hirae* ATCC 10541
- (ii) Agar medium for transferring test organism

|                                |        |
|--------------------------------|--------|
| Glucose                        | 10.0 g |
| Casein peptone                 | 5.0 g  |
| Yeast extract                  | 20.0 g |
| Potassium dihydrogen phosphate | 2.0 g  |
| Polysorbate 80                 | 0.1 g  |

|       |         |
|-------|---------|
| Agar  | 15.0 g  |
| Water | 1000 mL |

Mix all the ingredients, and sterilize. Adjust the pH of the solution so that it will be 6.7 to 6.8 after sterilization.

(iii) Liquid medium for suspending test organism—Use the culture medium (2).

(iv) Preparation of the test organism suspension—Puncture the test organism in the medium, prepared by dispensing 10 mL of the agar medium for transferring test organism in a test tube about 16 mm in inside diameter, incubate at 36.5 to 37.5°C for 20 to 24 hours. After sub-culturing at least three times, keep between 1 to 5°C. Transfer the organism so obtained in 10 mL of the liquid medium for suspending test organism, incubate at 36.5 to 37.5°C for 20 to 24 hours, and use this medium as the test organism stock suspension. Before use, add the test organism stock suspension to the liquid medium for suspending test organism so that the transmittance at 580 nm is 50 to 60%. Mix one volume of this suspension and 200 volume of the liquid medium for suspending test organism, and use this as the test organism suspension.

(v) Standard solution—Weigh accurately an amount of Gramicidin Reference Standard, previously dried under reduced pressure not exceeding 0.67 kPa at 60°C for 3 hours, equivalent to about 10 mg (potency), dissolve in ethanol (99.5) to make exactly 100 mL, and use this solution as the standard stock solution. Keep the standard stock solution at not exceeding 5°C and use within 7 days. Take exactly a suitable amount of the standard stock solution before use, add the following diluting solution to make a solution so that each mL contains 0.02 µg (potency), and use this solution as the standard solution.

Diluting solution: To 390 mL of propylene glycol add 210 mL of a mixture of ethanol (99.5) and acetone (9:1) and Sterile Purified Water to make 1000 mL.


(vi) Sample solution—Weigh accurately an amount of Gramicidin, equivalent to about 10 mg (potency), and dissolve in ethanol (99.5) to make exactly 100 mL. Take exactly a suitable amount of this solution, add the diluting solution obtained in (v) to make a solution so that each mL contains 0.02 µg (potency), and use this solution as the sample solution.

(vii) Procedure—Transfer 0.155 mL, 0.125 mL, 0.100 mL, 0.080 mL and 0.065 mL each of the standard solution, 0.100 mL of the sample solution and 0.100 mL of the diluting solution, separately, in test tubes about 14 mm in inside diameter and about 15 cm in length, and make three sets for each. To each of the test tube add 10 mL of the test organism suspension, stopper the tube, incubate in a water bath at 36.5 to 37.5°C for 180 to 270 minutes, add 0.5 mL of a solution of formaldehyde (1 in 3), and determine their transmittances at 580 nm.

**Containers and storage** Containers—Tight containers.

## Griseofulvin

グリセオフルビン



$C_{17}H_{17}ClO_6$ : 352.77

(2*S*,6'*R*)-7-Chloro-2',4,6-trimethoxy-6'-methylspiro[benzo[*b*]furan-2(3*H*),1'-(cyclohex-2'-ene)]-3,4'-dione [126-07-8]

Griseofulvin is a substance having antifungal activity produced by the growth of *Penicillium griseofulvum* or *Penicillium janczewskii*.

It contains not less than 960  $\mu$ g (potency) and not more than 1020  $\mu$ g (potency) per mg, calculated on the dried basis. The potency of Griseofulvin is expressed as mass (potency) of griseofulvin ( $C_{17}H_{17}ClO_6$ ).

**Description** Griseofulvin occurs as white, crystals or crystalline powder.

It is soluble in *N,N*-dimethylformamide, sparingly soluble in acetone, slightly soluble in methanol and in ethanol (95), and practically insoluble in water.

**Identification** (1) Determine the absorption spectrum of a solution of Griseofulvin in ethanol (95) (1 in 100,000) as directed under Ultraviolet-visible Spectrophotometry <2.24>, and compare the spectrum with the Reference Spectrum or the spectrum of a solution of Griseofulvin Reference Standard prepared in the same manner as the sample solution: both spectra exhibit similar intensities of absorption at the same wavelengths.

(2) Determine the infrared absorption spectrum of Griseofulvin as directed in the potassium bromide disk method under Infrared Spectrophotometry <2.25>, and compare the spectrum with the Reference Spectrum or the spectrum of Griseofulvin Reference Standard: both spectra exhibit similar intensities of absorption at the same wave numbers.

**Optical rotation** <2.49>  $[\alpha]_D^{20}$ : +350 – +364° (0.25 g calculated on the dried basis, *N,N*-dimethylformamide, 25 mL, 100 mm).

**Melting point** <2.60> 218 – 222°C

**Purity** (1) Acidity—Dissolve 0.25 g of Griseofulvin in 20 mL of neutralized ethanol, and add 2 drops of phenolphthalein TS and 1.0 mL of 0.02 mol/L sodium hydroxide VS: the color of the solution is red.

(2) Heavy metals <1.07>—Proceed with 1.0 g of Griseofulvin according to Method 2, and perform the test. Prepare the control solution with 2.5 mL of Standard Lead Solution (not more than 25 ppm).

(3) Arsenic <1.11>—Prepare the test solution with 1.0 g of Griseofulvin according to Method 3, and perform the test (not more than 2 ppm).

(4) Related substances—To 0.10 g of Griseofulvin add exactly 1 mL of the internal standard solution and acetone to make 10 mL, and use this solution as the sample solution. Separately, to 5.0 mg of Griseofulvin Reference Standard add exactly 1 mL of the internal standard solution and acetone to make 10 mL, and use this solution as the standard solution. Perform the test with 2  $\mu$ L each of the sample solution and standard solution as directed under Gas Chromatography <2.02> according to the following conditions, determine each peak area by the automatic integration method, and calculate the ratio,  $Q_1$ , of the peak area of dechlorogriseofulvin, having the relative retention time of about 0.6 with respect to griseofulvin, to that of the internal standard obtained from the sample solution, the ratio,  $Q_2$ , of the peak area of dehydrogriseofulvin, having the relative retention time of about 1.2 with respect to griseofulvin, to that of the internal standard obtained from the sample solution and the ratio,  $Q_S$ , of the peak area of griseofulvin to that of the internal standard obtained from the standard solution:  $Q_1/Q_S$  is not more than 0.6, and  $Q_2/Q_S$  is not more than 0.15. *Internal standard solution*—A solution of 9,10-diphenylanthracene in acetone (1 in 500).

*Operating conditions*—

Detector: An hydrogen flame-ionization detector.

Column: A glass column 4 mm in inside diameter and 1 m in length, packed with siliceous earth for gas chromatography coated with 25% phenyl-25% cyanopropyl-methylsilicone polymer for gas chromatography in the ratio of 1% (150 – 180  $\mu$ m in particle diameter).

Column temperature: A constant temperature of about 250°C.

Temperature of injection port: A constant temperature of about 270°C.

Temperature of detector: A constant temperature of about 300°C.

Carrier gas: Nitrogen

Flow rate: Adjust the flow rate so that the retention time of griseofulvin is about 10 minutes.

*System suitability*—

Test for required detectability: Measure exactly 1 mL of the standard solution, and add the internal standard solution diluted with acetone (1 in 10) to make exactly 10 mL. Confirm that the ratio of the peak area of griseofulvin to that of the internal standard obtained from 2  $\mu$ L of this solution is equivalent to 7 to 13% of that obtained from 2  $\mu$ L of the standard solution.

System performance: When the procedure is run with 2  $\mu$ L of the standard solution under the above operating conditions, the internal standard and griseofulvin are eluted in this order with the resolution between these peaks being not less than 5.

System repeatability: When the test is repeated 6 times with 2  $\mu$ L of the standard solution under the above operating conditions, the relative standard deviation of the ratios of the peak area of griseofulvin to that of the internal standard is not more than 5.0%.

(5) Petroleum ether soluble substances—To 1.0 g of Griseofulvin add 20 mL of petroleum ether, shake, and boil for 10 minutes under a reflux condenser. After cooling, filter through a dried filter paper, wash the filter paper with two 15-mL portions of petroleum ether, combine the washings to the filtrate, evaporate the petroleum ether on a water bath, and dry the residue at 105°C for 1 hour: the amount of the

residue is not more than 0.2%.

**Loss on drying** <2.41> Not more than 1.0% (1 g, reduced pressure not exceeding 0.67 kPa, 60°C, 3 hours).

**Residue on ignition** <2.44> Not more than 0.2% (1 g).

**Assay** Weigh accurately an amount of Griseofulvin and Griseofulvin Reference Standard, equivalent to about 50 mg (potency), dissolve each in 50 mL of *N,N*-dimethylformamide, add exactly 20 mL of the internal standard solution and water to make 250 mL, and use these solutions as the sample solution and standard solution. Perform the test with 10  $\mu$ L each of the sample solution and standard solution as directed under Liquid Chromatography <2.01> according to the following conditions, and determine the ratios,  $Q_T$  and  $Q_S$ , of the peak area of griseofulvin to that of the internal standard.

$$\begin{aligned} &\text{Amount } [\mu\text{g (potency)}] \text{ of } \text{C}_{17}\text{H}_{17}\text{ClO}_6 \\ &= W_S \times (Q_T/Q_S) \times 1000 \end{aligned}$$

$W_S$ : Amount [mg (potency)] of Griseofulvin Reference Standard

**Internal standard solution**—A solution of butyl parahydroxybenzoate in acetonitrile (1 in 400).

**Operating conditions**—

**Detector**: An ultraviolet absorption photometer (wavelength: 254 nm).

**Column**: A stainless steel column 4.6 mm in inside diameter and 25 cm in length, packed with octadecylsilanized silica gel for liquid chromatography (10  $\mu$ m in particle diameter).

**Column temperature**: A constant temperature of about 25°C.

**Mobile phase**: A mixture of water and acetonitrile (3:2).

**Flow rate**: Adjust the flow rate so that the retention time of griseofulvin is about 6 minutes.

**System suitability**—

**System performance**: When the procedure is run with 10  $\mu$ L of the standard solution under the above operating conditions, griseofulvin and the internal standard are eluted in this order with the resolution between these peaks being not less than 4.


**System repeatability**: When the test is repeated 6 times with 10  $\mu$ L of the standard solution under the above operating conditions, the relative standard deviation of the ratios of the peak area of griseofulvin to that of the internal standard is not more than 1.0%.

**Containers and storage** Containers—Tight containers.

## Guaifenesin

### Guaiacol Glyceryl Ether

グアイフェネシン



$\text{C}_{10}\text{H}_{14}\text{O}_4$ : 198.22

(2*RS*)-3-(2-Methoxyphenoxy)propane-1,2-diol [93-14-1]

Guaifenesin, when dried, contains not less than 98.0% and not more than 102.0% of  $\text{C}_{10}\text{H}_{14}\text{O}_4$ .

**Description** Guaifenesin occurs as a white crystals or crystalline powder.

It is freely soluble in ethanol (95), and sparingly soluble in water.

A solution of ethanol (95) (1 in 20) shows no optical rotation.

**Identification** (1) Determine the absorption spectrum of a solution of Guaifenesin (1 in 50,000) as directed under Ultraviolet-visible Spectrophotometry <2.24>, and compare the spectrum with the Reference Spectrum or the spectrum of a solution of Guaifenesin Reference Standard prepared in the same manner as the sample solution: both spectra exhibit similar intensities of absorption at the same wavelengths.

(2) Determine the infrared absorption spectrum of Guaifenesin, previously dried, as directed in the potassium bromide disk method under Infrared Spectrophotometry <2.25>, and compare the spectrum with the Reference Spectrum or the spectrum of previously dried Guaifenesin Reference Standard: both spectra exhibit similar intensities of absorption at the same wave numbers.

**Melting point** <2.60> 80 – 83°C

**pH** <2.54> Dissolve 1.0 g of Guaifenesin in 100 mL of water: the pH of the solution is between 5.0 and 7.0.

**Purity** (1) Clarity and color of solution—Dissolve 0.20 g of Guaifenesin in 10 mL of water: the solution is clear and colorless.

(2) Chloride <1.03>—Dissolve 0.7 g of Guaifenesin in 25 mL of water by warming. Cool, add 6 mL of dilute nitric acid and water to make 50 mL, and perform the test using this solution as the test solution. Prepare the control solution with 0.40 mL of 0.01 mol/L hydrochloric acid VS (not more than 0.020%).

(3) Heavy metals <1.07>—Dissolve 2.0 g of Guaifenesin in 25 mL of water by warming. Cool, add 2 mL of dilute acetic acid and water to make 50 mL, and perform the test using this solution as the test solution. Prepare the control solution with 2.0 mL of Standard Lead Solution (not more than 10 ppm).

(4) Arsenic <1.11>—Prepare the test solution with 1.0 g of Guaifenesin according to Method 3, and perform the test (not more than 2 ppm).

(5) Free guaiacol—To 1.0 g of Guaifenesin add exactly 25 mL of water, dissolve by warming, cool, and use this solu-

tion as the sample solution. Separately, dissolve 0.100 g of guaiacol in water to make exactly 1000 mL. Pipet 3 mL of this solution, add exactly 22 mL of water, and use this solution as the standard solution. To each of the sample solution and the standard solution add 1.0 mL of potassium hexacyanoferrate (III) TS and 5.0 mL of a solution of 4-aminoantipyrine (1 in 200), and immediately after shaking for exactly 5 seconds add a solution of sodium hydrogen carbonate (1 in 1200) to make exactly 100 mL. Determine the absorbances of these solutions at 500 nm exactly 15 minutes after the addition of the 4-aminoantipyrine solution as directed under Ultraviolet-visible Spectrophotometry <2.24>, using a solution, prepared in the same manner with 25 mL of water, as the blank: the absorbance of the solution obtained from the sample solution is not greater than that from the standard solution.

(6) Related substances—Dissolve 1.0 g of Guaifenesin in 100 mL of ethanol (95), and use this solution as the sample solution. Pipet 1 mL of the sample solution, add water to make exactly 200 mL, and use this solution as the standard solution. Perform the test with these solutions as directed under Thin-layer Chromatography <2.03>. Spot 10  $\mu$ L each of the sample solution and standard solution on a plate of silica gel for thin-layer chromatography. Develop the plate with a mixture of diethyl ether, ethanol (95), and ammonia solution (28) (40:10:1) to a distance of about 10 cm, and air-dry the plate. Spray evenly 4-dimethylaminobenzaldehyde TS for spraying on the plate, and heat at 110°C for 10 minutes: the spots other than the principal spot from the sample solution are not more intense than the spot from the standard solution.

**Loss on drying** <2.41> Not more than 0.5% (1 g, in vacuum, 60°C, 3 hours).

**Residue on ignition** <2.44> Not more than 0.1% (1 g).

**Assay** Weigh accurately about 60 mg of Guaifenesin and Guaifenesin Reference Standard, previously dried, and dissolve each then in water to make exactly 100 mL. Pipet 5 mL of these solutions, and add water to make exactly 100 mL, and use these solutions as the sample solution and standard solution. Determine the absorbances,  $A_T$  and  $A_S$ , of the sample solution and standard solution at 273 nm as directed under Ultraviolet-visible Spectrophotometry <2.24>.


$$\begin{aligned} \text{Amount (mg) of } C_{10}H_{14}O_4 \\ = W_S \times (A_T/A_S) \end{aligned}$$

$W_S$ : Amount (mg) of Guaifenesin Reference Standard

**Containers and storage** Containers—Tight containers.

## Guanabenz Acetate

グアナベンズ酢酸塩



$C_8H_8Cl_2N_4 \cdot C_2H_4O_2$ : 291.13  
(*E*)-1-(2,6-Dichlorobenzylideneamino)guanidine monoacetate [23256-50-0]

Guanabenz Acetate, when dried, contains not less than 98.5% of  $C_8H_8Cl_2N_4 \cdot C_2H_4O_2$ .

**Description** Guanabenz Acetate occurs as white crystals or crystalline powder.

It is freely soluble in acetic acid (100), soluble in methanol and in ethanol (95), slightly soluble in water, and practically insoluble in diethyl ether.

It is gradually affected by light.

Melting point: about 190°C (with decomposition).

**Identification** (1) To 5 mL of a solution of Guanabenz Acetate (1 in 1000) add 0.5 mL of a diluted ethanol (95) (5 in 6) which contains 16 g of urea and 0.2 g of 1-naphthol in 100 mL, and add 1 mL of *N*-bromosuccinimide TS: a purple color develops.

(2) Determine the absorption spectrum of a solution of Guanabenz Acetate in methanol (1 in 100,000) as directed under Ultraviolet-visible Spectrophotometry <2.24>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wavelengths.

(3) Determine the infrared absorption spectrum of Guanabenz Acetate, previously dried, as directed in the potassium bromide disk method under Infrared Spectrophotometry <2.25>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wave numbers.

(4) To 0.1 g of Guanabenz Acetate add 5 mL of water and 1 mL of ammonia TS, shake, filter, and neutralize the filtrate with dilute hydrochloric acid: the solution responds to the Qualitative Tests <1.09> (3) for acetate.

**Purity** (1) Heavy metals <1.07>—Proceed with 2.0 g of Guanabenz Acetate according to Method 2, and perform the test. Prepare the control solution with 2.0 mL of Standard Lead Solution (not more than 10 ppm).

(2) Related substances—Conduct this procedure without exposure to daylight, using light-resistant vessels. Dissolve 0.05 g of Guanabenz Acetate in 5 mL of methanol, and use this solution as the sample solution. Pipet 1 mL of the sample solution, add methanol to make exactly 10 mL, then pipet 1 mL of this solution, add methanol to make exactly 20 mL, and use this solution as the standard solution. Perform the test with these solutions as directed under Thin-layer Chromatography <2.03>. Spot 5  $\mu$ L each of the sample solution and standard solution on a plate of silica gel with fluorescent indicator for thin-layer chromatography. Develop the plate with a mixture of chloroform, methanol and ammonia solution (28) (80:20:1) to a distance of about 10 cm, and air-dry the plate. Examine under ultraviolet light (main wavelength: 254 nm): the spots other than the principal spot from the sample solution are not more intense than the spot from the standard solution. Place the plate in a chamber filled with iodine vapor for 10 minutes: the spots other than the principal spot from the sample solution are not more intense than the spot from the standard solution.

**Loss on drying** <2.41> Not more than 0.5% (1 g, in vacuum, phosphorus (V) oxide, 50°C, 3 hours).

**Residue on ignition** <2.44> Not more than 0.2% (1 g).

**Assay** Weigh accurately about 0.25 g of Guanabenz Acetate, previously dried, dissolve in 50 mL of acetic acid


(100), and titrate <2.50> with 0.1 mol/L perchloric acid VS (potentiometric titration). Perform a blank determination, and make any necessary correction.

Each mL of 0.1 mol/L perchloric acid VS  
= 29.11 mg of  $C_8H_8Cl_2N_4 \cdot C_2H_4O_2$

**Containers and storage** Containers—Tight containers.  
Storage—Light-resistant.

## Guanethidine Sulfate

グアネチジン硫酸塩



$C_{10}H_{22}N_4 \cdot H_2SO_4$ : 296.39  
1-[2-(Hexahydroazocin-1(2*H*)-yl)ethyl]guanidine  
monosulfate [645-43-2]

Guanethidine Sulfate, when dried, contains not less than 98.5% of  $C_{10}H_{22}N_4 \cdot H_2SO_4$ .

**Description** Guanethidine Sulfate occurs as white crystals or crystalline powder. It is odorless or has a slight, characteristic odor and a bitter taste.

It is very soluble in formic acid, freely soluble in water, and practically insoluble in ethanol (95) and in diethyl ether.

Melting point: 251 – 256°C (an evacuated sealed capillary tube, with decomposition).

**Identification** (1) To 4 mL of a solution of Guanethidine Sulfate (1 in 4000) add 2 mL of 1-naphthol TS, 1 mL of diacetyl TS and 15 mL of water, and allow to stand for 30 minutes: a red color develops.

(2) Determine the infrared absorption spectrum of Guanethidine Sulfate, previously dried, as directed in the potassium bromide disk method under Infrared Spectrophotometry <2.25>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wave numbers.

(3) A solution of Guanethidine Sulfate (1 in 10) responds to the Qualitative Tests <1.09> for sulfate.

**pH** <2.54> Dissolve 1.0 g of Guanethidine Sulfate in 50 mL of water: the pH of the solution is between 4.7 and 5.7.

**Purity** (1) Clarity and color of solution—Dissolve 1.0 g of Guanethidine Sulfate in 50 mL of water: the solution is clear and colorless.

(2) Methylisothiourrea sulfate—Dissolve 2.0 g of Guanethidine Sulfate in 80 mL of sodium hydroxide TS, and allow to stand for 10 minutes. Add 60 mL of hydrochloric acid, 2 g of sodium bromide and water to make 200 mL. Then, to this solution add 0.70 mL of 1/60 mol/L potassium bromate VS and 2 mL of zinc iodide-starch paste TS: a blue color develops.

(3) Heavy metals <1.07>—Proceed with 2.0 g of Guanethidine Sulfate according to Method 4, and perform the test. Prepare the control solution with 2.0 mL of Standard Lead Solution (not more than 10 ppm).

**Loss on drying** <2.41> Not more than 0.5% (1 g, 105°C,

4 hours).

**Residue on ignition** <2.44> Not more than 0.2% (1 g).

**Assay** Weigh accurately about 0.5 g of Guanethidine Sulfate, previously dried, dissolve in 2 mL of formic acid, add 70 mL of a mixture of acetic anhydride and acetic acid (100) (6:1), and titrate <2.50> with 0.1 mol/L perchloric acid VS (potentiometric titration). Perform a blank determination, and make any necessary correction.

Each mL of 0.1 mol/L perchloric acid VS  
= 29.64 mg of  $C_{10}H_{22}N_4 \cdot H_2SO_4$

**Containers and storage** Containers—Tight containers.  
Storage—Light-resistant.

## Adsorbed Habu-venom Toxoid

沈降はぶトキソイド

Adsorbed Habu-venom Toxoid is a liquid for injection containing habu toxoid prepared by treating toxic substances produced by habu (*Trimeresurus flavoviridis*) with formaldehyde by a method involving no appreciable loss of the immunogenicity and rendered insoluble by the addition of aluminum salt.

It conforms to the requirements of Adsorbed Habu-venom Toxoid in the Minimum Requirements for Biological Products.

**Description** Adsorbed Habu-venom Toxoid becomes a uniform whitish turbid liquid on shaking.

## Freeze-dried Habu Antivenom, Equine

乾燥はぶウマ抗毒素

Freeze-dried Habu Antivenom, Equine, is a preparation for injection which is dissolved before use.

It contains *Trimeresurus flavoviridis* antivenom in immunoglobulin of horse origin.

It conforms to the requirements of Freeze-dried Habu Anti-venom, Equine, in the Minimum Requirements for Biological Products.

**Description** Freeze-dried Habu Antivenom, Equine, becomes colorless or light yellow-brown, clear liquid or a slightly whitish turbid liquid on addition of solvent.

## Adsorbed Hepatitis B Vaccine

沈降 B 型肝炎ワクチン


Adsorbed Hepatitis B Vaccine is a liquid for injection prepared by adding an aluminum salt to a liquid containing a surface antigen of hepatitis B virus to make the HBs antigen insoluble.

It conforms to the requirements of Adsorbed Hepatitis B Vaccine in the Minimum Requirements for Biological Products.

**Description** Adsorbed Hepatitis B Vaccine becomes a homogeneous, whitish turbid liquid on shaking.

## Haloperidol

ハロペリドール



$C_{21}H_{23}ClFNO_2$ : 375.86

4-[4-(4-Chlorophenyl)-4-hydroxypiperidin-1-yl]-1-(4-fluorophenyl)butan-1-one [52-86-8]

Haloperidol, when dried, contains not less than 99.0% and not more than 101.0% of  $C_{21}H_{23}ClFNO_2$ .

**Description** Haloperidol occurs as white to pale yellow crystals or powder.

It is freely soluble in acetic acid (100), sparingly soluble in methanol, slightly soluble in 2-propanol and in ethanol (99.5), and practically insoluble in water.

**Identification** (1) Dissolve 30 mg of Haloperidol in 100 mL of 2-propanol. To 5 mL of the solution add 10 mL of 0.1 mol/L hydrochloric acid TS and 2-propanol to make 100 mL. Determine the absorption spectrum of the solution as directed under Ultraviolet-visible Spectrophotometry <2.24>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wavelengths.

(2) Determine the infrared absorption spectrum of Haloperidol as directed in the potassium bromide disk method under Infrared Spectrophotometry <2.25>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wave numbers.

**Melting point** <2.60> 149 – 153°C

**Purity** (1) Sulfate <1.14>—To 1.0 g of Haloperidol add 50 mL of water, shake, and filter. To 25 mL of the filtrate add 1 mL of dilute hydrochloric acid and water to make 50 mL, and perform the test using this solution as the test solution. Prepare the control solution with 0.50 mL of 0.005 mol/L sulfuric acid VS (not more than 0.048%).

(2) Heavy metals <1.07>—Proceed with 1.0 g of Haloperidol according to method 2, and perform the test. Prepare the control solution with 2.0 mL of Standard Lead Solution (not more than 20 ppm).

(3) Related substances—Dissolve 25 mg of Haloperidol in 50 mL of the mobile phase, and use this solution as the sample solution. Pipet 1 mL of the sample solution, add the mobile phase to make exactly 200 mL, and use this solution as the standard solution. Perform the test with exactly 10  $\mu$ L each of the sample solution and standard solution as directed under Liquid Chromatography <2.01> according to the fol-

lowing conditions, and determine each peak area by the automatic integration method: the area of the peak other than haloperidol is not larger than the peak area of haloperidol from the standard solution, and the total area of these peaks is not larger than 2 times the peak area of haloperidol from the standard solution. For this calculation, use the peak areas for the related substances, having the relative retention time of about 0.5, about 1.2 and about 2.6 with respect to haloperidol, after multiplying by their relative response factors, 0.75, 1.47 and 0.76, respectively.

**Operating conditions**—

**Detector:** An ultraviolet absorption photometer (wavelength: 220 nm).

**Column:** A stainless steel column 4.6 mm in inside diameter and 15 cm in length, packed with octadecylsilanized silica gel for liquid chromatography (5  $\mu$ m in particle diameter).

**Column temperature:** A constant temperature of about 40°C.

**Mobile phase:** Dissolve 2.95 g of trisodium citrate dihydrate in 900 mL of water, adjust to pH 3.5 with dilute hydrochloric acid, and add water to make 1000 mL. To 300 mL of this solution add 700 mL of methanol and 1.0 g of sodium lauryl sulfate.

**Flow rate:** Adjust the flow rate so that the retention time of haloperidol is about 9 minutes.

**Time span of measurement:** About 2 times as long as the retention time of haloperidol beginning after the solvent peak.

**System suitability**—

**Test for required detectability:** To exactly 5 mL of the standard solution add the mobile phase to make exactly 25 mL. Confirm that the peak area of haloperidol obtained with 10  $\mu$ L of this solution is equivalent to 15 to 25% of that with 10  $\mu$ L of the standard solution.

**System performance:** When the procedure is run with 10  $\mu$ L of the standard solution under the above operating conditions, the number of theoretical plates and the symmetry factor of the peak of haloperidol are not less than 4000 and not more than 2.0, respectively.

**System repeatability:** When the test is repeated 6 times with 10  $\mu$ L of the standard solution under the above operating conditions, the relative standard deviation of the peak area of haloperidol is not more than 2.0%.

**Loss on drying** <2.41> Not more than 0.5% (1 g, in vacuum, 60°C, phosphorus (V) oxide, 3 hours).

**Residue on ignition** <2.44> Not more than 0.1% (1 g).

**Assay** Weigh accurately about 0.6 g of haloperidol, previously dried, and dissolve in 40 mL of acetic acid (100), and titrate <2.50> with 0.1 mol/L perchloric acid VS (indicator: 1 drop of crystal violet TS). Perform a blank determination.

Each mL of 0.1 mol/L perchloric acid VS  
= 37.59 mg of  $C_{21}H_{23}ClFNO_2$

**Containers and storage** Containers—Tight containers.

Storage—Light-resistant.

## Haloperidol Tablets

ハロペリドール錠

Haloperidol Tablets contain not less than 93.0% and not more than 107.0% of the labeled amount of haloperidol ( $C_{21}H_{23}ClFNO_2$ ; 375.86).

**Method of preparation** Prepare as directed under Tablets, with Haloperidol.

**Identification** To pulverized Haloperidol Tablets, equivalent to 6 mg of Haloperidol according to the labeled amount, add 70 mL of 2-propanol, and heat on a water bath until to boiling while shaking. After cooling, add 2-propanol to make 100 mL, and centrifuge. To 5 mL of the supernatant liquid add 2 mL of 0.1 mol/L hydrochloric acid TS and 2-propanol to make 20 mL. Determine the absorption spectrum of the solution as directed under Ultraviolet-visible Spectrophotometry <2.24>: it exhibits maxima between 219 nm and 223 nm and between 243 nm and 247 nm.

**Uniformity of dosage units** <6.02> Perform the test according to the following method: it meets the requirement of the Content uniformity test.

To 1 tablet of Haloperidol Tablets add 5 mL of the mobile phase, disperse the particle with the aid of ultrasonic waves, add 30 mL of the mobile phase, and extract for 30 minutes with the aid of ultrasonic waves with occasional shaking. Shake for more 30 minutes, and add the mobile phase to make exactly 50 mL. Centrifuge the solution, pipet  $V$  mL of the supernatant liquid, equivalent to about 0.3 mg of haloperidol ( $C_{21}H_{23}ClFNO_2$ ), add exactly 2 mL of the internal standard solution and the mobile phase to make 25 mL, and use this solution as the sample solution. Separately, weigh accurately about 20 mg of haloperidol for assay, previously dried in vacuum at 60°C for 3 hours on phosphorus (V) oxide, and dissolve in the mobile phase to make exactly 100 mL. Pipet 15 mL of this solution, and add the mobile phase to make exactly 50 mL. Pipet 5 mL of this solution, add exactly 2 mL of the internal standard solution and the mobile phase to make 25 mL, and use this solution as the standard solution. Perform the test with 10  $\mu$ L each of the sample solution and standard solution as directed under Liquid Chromatography <2.01> according to the following conditions, and determine the ratios,  $Q_T$  and  $Q_S$ , of the peak area of haloperidol to that of the internal standard.

$$\begin{aligned} &\text{Amount (mg) of haloperidol (C}_{21}\text{H}_{23}\text{ClFNO}_2\text{)} \\ &= W_S \times (Q_T/Q_S) \times (1/V) \times (3/4) \end{aligned}$$

$W_S$ : Amount (mg) of haloperidol for assay

**Internal standard solution**—A solution of diphenyl in the mobile phase (1 in 6700).

**Operating conditions**—

Proceed as detected in the operating condition in the Assay.

**System suitability**—

**System performance**: When the procedure is run with 10  $\mu$ L of the standard solution under the above operating conditions, haloperidol and diphenyl are eluted in this order with the resolution between these peaks being not less than 5.

**System repeatability**: When the test is repeated 6 times with 10  $\mu$ L of the standard solution under the above operating conditions, the relative standard deviation of the ratio of the peak area of haloperidol to that of the internal standard is not more than 1.0%.

**Dissolution** Being specified separately.

**Assay** Weigh accurately, and powder not less than 20 Haloperidol Tablets. Weigh accurately a portion of the powder, equivalent to about 10 mg of haloperidol ( $C_{21}H_{23}ClFNO_2$ ), add 10 mL of water, disperse the particle with the aid of ultrasonic waves, add exactly 20 mL of the internal standard solution, extract for 30 minutes with the aid of ultrasonic waves with occasional shaking, and add the mobile phase to make 100 mL. Centrifuge after shaking for more 30 minutes, and use the supernatant liquid as the sample solution. Separately, weigh accurately about 25 mg of haloperidol for assay, previously dried in vacuum at 60°C for 3 hours on phosphorus (V) oxide, and dissolve in methanol to make exactly 25 mL. Pipet 10 mL of this solution, add exactly 20 mL of the internal standard solution and the mobile phase to make exactly 100 mL, and use this solution as the standard solution. Perform the test with 10  $\mu$ L each of the sample solution and standard solution as directed under Liquid Chromatography <2.01> according to the following conditions, and determine the ratios,  $Q_T$  and  $Q_S$ , of the peak area of haloperidol to that of the internal standard.

$$\begin{aligned} &\text{Amount (mg) of haloperidol (C}_{21}\text{H}_{23}\text{ClFNO}_2\text{)} \\ &= W_S \times (Q_T/Q_S) \times (2/5) \end{aligned}$$

$W_S$ : Amount (mg) of haloperidol for assay

**Internal standard solution**—A solution of diphenyl in methanol (1 in 2000).

**Operating conditions**—

**Detector**: An ultraviolet absorption photometer (wavelength: 220 nm).

**Column**: A stainless steel column 4.6 mm in inside diameter and 15 cm in length, packed with octadecylsilanized silica gel for liquid chromatography (5  $\mu$ m in particle diameter).

**Column temperature**: A constant temperature of about 40°C.

**Mobile phase**: Dissolve 2.95 g of trisodium citrate dihydrate in 900 mL of water, adjust to pH 3.5 with dilute hydrochloric acid, and add water to make 1000 mL. To 250 mL of this solution add 750 mL of methanol and 1.0 g of sodium lauryl sulfate, and mix to dissolve.

**Flow rate**: Adjust the flow rate so that the retention time of haloperidol is about 9 minutes.

**System suitability**—

**System performance**: When the procedure is run with 10  $\mu$ L of the standard solution under the above operating conditions, haloperidol and diphenyl are eluted in this order with the resolution between these peaks being not less than 5.

**System repeatability**: When the test is repeated 6 times with 10  $\mu$ L of the standard solution under the above operating conditions, the relative standard deviation of the ratio of the peak area of haloperidol to that of the internal standard is not more than 1.0%.

**Containers and storage** Containers—Tight containers.

Storage—Light-resistant for the tablets without coating.

## Halothane

ハロタン

C<sub>2</sub>HBrClF<sub>3</sub>: 197.38(2*RS*)-2-Bromo-2-chloro-1,1,1-trifluoroethane [151-67-7]

Halothane contains not less than 0.008% and not more than 0.012% of Thymol as a stabilizer.

**Description** Halothane is a clear, colorless, and mobile liquid.

It is miscible with ethanol (95), with diethyl ether and with isooctane.

It is slightly soluble in water.

It is a volatile, nonflammable liquid, and setting fire to its heated vapor does not support combustion.

It is affected by light.

Refractive index  $n_D^{20}$ : 1.369 – 1.371

**Identification** Transfer about 3  $\mu$ L of Halothane to a gas cell having light path 10 cm in length, and determine the infrared absorption spectrum as directed in the gas sampling method under Infrared Spectrophotometry <2.25>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wave numbers.

**Specific gravity** <2.56>  $d_{20}^{20}$ : 1.872 – 1.877

**Purity (1)** Acidity or alkalinity—Shake 60 mL of Halothane with 60 mL of freshly boiled and cooled water vigorously for 3 minutes. Separate the water layer, and use this as the sample solution. To 20 mL of the sample solution add 1 drop of bromocresol purple TS and 0.10 mL of 0.01 mol/L sodium hydroxide VS: a red-purple color develops. To 20 mL of the sample solution add 1 drop of bromocresol purple TS and 0.6 mL of 0.01 mol/L hydrochloric acid VS: a yellow color is produced.

(2) Halide and halogen—To 5 mL of the sample solution obtained in (1) add 1 drop of nitric acid and 0.20 mL of silver nitrate TS: no turbidity is produced. To 10 mL of the sample solution obtained in (1) add 1 mL of potassium iodide TS and 2 drops of starch TS, and allow to stand for 5 minutes: no blue color develops.

(3) Phosgene—Transfer 50 mL of Halothane to a dried 300-mL conical flask, suspend a strip of phosgene test paper vertically inside the flask with the lower end about 10 mm above the surface of the liquid, insert the stopper, and allow to stand at a dark place for 20 to 24 hours: the test paper shows no yellow color.

(4) Residue on evaporation—Pipet 50 mL of Halothane, evaporate on a water bath, and dry the residue at 105°C for 2 hours: the mass of the residue is not more than 1.0 mg.

(5) Volatile related substances—To 100 mL of Halothane add exactly 5.0  $\mu$ L of the internal standard, and use this solution as the sample solution. Perform the test with 5  $\mu$ L of the sample solution as directed under Gas Chromatography <2.02>, and determine each peak area by the automatic integration method: the total area of the peaks other than those

of halothane and the internal standard is not larger than the peak area of the internal standard.

**Internal standard**—1,1,2-Trichloro-1,2,2-trifluoroethane.

**Operating conditions**—

Detector: A hydrogen flame-ionization detector.

Column: A column about 3 mm in inside diameter and 3 m in length, at the first 2 m from the injection port, having macrogol 400 coated in the ratio of 30% on siliceous earth for gas chromatography (180 to 250  $\mu$ m in particle diameter), and at the remaining 1 m, having dinonyl phthalate coated in the ratio of 30% on siliceous earth for gas chromatography (180 to 250  $\mu$ m in particle diameter).

Column temperature: A constant temperature of about 50°C

Carrier gas: Nitrogen

Flow rate: Adjust the flow rate so that the retention time of the internal standard is 2 to 3 minutes.

Selection of column: Mix 3 mL of Halothane and 1 mL of the internal standard. Proceed with 1  $\mu$ L of this solution under the above operating conditions, and calculate the resolution. Use a column giving elution of the internal standard and halothane in this order with the resolution between these peaks being not less than 10.

Detection sensitivity: Adjust the detection sensitivity so that the peak height of the internal standard obtained from 5  $\mu$ L of the sample solution composes 30 to 70% of the full scale.

Time span of measurement: About 3 times as long as the retention time of halothane.

**Distilling range** <2.57> Not less than 95 vol distills within a 1°C range between 49°C and 51°C.

**Thymol** To 0.50 mL of Halothane add 5.0 mL of isooctane and 5.0 mL of titanium (IV) oxide TS, shake vigorously for 30 seconds, and allow to stand: the separated upper layer has more color than the following control solution A, and has no more color than the following control solution B.


Control solution: Dissolve 0.225 g of thymol for assay in isooctane to make exactly 100 mL. To 10 mL each of this solution, accurately measured, add isooctane to make exactly 150 mL and 100 mL, respectively. Proceed with 0.50 mL each of these solutions in the same manner as Halothane, and use the separated upper layers so obtained as the control solution A and B, respectively.

**Containers and storage** Containers—Tight containers.

Storage—Light-resistant, and not exceeding 30°C.

## Haloxazolam

ハロキサゾラム

C<sub>17</sub>H<sub>14</sub>BrFN<sub>2</sub>O<sub>2</sub>: 377.21

(11*bRS*)-10-Bromo-11*b*-(2-fluorophenyl)-2,3,7,11*b*-tetrahydro[1,3]oxazolo[3,2-*d*][1,4]benzodiazepin-6(5*H*)-one [59128-97-1]



Haloxazolam, when dried, contains not less than 99.0% of  $C_{17}H_{14}BrFN_2O_2$ .

**Description** Haloxazolam occurs as white crystals or crystalline powder. It is odorless and tasteless.

It is freely soluble in acetic acid (100), sparingly soluble in acetonitrile, in methanol and in ethanol (99.5), slightly soluble in diethyl ether, and practically insoluble in water.

Melting point: about 183°C (with decomposition).

**Identification (1)** Dissolve 0.01 g of Haloxazolam in 10 mL of methanol, add 1 drop of hydrochloric acid: the solution shows a yellow-green fluorescence under ultraviolet light (main wavelength: 365 nm). To this solution add 1 mL of sodium hydroxide TS: the fluorescence disappears immediately.

**(2)** Prepare the test solution with 0.05 g of Haloxazolam as directed under Oxygen Flask Combustion Method <1.06>, using a mixture of 20 mL of dilute sodium hydroxide TS and 1 mL of hydrogen peroxide (30) as an absorbing liquid: the test solution responds to the Qualitative Tests <1.09> for bromide and for fluoride.

**(3)** Determine the absorption spectrum of a solution of Haloxazolam in methanol (1 in 100,000) as directed under Ultraviolet-visible Spectrophotometry <2.24>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wavelengths.

**(4)** Determine the infrared absorption spectrum of Haloxazolam, previously dried, as directed in the potassium bromide disk method under Infrared Spectrophotometry <2.25>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wave numbers.

**Absorbance** <2.24>  $E_{1cm}^{1\%}$  (247 nm): 390 – 410 (10 mg, methanol, 1000 mL).

**Purity (1)** Clarity and color of solution—Dissolve 0.10 g of Haloxazolam in 20 mL of ethanol (99.5): the solution is clear and colorless.

**(2)** Soluble halides—To 1.0 g of Haloxazolam add 50 mL of water, allow to stand for 1 hour with occasional shaking, and filter. To 25 mL of the filtrate add 6 mL of dilute nitric acid and water to make 50 mL. Perform the test with this solution as directed under Chloride Limit Test <1.03>. Prepare the control solution with 0.10 mL of 0.01 mol/L hydrochloric acid VS.

**(3)** Heavy metals <1.07>—Proceed with 1.0 g of Haloxazolam according to Method 2, and perform the test. Prepare the control solution with 2.0 mL of Standard Lead Solution (not more than 20 ppm).

**(4)** Arsenic <1.11>—To 1.0 g of Haloxazolam in a decomposition flask add 5 mL of nitric acid and 2 mL of sulfuric acid, place a small funnel on the mouth of the flask, and heat carefully until white fumes are evolved. After cooling, add 2 mL of nitric acid, heat, repeat this procedure twice, add several 2-mL portions of hydrogen peroxide (30), and heat until the solution is colorless to pale yellow. After cooling, add 2 mL of a saturated solution of ammonium oxalate monohydrate, and heat until white fumes are evolved. After cooling, add water to make 5 mL, and perform the test with this solution: the solution has no more color than the following control solution (not more than 2 ppm).

Control solution: Proceed in the same manner as above

without using Haloxazolam, add 2.0 mL of Standard Arsenic Solution and water to make 5 mL, and proceed in the same manner as the test solution.

**(5)** Related substances—Dissolve 0.10 g of Haloxazolam in 100 mL of acetonitrile, and use this solution as the sample solution. Pipet 1 mL of the sample solution, add acetonitrile to make exactly 100 mL, and use this solution as the standard solution. Perform the test with exactly 10  $\mu$ L each of the sample solution and standard solution as directed under Liquid Chromatography <2.01> according to the following conditions. Determine each peak area of both solutions by the automatic integration method: the total area of all peaks other than the area of the haloxazolam from the sample solution is not larger than the peak area of the haloxazolam from the standard solution.

**Operating conditions—**

**Detector:** An ultraviolet absorption photometer (wavelength: 250 nm).

**Column:** A stainless steel column 4.6 mm in inside diameter and 15 cm in length, packed with octadecylsilanized silica gel for liquid chromatography (5  $\mu$ m in particle diameter).

**Column temperature:** A constant temperature of about 25°C.

**Mobile phase:** Dissolve 6.2 g of boric acid and 7.5 g of potassium chloride in 900 mL of water, adjust the pH with triethylamine to 8.5, and add water to make 1000 mL. To 300 mL of this solution add 200 mL of acetonitrile.

**Flow rate:** Adjust the flow rate so that the retention time of haloxazolam is about 10 minutes.

**Time span of measurement:** About 3 times as long as the retention time of haloxazolam beginning after the solvent peak.

**System suitability—**

**Test for required detection:** To exactly 5 mL of the standard solution add acetonitrile to make exactly 50 mL. Confirm that the peak area of haloxazolam obtained from 10  $\mu$ L of this solution is equivalent to 8 to 12% of that of haloxazolam obtained from 10  $\mu$ L of the standard solution.

**System performance:** Dissolve 10 mg each of Haloxazolam and cloxazolam in 200 mL of acetonitrile. When the procedure is run with 10  $\mu$ L of this solution under the above operating conditions, haloxazolam and cloxazolam are eluted in this order with the resolution between these peaks being not less than 1.5.

**System repeatability:** When the test is repeated 6 times with 10  $\mu$ L of the standard solution under the above operating conditions, the relative standard deviation of the peak area of haloxazolam is not more than 1.0%.

**Loss on drying** <2.41> Not more than 0.5% (1 g, 105°C, 3 hours).

**Residue on ignition** <2.44> Not more than 0.1% (1 g, platinum crucible).

**Assay** Weigh accurately about 0.5 g of Haloxazolam, previously dried, dissolve in 50 mL of acetic acid (100), and titrate <2.50> with 0.1 mol/L perchloric acid VS (potentiometric titration). Perform a blank determination, and make any necessary correction.

Each mL of 0.1 mol/L perchloric acid VS  
= 37.72 mg of  $C_{17}H_{14}BrFN_2O_2$

**Containers and storage** Containers—Tight containers.

Storage—Light-resistant.

## Heparin Sodium

ヘパリンナトリウム

Heparin Sodium is obtained from the livers, the lungs and the intestinal mucosa of healthy edible animals, and prolongs the clotting time of blood. Heparin Sodium obtained from the livers and the lungs contains not less than 110 Heparin Units per mg, and that obtained from the intestinal mucosa contains not less than 130 Heparin Units per mg.

Heparin Sodium, calculated on the dried basis, contains not less than 90% and not more than 110% of the labeled Units.

Label the name of the organ used as the starting material.

**Description** Heparin Sodium occurs as a white to grayish brown powder or grains. It is odorless.

It is soluble in water, and practically insoluble in ethanol (95) and in diethyl ether.

It is hygroscopic.

**pH** <2.54> The pH of a solution of 1.0 g of Heparin Sodium in 100 mL of water is between 6.0 and 8.0.

**Purity** (1) Clarity and color of solution—Dissolve 0.5 g of Heparin Sodium in 20 mL of water: the solution is clear and colorless to light yellow.

(2) Barium—Dissolve 0.03 g of Heparin Sodium in 3.0 mL of water, and use this solution as the sample solution. To 1.0 mL of the sample solution add 3 drops of dilute sulfuric acid, and allow to stand for 10 minutes: no turbidity is produced.

(3) Total nitrogen—Weigh accurately about 0.1 g of Heparin Sodium, previously dried at 60°C for 3 hours under reduced pressure, and perform the test as directed under Nitrogen Determination <1.08>: the amount of nitrogen (N: 14.01) is not more than 3.0%.

(4) Protein—To 1.0 mL of the sample solution obtained in (2) add 5 drops of a solution of trichloroacetic acid (1 in 5): neither a precipitate nor turbidity is produced.

**Loss on drying** <2.41> Not more than 10% (20 mg, in vacuum, 60°C, 3 hours).

**Residue on ignition** <2.44> Not more than 40% (after drying, 20 mg).

**Pyrogen** <4.04> Dissolve Heparin Sodium in isotonic sodium chloride solution so as to contain 1000 Units per mL according to the labeled Units. Inject into rabbits 2 mL of this solution per kg: it meets the requirement.

**Assay** (i) Substrate solution: Dissolve 15 mg of *N*-benzoyl-L-isoleucyl-L-glutamyl( $\gamma$ -OR)-glycyl-L-arginyl-*p*-nitroanilide hydrochloride in 20 mL of water.

(ii) Antithrombin III solution: Dissolve human antithrombin III in water to make a solution containing 1 Unit per mL.

(iii) Activated blood coagulation factor X solution: Dissolve bovine activated blood coagulation factor X in water to

make a solution containing 0.426 Units per mL.

(iv) Buffer solution: Dissolve 6.06 g of 2-amino-2-hydroxymethyl-1,3-propanediol in 750 mL of water, adjust the pH to 8.4 with 1 mol/L hydrochloric acid TS, and add water to make 1000 mL.

(v) Reaction stop solution: To 20 mL of acetic acid (100) add water to make 40 mL.

(vi) Heparin standard solutions: Dissolve Heparin Sodium Reference Standard in isotonic sodium chloride solution to make a solution containing 10 Units per mL, and use as the standard stock solution. To 100  $\mu$ L of the standard stock solution add the buffer solution to make exactly 5 mL, and use this solution as the standard solution. Prepare the heparin standard solutions (1), (2), (3), (4) and (5) by addition of antithrombin III solution, human normal plasma and the buffer solution to the standard solution as directed in the following table.

| Heparin standard solution No. | Heparin concentration (Unit/mL) | Buffer solution ( $\mu$ L) | Antithrombin III solution ( $\mu$ L) | Human normal plasma ( $\mu$ L) | Standard solution ( $\mu$ L) |
|-------------------------------|---------------------------------|----------------------------|--------------------------------------|--------------------------------|------------------------------|
|                               |                                 |                            |                                      |                                |                              |
| (1)                           | 0                               | 800                        | 100                                  | 100                            | 0                            |
| (2)                           | 0.02                            | 700                        | 100                                  | 100                            | 100                          |
| (3)                           | 0.04                            | 600                        | 100                                  | 100                            | 200                          |
| (4)                           | 0.06                            | 500                        | 100                                  | 100                            | 300                          |
| (5)                           | 0.08                            | 400                        | 100                                  | 100                            | 400                          |

(vii) Sample solution: Weigh accurately an adequate amount of Heparin Sodium, dissolve in isotonic sodium chloride solution so that each mL contains about 0.5 Units according to the labeled amount. To 100  $\mu$ L of this solution add 100  $\mu$ L of antithrombin III solution, 100  $\mu$ L of human normal plasma and 700  $\mu$ L of the buffer solution, and use this solution as the sample solution.

(viii) Procedure: Transfer 400  $\mu$ L of the sample solution to a test tube, and warm at 37°C for 4 minutes. Add 200  $\mu$ L of the activated blood coagulation factor X solution, mix well, warm at 37°C for exactly 30 seconds, add 400  $\mu$ L of the substrate solution, previously warmed at 37°C, and mix well. Allow the tube to stand at 37°C for exactly 3 minutes, add 600  $\mu$ L of the reaction stop solution, mix immediately, and determine the absorbance at 405 nm, using the blank solution prepared by addition of 600  $\mu$ L of the reaction stop solution and 600  $\mu$ L of water to 400  $\mu$ L of the sample solution. Proceed the same way with the heparin standard solution (1), the heparin standard solution (2), the heparin standard solution (3), the heparin standard solution (4) and the heparin standard solution (5), and determine their absorbances.

(ix) Calculation: Plot the absorbances of the standard solutions on the vertical axis and their heparin concentrations on the horizontal axis to prepare a calibration curve. Determine the heparin concentration, *C*, of the sample solution from its absorbance by using the curve, and calculate heparin Units per mg of Heparin Sodium from the following formula.

$$\text{Units per mg of Heparin Sodium} = C \times 10 \times (b/a)$$

*a*: Amount of sample (mg)

*b*: Total volume (mL) of isotonic sodium chloride solution used to dissolve the sample to make the solution containing about 0.5 Units per mL

**Containers and storage** Containers—Tight containers.

## Heparin Sodium Injection

ヘパリンナトリウム注射液

Heparin Sodium Injection is an aqueous solution for injection. It contains not less than 90% and not more than 110% of the labeled heparin Units.

Label the name of organ used as the starting material of Heparin Sodium supplied for preparing Heparin Sodium Injection.

**Method of preparation** Dissolve Heparin Sodium in Isotonic Sodium Chloride Solution and prepare as directed under Injections.

**Description** Heparin Sodium Injection is a clear, colorless to light yellow liquid.

**pH** <2.54> 5.5 – 8.0

**Purity** (1) Barium—Measure exactly a portion of Heparin Sodium Injection, equivalent to 3000 Units of Heparin Sodium according to the labeled Unit. Add water to make 3.0 mL and use this solution as the sample solution. To 1.0 mL of the sample solution add 3 drops of dilute sulfuric acid, and allow to stand for 10 minutes: no turbidity is produced.

(2) Protein—Proceed as directed in the Purity (4) under Heparin Sodium.

**Bacterial endotoxins** <4.01> Less than 0.0030 EU/unit.

**Extractable volume** <6.05> It meets the requirement.

**Foreign insoluble matter** Perform the test according to Method 1: it meets the requirement.

**Insoluble particulate matter** Perform the test according to Method 1: it meets the requirement.

**Sterility** Perform the test according to the Membrane filtration method: it meets the requirement.

**Assay** Proceed as directed in the Assay under Heparin Sodium, replacing the sample solution indicated in (vii) and the calculation in (ix) with the following.

Sample solution: Measure exactly an adequate portion of Heparin Sodium Injection according to the labeled Units, dilute it with isotonic sodium chloride solution so that each mL contains about 0.5 Units. To 100  $\mu$ L of this solution add 100  $\mu$ L of antithrombin III solution, 100  $\mu$ L of human normal plasma and 700  $\mu$ L of the buffer solution, and use this solution as the sample solution.

Calculation: Plot the absorbances of the standard solutions on the vertical axis and their heparin concentrations on the horizontal axis to prepare a calibration curve. Determine the heparin concentration, *C*, of the sample solution from its absorbance by using the curve, and calculate heparin Units per mL of Heparin Sodium Injection from the following formula.

$$\begin{aligned} &\text{Units per mL of Heparin Sodium Injection} \\ &= C \times 10 \times (b/a) \end{aligned}$$


*a*: Amount of sample (mL)

*b*: Total volume (mL) of isotonic sodium chloride solution used to dilute the sample to make the solution containing about 0.5 Units per mL

**Containers and storage** Containers—Hermetic containers.  
Storage—Light-resistant.

## Homatropine Hydrobromide

ホマトロピン臭化水素酸塩



$C_{16}H_{21}NO_3 \cdot HBr$ : 356.25

(1*R*,3*r*,5*S*)-8-Methyl-8-azabicyclo[3.2.1]oct-3-yl [(2*RS*)-2-hydroxy-2-phenyl]acetate monohydrobromide [51-56-9]

Homatropine Hydrobromide contains not less than 99.0% of  $C_{16}H_{21}NO_3 \cdot HBr$ , calculated on the dried basis.

**Description** Homatropine Hydrobromide occurs as white crystals or crystalline powder. It is odorless.

It is freely soluble in water, sparingly soluble in ethanol (95), slightly soluble in acetic acid (100), very slightly soluble in acetic anhydride, and practically insoluble in diethyl ether.

It is affected by light.

Melting point: about 214°C (with decomposition).

**Identification** (1) To 5 mL of a solution of Homatropine Hydrobromide (1 in 20) add 2 to 3 drops of iodine TS: a brown precipitate is produced.

(2) Dissolve 0.05 g of Homatropine Hydrobromide in 5 mL of water, and add 3 mL of 2,4,6-trinitrophenol TS: a yellow precipitate is produced. Filter the precipitate, wash with five 10-mL portions of water, and dry at 105°C for 2 hours: it melts <2.60> between 184°C and 187°C.

(3) A solution of Homatropine Hydrobromide (1 in 20) responds to the Qualitative Tests <1.09> for bromide.

**Purity** (1) Acidity—Dissolve 1.0 g of Homatropine Hydrobromide in 20 mL of water, and add 0.40 mL of 0.01 mol/L sodium hydroxide VS and 1 drop of methyl red-methylene blue TS: a green color develops.

(2) Atropine, hyoscyamine and scopolamine—To 10 mg of Homatropine Hydrobromide add 5 drops of nitric acid, evaporate on a water bath to dryness, and cool. Dissolve the residue in 1 mL of *N,N*-dimethylformamide, and add 5 to 6 drops of tetraethylammonium hydroxide TS: no red-purple color is produced.

(3) Related substances—Dissolve 0.15 g of Homatropine Hydrobromide in 3 mL of water, and use this solution as the sample solution.

(i) To 1 mL of the sample solution add 2 to 3 drops of tannic acid TS: no precipitate is produced.

(ii) To 1 mL of the sample solution add 2 to 3 drops each

of dilute hydrochloric acid and platinic chloride TS: no precipitate is produced.

**Loss on drying** <2.41> Not more than 1.5% (0.5 g, 105°C, 2 hours).

**Residue on ignition** <2.44> Not more than 0.2% (0.2 g).


**Assay** Dissolve by warming about 0.4 g of Homatropine Hydrobromide in 60 mL of a mixture of acetic anhydride and acetic acid (100) (7:3). Cool, and titrate <2.50> with 0.1 mol/L perchloric acid VS (potentiometric titration). Perform a blank determination.

Each mL of 0.1 mol/L perchloric acid VS  
= 35.63 mg of  $C_{19}H_{23}ClN_2 \cdot 2HCl$

**Containers and storage** Containers—Tight containers.  
Storage—Light-resistant.

## Homochlorcyclizine Hydrochloride

ホモクロルシクリジン塩酸塩



$C_{19}H_{23}ClN_2 \cdot 2HCl$ ; 387.77

1-[(*RS*)-(4-Chlorophenyl)(phenyl)methyl]-  
4-methylhexahydro-1*H*-1,4-diazepine dihydrochloride  
[1982-36-1]

Homochlorcyclizine Hydrochloride, when dried, contains not less than 98.0% of  $C_{19}H_{23}ClN_2 \cdot 2HCl$ .

**Description** Homochlorcyclizine Hydrochloride occurs as white to pale brown, crystals or powder.

It is very soluble in water, freely soluble in acetic acid (100), slightly soluble in ethanol (99.5), and very slightly soluble in acetonitrile and in acetic anhydride.

It dissolves in 0.1 mol/L hydrochloric acid TS.

It is hygroscopic.

It is colored slightly by light.

A solution of Homochlorcyclizine Hydrochloride (1 in 10) shows no optical rotation.

Melting point: about 227°C (with decomposition).

**Identification** (1) Determine the absorption spectrum of a solution of Homochlorcyclizine Hydrochloride in 0.1 mol/L hydrochloric acid TS (1 in 100,000) as directed under Ultraviolet-visible Spectrophotometry <2.24>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wavelengths.

(2) Determine the infrared absorption spectrum of Homochlorcyclizine Hydrochloride, previously dried, as directed in the potassium chloride disk method under Infrared Spectrophotometry <2.25>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wave numbers.

(3) A solution of Homochlorcyclizine Hydrochloride (1 in 100) responds to the Qualitative Tests <1.09> for chloride.

**Purity** (1) Heavy metals <1.07>—Proceed with 1.0 g of

Homochlorcyclizine Hydrochloride according to Method 2, and perform the test. Prepare the control solution with 2.0 mL of Standard Lead Solution (not more than 20 ppm).

(2) Related substances—Dissolve 0.10 g of Homochlorcyclizine Hydrochloride in 100 mL of the mobile phase, and use this solution as the sample solution. Measure exactly 1 mL of this solution, add the mobile phase to make exactly 100 mL, and use this solution as the standard solution. Perform the test with exactly 10  $\mu$ L each of the sample solution and standard solution as directed under Liquid Chromatography <2.01> according to the following conditions, and determine the peak areas by the automatic integration method: the areas of the peaks other than homochlorcyclizine obtained from the sample solution are not more than 1/2 times the peak area of homochlorcyclizine from the standard solution, and the total area of the peaks other than homochlorcyclizine from the sample solution is not more than the peak area of homochlorcyclizine from the standard solution.

**Operating conditions**—

**Detector:** An ultraviolet absorption photometer (wavelength: 223 nm).

**Column:** A stainless steel column 4.6 mm in inside diameter and 25 cm in length, packed with octadecylsilanized silica gel for liquid chromatography (5  $\mu$ m in particle diameter).

**Column temperature:** A constant temperature of about 40°C.

**Mobile phase:** A mixture of water, acetonitrile and perchloric acid (134:66:1).

**Flow rate:** Adjust the flow rate so that the retention time of homochlorcyclizine is about 10 minutes.

**Time span of measurement:** About 2 times as long as the retention time of homochlorcyclizine.

**System suitability**—

**Test for required detection:** To exactly 5 mL of the standard solution add the mobile phase to make exactly 50 mL. Confirm that the peak area of homochlorcyclizine obtained from 10  $\mu$ L of this solution is equivalent to 7 to 13% of that of homochlorcyclizine obtained from 10  $\mu$ L of the standard solution.

**System performance:** Dissolve 5 mg each of Homochlorcyclizine Hydrochloride and methyl parahydroxybenzoic acid in 100 mL of the mobile phase. When the procedure is run with 10  $\mu$ L of this solution under the above operating conditions, methyl parahydroxybenzoic acid and homochlorcyclizine are eluted in this order with the resolution between these peaks being not less than 5.

**System repeatability:** When the test is repeated 6 times with 10  $\mu$ L of the standard solution under the above operating conditions, the relative standard deviation of the peak area of homochlorcyclizine is not more than 1.0%.

**Loss on drying** <2.41> Not more than 2.0% (1 g, 110°C, 4 hours).

**Residue on ignition** <2.44> Not more than 0.2% (1 g).

**Assay** Weigh accurately about 0.3 g of Homochlorcyclizine Hydrochloride, previously dried, dissolve in 50 mL of a mixture of acetic anhydride and acetic acid (100) (7:3), and titrate <2.50> with 0.1 mol/L perchloric acid VS (potentiometric titration). Perform a blank determination.

Each mL of 0.1 mol/L perchloric acid VS

= 19.39 mg of  $C_{19}H_{23}ClN_2 \cdot 2HCl$

**Containers and storage** Containers—Tight containers.  
Storage—Light-resistant.

## Human Normal Immunoglobulin

人免疫グロブリン


Human Normal Immunoglobulin is a liquid for injection containing immunoglobulin G in serum globulins of humans.

It conforms to the requirements of Human Normal Immunoglobulin in the Minimum Requirements for Biological Products.

**Description** Human Normal Immunoglobulin is a clear, colorless or yellow-brown liquid.

## Hydralazine Hydrochloride

ヒドララジン塩酸塩



$C_8H_8N_4 \cdot HCl$ : 196.64

Phthalazin-1-ylhydrazine monohydrochloride [304-20-1]

Hydralazine Hydrochloride, when dried, contains not less than 98.0% of  $C_8H_8N_4 \cdot HCl$ .

**Description** Hydralazine Hydrochloride occurs as a white, crystalline powder. It is odorless, and has a bitter taste.

It is soluble in water, slightly soluble in ethanol (95), and practically insoluble in diethyl ether.

Melting point: about 275°C (with decomposition).

**Identification (1)** Determine the absorption spectrum of a solution of Hydralazine Hydrochloride (1 in 100,000) as directed under Ultraviolet-visible Spectrophotometry <2.24>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wavelengths.

(2) Determine the infrared absorption spectrum of Hydralazine Hydrochloride, previously dried, as directed in the potassium bromide disk method under Infrared Spectrophotometry <2.25>, and compare the spectrum with the Reference spectrum: both spectra exhibit similar intensities of absorption at the same wave numbers.

(3) A solution of Hydralazine Hydrochloride (1 in 4000) responds to the Qualitative Tests <1.09> for chloride.

**pH** <2.54> Dissolve 1.0 g of Hydralazine Hydrochloride in 50 mL of water: the pH of the solution is between 3.5 and 4.5.

**Purity (1)** Clarity and color of solution—Dissolve 1.0 g of Hydralazine Hydrochloride in 50 mL of water: the solution is clear, and colorless or pale yellow.

(2) Heavy metals <1.07>—Proceed with 1.0 g of Hydrala-

zine Hydrochloride according to Method 2, and perform the test. Prepare the control solution with 2.0 mL of Standard Lead Solution (not more than 20 ppm).

**Loss on drying** <2.41> Not more than 0.5% (0.5 g, in vacuum, phosphorus (V) oxide, 8 hours).

**Residue on ignition** <2.44> Not more than 0.1% (1 g).

**Assay** Weigh accurately about 0.15 g of Hydralazine Hydrochloride, previously dried, transfer it to a glass-stoppered flask, dissolve in 25 mL of water, add 25 mL of hydrochloric acid, cool to room temperature, add 5 mL of chloroform, and titrate <2.50> with 0.05 mol/L potassium iodate VS while shaking until the purple color of the chloroform layer disappears. The end point is reached when the red-purple color no more reappears in the chloroform layer within 5 minutes after the layer has been decolorized.

Each mL of 0.05 mol/L potassium iodate VS  
= 9.832 mg of  $C_8H_8N_4 \cdot HCl$

**Containers and storage** Containers—Tight containers.

## Hydralazine Hydrochloride for Injection

注射用ヒドララジン塩酸塩

Hydralazine Hydrochloride for Injection is a preparation for injection which is dissolved before use. It contains not less than 99% and not more than 113% of the labeled amount of hydralazine hydrochloride ( $C_8H_8N_4 \cdot HCl$ : 196.64).

**Method of preparation** Prepare as directed under Injections, with Hydralazine Hydrochloride.

**Description** Hydralazine Hydrochloride for Injection occurs as a white to pale yellow powder or mass. It is odorless, and has a bitter taste.

**Identification** Determine the absorption spectrum of a solution of Hydralazine Hydrochloride for Injection (1 in 100,000) as directed under Ultraviolet-visible Spectrophotometry <2.24>: it exhibits maxima between 238 nm and 242 nm, between 258 nm and 262 nm, between 301 nm and 305 nm, and between 313 nm and 317 nm.

**pH** <2.54> Dissolve 1.0 g of Hydralazine Hydrochloride for Injection in 50 mL of water: the pH of this solution is between 3.5 and 4.5.

**Assay** Weigh accurately the contents of not less than 10 samples of Hydralazine Hydrochloride for Injection. Weigh accurately about 0.15 g of the contents, transfer it to a glass-stoppered flask, dissolve in 25 mL of water, add 25 mL of hydrochloric acid, cool to room temperature, and proceed as directed in the Assay under Hydralazine Hydrochloride.

Each mL of 0.05 mol/L potassium iodate VS  
= 9.832 mg of  $C_8H_8N_4 \cdot HCl$

**Containers and storage** Containers—Hermetic containers.

## Hydralazine Hydrochloride Powder

ヒドララジン塩酸塩散

Hydralazine Hydrochloride Powder contains not less than 95% and not more than 105% of the labeled amount of hydralazine hydrochloride ( $\text{C}_8\text{H}_8\text{N}_4\cdot\text{HCl}$ ; 196.64).

**Method of preparation** Prepare as directed under Powder, with Hydralazine Hydrochloride.

**Identification** Weigh a portion of Hydralazine Hydrochloride Powder, equivalent to 25 mg of Hydralazine Hydrochloride according to the labeled amount, add 100 mL of water, shake well, and filter, if necessary. Add water to 2 mL of the filtrate to make 50 mL and determine the absorption spectrum of this solution as directed under Ultraviolet-visible Spectrophotometry <2.24>: it exhibits maxima between 238 nm and 242 nm, between 258 nm and 262 nm, between 301 nm and 305 nm, and between 313 nm and 317 nm.

**Assay** Weigh accurately a portion of Hydralazine Hydrochloride Powder, equivalent to about 0.15 g of Hydralazine Hydrochloride, transfer it to a glass-stoppered flask, add 25 mL of water, shake well, add 25 mL of hydrochloric acid, cool to room temperature, and proceed as directed in the Assay under Hydralazine Hydrochloride.

Each mL of 0.05 mol/L potassium iodate VS  
= 9.832 mg of  $\text{C}_8\text{H}_8\text{N}_4\cdot\text{HCl}$

**Containers and storage** Containers—Tight containers.

## Hydralazine Hydrochloride Tablets

ヒドララジン塩酸塩錠

Hydralazine Hydrochloride Tablets contain not less than 95% and not more than 105% of the labeled amount of hydralazine hydrochloride ( $\text{C}_8\text{H}_8\text{N}_4\cdot\text{HCl}$ ; 196.64).

**Method of preparation** Prepare as directed under Tablets, with Hydralazine Hydrochloride.

**Identification** Weigh a quantity of powdered Hydralazine Hydrochloride Tablets, equivalent to 25 mg of Hydralazine Hydrochloride according to the labeled amount, add 100 mL of water, mix well, and filter if necessary. To 2 mL of this solution add water to make 50 mL, and determine the absorption spectrum of this solution as directed under Ultraviolet-visible Spectrophotometry <2.24>: it exhibits maxima between 238 nm and 242 nm, between 258 nm and 262 nm, between 301 nm and 305 nm and between 313 nm and 317 nm.

**Dissolution** <6.10> Perform the test according to the following method: it meets the requirement.

Perform the test with 1 tablet of Hydralazine Hydrochloride Tablets at 50 revolutions per minute according to the Paddle method, using 900 mL of water as the test solution. Take 30 mL or more of the dissolved solution 45 minutes af-

ter start of the dissolution test, and filter through a membrane filter with pore size of not more than  $0.8\ \mu\text{m}$ . Discard the first 10 mL of the filtrate, pipet the subsequent  $V$  mL, add water to make exactly  $V'$  mL so that each mL contains about  $11\ \mu\text{g}$  of hydralazine hydrochloride ( $\text{C}_8\text{H}_8\text{N}_4\cdot\text{HCl}$ ) according to the labeled amount, and use this solution as the sample solution. Separately, weigh accurately about 0.05 g of hydralazine hydrochloride for assay, previously dried at  $105^\circ\text{C}$  for 3 hours, and dissolve in water to make exactly 50 mL. Pipet 1 mL of this solution, add water to make exactly 100 mL, and use this solution as the standard solution. Determine the absorbances,  $A_T$  and  $A_S$ , of the sample solution and the standard solution at 260 nm as directed under Ultraviolet-visible Spectrophotometry <2.24>. The dissolution rate of Hydralazine Hydrochloride Tablets in 45 minutes is not less than 80%.

Dissolution rate (%) with respect to the labeled amount of hydralazine hydrochloride ( $\text{C}_8\text{H}_8\text{N}_4\cdot\text{HCl}$ )  
=  $W_S \times (A_T/A_S) \times (V'/V) \times (1/C) \times 18$

$W_S$ : Amount (mg) of hydralazine hydrochloride for assay.

$C$ : Labeled amount (mg) of hydralazine hydrochloride ( $\text{C}_8\text{H}_8\text{N}_4\cdot\text{HCl}$ ) in 1 tablet.

**Assay** Weigh accurately not less than 20 Hydralazine Hydrochloride Tablets, and powder. Weigh accurately a portion of the powder, equivalent to about 0.15 g of hydralazine hydrochloride ( $\text{C}_8\text{H}_8\text{N}_4\cdot\text{HCl}$ ), transfer it to a glass-stoppered flask, and proceed as directed in the Assay under Hydralazine Hydrochloride.

Each mL of 0.05 mol/L potassium iodate VS  
= 9.832 mg of  $\text{C}_8\text{H}_8\text{N}_4\cdot\text{HCl}$

**Containers and storage** Containers—Tight containers.

## Hydrochloric Acid

塩酸

Hydrochloric Acid contains not less than 35.0% and not more than 38.0% of hydrogen chloride ( $\text{HCl}$ ; 36.46).

**Description** Hydrochloric Acid is a colorless liquid having a pungent odor.

It is fuming but ceases to fume when it is diluted with 2 volumes of water.

Specific gravity  $d_{20}^{20}$ , about 1.18

**Identification** (1) Allow a glass stick wet with ammonia TS to come near the surface of Hydrochloric Acid: a remarkable white smoke evolves.

(2) A solution of Hydrochloric Acid (1 in 100) changes blue litmus paper to red, and responds to the Qualitative Tests <1.09> for chloride.

**Purity** (1) Sulfate <1.14>—To 15 mL of Hydrochloric Acid add water to make 50 mL, and use this solution as the sample solution. To 3.0 mL of the sample solution add 5 mL of water and 5 drops of barium chloride TS, and allow to stand for 1 hour: no turbidity is produced.

(2) Sulfite—To 3.0 mL of the sample solution obtained in (1) add 5 mL of water and 1 drop of iodine TS: the color of io-

dine TS does not disappear.

(3) Bromide or iodide—Place 10 mL of the sample solution obtained in (1) in a glass-stoppered test tube, add 1 mL of chloroform and 1 drop of 0.002 mol/L potassium permanganate VS, and shake well: the chloroform layer remains colorless.

(4) Bromine or chlorine—Place 10 mL of the sample solution obtained in (1) in a glass-stoppered test tube, add 5 drops of potassium iodide TS and 1 mL of chloroform, and shake for 1 minute: the chloroform layer remains free from a purple color.

(5) Heavy metals <1.07>—Evaporate 5 mL of Hydrochloric Acid on a water bath to dryness, and add 2 mL of dilute acetic acid and water to the residue to make 50 mL. Perform the test using this solution as the test solution. Prepare the control solution as follows: to 3.0 mL of Standard Lead Solution add 2 mL of dilute acetic acid and water to make 50 mL (not more than 5 ppm).

(6) Arsenic <1.11>—Prepare the test solution with 1.7 mL of Hydrochloric Acid according to Method 1, and perform the test (not more than 1 ppm).

(7) Mercury—Dilute 20 mL of Hydrochloric Acid with water to make exactly 100 mL, and use the solution as the sample solution. Perform the test with this sample solution as directed under Atomic Absorption Spectrophotometry <2.23> (cold vapor type). Place the sample solution in a sample bottle of the atomic absorption spectrophotometer, add 10 mL of tin (II) chloride-sulfuric acid TS, connect the bottle immediately to the spectrophotometer, circulate air, and determine the absorbance  $A_T$  of the sample solution after the recorder reading has risen rapidly, and becomes constant at a wavelength of 253.7 nm. On the other hand, to 8 mL of Standard Mercury Solution add water to make exactly 100 mL, and determine the absorbance  $A_S$  of the solution obtained by the same procedure as used for the sample solution:  $A_T$  is smaller than  $A_S$  (not more than 0.04 ppm).

**Residue on ignition** <2.44> Pipet 10 mL of Hydrochloric Acid, add 2 drops of sulfuric acid, evaporate to dryness, and ignite: not more than 1.0 mg of residue remains.

**Assay** Weigh accurately a glass-stoppered flask containing 20 mL of water, add about 3 mL of Hydrochloric Acid, and weigh accurately again. Dilute with 25 mL of water, and titrate <2.50> with 1 mol/L sodium hydroxide VS (indicator: 2 to 3 drops of methyl red TS).

Each mL of 1 mol/L sodium hydroxide VS  
= 36.46 mg of HCl

**Containers and storage** Containers—Tight containers.

## Dilute Hydrochloric Acid

希塩酸

Dilute Hydrochloric Acid contains not less than 9.5 w/v% and not more than 10.5 w/v% of hydrogen chloride (HCl: 36.46).

**Description** Dilute Hydrochloric Acid is a colorless liquid. It is odorless and has a strong acid taste.

Specific gravity  $d_{20}^{20}$ : about 1.05

**Identification** A solution of Dilute Hydrochloric Acid (1 in 30) changes blue litmus paper to red and responds to the Qualitative Tests <1.09> for chloride.

**Purity** (1) Sulfate—To 3.0 mL of Dilute Hydrochloric Acid add 5 mL of water and 5 drops of barium chloride TS, and allow to stand for 1 hour: no turbidity is produced.

(2) Sulfite—To 3.0 mL of Dilute Hydrochloric Acid add 5 mL of water and 1 drop of iodine TS: the color of iodine TS does not disappear.

(3) Bromide or iodide—Place 10 mL of Dilute Hydrochloric Acid in a glass-stoppered test tube, add 1 mL of chloroform and 1 drop of 0.002 mol/L potassium permanganate VS, and shake well: the chloroform layer remains colorless.

(4) Bromine or chlorine—Place 10 mL of Dilute Hydrochloric Acid in a glass-stoppered test tube, add 5 drops of potassium iodide TS and 1 mL of chloroform, and shake for 1 minute: the chloroform layer remains free from a purple color.

(5) Heavy metals <1.07>—Evaporate 9.5 mL of Dilute Hydrochloric Acid on a water bath to dryness, add 2 mL of dilute acetic acid and water to make 50 mL, and perform the test using this solution as the test solution. Prepare the control solution as follows: to 3.0 mL of Standard Lead Solution add 2 mL of dilute acetic acid and water to make 50 mL (not more than 3 ppm).

(6) Arsenic <1.11>—Prepare the test solution with 4.0 mL of Dilute Hydrochloric Acid according to Method 1, and perform the test (not more than 0.5 ppm).

(7) Mercury—Dilute 80 mL of Dilute Hydrochloric Acid with water to make exactly 100 mL, and use this solution as the sample solution. Perform the test with this solution according to the Atomic Absorption Spectrophotometry <2.23> (cold vapor type). Place the sample solution in a sample bottle of the atomic absorption spectrophotometer, add 10 mL of tin (II) chloride-sulfuric acid TS, connect the bottle immediately to the spectrophotometer, circulate air, and read the absorbance  $A_T$  of the sample solution after the recorder reading has risen rapidly and become constant at a wavelength of 253.7 nm. On the other hand, to 8 mL of Standard Mercury Solution add water to make exactly 100 mL, and read the absorbance  $A_S$  of the solution obtained by the same procedure as used for the sample solution:  $A_T$  is smaller than  $A_S$  (not more than 0.01 ppm).

**Residue on ignition** <2.44> Pipet 10 mL of Dilute Hydrochloric Acid, add 2 drops of sulfuric acid, evaporate to dryness, and ignite: the mass of the residue is not more than 1.0 mg.

**Assay** Measure exactly 10 mL of Dilute Hydrochloric Acid, and dilute with 20 mL of water. Titrate <2.50> with 1 mol/L sodium hydroxide VS (indicator: 2 to 3 drops of methyl red TS).

Each mL of 1 mol/L sodium hydroxide VS  
= 36.46 mg of HCl

**Containers and storage** Containers—Tight containers.

## Hydrochloric Acid Lemonade

塩酸リモナーデ

### Method of preparation

|                          |                       |
|--------------------------|-----------------------|
| Dilute Hydrochloric Acid | 5 mL                  |
| Simple Syrup             | 80 mL                 |
| Purified Water           | a sufficient quantity |
| To make 1000 mL          |                       |


Prepare before use as directed under Lemonades, with the above ingredients.

**Description** Hydrochloric Acid Lemonade is a clear, colorless liquid. It has a sweet, cool, acid taste.

**Containers and storage** Containers—Tight containers.

## Hydrochlorothiazide

ヒドロクロロチアジド



$C_7H_8ClN_3O_4S_2$ : 297.74

6-Chloro-3,4-dihydro-2H-1,2,4-benzothiadiazine-7-sulfonamide 1,1-dioxide [58-93-5]

Hydrochlorothiazide, when dried, contains not less than 99.0% of  $C_7H_8ClN_3O_4S_2$ .

**Description** Hydrochlorothiazide occurs as white crystals or crystalline powder. It is odorless, and has a slightly bitter taste.

It is freely soluble in acetone, sparingly soluble in acetonitrile, very slightly soluble in water and in ethanol (95), and practically insoluble in diethyl ether.

It dissolves in sodium hydroxide TS.

Melting point: about 267°C (with decomposition).

**Identification** (1) To 5 mg of Hydrochlorothiazide add 5 mL of disodium chlomotropate TS, and allow to stand for 5 minutes: a purple color develops.

(2) Fuse a mixture of 0.1 g of Hydrochlorothiazide and 0.5 g of sodium carbonate decahydrate cautiously: the gas evolved changes moistened red litmus paper to blue. After cooling, crush with a glass rod, add 10 mL of water, stir, and filter. To 4 mL of the filtrate add 2 drops of hydrogen peroxide (30), 5 mL of diluted hydrochloric acid (1 in 5) and 2 to 3 drops of barium chloride TS: a white precipitate is produced.

(3) To 4 mL of the filtrate obtained in (2) add 5 mL of dilute nitric acid and 3 drops of silver nitrate TS: a white precipitate is produced.

(4) Dissolve 12 mg of Hydrochlorothiazide in 100 mL of sodium hydroxide TS. Dilute 10 mL of the solution with water to make 100 mL. Determine the absorption spectrum of the solution as directed under Ultraviolet-visible Spec-

trophotometry <2.24>, and compare the spectrum with the Reference Spectrum or the spectrum of a solution of Hydrochlorothiazide Reference Standard prepared in the same manner as the sample solution: both spectra exhibit similar intensities of absorption at the same wavelengths.

**Purity** (1) Chloride <1.03>—Dissolve 1.0 g of Hydrochlorothiazide in 30 mL of acetone, add 6 mL of dilute nitric acid and water to make 50 mL, and perform the test using this solution as the test solution. Prepare the control solution as follows: to 1.0 mL of 0.01 mol/L hydrochloric acid VS add 30 mL of acetone, 6 mL of dilute nitric acid and water to make 50 mL (not more than 0.036%).

(2) Sulfate <1.14>—Dissolve 1.0 g of Hydrochlorothiazide in 30 mL of acetone, add 1 mL of dilute hydrochloric acid and water to make 50 mL, and perform the test using this solution as the test solution. Prepare the control solution as follows: to 1.0 mL of 0.005 mol/L sulfuric acid VS add 30 mL of acetone, 1 mL of dilute hydrochloric acid and water to make 50 mL (not more than 0.048%).

(3) Heavy metals <1.07>—Proceed with 1.0 g of Hydrochlorothiazide according to Method 2, and perform the test. Prepare the control solution with 2.0 mL of Standard Lead Solution (not more than 20 ppm).

(4) Primary aromatic amines—Dissolve 80 mg of Hydrochlorothiazide in acetone to make exactly 100 mL. Measure exactly 1 mL of the solution, add 3.0 mL of dilute hydrochloric acid, 3.0 mL of water and 0.15 mL of sodium nitrite TS, shake, and allow to stand for 1 minute. Shake this solution with 1.0 mL of ammonium amidosulfate TS, allow to stand for 3 minutes, then add 1.0 mL of *N*-(1-naphthyl)-*N'*-diethylethylenediamine oxalate TS, shake, and allow to stand for 5 minutes. Perform the test with this solution as directed under Ultraviolet-visible Spectrophotometry <2.24>, using a solution prepared with 1.0 mL of acetone in the same manner as the blank: the absorbance at 525 nm is not more than 0.10.

**Loss on drying** <2.41> Not more than 1.0% (1 g, 105°C, 2 hours).

**Residue on ignition** <2.44> Not more than 0.1% (1 g).

**Assay** Weigh accurately about 30 mg each of Hydrochlorothiazide and Hydrochlorothiazide Reference Standard, previously dried, and dissolve in 150 mL of the mobile phase, add exactly 10 mL each of the internal standard solution, then add the mobile phase to make 200 mL, and use these solutions as the sample solution and the standard solution, respectively. Perform the test with 20  $\mu$ L each of the sample solution and standard solution as directed under Liquid Chromatography <2.01> according to the following conditions, and calculate the ratios,  $Q_T$  and  $Q_S$ , of the peak area of hydrochlorothiazide to that of the internal standard.

$$\text{Amount (mg) of } C_7H_8ClN_3O_4S_2 = W_S \times (Q_T/Q_S)$$

$W_S$ : Amount (mg) of Hydrochlorothiazide Reference Standard

**Internal standard solution**—A solution of 4-aminoacetophenone in acetonitrile (9 in 2000).

**Operating conditions**—

Detector: An ultraviolet absorption photometer (wavelength: 254 nm).

Column: A stainless steel column 4.6 mm in inside di-



ameter and 25 cm in length, packed with octadecylsilanized silica gel for liquid chromatography (5  $\mu$ m in particle diameter).

Column temperature: A constant temperature of about 25°C.

Mobile phase: A mixture of 0.1 mol/L sodium dihydrogen phosphate TS, pH 3.0 and acetonitrile (9:1).

Flow rate: Adjust the flow rate so that the retention time of hydrochlorothiazide is about 10 minutes.

*System suitability—*


System performance: When the procedure is run with 20  $\mu$ L of the standard solution under the above operating conditions, hydrochlorothiazide and the internal standard are eluted in this order with the resolution between these peaks being not less than 4.

System repeatability: When the test is repeated 6 times with 20  $\mu$ L of the standard solution under the above operating conditions, the relative standard deviation of the ratios of the peak area of hydrochlorothiazide to that of the internal standard is not more than 1.0%.

**Containers and storage** Containers—Well-closed containers.

## Hydrocortisone

ヒドロコルチゾン



$C_{21}H_{30}O_5$ : 362.46

11 $\beta$ ,17,21-Trihydroxypregn-4-ene-3,20-dione [50-23-7]

Hydrocortisone, when dried, contains not less than 97.0% and not more than 102.0% of  $C_{21}H_{30}O_5$ .

**Description** Hydrocortisone occurs as a white, crystalline powder. It is odorless.

It is sparingly soluble in methanol, in ethanol (95) and in 1,4-dioxane, slightly soluble in chloroform, and very slightly soluble in diethyl ether and in water.

Melting point: 212 – 220°C (with decomposition).

**Identification (1)** Add 2 mL of sulfuric acid to 2 mg of Hydrocortisone: the solution shows a yellow-green fluorescence immediately, and the color of the solution changes gradually from orange to dark red. Dilute carefully the solution with 10 mL of water: the color changes through yellow to orange-yellow with green fluorescence, and a small amount of a flocculent precipitate is formed.

(2) Dissolve 0.01 g of Hydrocortisone in 1 mL of methanol, add 1 mL of Fehling's TS, and heat: a red precipitate is formed.

(3) Determine the infrared absorption spectrum of Hydrocortisone, previously dried, as directed in the potassium bromide disk method under Infrared Spectrophotometry <2.25>, and compare the spectrum with the Reference Spectrum or the spectrum of previously dried Hydrocortisone Reference Standard: both spectra exhibit similar intensities

of absorption at the same wave numbers. If any difference appears between the spectra, dissolve Hydrocortisone and Hydrocortisone Reference Standard in ethanol (95), respectively, then evaporate the ethanol to dryness, and repeat the test on the residues.

**Optical rotation** <2.49>  $[\alpha]_D^{20}$ : +150 – +156° (after drying, 0.1 g, 1,4-dioxane, 10 mL, 100 mm).

**Purity** Related substances—Dissolve 20 mg of Hydrocortisone in 10 mL of a mixture of chloroform and methanol (9:1), and use this solution as the sample solution. Pipet 1 mL of this solution, add a mixture of chloroform and methanol (9:1) to make exactly 50 mL, and use this solution as the standard solution. Perform the test with these solutions as directed under Thin-layer Chromatography <2.03>. Spot 10  $\mu$ L each of the sample solution and standard solution on a plate of silica gel with fluorescent indicator for thin-layer chromatography. Develop the plate with a mixture of chloroform and ethanol (95) (17:3) to a distance of about 10 cm, and air-dry the plate. Examine under ultraviolet light (main wavelength: 254 nm): the spots other than the principal spot from the sample solution are not more intense than the spot from the standard solution.

**Loss on drying** <2.41> Not more than 1.0% (0.5 g, 105°C, 3 hours).

**Residue on ignition** <2.44> Not more than 0.1% (0.5 g).

**Assay** Dissolve about 20 mg each of Hydrocortisone and Hydrocortisone Reference Standard, previously dried and accurately weighed, in 20 mL each of a mixture of chloroform and methanol (9:1), add 10 mL each of the internal standard solution, then add a mixture of chloroform and methanol (9:1) to make 50 mL, and use these solutions as the sample solution and the standard solution. Perform the test with 5  $\mu$ L each of these solutions as directed under Liquid Chromatography <2.01> according to the following conditions, and calculate the ratios,  $Q_T$  and  $Q_S$ , of the peak area of hydrocortisone to that of the internal standard, respectively.

$$\begin{aligned} \text{Amount (mg) of } C_{21}H_{30}O_5 \\ = W_S \times (Q_T/Q_S) \end{aligned}$$

$W_S$ : Amount (mg) of Hydrocortisone Reference Standard

**Internal standard solution**—A solution of prednisone in a mixture of chloroform and methanol (9:1) (9 in 10,000).

**Operating conditions**—

Detector: An ultraviolet absorption photometer (wavelength: 254 nm).

Column: A stainless steel column 4.6 mm in inside diameter and 25 cm in length, packed with silica gel for liquid chromatography (5  $\mu$ m in particle diameter).

Column temperature: A constant temperature of about 20°C.

Mobile phase: A mixture of chloroform, methanol and acetic acid (100) (1000:20:1).

Flow rate: Adjust the flow rate so that the retention time of hydrocortisone is about 15 minutes.

*System suitability—*


System performance: When the procedure is run with 5  $\mu$ L of the standard solution under the above operating conditions, the internal standard and hydrocortisone are eluted in this order with the resolution between these peaks being not less than 7.

**System repeatability:** When the test is repeated 6 times with 5  $\mu\text{L}$  of the standard solution under the above operating conditions, the relative standard deviation of the ratios of the peak area of hydrocortisone to that of the internal standard is not more than 1.0%.

**Containers and storage** Containers—Tight containers.

## Hydrocortisone Acetate

ヒドロコルチゾン酢酸エステル



$\text{C}_{23}\text{H}_{32}\text{O}_6$ : 404.50

11 $\beta$ ,17,21-Trihydroxypregn-4-ene-3,20-dione 21-acetate  
[50-03-3]

Hydrocortisone Acetate, when dried, contains not less than 97.0% and not more than 102.0% of  $\text{C}_{23}\text{H}_{32}\text{O}_6$ .

**Description** Hydrocortisone Acetate occurs as white crystals or crystalline powder. It is odorless.

It is sparingly soluble in 1,4-dioxane, slightly soluble in methanol, in ethanol (95) and in chloroform, very slightly soluble in diethyl ether, and practically insoluble in water.

Melting point: about 220°C (with decomposition).

**Identification (1)** Add 2 mL of sulfuric acid to 2 mg of Hydrocortisone Acetate: the solution shows a yellowish green fluorescence immediately, and the color of the solution gradually changes through orange-yellow to dark red. This solution shows a strong light green fluorescence under ultraviolet light. Add carefully 10 mL of water to this solution: the color changes from yellow to orange-yellow with a light green fluorescence, and a yellow-brown, flocculent precipitate is formed.

(2) Dissolve 0.01 g of Hydrocortisone Acetate in 1 mL of methanol by warming, add 1 mL of Fehling's TS, and heat: an orange to red precipitate is formed.

(3) To 0.05 g of Hydrocortisone Acetate add 2 mL of potassium hydroxide-ethanol TS, and heat on a water bath for 5 minutes. Cool, add 2 mL of diluted sulfuric acid (2 in 7), and boil gently for 1 minute: the odor of ethyl acetate is perceptible.

(4) Determine the infrared absorption spectra of Hydrocortisone Acetate and Hydrocortisone Acetate Reference Standard, previously dried, as directed in the potassium bromide disk method under Infrared Spectrophotometry <2.25>: both the sample and the Reference Standard exhibit similar intensities of absorption at the same wave numbers. If any difference appears, dissolve the sample and Reference Standard in ethanol (95), respectively, evaporate to dryness, and repeat the test on the residues.

**Optical rotation** <2.49>  $[\alpha]_D^{20}$ : +158 – +165° (after drying, 50 mg, 1,4-dioxane, 10 mL, 100 mm).

**Purity** Related substances—Dissolve 40 mg of Hydrocortisone Acetate in 25 mL of a mixture of chloroform and methanol (9:1), and use this solution as the sample solution. Pipet 2 mL of the sample solution, add a mixture of chloroform and methanol (9:1) to make exactly 100 mL, and use this solution as the standard solution. Perform the test with these solutions as directed under Thin-layer Chromatography <2.03>. Spot 5  $\mu\text{L}$  each of the sample solution and standard solution on a plate of silica gel for thin-layer chromatography. Develop the plate with a mixture of dichloromethane, diethyl ether, methanol and water (160:30:8:1) to a distance of about 12 cm, and air-dry the plate. Spray evenly alkaline blue tetrazolium TS on the plate: the spots other than the principal spot from the sample solution are not more intense than the spot from the standard solution.

**Loss on drying** <2.41> Not more than 1.0% (0.5 g, 105°C, 3 hours).

**Residue on ignition** <2.44> Not more than 0.1% (0.5 g).

**Assay** Dissolve about 20 mg each of Hydrocortisone Acetate and Hydrocortisone Acetate Reference Standard, previously dried and accurately weighed, in methanol, add exactly 10 mL each of the internal standard solution, then add methanol to make 100 mL, and use these solutions as the sample solution and the standard solution. Perform the test with 20  $\mu\text{L}$  each of the sample solution and standard solution as directed under Liquid Chromatography <2.01> according to the following conditions, and calculate the ratios,  $Q_T$  and  $Q_S$ , of the peak area of hydrocortisone acetate to that of the internal standard, respectively.

$$\text{Amount (mg) of } \text{C}_{23}\text{H}_{32}\text{O}_6 = W_S \times (Q_T/Q_S)$$

$W_S$ : Amount (mg) of Hydrocortisone Acetate Reference Standard

**Internal standard solution**—A solution of benzyl parahydroxybenzoate in methanol (1 in 1000).

**Operating conditions**—

**Detector:** An ultraviolet absorption photometer (wavelength: 254 nm).

**Column:** A stainless steel column 3.9 mm in inside diameter and 30 cm in length, packed with octadecylsilanized silica gel for liquid chromatography (10  $\mu\text{m}$  in particle diameter).

**Column temperature:** A constant temperature of about 25°C.

**Mobile phase:** A mixture of water and acetonitrile (13:7).

**Flow rate:** Adjust the flow rate so that the retention time of hydrocortisone acetate is about 8 minutes.

**System suitability**—

**System performance:** When the procedure is run with 20  $\mu\text{L}$  of the standard solution under the above operating conditions, hydrocortisone acetate and the internal standard are eluted in this order with the resolution between these peaks being not less than 4.

**System repeatability:** When the test is repeated 6 times with 20  $\mu\text{L}$  of the standard solution under the above operating conditions, the relative standard deviation of the ratios of the peak area of hydrocortisone acetate to that of the internal standard is not more than 1.0%.

**Containers and storage** Containers—Tight containers.

## Hydrocortisone and Diphenhydramine Ointment

ヒドロコルチゾン・ジフェンヒドラミン軟膏

### Method of preparation

|                        |                       |
|------------------------|-----------------------|
| Hydrocortisone Acetate | 5 g                   |
| Diphenhydramine        | 5 g                   |
| White Petrolatum       | a sufficient quantity |

To make 1000 g

Prepare as directed under Ointments, with the above ingredients.

**Description** Hydrocortisone and Diphenhydramine Ointment is white to pale yellow in color.

**Identification (1)** To 1 g of Hydrocortisone and Diphenhydramine Ointment add 10 mL of ethanol (95), heat on a water bath for 5 minutes with occasional shaking, cool, and filter. Take 5 mL of the filtrate, distill off the ethanol, and to the residue add 2 mL of sulfuric acid: the solution shows a yellow-green fluorescence immediately and the color of the solution gradually changes through yellow to yellow-brown. Add carefully 10 mL of water to this solution: the color changes to yellow with green fluorescence, and a light yellow, flocculent precipitate is formed (hydrocortisone acetate).

(2) To 1 mL of the filtrate obtained in (1) add 5 mL of potassium hydrogen phthalate buffer solution, pH 4.6, and 2 mL of bromophenol blue TS, and add further 5 mL of chloroform. Shake well, and allow to stand: a yellow color develops in the chloroform layer (diphenhydramine).


(3) To 1 g of Hydrocortisone and Diphenhydramine Ointment add 5 mL of methanol, warm, and shake. After cooling, separate the methanol layer, and use this layer as the sample solution. Dissolve 0.01 g each of hydrocortisone acetate and diphenhydramine in 10 mL each of methanol, and use these solutions as standard solutions (1) and (2). Perform the test with the sample solution and standard solutions (1) and (2) as directed under Thin-layer Chromatography <2.03>. Spot 5  $\mu$ L each of these solutions on a plate of silica gel with a complex fluorescent indicator for thin-layer chromatography. Develop the plate with a mixture of ethyl acetate and diethyl ether (4:1) to a distance of about 10 cm, and air-dry the plate. Examine under ultraviolet light (broad spectrum wavelength): two spots from the sample solution show the same *R<sub>f</sub>* value as the corresponding spots from standard solutions (1) and (2).

**Containers and storage** Containers—Tight containers.

Storage—Light-resistant.

## Hydrocortisone Butyrate

ヒドロコルチゾン酪酸エステル



$C_{25}H_{36}O_6$ : 432.55

11 $\beta$ ,17,21-Trihydroxypregn-4-ene-3,20-dione 17-butyrate  
[13609-67-1]

Hydrocortisone Butyrate, when dried, contains not less than 96.0% and not more than 104.0% of  $C_{25}H_{36}O_6$ .

**Description** Hydrocortisone Butyrate occurs as a white powder. It is odorless.

It is freely soluble in tetrahydrofuran, in chloroform and in 1,2-dichloroethane, soluble in methanol, sparingly soluble in ethanol (99.5), slightly soluble in diethyl ether, and practically insoluble in water.

Melting point: about 200°C (with decomposition).

**Identification (1)** Add 2 mL of sulfuric acid to 2 mg of Hydrocortisone Butyrate: the solution shows a yellowish green fluorescence immediately, and the color of the solution gradually changes through orange-yellow to dark red. This solution shows a strong light green fluorescence under ultraviolet light (main wavelength: 254 nm). Add carefully 10 mL of water to this solution: the color changes from yellow to orange-yellow with a light green fluorescence, and a yellow-brown, flocculent precipitate is formed.

(2) Dissolve 0.01 g of Hydrocortisone Butyrate in 1 mL of methanol by warming, add 1 mL of Fehling's TS, and heat: an orange to red precipitate is formed.

(3) To 0.05 g of Hydrocortisone Butyrate add 2 mL of potassium hydroxide-ethanol TS, and heat on a water bath for 5 minutes. Cool, add 2 mL of diluted sulfuric acid (2 in 7), and boil gently for 1 minute: the odor of ethyl butyrate is perceptible.

(4) Determine the infrared absorption spectrum of Hydrocortisone Butyrate, previously dried, as directed in the potassium bromide disk method under Infrared Spectrophotometry <2.25>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wave numbers.

**Optical rotation** <2.49>  $[\alpha]_D^{25}$ : +48 – +52° (after drying, 0.1 g, chloroform, 10 mL, 100 mm).

**Purity (1)** Heavy metals <1.07>—Proceed with 1.0 g of Hydrocortisone Butyrate according to method 2, and perform the test. Prepare the control solution with 2.0 mL of Standard Lead Solution (not more than 20 ppm).

(2) Related substances—Dissolve 25 mg of Hydrocortisone Butyrate in 5 mL of tetrahydrofuran, and use this solution as the sample solution. Pipet 2 mL of this solution, and add tetrahydrofuran to make exactly 50 mL. Pipet 5 mL of

this solution, add tetrahydrofuran to make exactly 20 mL, and use this solution as the standard solution. Perform the test with these solutions as directed under Thin-layer Chromatography <2.03>. Spot 10  $\mu$ L each of the sample solution and standard solution on a plate of silica gel for thin-layer chromatography. Develop the plate with a mixture of 1,2-dichloroethane, methanol and water (470:30:1) to a distance of about 15 cm, and air-dry the plate. Spray evenly alkaline blue tetrazolium TS on the plate: the spots other than the principal spot from the sample solution are not more than two in number, and not more intense than those from the standard solution in color.

**Loss on drying** <2.41> Not more than 1.0% (1 g, 105°C, 3 hours).

**Residue on ignition** <2.44> Not more than 0.1% (1 g).


**Assay** Weigh accurately about 50 mg of Hydrocortisone Butyrate, previously dried, and dissolve in ethanol (99.5) to make exactly 100 mL. Pipet 2 mL of this solution, and add ethanol (99.5) to make exactly 50 mL. Determine the absorbance *A* of this solution at the wavelength of maximum absorption at about 241 nm as directed under Ultraviolet-visible Spectrophotometry <2.24>.

$$\text{Amount (mg) of } C_{25}H_{36}O_6 = (A/375) \times 25,000$$

**Containers and storage** Containers—Tight containers.

## Hydrocortisone Sodium Phosphate

ヒドロコルチゾンリン酸エステルナトリウム



$C_{21}H_{29}Na_2O_8P$ : 486.40

Disodium 11 $\beta$ ,17,21-trihydroxypregn-4-ene-3,20-dione 21-phosphate [6000-74-4]

Hydrocortisone Sodium Phosphate contains not less than 96.0% and not more than 102.0% of  $C_{21}H_{29}Na_2O_8P$ , calculated on the dried basis.

**Description** Hydrocortisone Sodium Phosphate occurs as a white to light yellow powder. It is odorless.

It is freely soluble in water, sparingly soluble in methanol, very slightly soluble in ethanol (95), and practically insoluble in diethyl ether.

It is hygroscopic.

**Identification** (1) To 2 mg of Hydrocortisone Sodium Phosphate add 2 mL of sulfuric acid: a yellowish green fluorescence is exhibited initially, then gradually changes through orange-yellow to dark red. Examine the solution under ultraviolet light (main wavelength: 254 nm): an intense, light green fluorescence is exhibited. To this solution add carefully 10 mL of water: the color changes from yellow to orange-yellow with a light green fluorescence and a yellow-brown, flocculent floating substance is formed.

(2) Determine the infrared absorption spectrum of

Hydrocortisone Sodium Phosphate as directed in the paste method under Infrared Spectrophotometry <2.25>, and compare the spectrum with the Reference Spectrum or the spectrum of Hydrocortisone Sodium Phosphate Reference Standard: both spectra exhibit similar intensities of absorption at the same wave numbers. If any difference appears between the spectra, dissolve Hydrocortisone Sodium Phosphate and Hydrocortisone Sodium Phosphate Reference Standard in methanol, respectively, then evaporate the methanol to dryness, and repeat the test on the residues.

(3) Moisten 1.0 g of Hydrocortisone Sodium Phosphate with a small quantity of sulfuric acid, and incinerate by gradual heating. After cooling, dissolve the residue in 10 mL of dilute nitric acid, and heat in a water bath for 30 minutes. After cooling, filter if necessary. This solution responds to the Qualitative Tests <1.09> for sodium salt and for phosphate.

**Optical rotation** <2.49>  $[\alpha]_D^{20}$ : +121 – +129° (1 g, calculated on the dried basis, phosphate buffer solution, pH 7.0, 100 mL, 100 mm).

**pH** <2.54> Dissolve 1.0 g of Hydrocortisone Sodium Phosphate in 100 mL of water: the pH of this solution is between 7.5 and 9.5.

**Purity** (1) Clarity and color of solution—Dissolve 1.0 g of Hydrocortisone Sodium Phosphate in 10 mL of water: the solution is clear and colorless to pale yellow.

(2) Chloride <1.03>—Dissolve 0.30 g of Hydrocortisone Sodium Phosphate in 20 mL of water, and add 6 mL of dilute nitric acid and water to make 100 mL. To 5 mL of this solution add water to make 50 mL. Perform the test using this solution as the test solution. Prepare the control solution with 0.25 mL of 0.01 mol/L hydrochloric acid VS (not more than 0.600%).

(3) Heavy metals <1.07>—Proceed with 0.5 g of Hydrocortisone Sodium Phosphate according to Method 2, and perform the test. Prepare the control solution with 2.0 mL of Standard Lead Solution (not more than 40 ppm).

(4) Arsenic <1.11>—Prepare the test solution with 1.0 g of Hydrocortisone Sodium Phosphate according to Method 3, and perform the test (not more than 2 ppm).

(5) Free phosphoric acid—Weigh accurately about 0.25 g of Hydrocortisone Sodium Phosphate, dissolve in water to make exactly 100 mL, and use this solution as the sample solution. Pipet 5 mL each of the sample solution and Standard Phosphoric Acid Solution into separate 25-mL volumetric flasks, add 2.5 mL of hexaammonium heptamolybdate-sulfuric acid TS and 1 mL of 1-amino-2-naphthol-4-sulfonic acid TS, shake, add water to make exactly 25 mL, and allow to stand at  $20 \pm 1^\circ\text{C}$  for 30 minutes. Perform the test with these solutions as directed under Ultraviolet-visible Spectrophotometry <2.24>, using a solution prepared with 5 mL of water in the same manner as the blank. Determine the absorbances,  $A_T$  and  $A_S$ , of each solution from the sample solution and Standard Phosphoric Acid Solution at 740 nm: the amount of free phosphoric acid is not more than 1.0%.

Content (%) of free phosphoric acid ( $H_3PO_4$ )

$$= (A_T/A_S) \times (1/W) \times 257.8$$

*W*: Amount (mg) of Hydrocortisone Sodium Phosphate, calculated on the dried basis.

(6) Free hydrocortisone—Dissolve 25 mg of Hydrocorti-

sone Sodium Phosphate in the mobile phase to make exactly 20 mL, and use this solution as the sample solution. Separately, weigh 25 mg of Hydrocortisone Reference Standard, previously dried at 105°C for 3 hours, and dissolve in the mobile phase to make exactly 100 mL. Pipet 10 mL of this solution, add the mobile phase to make exactly 200 mL, and use this solution as the standard solution. Perform the test with exactly 20  $\mu$ L each of the sample solution and standard solution as directed under Liquid Chromatography <2.01> according to the following conditions. Determine the peak areas,  $A_T$  and  $A_S$ , of hydrocortisone from each solution:  $A_T$  is not larger than  $A_S$ .

**Operating conditions—**

Detector, column, column temperature, mobile phase, and flow rate: Proceed as directed in the operating conditions in the Assay.

**System suitability—**

System performance: Proceed as directed in the system suitability in the Assay.

System repeatability: When the test is repeated 6 times with 20  $\mu$ L of the standard solution under the above operating conditions, the relative standard deviation of the peak area of hydrocortisone is not more than 1.0%.

**Loss on drying** <2.41> Not more than 5.0% (1 g, in vacuum, 80°C, 5 hours).

**Assay** Weigh accurately about 20 mg each of Hydrocortisone Sodium Phosphate and Hydrocortisone Sodium Phosphate Reference Standard (previously determine the loss on drying <2.41> in the same manner as Hydrocortisone Sodium Phosphate), dissolve each in 50 mL of the mobile phase, add exactly 10 mL of the internal standard solution, then add the mobile phase to make 200 mL, and use these solutions as the sample solution and the standard solution, respectively. Perform the test with 20  $\mu$ L each of the sample solution and standard solution as directed under Liquid Chromatography <2.01> according to the following conditions, and calculate the ratios,  $Q_T$  and  $Q_S$ , of the peak area of hydrocortisone phosphate to that of the internal standard, respectively.

$$\text{Amount (mg) of } C_{21}H_{29}Na_2O_8P = W_S \times (Q_T/Q_S)$$

$W_S$ : Amount (mg) of Hydrocortisone Sodium Phosphate Reference Standard, calculated on the dried basis

**Internal standard solution—**A solution of isopropyl parahydroxybenzoate in the mobile phase (3 in 5000).

**Operating conditions—**

Detector: An ultraviolet absorption photometer (wavelength: 254 nm).

Column: A stainless steel column 4.6 mm in inside diameter and 25 cm in length, packed with octadecylsilanized silica gel for liquid chromatography (7  $\mu$ m in particle diameter).

Column temperature: A constant temperature of about 25°C.

Mobile phase: A mixture of 0.05 mol/L sodium dihydrogen phosphate TS, pH 2.6 and methanol (1:1).

Flow rate: Adjust the flow rate so that the retention time of hydrocortisone phosphate is about 10 minutes.

**System suitability—**

System performance: When the procedure is run with 20  $\mu$ L of the standard solution under the above operating conditions, hydrocortisone phosphate and isopropyl para-


hydroxybenzoate are eluted in this order with the resolution between these peaks being not less than 8.

System repeatability: When the test is repeated 6 times with 20  $\mu$ L of the standard solution under the above operating conditions, the relative standard deviation of the ratios of the peak area of hydrocortisone phosphate to that of the internal standard is not more than 1.0%.

**Containers and storage** Containers—Tight containers.

## Hydrocortisone Sodium Succinate

ヒドロコルチゾンコハク酸エステルナトリウム



$C_{25}H_{33}NaO_8$ : 484.51

Monosodium 11 $\beta$ ,17,21-trihydroxypregn-4-ene-3,20-dione 21-succinate [125-04-2]

Hydrocortisone Sodium Succinate, calculated on the dried basis, contains not less than 97.0% and not more than 103.0% of  $C_{25}H_{33}NaO_8$ .

**Description** Hydrocortisone Sodium Succinate occurs as white powder or masses. It is odorless.

It is freely soluble in water, in methanol and in ethanol (95), and practically insoluble in diethyl ether.

It is hygroscopic.

It is gradually colored by light.

**Identification (1)** Dissolve 0.2 g of Hydrocortisone Sodium Succinate in 20 mL of water, and add 0.5 mL of dilute hydrochloric acid with stirring: a white precipitate is formed. Collect the precipitate, wash it with two 10-mL portions of water, and dry at 105°C for 3 hours. To 3 mg of this dried matter add 2 mL of sulfuric acid: the solution shows a yellowish green fluorescence immediately, and the color of the solution gradually changes through orange-yellow to dark red. This solution shows a strong light green fluorescence under ultraviolet light. Add carefully 10 mL of water to this solution: the color changes from yellow to orange-yellow with a light green fluorescence, and a yellow-brown flocculent precipitate is formed.

(2) Dissolve 0.01 g of the dried matter obtained in (1) in 1 mL of methanol, add 1 mL of Fehling's TS, and heat: an orange to red precipitate is formed.

(3) To 0.1 g of the dried matter obtained in (1) add 2 mL of sodium hydroxide TS, and allow to stand for 10 minutes. Filter the solution to remove the precipitate formed, mix the filtrate with 1 mL of dilute hydrochloric acid, filter if necessary, then adjust the solution to a pH of about 6 with diluted ammonia TS (1 in 10), and add 2 to 3 drops of iron (III) chloride TS: a brown precipitate is formed.

(4) Determine the infrared absorption spectrum of the dried matter obtained in (1) as directed in the potassium bromide disk method under Infrared Spectrophotometry <2.25>, and compare the spectrum with the Reference Spec-

trum or the spectrum of previously dried Hydrocortisone Succinate Reference Standard: both spectra exhibit similar intensities of absorption at the same wave numbers. If any difference appears between the spectra, dissolve Hydrocortisone Sodium Succinate and Hydrocortisone Succinate Reference Standard in methanol, respectively, then evaporate the methanol to dryness, and repeat the test on the residues.

(5) Hydrocortisone Sodium Succinate responds to the Qualitative Tests <1.09> (1) for sodium salt.

**Optical rotation** <2.49>  $[\alpha]_D^{20}$ : +135 – +145° (0.1 g, calculated on the dried basis, ethanol (95), 10 mL, 100 mm).

**Purity** (1) Clarity and color of solution—Dissolve 0.5 g of Hydrocortisone Sodium Succinate in 10 mL of water: the solution is clear and colorless.

(2) Other steroids—Dissolve 25 mg of Hydrocortisone Sodium Succinate in methanol to make exactly 10 mL, and use this solution as the sample solution. Separately, dissolve 0.025 g of hydrocortisone in methanol to make exactly 10 mL. Pipet 1 mL of this solution, add methanol to make exactly 20 mL, and use this solution as the standard solution (1). Pipet 6 mL of the standard solution (1), add methanol to make exactly 10 mL, and use this solution as the standard solution (2). Perform the test with these solutions as directed under Thin-layer Chromatography <2.03>. Spot 3  $\mu$ L each of the sample solution and standard solutions (1) and (2) on a plate of silica gel with fluorescent indicator for thin-layer chromatography. Develop the plate with a mixture of chloroform, ethanol (99.5) and formic acid (150:10:1) to a distance of about 10 cm, and air-dry the plate. Examine under ultraviolet light (main wavelength: 254 nm): the spot from the sample solution corresponding to the spot from the standard solution (1) is not more intense than the spot from the standard solution (1). Any spot other than the principal spot and the above spot obtained from the sample solution is not more than one, and is not more intense than the spot from the standard solution (2).

**Loss on drying** <2.41> Not more than 2.0% (0.5 g, 105°C, 3 hours).

**Assay** Weigh accurately about 10 mg of Hydrocortisone Sodium Succinate, and dissolve in methanol to make exactly 100 mL. Pipet 5 mL of this solution, add methanol to make exactly 50 mL, and use this solution as the sample solution. Separately, weigh accurately about 0.01 g of Hydrocortisone Succinate Reference Standard, previously dried at 105°C for 3 hours, proceed in the same manner as directed for the sample solution, and use this solution as the standard solution. Determine the absorbances,  $A_T$  and  $A_S$ , of the sample solution and standard solution at 240 nm as directed under Ultraviolet-visible Spectrophotometry <2.24>, respectively.


$$\begin{aligned} \text{Amount (mg) of } C_{25}H_{33}NaO_8 \\ = W_S \times (A_T/A_S) \times 1.0475 \end{aligned}$$

$W_S$ : Amount (mg) of Hydrocortisone Succinate Reference Standard

**Containers and storage** Containers—Tight containers.  
Storage—Light-resistant.

## Hydrocortisone Succinate

ヒドロコルチゾンコハク酸エステル



$C_{25}H_{34}O_8$ : 462.53

11 $\beta$ ,17,21-Trihydroxypregn-4-ene-3,20-dione  
21-(hydrogen succinate) [2203-97-6]

Hydrocortisone Succinate, when dried, contains not less than 97.0% and not more than 103.0% of  $C_{25}H_{34}O_8$ .

**Description** Hydrocortisone Succinate occurs as a white crystalline powder.

It is very soluble in methanol, freely soluble in ethanol (99.5), sparingly soluble in ethanol (95), and practically insoluble in water.

**Identification** (1) To 3 mg of Hydrocortisone Succinate add 2 mL of sulfuric acid: the solution shows a yellowish green fluorescence immediately, and the color of the solution gradually changes through orange-yellow to dark red. This solution shows a strong light green fluorescence under ultraviolet light. Add carefully 10 mL of water to this solution: the color changes from yellow to orange-yellow with a light green fluorescence, and a yellow-brown flocculent precipitate is formed.

(2) Determine the infrared absorption spectrum of Hydrocortisone Succinate, previously dried, as directed in the potassium bromide disk method under Infrared Spectrophotometry <2.25>, and compare the spectrum with the Reference Spectrum or the spectrum of previously dried Hydrocortisone Succinate Reference Standard: both spectra exhibit similar intensities of absorption at the same wave numbers. If any difference appears between the spectra, dissolve Hydrocortisone Succinate and Hydrocortisone Succinate Reference Standard in methanol, respectively, then evaporate the methanol to dryness, and repeat the test on the residues.

**Optical rotation** <2.49>  $[\alpha]_D^{20}$ : +147 – +153° (after drying, 0.1 g, ethanol (99.5), 10 mL, 100 mm).

**Purity** Related substances—Dissolve 25 mg of Hydrocortisone Succinate in exactly 10 mL of methanol, and use this solution as the sample solution. Separately, dissolve 0.025 g of hydrocortisone in exactly 10 mL of methanol. Pipet 1 mL of this solution, dilute with methanol to exactly 50 mL, and use this solution as the standard solution. Perform the test with these solutions as directed under Thin-layer Chromatography <2.03>. Spot 3  $\mu$ L each of the sample solution and standard solution on a plate of silica gel with fluorescent indicator for thin-layer chromatography. Develop the plate with a mixture of chloroform, ethanol (99.5) and formic acid (150:10:1) to a distance of about 10 cm, and air-dry the plate. Examine un-

der ultraviolet light (main wavelength: 254 nm): the spots other than the principal spot from the sample solution are not more intense than the spot from the standard solution.

**Loss on drying** <2.41> Not more than 2.0% (0.5 g, 105°C, 3 hours).

**Residue on ignition** <2.44> Not more than 0.1% (0.5 g).

**Assay** Weigh accurately about 50 mg each of Hydrocortisone Succinate and Hydrocortisone Succinate Reference Standard, previously dried, and dissolve in methanol to make exactly 50 mL. Pipet 5 mL each of these solutions, add exactly 5 mL each of the internal standard solution, then add methanol to make 50 mL, and use these solutions as the sample solution and standard solution. Perform the test with 10  $\mu$ L each of the sample solution and standard solution as directed under Liquid Chromatography <2.01> according to the following operating conditions, and calculate the ratios,  $Q_T$  and  $Q_S$ , of the peak area of hydrocortisone succinate to that of the internal standard, respectively.

$$\begin{aligned} &\text{Amount (mg) of } C_{25}H_{34}O_8 \\ &= W_S \times (Q_T/Q_S) \end{aligned}$$

$W_S$ : Amount (mg) of Hydrocortisone Succinate Reference Standard

**Internal standard solution**—A solution of butyl parahydroxy benzoate in methanol (1 in 2500).

**Operating conditions**—

**Detector:** An ultraviolet absorption photometer (wavelength: 254 nm).

**Column:** A stainless steel column 4 mm in inside diameter and 30 cm in length, packed with octadecylsilanized silica gel (10  $\mu$ m in particle diameter).

**Column temperature:** A constant temperature of about 25°C.

**Mobile phase:** A mixture of acetic acid-sodium acetate buffer solution, pH 4.0 and acetonitrile (3:2).

**Flow rate:** Adjust the flow rate so that the retention time of hydrocortisone succinate is about 5 minutes.

**System suitability**—

**System performance:** When the procedure is run with 10  $\mu$ L of the standard solution under the above operating conditions, hydrocortisone succinate and the internal standard are eluted in this order with the resolution between these peaks being not less than 9.


**System repeatability:** When the test is repeated 6 times with 10  $\mu$ L of the standard solution under the above operating conditions, the relative standard deviation of the ratios of the peak area of hydrocortisone succinate to that of the internal standard is not more than 1.0%.

**Containers and storage** Containers—Tight containers.

Storage—Light-resistant.

## Hydrocotarnine Hydrochloride Hydrate

ヒドロコタルニン塩酸塩水和物



$C_{12}H_{15}NO_3 \cdot HCl \cdot H_2O$ : 275.73

4-Methoxy-6-methyl-5,6,7,8-

tetrahydro[1,3]dioxolo[4,5-g]isoquinoline

monohydrochloride monohydrate [5985-55-7, anhydride]

Hydrocotarnine Hydrochloride Hydrate, when dried, contains not less than 98.0% of hydrocotarnine-hydrochloride ( $C_{12}H_{15}NO_3HCl$ : 257.72).

**Description** Hydrocotarnine Hydrochloride Hydrate occurs as white to pale yellow crystals or crystalline powder.

It is freely soluble in water, sparingly soluble in ethanol (95) and in acetic acid (100), and slightly soluble in acetic anhydride.

**Identification (1)** Determine the absorption spectrum of a solution of Hydrocotarnine Hydrochloride Hydrate (1 in 10,000) as directed under Ultraviolet-visible Spectrophotometry <2.24>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wavelengths.

**(2)** Determine the infrared absorption spectrum of Hydrocotarnine Hydrochloride Hydrate as directed in the potassium bromide disk method under Infrared Spectrophotometry <2.25>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wave numbers.

**(3)** A solution of Hydrocotarnine Hydrochloride Hydrate (1 in 50) responds to the Qualitative Tests <1.09> (2) for chloride.

**pH** <2.54> Dissolve 1.0 g of Hydrocotarnine Hydrochloride Hydrate in 20 mL of water: the pH of the solution is between 4.0 and 6.0.

**Purity (1)** Clarity and color of solution—Dissolve 0.5 g of Hydrocotarnine Hydrochloride Hydrate in 10 mL of water: the solution is clear, and when perform the test with this solution as directed under Ultraviolet-visible Spectrophotometry <2.24>, using water as the blank, the absorbance at 400 nm is not more than 0.17.

**(2)** Heavy metals <1.07>—Proceeds with 1.0 g of Hydrocotarnine Hydrochloride Hydrate according to Method 1, and perform the test. Prepare the control solution with 2.0 mL of Standard Lead Solution (not more than 20 ppm).

**(3)** Related substances—Dissolve 0.10 g of Hydrocotarnine Hydrochloride Hydrate in 10 mL of diluted ethanol (99.5) (1 in 2), and use this solution as the sample solution. Pipet 1 mL of the sample solution, add diluted ethanol (99.5) (1 in 2) to make exactly 100 mL, and use this solution as the standard solution. Perform the test with these

solutions as directed under Thin-layer Chromatography <2.03>. Spot 10  $\mu$ L each of the sample solution and standard solution on a plate of silica gel with fluorescent indicator for thin-layer chromatography. Develop the plate with a mixture of acetone, toluene, ethanol (99.5) and ammonia water (28) (20:20:3:1) to a distance of about 15 cm, and air-dry the plate. Examine under ultraviolet light (main wavelength: 365 nm): the spots other than the principal spot from the sample solution are not more intense than the spot from the standard solution.

**Loss on drying** <2.41> Not more than 7.0% (1 g, 105°C, 3 hours).

**Residue on ignition** <2.44> Not more than 0.2% (1 g).

**Assay** Weigh accurately about 0.5 g of Hydrocotarnine Hydrochloride Hydrate, previously dried. Dissolve in 50 mL of a mixture of acetic anhydride and acetic acid (100) (7:3) by warming. Cool, and titrate <2.50> with 0.1 mol/L perchloric acid VS (potentiometric titration). Perform a blank determination.

Each mL of 0.1 mol/L perchloric acid VS  
= 25.77 mg of  $C_{12}H_{15}NO_3HCl$

**Containers and storage** Containers—Tight containers.

## Hydrogenated Oil

硬化油

Hydrogenated Oil is the fat obtained by hydrogenation of fish oil or of other oils originating from animal or vegetable.

**Description** Hydrogenated Oil occurs as a white mass or powder and has a characteristic odor and a mild taste.

It is freely soluble in diethyl ether, very slightly soluble in ethanol (95), and practically insoluble in water.

The oil obtained by hydrogenation of castor oil is slightly soluble in diethyl ether, very slightly soluble in ethanol (95), and practically insoluble in water.

**Acid value** <1.13> Not more than 2.0.

**Purity** (1) Moisture and coloration—Hydrogenated Oil (5.0 g), melted by heating on a water bath, forms a clear liquid, from which no water separates. In a 10-mm thick layer of the liquid, it is colorless or slightly yellow.

(2) Alkalinity—To 2.0 g of Hydrogenated Oil add 10 mL of water, melt by heating on a water bath, and shake vigorously. After cooling, add 1 drop of phenolphthalein TS to the separated water layer: no color develops.

(3) Chloride—To 1.5 g of Hydrogenated Oil add 30 mL of ethanol (95), boil for 10 minutes under a reflux condenser, and filter after cooling. To 20 mL of the filtrate add 5 drops of a solution of silver nitrate in ethanol (95) (1 in 50): the turbidity of the solution does not exceed that of the following control solution.

Control solution: To 1.0 mL of 0.01 mol/L hydrochloric acid VS add ethanol (95) to make 20 mL, then add 5 drops of a solution of silver nitrate in ethanol (95) (1 in 50).

(4) Heavy metals—Heat 2.0 g of Hydrogenated Oil with 5 mL of dilute hydrochloric acid and 10 mL of water on a

water bath for 5 minutes with occasional shaking. After cooling, filter, and make 5 mL of the filtrate weakly alkaline with ammonia TS, then add 3 drops of sodium sulfide TS: the solution remains unchanged.

(5) Nickel—Place 5.0 g of Hydrogenated Oil in a quartz or porcelain crucible, heat slightly with caution at the beginning, and, after carbonization, incinerate by strong heating ( $500 \pm 20^\circ\text{C}$ ). Cool, add 1 mL of hydrochloric acid, evaporate on a water bath to dryness, dissolve the residue in 3 mL of dilute hydrochloric acid, and add 7 mL of water. Then add 1 mL of bromine TS and 1 mL of a solution of citric acid monohydrate (1 in 5), make alkaline with 5 mL of ammonia TS, and cool in running water. To this solution add 1 mL of dimethylglyoxime TS, add water to make 20 mL, and use this solution as the test solution. Allow to stand for 5 minutes: the solution has no more color than the following control solution.

Control solution: Evaporate 1 mL of hydrochloric acid on a water bath to dryness, add 1 mL of Standard Nickel Solution and 3 mL of dilute hydrochloric acid, and add 6 mL of water. Then proceed as directed in the test solution, add water to make 20 mL, and allow to stand for 5 minutes.

**Residue on ignition** <2.44> Not more than 0.1% (5 g).

**Containers and storage** Containers—Well-closed containers.

## Hydrophilic Ointment

親水軟膏

### Method of preparation

|                              |                       |
|------------------------------|-----------------------|
| White Petrolatum             | 250 g                 |
| Stearyl Alcohol              | 200 g                 |
| Propylene Glycol             | 120 g                 |
| Polyoxyethylene hydrogenated |                       |
| castor oil 60                | 40 g                  |
| Glycerin Monostearate        | 10 g                  |
| Methyl Parahydroxybenzoate   | 1 g                   |
| Propyl Parahydroxybenzoate   | 1 g                   |
| Purified Water               | a sufficient quantity |

To make 1000 g

Melt White Petrolatum, Stearyl Alcohol, polyoxyethylene hydrogenated castor oil 60 and Glycerin Monostearate by heating on a water bath, stir, and keep temperature of the mixture at about  $75^\circ\text{C}$ . To Propylene Glycol add Methyl Parahydroxybenzoate and Propyl Parahydroxybenzoate, melt by warming if necessary, dissolve in Purified Water, and warm to about  $75^\circ\text{C}$ . Add this solution to the above mixture, stir to form emulsion, cool, and stir thoroughly until it congeals.


**Description** Hydrophilic Ointment is white in color. It has a slight, characteristic odor.

**Containers and storage** Containers—Tight containers.



## Hydroxocobalamin Acetate

ヒドロキソコバラミン酢酸塩



$C_{62}H_{89}CoN_{13}O_{15}P \cdot C_2H_4O_2$ : 1406.41  
 $Co\alpha$ -[ $\alpha$ -(5,6-Dimethyl-1H-benzimidazol-1-yl)]- $Co\beta$ -hydroxocobamide monoacetate  
 [13422-51-0, Hydroxocobalamin]

Hydroxocobalamin Acetate contains not less than 95.0% of  $C_{62}H_{89}CoN_{13}O_{15}P \cdot C_2H_4O_2$ , calculated on the dried basis.

**Description** Hydroxocobalamin Acetate occurs as dark red crystals or powder. It is odorless.

It is freely soluble in water, slightly soluble in ethanol (95), and practically insoluble in diethyl ether.

It is hygroscopic.

**Identification (1)** Determine the absorption spectrum of a solution of Hydroxocobalamin Acetate in acetic acid-sodium acetate buffer solution, pH 4.5 (1 in 50,000) as directed under Ultraviolet-visible Spectrophotometry <2.24>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wavelengths.

(2) Mix 1 mg of Hydroxocobalamin Acetate with 0.05 g of potassium hydrogen sulfate, and fuse by igniting. Cool, break up the mass with a glass rod, add 3 mL of water, and dissolve by boiling. Add 1 drop of phenolphthalein TS, and add dropwise sodium hydroxide TS until the solution develops a light red. Then add 0.5 g of sodium acetate trihydrate, 0.5 mL of dilute acetic acid and 0.5 mL of a solution of disodium 1-nitroso-2-naphthol-3,6-disulfonate (1 in 500): a red to orange-red color develops immediately. Then add 0.5 mL of hydrochloric acid, and boil for 1 minute: the red color does not disappear.

(3) Add 0.5 mL of ethanol (99.5) and 1 mL of sulfuric acid to 0.02 g of Hydroxocobalamin Acetate, and heat the mixture: the odor of ethyl acetate is perceptible.

**Purity** Cyanocobalamin and colored impurities—Dissolve 50 mg of Hydroxocobalamin Acetate in exactly 5 mL each of

acetic acid-sodium acetate buffer solution, pH 5.0, in two tubes. To one tube add 0.15 mL of potassium thiocyanate TS, allow to stand for 30 minutes, and use this solution as the sample solution (1). To the other tube add 0.10 mL of potassium cyanide TS, allow to stand for 30 minutes, and use this solution as the sample solution (2). Separately, dissolve 3.0 mg of Cyanocobalamin Reference Standard in exactly 10 mL of acetic acid-sodium acetate buffer solution, pH 5.0, and use this solution as the standard solution. Perform the test with these solutions as directed under Thin-layer Chromatography <2.03>. Apply 20  $\mu$ L each of the sample solution and standard solution 25 mm in length along the starting line, 10 mm apart from each other, on a plate of silica gel for thin-layer chromatography. Develop the plate for 18 hours with 2-butanol saturated with water, while supporting the plate at an angle of about 15° to a horizontal plane, and air-dry the plate: the spot from the sample solution (1) corresponding to that from the standard solution is not more intense than the spot from the standard solution, and the spots other than the principal spot from the sample solution (2) are not more intense than the spot from the standard solution.

**Loss on drying** <2.41> Not more than 12% (50 mg, in vacuum at a pressure not exceeding 0.67 kPa, phosphorus (V) oxide, 100°C, 6 hours).

**Assay** Weigh accurately about 20 mg of Hydroxocobalamin Acetate, and dissolve in acetic acid-sodium acetate buffer solution, pH 5.0, to make exactly 50 mL. Pipet 2 mL of this solution into a 50-mL volumetric flask, add 1 mL of a solution of potassium cyanide (1 in 1000), and allow to stand for 30 minutes at ordinary temperature. Add acetic acid-sodium acetate buffer solution, pH 5.0, to make exactly 50 mL, and use this solution as the sample solution. Separately, weigh accurately about 20 mg of Cyanocobalamin Reference Standard after determining the loss on drying in the same manner as for Cyanocobalamin, and dissolve in water to make exactly 50 mL. To 2 mL of this solution, exactly measured, add acetic acid-sodium acetate buffer solution, pH 5.0, to make exactly 50 mL, and use this solution as the standard solution. Determine the absorbances,  $A_T$  and  $A_S$ , of the sample solution and standard solution at 361 nm as directed under Ultraviolet-visible Spectrophotometry <2.24>.

$$\begin{aligned} \text{Amount (mg) of } C_{62}H_{89}CoN_{13}O_{15}P \cdot C_2H_4O_2 \\ = W_S \times (A_T/A_S) \times 1.0377 \end{aligned}$$

$W_S$ : Amount (mg) of Cyanocobalamin Reference Standard, calculated on the dried basis

**Containers and storage** Containers—Tight containers.

Storage—Light-resistant, and in a cold place.

## Hydroxypropylcellulose

ヒドロキシプロピルセルロース

Hydroxypropylcellulose is a hydroxypropyl ether of cellulose.

Hydroxypropylcellulose, when dried, contains not less than 53.4% and not more than 77.5% of hydroxypropoxy group ( $-OC_3H_6OH$ : 75.09).

**Description** Hydroxypropylcellulose occurs as a white to

yellowish white powder.

It is practically insoluble in diethyl ether.

It forms a viscous liquid upon addition of water or ethanol (95).

**Identification (1)** To 1 g of Hydroxypropylcellulose add 100 mL of water, heat in a water bath at 70°C for 5 minutes with stirring, and cool while shaking. Allow to stand at room temperature until it becomes more homogeneous and viscous, and use this solution as the sample solution. To 2 mL of the sample solution add 1 mL of anthrone TS gently: a blue to green color develops at the zone of contact.

**(2)** Heat the sample solution obtained in (1): a white turbidity or precipitate is produced, and the turbidity or precipitate disappears when cooled.

**(3)** To 1 g of Hydroxypropylcellulose add 100 mL of ethanol (95), and allow to stand after stirring: a homogeneous and viscous liquid is produced.

**pH** <2.54> Dissolve 1.0 g of Hydroxypropylcellulose in 50 mL of freshly boiled and cooled water: the pH of the solution is between 5.0 and 7.5.

**Purity (1)** Clarity of solution—Use an outer glass cylinder, 250 mm in height, 25 mm in internal diameter, 2 mm in thickness, with a high-quality glass plate 2 mm thick at the bottom, and inner glass cylinder, 300 mm in height, 15 mm in internal diameter, 2 mm in thickness, with a high-quality glass plate 2 mm thick at the bottom. In the outer cylinder place a solution prepared by adding 1.0 g of Hydroxypropylcellulose to 100 mL of water, heat while stirring in a water bath at 70°C, and then cool to room temperature. Place this cylinder on a sheet of white paper on which 15 parallel, black, 1-mm width lines are drawn at 1-mm intervals. Place the inner cylinder, and move it up and down while viewing downward through the bottom of the inner cylinder, and measure the minimum height of the solution between the bottom of the outer cylinder and the lower end of the inner cylinder at the time when the lines on the paper cannot be differentiated. The average value obtained from three repeated procedures is greater than that obtained from the following control solution treated in the same manner.

Control solution: To 5.50 mL of 0.005 mol/L sulfuric acid VS add 1 mL of dilute hydrochloric acid, 5 mL of ethanol (95) and water to make 50 mL. To this solution add 2 mL of barium chloride TS, mix, allow to stand for 10 minutes, and shake well before use.

**(2)** Chloride <1.03>—Add 1.0 g of Hydroxypropylcellulose to 30 mL of water, heat in a water bath with stirring for 30 minutes, and filter while being hot. Wash the residue with three 15-mL portions of hot water, combine the washings with the filtrate, and add water to make 100 mL after cooling. To 10 mL of the sample solution add 6 mL of dilute nitric acid and water to make 50 mL, and perform the test using this solution as the test solution. Prepare the control solution with 0.40 mL of 0.01 mol/L hydrochloric acid VS (not more than 0.142%).

**(3)** Sulfate <1.14>—To 40 mL of the sample solution obtained in (2) add 1 mL of dilute hydrochloric acid and water to make 50 mL, and perform the test using this solution as the test solution. Prepare the control solution with 0.40 mL of 0.005 mol/L sulfuric acid VS (not more than 0.048%).

**(4)** Heavy metals <1.07>—Proceed with 1.0 g of Hydroxypropylcellulose according to Method 2 and perform the test.

Prepare the control solution with 2.0 mL of Standard Lead Solution (not more than 20 ppm).

**(5)** Arsenic <1.11>—Prepare the test solution with 1.0 g of Hydroxypropylcellulose according to Method 3, and perform the test (not more than 2 ppm).

**Loss on drying** <2.41> Not more than 5.0% (1 g, 105°C, 4 hours).

**Residue on ignition** <2.44> Not more than 0.5% (1 g).

**Assay (i)** Apparatus—Reaction flask: A 5-mL screw-cap pressure-tight glass bottle, having an inverted conical bottom inside, 20 mm in outside diameter, 50 mm in height up to the neck, 2 mL in capacity up to a height of about 30 mm, equipped with a pressure-tight septum of heat-resisting resin and also with an inside stopper or sealer of fluoroplastic.

Heater: A square aluminum block 60 to 80 mm thick, having holes 20.6 mm in diameter and 32 mm in depth, capable of maintaining the inside temperature within  $\pm 1^\circ\text{C}$ .

**(ii)** Procedure—Weigh accurately about 65 mg of Hydroxypropylcellulose, previously dried, transfer to the reaction flask, add 65 mg of adipic acid, 2.0 mL of the internal standard solution and 2.0 mL of hydroiodic acid, stopper the flask tightly, and weigh accurately. Shake the flask for 30 seconds, heat at 150°C on the heater for 30 minutes with repeated shaking at 5-minute intervals, and continue heating for an additional 30 minutes. Allow the flask to cool, and again weigh accurately. If the mass loss is less than 10 mg, use the upper layer of the mixture as the sample solution. Separately, take 65 mg of adipic acid, 2.0 mL of the internal standard solution and 2.0 mL of hydroiodic acid in another reaction flask, stopper tightly, and weigh accurately. Add 50  $\mu\text{L}$  of isopropyl iodide for assay, and again weigh accurately. Shake the reaction flask for 30 seconds, and use the upper layer of the content as the standard solution. Perform the test as directed under Gas Chromatography <2.02> with 1  $\mu\text{L}$  each of the sample solution and standard solution according to the following conditions, and calculate the ratios,  $Q_T$  and  $Q_S$ , of the peak area of isopropyl iodide to that of the internal standard.

$$\begin{aligned} \text{Amount (\% of hydroxypropoxy group (C}_3\text{H}_7\text{O}_2\text{))} \\ = (W_S/W_T) \times (Q_T/Q_S) \times 44.17 \end{aligned}$$

$W_S$ : Amount (mg) of isopropyl iodide in the standard solution.

$W_T$ : Amount (mg) of the sample

**Internal standard solution**—A solution of *n*-octane in *o*-xylene (4 in 100).

**Operating conditions**—

Detector: A thermal conductivity detector or hydrogen flame-ionization detector.

Column: A glass column about 3 mm in inside diameter and about 3 m in length, packed with siliceous earth for gas chromatography, 180 to 250  $\mu\text{m}$  in particle diameter, coated with methyl silicone polymer for gas chromatography at the ratio of 20%.

Column temperature: A constant temperature of about 100°C.

Carrier gas: Helium (for thermal-conductivity detector); helium or nitrogen (for hydrogen flame-ionization detector).

Flow rate: Adjust the flow rate so that the retention time of the internal standard is about 10 minutes.

Selection of column: Proceed with 1  $\mu\text{L}$  of the standard so-

lution under the above operating conditions. Use a column giving well-resolved peaks of isopropyl iodide and the internal standard in this order.

**Containers and storage** Containers—Well-closed containers.

## Low Substituted Hydroxypropylcellulose

低置換度ヒドロキシプロピルセルロース

Low Substituted Hydroxypropylcellulose is a low substituted hydroxypropyl ether of cellulose.

Low Substituted Hydroxypropylcellulose, when dried, contains not less than 5.0% and not more than 16.0% of hydroxypropoxy group ( $-\text{OC}_3\text{H}_6\text{OH}$ : 75.09).

**Description** Low Substituted Hydroxypropylcellulose occurs as a white to yellowish white powder or granules. It is odorless or has a slight, characteristic odor. It is tasteless.

It is practically insoluble in ethanol (95) and in diethyl ether.

It dissolves in a solution of sodium hydroxide (1 in 10), and produces a viscous solution.

It swells in water, in sodium carbonate TS and in 2 mol/L hydrochloric acid TS.

**Identification** (1) To 0.02 g of Low Substituted Hydroxypropylcellulose add 2 mL of water, shake, and produce a turbid solution. Add 1 mL of anthrone TS gently: a blue to blue-green color develops at the zone of contact.

(2) To 0.1 g of Low Substituted Hydroxypropylcellulose add 10 mL of water, stir and produce a turbid solution. Add 1 g of sodium hydroxide, shake until it becomes homogeneous, and use this solution as the sample solution. To 0.1 mL of the sample solution add 9 mL of diluted sulfuric acid (9 in 10), shake well, heat in a water bath for exactly 3 minutes, immediately cool in an ice bath, add carefully 0.6 mL of ninhydrin TS, shake well, and allow to stand at 25°C: a red color develops at first, and it changes to purple within 100 minutes.

(3) To 5 mL of the sample solution obtained in (2) add 10 mL of a mixture of acetone and methanol (4:1), and shake: a white, flocculent precipitate is produced.

**pH** <2.54> To 1.0 g of Low Substituted Hydroxypropylcellulose add 100 mL of freshly boiled and cooled water, and shake: the pH of the solution is between 5.0 and 7.5.

**Purity** (1) Chloride <1.03>—To 0.5 g of Low Substituted Hydroxypropylcellulose add 30 mL of hot water, stir well, heat on a water bath for 10 minutes, and filter the supernatant liquid by decantation while being hot. Wash the residue thoroughly with 50 mL of hot water, combine the washings with the filtrate, and add water to make 100 mL after cooling. To 5 mL of the sample solution add 6 mL of dilute nitric acid and water to make 50 mL, and perform the test using this solution as the test solution. Prepare the control solution with 0.25 mL of 0.01 mol/L hydrochloric acid VS (not more than 0.335%).

(2) Heavy metals <1.07>—Proceed with 2.0 g of Low Substituted Hydroxypropylcellulose according to Method 2, and perform the test. Prepare the control solution with 2.0

mL of Standard Lead Solution (not more than 10 ppm).

(3) Arsenic <1.11>—Prepare the test solution with 1.0 g of Low Substituted Hydroxypropylcellulose, according to Method 3, and perform the test (not more than 2 ppm).

**Loss on drying** <2.41> Not more than 6.0% (1 g, 105°C, 1 hour).

**Residue on ignition** <2.44> Not more than 1.0% (1 g).

**Assay** (i) Apparatus—Reaction flask: A 5-mL screw-cap pressure-tight glass bottle, having an inverted conical bottom inside, 20 mm in outside diameter, 50 mm in height up to the neck, 2 mL in capacity up to a height of about 30 mm, equipped with a pressure-tight septum of heat-resisting resin and also with an inside stopper or sealer of fluoroplastic.

Heater: A square-shaped aluminum block 60 to 80 mm thick, having holes 20.6 mm in diameter and 32 mm in depth, capable of maintaining the inside temperature within  $\pm 1^\circ\text{C}$ .

(ii) Procedure—Weigh accurately about 65 mg of Low Substituted Hydroxypropylcellulose, previously dried, transfer to the reaction flask, add 65 mg of adipic acid, 2.0 mL of the internal standard solution and 2.0 mL of hydroiodic acid, stopper the flask tightly, and weigh accurately. Shake the flask for 30 seconds, heat at 150°C on the heater for 30 minutes with repeated shaking at 5-minute intervals, and continue heating for an additional 30 minutes. Allow the flask to cool, and again weigh accurately. If the mass loss is less than 10 mg, use the upper layer of the mixture as the sample solution. Separately, take 65 mg of adipic acid, 2.0 mL of the internal standard solution and 2.0 mL of hydroiodic acid in another reaction flask, stopper tightly, and weigh accurately. Add 15  $\mu\text{L}$  of isopropyl iodide for assay, and again weigh accurately. Shake the reaction flask for 30 seconds, and use the upper layer of the content as the standard solution. Perform the test as directed under Gas Chromatography <2.02> with 2  $\mu\text{L}$  each of the sample solution and standard solution according to the following conditions, and calculate the ratios,  $Q_T$  and  $Q_S$ , of the peak area of isopropyl iodide to that of the internal standard.

$$\begin{aligned} \text{Amount (\% of hydroxypropoxy group (C}_3\text{H}_7\text{O}_2\text{))} \\ = (W_S/W_T) \times (Q_T/Q_S) \times 44.17 \end{aligned}$$

$W_S$ : Amount (mg) of isopropyl iodide in the standard solution.

$W_T$ : amount (mg) of the sample

**Internal standard solution**—A solution of *n*-octane in *o*-xylene (1 in 50).

**Operating conditions**—

Detector: A thermal conductivity detector or hydrogen flame-ionization detector.

Column: A glass column about 3 mm in inside diameter and about 3 m in length, packed with siliceous earth for gas chromatography, 180 to 250  $\mu\text{m}$  in particle diameter, coated with methyl silicone polymer for gas chromatography at the ratio of 20%.

Column temperature: A constant temperature of about 100°C.

Carrier gas: Helium (for thermal-conductivity detector); helium or nitrogen (for hydrogen flame-ionization detector).

Flow rate: Adjust the flow rate so that the retention time of the internal standard is about 10 minutes.


Selection of column: Proceed with 2  $\mu\text{L}$  of the standard solution under the above operating conditions. Use a column

giving well-resolved peaks of isopropyl iodide and the internal standard in this order.

**Containers and storage** Containers—Tight containers.

## Hydroxyzine Hydrochloride

ヒドロキシジン塩酸塩



$C_{21}H_{27}ClN_2O_2 \cdot 2HCl$ : 447.83

2-(2- {4-[(*RS*)-(4-Chlorophenyl)phenylmethyl]piperazin-1-yl} ethoxy)ethanol dihydrochloride [2192-20-3]

Hydroxyzine Hydrochloride, when dried, contains not less than 98.5% of  $C_{21}H_{27}ClN_2O_2 \cdot 2HCl$ .

**Description** Hydroxyzine Hydrochloride occurs as a white, crystalline powder. It is odorless, and has a bitter taste.

It is very soluble in water, freely soluble in methanol, in ethanol (95) and in acetic acid (100), very slightly soluble in acetic anhydride, and practically insoluble in diethyl ether.

Melting point: about 200°C (with decomposition).

**Identification (1)** To 5 mL of a solution of Hydroxyzine Hydrochloride (1 in 100) add 2 to 3 drops of ammonium thiocyanate-cobaltous nitrate TS: a blue precipitate is formed.

**(2)** Determine the absorption spectrum of a solution of Hydroxyzine Hydrochloride in methanol (1 in 100,000) as directed under Ultraviolet-visible Spectrophotometry <2.24>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wavelengths.

**(3)** A solution of Hydroxyzine Hydrochloride (1 in 10) responds to the Qualitative Tests <1.09> for chloride.

**pH <2.54>** Dissolve 1.0 g of Hydroxyzine Hydrochloride in 20 mL of water: the pH of this solution is between 1.3 and 2.5.

**Purity (1)** Clarity and color of solution—Dissolve 1.0 g of Hydroxyzine Hydrochloride in 10 mL of water: the solution is clear and colorless.

**(2)** Heavy metals <1.07>—Proceed with 1.0 g of Hydroxyzine Hydrochloride according to Method 2, and perform the test. Prepare the control solution with 2.0 mL of Standard Lead Solution (not more than 20 ppm).

**(3)** Related substances—Dissolve 0.20 g of Hydroxyzine Hydrochloride in 10 mL of methanol, and use this solution as the sample solution. Pipet 1 mL of the sample solution, add methanol to make exactly 200 mL, and use this solution as the standard solution. Perform the test with these solutions as directed under Thin-layer Chromatography <2.03>. Spot 5  $\mu$ L each of the sample solution and standard solution on a plate of silica gel for thin-layer chromatography. Develop the plate with a mixture of ethyl acetate, ethanol (95) and ammonia solution (28) (150:95:1) to a distance of about 10 cm, and air-dry the plate. Allow the plate to stand in iodine vapor: the

spots other than the principal spot from the sample solution are not more intense than the spot from the standard solution.

**Loss on drying <2.41>** Not more than 3.0% (1 g, 105°C, 2 hours).

**Residue on ignition <2.44>** Not more than 0.2% (1 g).

**Assay** Weigh accurately about 0.1 g of Hydroxyzine Hydrochloride, previously dried, dissolve in 60 mL of a mixture of acetic anhydride and acetic acid (100) (7:3), and titrate <2.50> with 0.1 mol/L perchloric acid VS (potentiometric titration). Perform a blank determination, and make any necessary correction.


Each mL of 0.1 mol/L perchloric acid VS

= 22.39 mg of  $C_{21}H_{27}ClN_2O_2 \cdot 2HCl$

**Containers and storage** Containers—Tight containers.

## Hydroxyzine Pamoate

ヒドロキシジンパモ酸塩



$C_{21}H_{27}ClN_2O_2 \cdot C_{23}H_{16}O_6$ : 763.27

2-(2- {4-[(*RS*)-(4-Chlorophenyl)phenylmethyl]piperazin-1-yl} ethoxy)ethanol mono[4,4'-methylenebis(3-hydroxy-2-naphthoate)] (1/1) [10246-75-0]

Hydroxyzine Pamoate contains not less than 98.0% of  $C_{21}H_{27}ClN_2O_2 \cdot C_{23}H_{16}O_6$ , calculated on the anhydrous basis.

**Description** Hydroxyzine Pamoate occurs as a light yellow, crystalline powder. It is odorless, and has a slightly bitter taste.

It is freely soluble in *N,N*-dimethylformamide, slightly soluble in acetone, and practically insoluble in water, in methanol, in ethanol (95) and in diethyl ether.

**Identification (1)** To 0.1 g of Hydroxyzine Pamoate add 25 mL of sodium hydroxide TS, and shake well. Extract with 20 mL of chloroform, and use the chloroform layer as the sample solution. Use the water layer for test (4). To 5 mL of the sample solution add 2 mL of ammonium thiocyanate-cobaltous nitrate TS, shake well, and allow to stand: a blue color is produced in the chloroform layer.

**(2)** Evaporate 2 mL of the sample solution obtained in (1) on a water bath to dryness, and dissolve the residue in 0.1 mol/L hydrochloric acid TS to make 500 mL. Determine the absorption spectrum of the solution as directed under Ultraviolet-visible Spectrophotometry <2.24>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wavelengths.

**(3)** Perform the test with Hydroxyzine Pamoate as direct-

ed under Flame Coloration Test <1.04> (2): a green color appears.

(4) To 1 mL of the water layer obtained in (1), add 2 mL of 1 mol/L hydrochloric acid TS: a yellow precipitate is produced. Collect the precipitate, dissolve the precipitate in 5 mL of methanol, and add 1 drop of iron (III) chloride TS: a green color is produced.

**Purity** (1) Clarity and color of solution—Dissolve 1.0 g of Hydroxyzine Pamoate in 10 mL of *N,N*-dimethylformamide: the solution is clear, and shows a slightly greenish, light yellow-brown color.

(2) Chloride <1.03>—To 0.3 g of Hydroxyzine Pamoate add 6 mL of dilute nitric acid and 10 mL of water, shake for 5 minutes, and filter. Wash the residue with two 10-mL portions of water, combine the washings with the filtrate, and add water to make 50 mL. Perform the test using this solution as the test solution. Prepare the control solution with 0.80 mL of 0.01 mol/L hydrochloric acid VS (not more than 0.095%).

(3) Heavy metals <1.07>—Proceed with 1.0 g of Hydroxyzine Pamoate according to Method 2, and perform the test. Prepare the control solution with 2.0 mL of Standard Lead Solution (not more than 20 ppm).

(4) Arsenic <1.11>—Prepare the test solution with 2.0 g of Hydroxyzine Pamoate according to Method 3, and perform the test (not more than 1 ppm).

(5) Related substances—Dissolve 0.40 g of Hydroxyzine Pamoate in 10 mL of a mixture of sodium hydroxide TS and acetone (1:1), and use the solution as the sample solution. Pipet 1 mL of this solution, add a mixture of sodium hydroxide TS and acetone (1:1) to make exactly 20 mL. Pipet 5 mL of this solution, add a mixture of sodium hydroxide TS and acetone (1:1) to make exactly 50 mL, and use the solution as the standard solution. Perform the test with these solutions as directed under Thin-layer Chromatography <2.03>. Spot 5  $\mu$ L each of the sample solution and standard solution on a plate of silica gel for thin-layer chromatography. Develop the plate with a mixture of ethyl acetate, ethanol (95) and ammonia TS (150:95:1) to a distance of about 10 cm, and air-dry the plate. Spray evenly hydrogen hexachloroplatinate (IV)-potassium iodide TS on the plate: the spots other than the spots from hydroxyzine and pamoic acid obtained from the sample solution are not more intense than the spot from the standard solution.

**Water** <2.48> Not more than 3.0% (1 g, direct titration).

**Residue on ignition** <2.44> Not more than 0.5% (1 g).


**Assay** Weigh accurately about 0.6 g of Hydroxyzine Pamoate, add 25 mL of sodium hydroxide TS, shake well, and extract with six 25-mL portions of chloroform. Filter each extract through 5 g of anhydrous sodium sulfate on a pledget of absorbent cotton. Combine the chloroform extracts, and evaporate the combined chloroform extracts on a water bath to about 30 mL. Add 30 mL of acetic acid (100), and titrate <2.50> with 0.1 mol/L perchloric acid VS until the color of the solution changes from purple through blue to blue-green (indicator: 2 drops of crystal violet TS). Perform a blank determination.

Each mL of 0.1 mol/L perchloric acid VS  
= 38.16 mg of  $C_{21}H_{27}ClN_2O_2 \cdot C_{23}H_{16}O_6$

**Containers and storage** Containers—Tight containers.

## Hymecromone

ヒメクロモン



$C_{10}H_8O_3$ : 176.17

7-Hydroxy-4-methylchromen-2-one [90-33-5]

Hymecromone, when dried, contains not less than 98.0% of  $C_{10}H_8O_3$ .

**Description** Hymecromone occurs as white crystals or crystalline powder. It is odorless and tasteless.

It is freely soluble in *N,N*-dimethylformamide, sparingly soluble in ethanol (95), in ethanol (99.5) and in acetone, slightly soluble in diethyl ether, and practically insoluble in water.

**Identification** (1) Dissolve 2 mg of Hymecromone in 5 mL of ammonia-ammonium chloride buffer solution, pH 11.0: the solution shows an intense blue-purple fluorescence.

(2) Dissolve 0.025 g of Hymecromone in 5 mL of diluted ethanol (95) (1 in 2), and add 1 drop of iron (III) chloride TS: initially a blackish brown color develops, and when allowed to stand the color changes to yellow-brown.

(3) Determine the absorption spectrum of a solution of Hymecromone in ethanol (99.5) (1 in 250,000) as directed under Ultraviolet-visible Spectrophotometry <2.24>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wavelengths.

(4) Determine the infrared absorption spectrum of Hymecromone, previously dried, as directed in the potassium bromide disk method under Infrared Spectrophotometry <2.25>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wave numbers.

**Melting point** <2.60> 187 – 191°C

**Purity** (1) Chloride <1.03>—Dissolve 0.8 g of Hymecromone in 40 mL of a mixture of acetone and water (2:1), and add 6 mL of dilute nitric acid and a mixture of acetone and water (2:1) to make 50 mL. Perform the test using this solution as the test solution. Prepare the control solution as follows: to 0.25 mL of 0.01 mol/L hydrochloric acid VS add 6 mL of dilute nitric acid and a mixture of acetone and water (2:1) to make 50 mL (not more than 0.011%).

(2) Sulfate <1.14>—Dissolve 0.8 g of Hymecromone in 40 mL of a mixture of acetone and water (2:1), and add 1 mL of dilute hydrochloric acid and a mixture of acetone and water (2:1) to make 50 mL. Perform the test using this solution as the test solution. Prepare the control solution as follows: to 0.40 mL of 0.005 mol/L sulfuric acid VS add 1 mL of dilute hydrochloric acid and a mixture of acetone and water (2:1) to make 50 mL (not more than 0.024%).

(3) Heavy metals <1.07>—Proceed with 2.0 g of Hymecromone according to Method 2, and perform the test. Prepare the control solution with 2.0 mL of Standard Lead

Solution (not more than 10 ppm).

(4) Arsenic <1.11>—Prepare the test solution with 1.0 g of Hymecromone according to Method 3, and perform the test (not more than 2 ppm).

(5) Related substances—Dissolve 80 mg of Hymecromone in 10 mL of ethanol (95), and use this solution as the sample solution. Pipet 1 mL of the sample solution, and add ethanol (95) to make exactly 50 mL. Pipet 1 mL of this solution, add ethanol (95) to make exactly 20 mL, and use this solution as the standard solution. Perform the test with these solutions as directed under Thin-layer Chromatography <2.03>. Spot 10  $\mu$ L each of the sample solution and standard solution on a plate of silica gel for thin-layer chromatography. Develop the plate with a mixture of chloroform and ethanol (95) (10:1) to a distance of about 10 cm, and air-dry the plate. Allow the plate to stand in iodine vapor for 5 minutes: the spots other than the principal spot from the sample solution are not more intense than the spot from the standard solution.

**Loss on drying** <2.41> Not more than 0.5% (1 g, 105°C, 4 hours).

**Residue on ignition** <2.44> Not more than 0.1% (1 g).

**Assay** Weigh accurately about 0.25 g of Hymecromone, previously dried, dissolve in 90 mL of *N,N*-dimethylformamide, and titrate <2.50> with 0.1 mol/L tetramethylammonium hydroxide VS (potentiometric titration). Separately, perform a blank determination with a solution prepared by adding 14 mL of water to 90 mL of *N,N*-dimethylformamide, and make any necessary correction.

Each mL of 0.1 mol/L tetramethylammonium hydroxide VS  
= 17.62 mg of C<sub>10</sub>H<sub>8</sub>O<sub>3</sub>

**Containers and storage** Containers—Tight containers.

## Hypromellose

### Hydroxypropylmethylcellulose

ヒプロメロース

Cellulose, 2-hydroxypropyl methyl ether  
[9004-65-3]

This monograph is harmonized with the European Pharmacopoeia and the U.S. Pharmacopoeia. The parts of the text that are not harmonized are marked with symbol (♦ ♦).

Hypromellose is a methyl and hydroxypropyl mixed ether of cellulose.

There are four substitution types of Hypromellose, 1828, 2208, 2910 and 2910. They contain methoxy (-OCH<sub>3</sub>; 31.03) and hydroxypropoxy (-OC<sub>3</sub>H<sub>6</sub>OH; 75.09) groups conforming to the limits for the types of Hypromellose shown in the table below, calculated on the dried basis.

The viscosity is shown in millipascal second (mPa·s) on the label, together with the substitution type.

| Substitution Type | Methoxy Group (%) |      | Hydroxypropoxy Group (%) |      |
|-------------------|-------------------|------|--------------------------|------|
|                   | Min.              | Max. | Min.                     | Max. |
| 1828              | 16.5              | 20.0 | 23.0                     | 32.0 |
| 2208              | 19.0              | 24.0 | 4.0                      | 12.0 |
| 2906              | 27.0              | 30.0 | 4.0                      | 7.5  |
| 2910              | 28.0              | 30.0 | 7.0                      | 12.0 |

♦**Description** Hypromellose occurs as a white to yellowish white, powder or granules.

It is practically insoluble in ethanol (99.5).

It swells with water and becomes a clear or slightly turbid, viscous solution.♦

**Identification** (1) Disperse evenly 1.0 g of Hypromellose over the surface of 100 mL of water in a beaker, while gently tapping the top of the container, if necessary, and allow the beaker to stand: it aggregates on the surface of water.

(2) Add 1.0 g of Hypromellose to 100 mL of hot water, and stir: it becomes a suspension. Cool the suspension to 10°C, and stir: the resulting liquid is a clear or a slightly cloudy, viscous fluid.

(3) To 0.1 mL of the viscous fluid obtained in (2) add 9 mL of diluted sulfuric acid (9 in 10), stir, heat in a water bath for exactly 3 minutes, and immediately cool in ice water. Add carefully 0.6 mL of ninhydrin TS, stir, and allow to stand at 25°C: the solution shows a light red color first, then changes to purple color within 100 minutes.

(4) Pour and spread out 2 to 3 mL of the viscous fluid obtained in (2) onto a glass plate, and allow the water to evaporate: a transparent film results.

(5) Pipet 50 mL of water, add exactly 50 mL of the viscous fluid obtained in (2), and warm to rise the temperature at a rate of 2 to 5°C per minute while stirring: the temperature, when a white turbidity of the solution starts to increase, is not less than 50°C.

**Viscosity** <2.53> Method I: Apply to Hypromellose having a labeled viscosity of less than 600 mPa·s. Put exactly an amount of Hypromellose, equivalent to 4.000 g on the dried basis, in a tared, wide-mouth bottle, add hot water to make 200.0 g, stopper the bottle, stir by mechanical means at 350- to 450-revolutions per minute for 10 to 20 minutes to get a homogeneous dispersion. If necessary, take off the sample attached on the walls of the bottle, put them in the dispersed solution, and dissolve by continuing the stirring in a water bath not exceeding 10°C for 20 to 40 minutes while stirring. Add cooled water, if necessary, to make 200.0 g, and use this solution as the sample solution. Centrifuge the solution if necessary to expel any entrapped air bubbles. Perform the test with the sample solution at 20 ± 0.1°C as directed in Method I under Viscosity Determination: not less than 80% and not more than 120% of the labeled viscosity.

Method II: Apply to Hypromellose having a labeled viscosity of not less than 600 mPa·s. Put exactly an amount of Hypromellose, equivalent to 10.00 g on the dried basis, in a tared, wide-mouth bottle, add hot water to make 500.0 g, stopper the bottle, and prepare the sample solution in the same manner as directed in Method I. Perform the test with the sample solution at 20 ± 0.1°C as directed in Method II (2)

under Viscosity Determination, using a single cylinder-type rotational viscometer, according to the following operating conditions: not less than 75% and not more than 140% of the labeled viscosity.

**Operating conditions—**

Apparatus: Brookfield type viscometer LV model

Rotor No., rotation frequency, and conversion factor: According to the following table, depending on the labeled viscosity.

| Labeled viscosity<br>(mPa·s)         | Rotor<br>No. | Rotation<br>frequency<br>/min | Conversion<br>factor |
|--------------------------------------|--------------|-------------------------------|----------------------|
| Not less than 600 and less than 1400 | 3            | 60                            | 20                   |
| " 1400 " 3500                        | 3            | 12                            | 100                  |
| " 3500 " 9500                        | 4            | 60                            | 100                  |
| " 9500 " 99500                       | 4            | 6                             | 1000                 |
| " 99500                              | 4            | 3                             | 2000                 |

Procedure of apparatus: Read value after 2 minutes of rotation, and stop the rotation for 2 minutes. Repeat this procedure more two times, and average three observed values.

**pH** <2.54> Allow the sample solution obtained in the Viscosity to stand at  $20 \pm 2^\circ\text{C}$  for 5 minutes: the pH of the solution thus obtained is between 5.0 and 8.0.

**Purity** Heavy metals—Put 1.0 g of Hypromellose in a 100-mL Kjeldahl flask, add a sufficient amount of a mixture of nitric acid and sulfuric acid (5:4) to wet the sample, and heat gently. Repeat this procedure until to use totally 18 mL of the mixture of nitric acid and sulfuric acid. Then boil gently until the solution changes to black. After cooling, add 2 mL of nitric acid, and heat until the solution changes to black. Repeat this procedure until the solution no longer changes to black, and heat strongly until dense white fumes are evolved. After cooling, add 5 mL of water, boil gently until dense white fumes are evolved, then heat until the volume of the solution becomes to 2 to 3 mL. After cooling, if the solution reveals yellow color by addition of 5 mL of water, add 1 mL of hydrogen peroxide (30), and heat until the volume of the solution becomes to 2 to 3 mL. After cooling, dilute the solution with 2 to 3 mL of water, transfer to a Nessler tube, add water to make 25 mL, and use this solution as the test solution. Separately, put 2.0 mL of Standard Lead Solution in a 100-mL kjeldahl flask, add 18 mL of the mixture of nitric acid and sulfuric acid (5:4) and an amount of nitric acid equal to that used for preparation of the test solution, and heat until white fumes are evolved. After cooling, add 10 mL of water. In the case where hydrogen peroxide (30) is added for the preparation of the test solution, add the same amount of hydrogen peroxide (30), then proceed in the same manner for preparation of the test solution, and use so obtained solution as the control solution. Adjust the test solution and the control solution to pH 3.0 to 4.0 with ammonia solution (28), and add water to make 40 mL, respectively. To these solutions add 1.2 mL of thioacetamide-alkaline glycerin TS, 2 mL of acetate buffer solution, pH 3.5 and water to make 50 mL, separately. After allowing to stand for 5 minutes, observe vertically both tubes on a white background: the color obtained with the test solution is not more intense than that with the control solution (not more than 20 ppm).

**Loss on drying** <2.41> Not more than 5.0% (1 g,  $105^\circ\text{C}$ , 1 hour).

**Residue on ignition** <2.44> Not more than 1.5% (1 g).

**Assay** (i) Apparatus—Reaction bottle: A 5-mL pressure-tight glass vial, having 20 mm in outside diameter and 50 mm in height, the neck 20 mm in outside diameter and 13 mm in inside diameter, equipped with a septum of butyl-rubber processed the surface with fluoroplastics, which can be fixed tightly to vial with aluminum cap, or equivalent.

Heater: A square-shaped aluminum block, having holes 20 mm in diameter and 32 mm in depth, adopted to the reaction bottle. Capable of stirring the content of the reaction bottle by means of magnetic stirrer or of reciprocal shaker about 100 times per minute.

(ii) Procedure—Weigh accurately about 65 mg of Hypromellose, transfer to the reaction bottle, add 0.06 to 0.10 g of adipic acid, 2.0 mL of the internal standard solution and 2.0 mL of hydroiodic acid, stopper the bottle immediately, and weigh accurately. Stir or shake for 60 minutes while heating so that the temperature of the bottle content is  $130 \pm 2^\circ\text{C}$ . In the case when the stirrer or shaker is not available, heat for 30 minutes with repeated shaking at 5-minute intervals by hand, and continue heating for an additional 30 minutes. Allow the bottle to cool, and again weigh accurately. If the mass loss is less than 0.50% or there is no evidence of a leak, use the upper layer of the mixture as the sample solution. Separately, put 0.06 to 0.10 g of adipic acid, 2.0 mL of the internal standard solution and 2.0 mL of hydroiodic acid in a reaction bottle, stopper the bottle immediately, and weigh accurately. Add 45  $\mu\text{L}$  of iodomethane for assay and 15 to 22  $\mu\text{L}$  of isopropyl iodide for assay through the septum using micro-syringe with weighing accurately every time, stir thoroughly, and use the upper layer of the mixture as the standard solution. Perform the test with 1 to 2  $\mu\text{L}$  each of the sample solution and standard solution as directed under Gas Chromatography <2.02> according to the following conditions, and calculate the ratios,  $Q_{\text{Ta}}$  and  $Q_{\text{Tb}}$ , of the peak area of iodomethane and isopropyl iodide to that of the internal standard obtained from the sample solution, and  $Q_{\text{Sa}}$  and  $Q_{\text{Sb}}$ , of the peak area of iodomethane and isopropyl iodide to that of the internal standard from the standard solution.

$$\begin{aligned} \text{Content (\%)} & \text{ of methoxy group } (-\text{CH}_3\text{O}) \\ & = (Q_{\text{Ta}}/Q_{\text{Sa}}) \times (W_{\text{Sa}}/W) \times 21.86 \end{aligned}$$

$$\begin{aligned} \text{Content (\%)} & \text{ of hydroxypropoxy group } (-\text{C}_3\text{H}_7\text{O}_2) \\ & = (Q_{\text{Tb}}/Q_{\text{Sb}}) \times (W_{\text{Sb}}/W) \times 44.17 \end{aligned}$$

$W_{\text{Sa}}$ : Amount (mg) of iodomethane for assay.

$W_{\text{Sb}}$ : Amount (mg) of isopropyl iodide for assay.

$W$ : Amount (mg) of sample, calculated on the dried basis

**Internal standard solution**—A solution of *n*-octane in *o*-xylene (3 in 100).

**Operating conditions—**

Detector: A thermal conductivity detector or hydrogen flame-ionization detector.

Column: A glass column 3–4 mm in inside diameter and 1.8–3 m in length, packed with siliceous earth for gas chromatography, 125 to 150  $\mu\text{m}$  in diameter, coated with methyl silicone polymer at the ratio of 10–20%.

Column temperature: A constant temperature of about  $100^\circ\text{C}$ .

Carrier gas: Helium for thermal conductivity detector, or Helium or Nitrogen for hydrogen, flame-ionization detector.

Flow rate: Adjust the flow rate so that the retention time of the internal standard is about 10 minutes.

*System suitability—*

System performance: When the procedure is run with 1 – 2  $\mu$ L of the standard solution under the above operating conditions, iodomethane, isopropyl iodide and the internal standard are eluted in this order, with complete separation of these peaks.

♦**Containers and storage** Containers—Well-closed containers. ♦

## Hypromellose Phthalate

ヒプロメロースフタル酸エステル

Hydroxypropyl methylcellulose benzene-1,2-dicarboxylate  
[9050-31-1]

Hypromellose Phthalate is a monophthalic acid ester of hypromellose.

It contains methoxy group ( $-\text{OCH}_3$ : 31.03), hydroxypropoxy group ( $-\text{OC}_3\text{H}_6\text{OH}$ : 75.09), and carboxybenzoyl group ( $-\text{COC}_6\text{H}_4\text{COOH}$ : 149.12).

There are two substitution types, 200731 and 220824, and each type contains indicated amount of carboxybenzoyl group in the accompanying table, calculated on the anhydrous basis.

| Substitution Type | Carboxybenzoyl group (%) |      |
|-------------------|--------------------------|------|
|                   | Min.                     | Max. |
| 200731            | 27.0                     | 35.0 |
| 220824            | 21.0                     | 27.0 |

Its substitution type and its kinematic viscosity are shown in square mm per second ( $\text{mm}^2/\text{s}$ ) on the label.

**Description** Hypromellose Phthalate occurs as white powder or granules. It is odorless and tasteless.

It is practically insoluble in acetonitrile, in ethanol (99.5), and in hexane.

It becomes a viscous liquid when a mixture of methanol and dichloromethane (1:1) or a mixture of ethanol (99.5) and acetone (1:1) is added.

It dissolves in sodium hydroxide TS.

**Identification** Determine the infrared absorption spectrum of Hypromellose Phthalate as directed in the potassium bromide disk method under Infrared Spectrophotometry <2.25>, and compare the spectrum with the Reference Spectrum or spectrum of Hypromellose Phthalate Reference Standard: both spectra exhibit similar intensities of absorption at the same wave numbers.

**Viscosity** <2.53> To 10 g of Hypromellose Phthalate, previously dried at  $105^\circ\text{C}$  for 1 hour, add 90 g of a mixture of methanol and dichloromethane in equal mass ratio, and stir to dissolve. Determine the viscosity at  $20 \pm 0.1^\circ\text{C}$  as directed in Method I under Viscosity Determination: the viscosity is not less than 80% and not more than 120% of the labeled unit.

**Purity** (1) Chloride <1.03>—Dissolve 1.0 g of Hypromellose Phthalate in 40 mL of 0.2 mol/L sodium hydroxide VS,

add 1 drop of phenolphthalein TS, and add dilute nitric acid dropwise with vigorous stirring until the red color is discharged. Further add 20 mL of dilute nitric acid with stirring. Heat on a water bath with stirring until the gelatinous precipitate formed turns to granular particles. After cooling, centrifuge, and take off the supernatant liquid. Wash the precipitate with three 20-mL portions of water by centrifuging each time, combine the supernatant liquid and the washings, add water to make 200 mL, and filter. Perform the test with 50 mL of the filtrate. Control solution: To 0.50 mL of 0.01 mol/L hydrochloric acid VS add 10 mL of 0.2 mol/L sodium hydroxide VS, 7 mL of dilute nitric acid and water to make 50 mL (not more than 0.07%).

(2) Heavy metals <1.07>—Proceed with 2.0 g of Hypromellose Phthalate according to Method 2, and perform the test. Prepare the control solution with 2.0 mL of Standard Lead Solution (not more than 10 ppm).

(3) Phthalic acid—Weigh accurately about 0.2 g of Hypromellose Phthalate, add about 50 mL of acetonitrile to dissolve partially with the aid of ultrasonicator, add 10 mL of water, and dissolve further with the ultrasonicator. After cooling, add acetonitrile to make exactly 100 mL, and use this solution as the sample solution. Separately, weigh accurately about 12.5 mg of phthalic acid, dissolve in about 125 mL of acetonitrile by stirring, add 25 mL of water and acetonitrile to make exactly 250 mL, and use this solution as the standard solution. Perform the test with 10  $\mu$ L each of the sample solution and standard solution as directed under Liquid Chromatography <2.01> according to the following conditions, and determine the peak areas of phthalic acid,  $A_T$  and  $A_S$ , of both solutions: amount of phthalic acid ( $\text{C}_8\text{H}_6\text{O}_4$ : 166.13) is not more than 1.0%.

$$\begin{aligned} &\text{Amount (\%)} \text{ of phthalic acid} \\ &= (C/W) \times (A_T/A_S) \times 10 \end{aligned}$$

$C$ : concentration of phthalic acid in the standard solution ( $\mu\text{g/mL}$ )

$W$ : amount (mg) of the sample, calculated on the anhydrous basis

*Operating conditions—*

Detector: An ultraviolet absorption photometer (wavelength: 235 nm).

Column: A stainless steel column about 4 mm in inside diameter and about 25 cm in length, packed with octadecylsilanized silica gel for liquid chromatography (3 to 10  $\mu\text{m}$  in particle diameter).

Column temperature: A constant temperature of about  $20^\circ\text{C}$ .

Mobile phase: A mixture of 0.1 mol/L cyanoacetic acid and acetonitrile (17:3).

Flow rate: About 2.0 mL per minute.

System repeatability: Repeat the test six times with the standard solution under the above operating conditions: the relative standard deviation of the peak area of phthalic acid is not more than 1.0%.

**Water** <2.48> Not more than 5.0% (1 g, direct titration, using a mixture of ethanol (99.5) and dichloromethane (3:2) instead of methanol for Karl Fischer method).

**Residue on ignition** <2.44> Not more than 0.2% (1 g).

**Assay** Weigh accurately about 1 g of Hypromellose Phthalate, dissolve in 50 mL of a mixture of ethanol (95), acetone



and water (2:2:1), and titrate <2.50> with 0.1 mol/L sodium hydroxide VS (indicator: 2 drops of phenolphthalein TS). Perform a blank determination, and make any necessary correction.

$$\text{Amount (\% of carboxybenzoyl group (C}_8\text{H}_5\text{O}_3\text{))} \\ = \{(0.01 \times 149.1 \times V)/W\} - \{(2 \times 149.1 \times P)/166.1\}$$

*P*: amount (%) of phthalic acid obtained in the Purity (3)


*V*: amount (mL) of 0.1 mol/L sodium hydroxide VS consumed

*W*: amount (g) of the sample, calculated on the anhydrous basis

**Containers and storage** Containers—Tight containers.

## Ibuprofen

イブプロフェン



$\text{C}_{13}\text{H}_{18}\text{O}_2$ : 206.28

(2*RS*)-2-[4-(2-Methylpropyl)phenyl]propanoic acid  
[15687-27-1]

Ibuprofen, when dried, contains not less than 98.5% of  $\text{C}_{13}\text{H}_{18}\text{O}_2$ .

**Description** Ibuprofen occurs as a white crystalline powder.

It is freely soluble in ethanol (95) and in acetone, and practically insoluble in water.

It dissolves in dilute sodium hydroxide TS.

**Identification** (1) Determine the absorption spectrum of a solution of Ibuprofen in dilute sodium hydroxide TS (3 in 20,000) as directed under Ultraviolet-visible Spectrophotometry <2.24>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wavelengths.

(2) Determine the infrared absorption spectrum of Ibuprofen, previously dried, as directed in the potassium bromide disk method under Infrared Spectrophotometry <2.25>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wave numbers.

**Melting point** <2.60> 75 – 77°C

**Purity** (1) Heavy metals <1.07>—Proceed with 3.0 g of Ibuprofen according to Method 2, and perform the test. Prepare the control solution with 3.0 mL of Standard Lead Solution (not more than 10 ppm).

(2) Arsenic <1.11>—Prepare the test solution with 1.0 g of Ibuprofen according to Method 3, and perform the test (not more than 2 ppm).

(3) Related substances—Dissolve 0.50 g of Ibuprofen in exactly 5 mL of acetone, and use this solution as the sample solution. Pipet 1 mL of the sample solution, add acetone to make exactly 100 mL, and use this solution as the standard solution. Perform the test with these solutions as directed under Thin-layer Chromatography <2.03>. Spot 5  $\mu\text{L}$  each of the sample solution and standard solution on a plate of silica

gel with fluorescent indicator for thin-layer chromatography. Develop the plate with a mixture of hexane, ethyl acetate and acetic acid (100) (15:5:1) to a distance of about 10 cm, and air-dry the plate. Examine the plate under ultraviolet light (main wavelength: 254 nm): the spots other than the principal spot from the sample solution are not more intense than the spot from the standard solution.

**Loss on drying** <2.41> Not more than 0.5% (1 g, in vacuum at a pressure not exceeding 0.67 kPa, phosphorus (V) oxide, 4 hours).

**Residue on ignition** <2.44> Not more than 0.1% (1 g).

**Assay** Weigh accurately about 0.5 g of Ibuprofen, previously dried, dissolve in 50 mL of ethanol (95), and titrate <2.50> with 0.1 mol/L sodium hydroxide VS (indicator: 3 drops of phenolphthalein TS). Perform a blank determination, and make any necessary correction.

Each mL of 0.1 mol/L sodium hydroxide VS  
= 20.63 mg of  $\text{C}_{13}\text{H}_{18}\text{O}_2$

**Containers and storage** Containers—Well-closed containers.

## Ichthammol

イクタモール

Ichthammol, calculated on the dried basis, contains not less than 2.5% of ammonia ( $\text{NH}_3$ : 17.030), not more than 8.0% of ammonium sulfate [ $(\text{NH}_4)_2\text{SO}_4$ : 132.14], and not less than 10.0% of total sulfur (as S: 32.07).

**Description** Ichthammol is a red-brown to blackish brown, viscous fluid. It has a characteristic odor.

It is miscible with water, and is partially soluble in ethanol (95) and in diethyl ether.

**Identification** (1) To 4 mL of a solution of Ichthammol (3 in 10) add 8 mL of hydrochloric acid: a yellow-brown to blackish brown, oily or resinous mass is produced. Cool the mass with ice to solidify, and discard the water layer. Wash the residue with diethyl ether: a part of the mass dissolves but it does not dissolve completely even when it is washed until almost no color develops in the washing. Perform the following tests with this residue.

(i) To 0.1 g of the residue add 1 mL of a mixture of diethyl ether and ethanol (95) (1:1): it dissolves.

(ii) To 0.1 g of the residue add 2 mL of water: it dissolves. To 1 mL of this solution add 0.4 mL of hydrochloric acid: a yellow-brown to blackish brown oily or resinous substance is produced.

(iii) To 1 mL of the solution obtained in (ii) add 0.3 g of sodium chloride: a yellow-brown or blackish brown oily or resinous substance is produced.

(2) Boil 2 mL of a solution of Ichthammol (1 in 10) with 2 mL of sodium hydroxide TS: the gas evolved changes moistened red litmus paper to blue.

**Loss on drying** <2.41> Not more than 50% (0.5 g, 105°C, 6 hours).

**Residue on ignition** <2.44> Not more than 0.5% (1 g).



$$= W_S \times (A_T/A_S) \times 1000$$

$W_S$ : Amount [mg (potency)] of Idarubicin Hydrochloride Reference Standard

**Operating conditions—**

**Detector:** An ultraviolet absorption photometer (wavelength: 254 nm).

**Column:** A stainless steel column 3.9 mm in inside diameter and 15 cm in length, packed with octadecylsilanized silica gel for liquid chromatography (4  $\mu$ m in particle diameter).

**Column temperature:** A constant temperature of about 35°C.

**Mobile phase:** Dissolve 10.2 g of potassium dihydrogenphosphate in a suitable amount of water, add 1 mL of phosphoric acid and water to make 750 mL, and add 250 mL of tetrahydrofuran. To 500 mL of this solution add 0.72 g of sodium lauryl sulfate and 0.5 mL of *N,N*-dimethyl-*n*-octylamine, and adjust to pH 4 with 2 mol/L sodium hydroxide TS.

**Flow rate:** Adjust the flow rate so that the retention time of idarubicin is about 15 minutes.

**System suitability—**

**System performance:** When the procedure is run with 20  $\mu$ L of the standard solution under the above operating conditions, the number of theoretical steps of the peak of idarubicin is not less than 3000 steps.

**System repeatability:** When the test is repeated 6 times with 20  $\mu$ L of the standard solution under the above operating conditions, the relative standard deviation of the peak areas of idarubicin is not more than 2.0%.

**Containers and storage** Containers—Tight containers.

## Idarubicin Hydrochloride for Injection

注射用イダルビシン塩酸塩

Idarubicin Hydrochloride for Injection is a preparation for injection, which is dissolved before use.

It contains not less than 90.0% and not more than 110.0% of the labeled amount of idarubicin hydrochloride ( $C_{26}H_{27}NO_9 \cdot HCl$ ; 533.95).

**Method of preparation** Prepare as directed under Injections, with Idarubicin Hydrochloride.

**Description** Idarubicin Hydrochloride for Injection occurs as yellow-red masses.

**Identification (1)** Dissolve an amount of Idarubicin Hydrochloride for Injection, equivalent to 2 mg (potency) of Idarubicin Hydrochloride according to the labeled amount, in 5 mL of sodium hydroxide TS: the solution shows a blue-purple color.

**(2)** Dissolve an amount of Idarubicin Hydrochloride for Injection, equivalent to 1 mg (potency) of Idarubicin Hydrochloride according to the labeled amount, in 1 mL of water, and add methanol to make 100 mL. Determine the absorption spectrum of the solution as directed under Ultraviolet-visible Spectrophotometry <2.24>: it exhibits maxima between 250 nm and 254 nm, between 285 nm and 289 nm,

between 480 nm and 484 nm, and between 510 nm and 520 nm.

**pH <2.54>** The pH of a solution prepared by dissolving an amount of Idarubicin Hydrochloride for Injection, equivalent to 5 mg (potency) of Idarubicin Hydrochloride according to the labeled amount, in 5 mL of water is between 5.0 and 7.0.

**Purity** Clarity and color of solution—Dissolve an amount of Idarubicin Hydrochloride for Injection, equivalent to 5 mg (potency) of Idarubicin Hydrochloride according to the labeled amount, in 5 mL of water: the solution is clear and yellow-red.

**Water <2.48>** Weigh accurately the mass of one Idarubicin Hydrochloride for Injection, add 5 mL of methanol for Karl Fischer method using a syringe, dissolve with thorough shaking, and perform the test with 4 mL of this solution as directed in the Volumetric titration (direct titration). Use 4 mL of methanol for Karl Fischer method as the blank. Determine the mass of the content from the difference between the mass of one Idarubicin Hydrochloride for Injection obtained above and the mass of its bottle and rubber stopper, which are weighed accurately after washing with water then with ethanol (95), drying at 105°C for 1 hour and allowing to cool to room temperature in a desiccator (not more than 4.0%).

**Bacterial endotoxins <4.01>** Less than 8.9 EU/mg (potency).

**Uniformity of dosage units <6.02>** Perform the test according to the following method: it meets the requirement of the Content uniformity test.

To one Idarubicin Hydrochloride for Injection add the mobile phase prepared without addition of sodium lauryl sulfate to make exactly  $V$  mL so that each mL contains 0.2 mg (potency) of idarubicin hydrochloride ( $C_{26}H_{27}NO_9 \cdot HCl$ ), and use this solution as the sample solution. Separately, weigh accurately an amount of Idarubicin Hydrochloride Reference Standard, equivalent to about 10 mg (potency), dissolve in the mobile phase without containing sodium lauryl sulfate to make exactly 50 mL, and use this solution as the standard solution. Proceed as directed in the Assay under Idarubicin Hydrochloride.

$$\text{Amount [mg (potency)] of idarubicin hydrochloride } (C_{26}H_{27}NO_9 \cdot HCl) = W_S \times (A_T/A_S) \times (V/50)$$

$W_S$ : Amount [mg (potency)] of Idarubicin Hydrochloride Reference Standard

**Foreign insoluble matter <6.06>** Perform the test according to the Method 2: it meets the requirement.

**Insoluble particulate matter <6.07>** Perform the test according to the Method 1: it meets the requirement.

**Sterility <4.06>** Perform the test according to the Membrane filtration method: it meets the requirement.

**Assay** Weigh accurately the mass of the contents of not less than 10 Idarubicin Hydrochloride for Injection. Weigh accurately an amount of the content, equivalent to about 5 mg (potency) according to the labeled amount, dissolve in the mobile phase prepared without addition of sodium lauryl sulfate to make exactly 25 mL, and use this solution as the sample solution. Separately, weigh accurately an amount of

Idarubicin Hydrochloride Reference Standard, equivalent to about 10 mg (potency), dissolve in the mobile phase without containing sodium lauryl sulfate to make exactly 50 mL, and use this solution as the standard solution. Proceed as directed in the Assay under Idarubicin Hydrochloride.


$$\begin{aligned} &\text{Amount [mg (potency)] of idarubicin hydrochloride} \\ &(\text{C}_{26}\text{H}_{27}\text{NO}_9 \cdot \text{HCl}) \\ &= W_S \times (A_T/A_S) \times (1/2) \end{aligned}$$

$W_S$ : Amount [mg (potency)] of Idarubicin Hydrochloride Reference Standard

**Containers and storage** Containers—Hermetic containers.

## Idoxuridine

イドクスウリジン



$\text{C}_9\text{H}_{11}\text{IN}_2\text{O}_5$ ; 354.10

5-Iodo-2'-deoxyuridine [54-42-2]

Idoxuridine, when dried, contains not less than 98.0% of  $\text{C}_9\text{H}_{11}\text{IN}_2\text{O}_5$ .

**Description** Idoxuridine occurs as colorless, crystals or a white, crystalline powder. It is odorless.

It is freely soluble in dimethylamide, slightly soluble in water, very slightly soluble in ethanol (95), and practically insoluble in diethyl ether.

It dissolves in sodium hydroxide TS.

Melting point: about 176°C (with decomposition).

**Identification (1)** Dissolve 0.01 g of Idoxuridine in 5 mL of water by warming, add 5 mL of diphenylamine-acetic acid TS, and heat for 5 minutes: a blue color develops.

(2) Heat 0.1 g of Idoxuridine: a purple gas evolves.

(3) Dissolve 2 mg of Idoxuridine in 50 mL of 0.01 mol/L sodium hydroxide. Determine the absorption spectrum of the solution as directed under Ultraviolet-visible Spectrophotometry <2.24>, and compare the spectrum with the Reference Spectrum or the spectrum of a solution of Idoxuridine Reference Standard prepared in the same manner as the sample solution: both spectra exhibit similar intensities of absorption at the same wavelengths.

**Optical rotation** <2.49>  $[\alpha]_D^{20}$ : +28 – +31° (after drying, 0.2 g, sodium hydroxide TS, 20 mL, 100 mm).

**Purity (1)** Clarity and color of solution—Dissolve 0.20 g of Idoxuridine in 5 mL of a solution of sodium hydroxide (1 in 200): the solution is clear and colorless.

(2) Heavy metals <1.07>—Proceed with 2.0 g of Idoxuridine according to Method 2, and perform the test. Prepare the control solution with 2.0 mL of Standard Lead Solution (not more than 10 ppm).

(3) Related substances—Dissolve 0.10 g of Idoxuridine in exactly 10 mL of a mixture of dilute ethanol and ammonia solution (28) (99:1), and use this solution as the sample solution. Perform the test with the sample solution as directed under Thin-layer Chromatography <2.03>. Spot 50  $\mu\text{L}$  of the sample solution on a plate of silica gel with fluorescent indicator for thin-layer chromatography. Develop the plate with a mixture of ethyl acetate and diluted 2-propanol (2 in 3) (4:1) to a distance of about 10 cm, and air-dry the plate. Then develop two-dimensionally at right angles to the first, and air-dry the plate. Examine under ultraviolet light (main wavelength: 254 nm): any spot other than the principal spot does not appear.

(4) Iodine and iodide—Dissolve 0.10 g of Idoxuridine in 20 mL of water and 5 mL of sodium hydroxide TS, and add immediately 5 mL of dilute sulfuric acid under ice-cooling. Allow to stand for 10 minutes with occasional shaking, and filter. Transfer the filtrate into a Nessler tube, add 10 mL of chloroform and 3 drops of a solution of potassium iodate (1 in 100), shake for 30 seconds, and allow to stand: the chloroform layer has no more color than the following control solution.

Control solution: Weigh accurately 0.111 g of potassium iodide, and dissolve in water to make 1000 mL. To exactly 1 mL of this solution add 19 mL of water, 5 mL of sodium hydroxide TS and 5 mL of dilute sulfuric acid, mix, and filter. Transfer the filtrate to a Nessler tube, and proceed in the same manner.

**Loss on drying** <2.41> Not more than 0.5% (2 g, in vacuum, 60°C, 3 hours).

**Residue on ignition** <2.44> Not more than 0.3% (1 g).

**Assay** Weigh accurately about 0.7 g of Idoxuridine, previously dried, dissolve in 80 mL of *N,N*-dimethylformamide, and titrate <2.50> with 0.1 mol/L tetramethylammonium hydroxide VS until the color of the solution changes from yellow through yellow-green to blue (indicator: 5 drops of thymol blue-dimethylformamide TS). Perform a blank determination, and make any necessary correction.

$$\begin{aligned} &\text{Each mL of 0.1 mol/L tetramethylammonium} \\ &\text{hydroxide VS} \\ &= 35.41 \text{ mg of } \text{C}_9\text{H}_{11}\text{IN}_2\text{O}_5 \end{aligned}$$

**Containers and storage** Containers—Tight containers. Storage—Light-resistant.

## Idoxuridine Ophthalmic Solution

イドクスウリジン点眼液

Idoxuridine Ophthalmic Solution contains not less than 90% and not more than 110% of the labeled amount of idoxuridine ( $\text{C}_9\text{H}_{11}\text{IN}_2\text{O}_5$ ; 354.10).

**Method of preparation** Prepare as directed under Ophthalmic Solutions, with Idoxuridine.

**Description** Idoxuridine Ophthalmic Solution is a clear, colorless liquid.

**Identification (1)** To a volume of Idoxuridine Ophthalmic Solution, equivalent to 5 mg of Idoxuridine according to the

labeled amount, add 5 mL of diphenylamine-acetic acid TS, and heat for 20 minutes: a light blue color develops.

(2) Place a volume of Idoxuridine Ophthalmic Solution, equivalent to 5 mg of Idoxuridine according to the labeled amount, in a porcelain crucible, add 0.1 g of anhydrous sodium carbonate, heat slowly, evaporate to dryness and ignite until the residue is incinerated. Dissolve the residue in 5 mL of water, acidify with hydrochloric acid, and add 2 to 3 drops of sodium nitrite TS: a yellow-brown color develops. Then add 2 to 3 drops of starch TS: a deep blue color develops.

(3) To a volume of Idoxuridine Ophthalmic Solution, equivalent to 2 mg of Idoxuridine according to the labeled amount, add 0.01 mol/L sodium hydroxide TS to make 50 mL. Determine the absorption spectrum of this solution as directed under Ultraviolet-visible Spectrophotometry <2.24>: it exhibits a maximum between 277 nm and 281 nm.

pH <2.54> 4.5 – 7.0

**Purity** 5-Iodouracil and 2'-deoxyuridine—To a volume of Idoxuridine Ophthalmic Solution, equivalent to 4.0 mg of Idoxuridine according to the labeled amount, add water to make exactly 5 mL, and use this solution as the sample solution. Separately, dissolve 12.0 mg of 5-iodouracil for liquid chromatography and 4.0 mg of 2'-deoxyuridine for liquid chromatography in water to make exactly 200 mL. Measure exactly 5 mL of this solution, add water to make exactly 25 mL, and use this solution as the standard solution. Perform the test with exactly 10  $\mu$ L each of the sample solution and standard solution as directed under Liquid Chromatography <2.01> according to the following conditions, and determine the peak areas of 5-iodouracil and 2'-deoxyuridine: the peak areas of 5-iodouracil and 2'-deoxyuridine of the sample solution are not more than the peak areas of 5-iodouracil and 2'-deoxyuridine of the standard solution.

**Operating conditions**—

**Detector:** An ultraviolet absorption photometer (wavelength: 254 nm).

**Column:** A stainless steel column 3.9 mm in inside diameter and 30 cm in length, packed with octadecylsilanized silica gel for liquid chromatography (10  $\mu$ m in particle diameter).

**Column temperature:** A constant temperature of about 25°C.

**Mobile phase:** A mixture of water and methanol (24:1).

**Flow rate:** Adjust the flow rate so that the retention time of 2'-deoxyuridine is about 6 minutes.

**System suitability**—

**System performance:** When the procedure is run with 10  $\mu$ L of the standard solution under the above operating conditions, 2'-deoxyuridine and 5-iodouracil are eluted in this order with the resolution between these peaks being not less than 2.0.

**System repeatability:** When the test is repeated 6 times with 10  $\mu$ L of the standard solution under the above operating conditions, the relative standard deviation of the peak area of 2'-deoxyuridine is not more than 1.0%.

**Assay** Measure exactly a volume of Idoxuridine Ophthalmic Solution, equivalent to 3 mg of idoxuridine ( $C_9H_{11}IN_2O_5$ ) according to the labeled amount, add exactly 2 mL of the internal standard solution, then add water to make 10 mL, and use this solution as the sample solution. Separately weigh accurately about 10 mg of Idoxuridine

Reference Standard, previously dried at 60°C for 3 hours, dissolve in water to make exactly 10 mL. Measure exactly 3 mL of this solution, add exactly 2 mL of the internal standard solution, then add water to make 10 mL, and use this solution as the standard solution. Perform the test with 10  $\mu$ L each of the sample solution and standard solution as directed under Liquid Chromatography <2.01> according to the following conditions, and calculate the ratios,  $Q_T$  and  $Q_S$ , of the peak area of idoxuridine to that of the internal standard, respectively.

$$\begin{aligned} &\text{Amount (mg) of idoxuridine (C}_9\text{H}_{11}\text{IN}_2\text{O}_5\text{)} \\ &= W_S \times (Q_T/Q_S) \times (3/10) \end{aligned}$$

$W_S$ : Amount (mg) of Idoxuridine Reference Standard

**Internal standard solution**—A solution of sulfathiazole in the mobile phase (1 in 4000).

**Operating conditions**—

**Detector:** An ultraviolet absorption photometer (wavelength: 254 nm).

**Column:** A stainless steel column 3.9 mm in inside diameter and 30 cm in length, packed with octadecylsilanized silica gel for liquid chromatography (10  $\mu$ m in particle diameter).

**Column temperature:** A constant temperature of about 25°C.

**Mobile phase:** A mixture of water and methanol (87:13).

**Flow rate:** Adjust the flow rate so that the retention time of idoxuridine is about 9 minutes.

**System suitability**—

**System performance:** When the procedure is run with 10  $\mu$ L of the standard solution under the above operating conditions, idoxuridine and the internal standard are eluted in this order with the resolution between these peaks being not less than 2.0.


**System repeatability:** When the test is repeated 6 times with 10  $\mu$ L of the standard solution under the above operating conditions, the relative standard deviation of the ratios of the peak area of idoxuridine to that of the internal standard is not more than 1.0%.

**Containers and storage** Containers—Tight containers.

**Storage**—Light-resistant, in a cold place, and avoid freezing.

## Ifenprodil Tartrate

イフェンプロジル酒石酸塩



( $C_{21}H_{27}NO_2$ ) $_2$ · $C_4H_6O_6$ : 800.98  
(1*RS*,2*SR*)-4-[2-(4-Benzylpiperidin-1-yl)-1-hydroxypropyl]phenol hemi-(2*R*,3*R*)-tartrate [23210-58-4]

Ifenprodil Tartrate contains not less than 98.5% of ( $C_{21}H_{27}NO_2$ ) $_2$ · $C_4H_6O_6$ , calculated on the anhydrous basis.

**Description** Ifenprodil Tartrate occurs as a white crystalline

powder. It is odorless.

It is freely soluble in acetic acid (100), soluble in ethanol (95), slightly soluble in water and in methanol, and practically insoluble in diethyl ether.

Optical rotation  $[\alpha]_D^{20}$ : +11 – +15° (1 g, calculated on the anhydrous basis, ethanol (95), 20 mL, 100 mm).

Melting point: about 148°C (with decomposition).

**Identification (1)** Determine the absorption spectrum of a solution of Ifenprodil Tartrate in methanol (1 in 100,000) as directed under Ultraviolet-visible Spectrophotometry <2.24>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wavelengths.

(2) Determine the infrared absorption spectrum of Ifenprodil Tartrate as directed in the potassium bromide disk method under Infrared Spectrophotometry <2.25>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wave numbers.

(3) Dissolve 0.4 g of Ifenprodil Tartrate in 40 mL of water by warming. After cooling, add 0.5 mL of ammonia TS to this solution, extract with two 40-mL portions of chloroform, and collect the water layer. Evaporate 30 mL of the water layer on a water bath to dryness, and after cooling, dissolve the residue in 6 mL of water: the solution responds to the Qualitative Tests <1.09> for tartrate.

**Purity (1)** Heavy metals <1.07>—Proceed with 2.0 g of Ifenprodil Tartrate according to Method 2, and perform the test. Prepare the control solution with 2.0 mL of Standard Lead Solution (not more than 10 ppm).

(2) Related substances—Dissolve 0.30 g of Ifenprodil Tartrate in 10 mL of diluted ethanol (95) (3 in 4), and use this solution as the sample solution. Pipet 1 mL of the sample solution, add diluted ethanol (95) (3 in 4) to make exactly 200 mL, and use this solution as the standard solution. Perform the test with these solutions as directed under Thin-layer Chromatography <2.03>. Spot 10  $\mu$ L each of the sample solution and standard solution on a plate of silica gel for thin-layer chromatography. Develop the plate with a mixture of ethyl acetate, hexane, 1-butanol and ammonia solution (28) (140:40:20:1) to a distance of about 10 cm, and air-dry the plate. Spray hydrogen hexachloroplatinate (IV)-potassium iodide TS evenly on the plate: the spots other than the principal spot from the sample solution are not more intense than the spot from the standard solution.

**Water** <2.48> Not more than 4.0% (0.5 g, direct titration).

**Residue on ignition** <2.44> Not more than 0.1% (1 g).

**Assay** Weigh accurately about 0.5 g of Ifenprodil Tartrate, dissolve in 50 mL of acetic acid (100), and titrate <2.50> with 0.1 mol/L perchloric acid VS (potentiometric titration). Perform a blank determination, and make any necessary correction.


$$\begin{aligned} \text{Each mL of 0.1 mol/L perchloric acid VS} \\ = 40.05 \text{ mg of } (\text{C}_{21}\text{H}_{27}\text{NO}_2)_2 \cdot \text{C}_4\text{H}_6\text{O}_6 \end{aligned}$$

**Containers and storage** Containers—Well-closed containers.

Storage—Light-resistant.

## Imipenem Hydrate

イミペネム水和物



$\text{C}_{12}\text{H}_{17}\text{N}_3\text{O}_4\text{S} \cdot \text{H}_2\text{O}$ : 317.36

(5*R*,6*S*)-3-[2-(Formimidoethylsulfanyl)-6-[(1*R*)-1-hydroxyethyl]-7-oxo-1-azabicyclo[3.2.0]hept-2-ene-2-carboxylic acid monohydrate [74431-23-5]

Imipenem Hydrate contains not less than 980  $\mu$ g (potency) and not more than 1010  $\mu$ g (potency) per mg, calculated on the anhydrous basis. The potency of Imipenem Hydrate is expressed as mass (potency) of imipenem ( $\text{C}_{12}\text{H}_{17}\text{N}_3\text{O}_4\text{S}$ : 299.35).

**Description** Imipenem Hydrate occurs as white to light yellow crystalline powder.

It is sparingly soluble in water, and practically insoluble in ethanol (99.5).

**Identification (1)** Determine the absorption spectrum of a solution of Imipenem Hydrate in 0.1 mol/L 3-(*N*-morpholino)propanesulfonic acid buffer solution, pH 7.0 (1 in 50,000) as directed under Ultraviolet-visible Spectrophotometry <2.24>, and compare the spectrum with the Reference Spectrum or the spectrum of a solution of Imipenem Reference Standard prepared in the same manner as the sample solution: both spectra exhibit similar intensities of absorption at the same wavelengths.

(2) Determine the infrared absorption spectrum of Imipenem Hydrate as directed in the potassium bromide disk method under Infrared Spectrophotometry <2.25>, and compare the spectrum with the Reference Spectrum or the spectrum of Imipenem Reference Standard: both spectra exhibit similar intensities of absorption at the same wave numbers.

**Optical rotation** <2.49>  $[\alpha]_D^{20}$ : +89 – +94° (50 mg calculated on the anhydrous basis, 0.1 mol/L 3-(*N*-morpholino)propanesulfonic acid buffer solution, pH 7.0, 10 mL, 100 mm).

**pH** <2.54> The pH of a solution obtained by dissolving 1.0 g of Imipenem Hydrate in 200 mL of water is between 4.5 and 7.0.

**Purity (1)** Heavy metals <1.07>—Proceed with 1.0 g of Imipenem Hydrate according to Method 2, and perform the test. Prepare the control solution with 2.0 mL of Standard Lead Solution (not more than 20 ppm).

(2) Arsenic <1.11>—Put 2.0 g of Imipenem Hydrate in a crucible, add 5 mL of nitric acid and 1 mL of sulfuric acid, and heat carefully until white fumes evolve. After cooling, add 2 mL of nitric acid, heat, and repeat this procedure once more. Then add 2 mL of hydrogen peroxide (30), heat, and repeat this procedure several times until the color of the solution changes to colorless to pale yellow. After cooling, heat again until white fumes evolve. After cooling, add water to make 5 mL, and perform the test with this solution as the test

solution (not more than 1 ppm).

(3) Related substances—Dissolve 50 mg of Imipenem Hydrate in 50 mL of 0.1 mol/L 3-(*N*-morpholino)propanesulfonic acid buffer solution, pH 7.0, and use this solution as the sample solution. Pipet 1 mL of the sample solution, add 0.1 mol/L 3-(*N*-morpholino)propanesulfonic acid buffer solution, pH 7.0 to make exactly 100 mL, and use this solution as the standard solution. Perform the test with exactly 10  $\mu$ L each of the sample solution and standard solution as directed under Liquid Chromatography <2.01> according to the following conditions, and determine each peak area by the automatic integration method: the peak area of thienamycin, having the relative retention time of about 0.8 with respect to imipenem, obtained from the sample solution is not more than 1.4 times the peak area of imipenem from the standard solution, the area of the peak other than imipenem and thienamycin from the sample solution is not more than 1/3 times the peak area of imipenem from the standard solution, and the total area of the peaks other than imipenem and thienamycin from the sample solution is not more than the peak area of imipenem from the standard solution.

#### Operating conditions—

Detector, column, column temperature, mobile phase, and flow rate: Proceed as directed in the operating conditions in the Assay.

Time span of measurement: About 2 times as long as the retention time of imipenem.

#### System suitability—

Test for required detectability: Measure exactly 5 mL of the standard solution, add 0.1 mol/L 3-(*N*-morpholino)propanesulfonic acid buffer solution, pH 7.0 to make exactly 50 mL. Confirm that the peak area of imipenem from 10  $\mu$ L of this solution is equivalent to 7 to 13% of that from the standard solution.

System performance: Proceed as directed in the system suitability in the Assay.

System repeatability: When the test is repeated 6 times with 10  $\mu$ L of the standard solution under the above operating conditions, the relative standard deviation of the peak area of imipenem is not more than 2.0%.

**Water** <2.48> Not less than 5.0% and not more than 8.0% (20 mg, coulometric titration, water evaporation temperature: 140°C).

**Residue on ignition** <2.44> Not more than 0.2% (1 g).

**Assay** Weigh accurately an amount of Imipenem Hydrate and Imipenem Reference Standard, equivalent to about 50 mg (potency), dissolve each in 0.1 mol/L 3-(*N*-morpholino)propanesulfonic acid buffer solution, pH 7.0 to make exactly 50 mL, and use these solutions as the sample solution and the standard solution. Perform the test with exactly 10  $\mu$ L each of the sample solution and standard solution, within 30 minutes after preparation of these solutions, as directed under Liquid Chromatography <2.01> according to the following conditions, and determine the peak areas,  $A_T$  and  $A_S$ , of imipenem of these solutions.

$$\text{Amount } [\mu\text{g (potency)}] \text{ of imipenem (C}_{12}\text{H}_{17}\text{N}_3\text{O}_4\text{S)} \\ = W_S \times (A_T/A_S) \times 1000$$

$W_S$ : Amount [mg (potency)] of Imipenem Reference Standard

#### Operating conditions—

Detector: An ultraviolet absorption photometer (wavelength: 280 nm).

Column: A stainless steel column 3.9 mm in inside diameter and 30 cm in length, packed with octadecylsilanized silica gel for liquid chromatography (10  $\mu$ m in particle diameter).

Column temperature: A constant temperature of about 25°C.

Mobile phase: A mixture of 0.1 mol/L 3-(*N*-morpholino)propanesulfonic acid buffer solution, pH 7.0 and acetonitrile (100:1).

Flow rate: Adjust the flow rate so that the retention time of imipenem is about 6 minutes.

#### System suitability—

System performance: Dissolve 50 mg of Imipenem and 75 mg of resorcinol in 50 mL of 0.1 mol/L 3-(*N*-morpholino)propanesulfonic acid buffer solution, pH 7.0. When the procedure is run with 10  $\mu$ L of this solution under the above operating conditions, imipenem and resorcinol are eluted in this order with the resolution between these peaks being not less than 4.

System repeatability: When the test is repeated 5 times with 10  $\mu$ L of the standard solution under the above operating conditions, the relative standard deviation of the peak area of imipenem is not more than 0.80%.

**Containers and storage** Containers—Hermetic containers.

## Imipenem and Cilastatin Sodium for Injection

注射用イミペネム・シラスタチンナトリウム

Imipenem and Cilastatin Sodium for Injection is a preparation for injection which is dissolved or suspended before use.

It contains not less than 93.0% and not more than 115.0% of the labeled amount of imipenem (C<sub>12</sub>H<sub>17</sub>N<sub>3</sub>O<sub>4</sub>S: 299.35) and an amount of cilastatin sodium (C<sub>16</sub>H<sub>25</sub>N<sub>2</sub>NaO<sub>5</sub>S: 380.43), equivalent to not less than 93.0% and not more than 115.0% of the labeled amount of cilastatin (C<sub>16</sub>H<sub>26</sub>N<sub>2</sub>O<sub>5</sub>S: 358.45).

**Method of preparation** Prepare as directed under Injections, with Imipenem Hydrate and Cilastatin Sodium.

**Description** Imipenem and Cilastatin Sodium for Injection occurs as a white to light yellowish white powder.

**Identification** (1) To 1 mL of a solution of Imipenem and Cilastatin Sodium for Injection (1 in 100) add 1 mL of ninhydrin TS, heat in a water bath for 5 minutes: a purple color appears (cilastatin).

(2) To 2 mL of a solution of Imipenem and Cilastatin Sodium for Injection (1 in 1000) add 0.1 mol/L 3-(*N*-morpholino)propanesulfonic acid buffer solution, pH 7.0 to make 50 mL, and determine the absorption spectrum of the solution as directed under Ultraviolet-visible Spectrophotometry <2.24>: it exhibits a maximum between 296 nm and 300 nm (imipenem).

**pH** <2.54> The pH of a solution prepared by dissolving an

amount of Imipenem and Cilastatin Sodium for Injection, equivalent to 0.5 g (potency) of Imipenem Hydrate according to the labeled amount, in 100 mL of isotonic sodium chloride solution is between 6.5 and 8.0. The pH of the Injection intended for intramuscular use is between 6.0 and 7.5.

**Purity** Clarity and color of solution—Dissolve an amount of Imipenem and Cilastatin Sodium for Injection, equivalent to 0.5 g (potency) of Imipenem Hydrate according to the labeled amount, in 100 mL of isotonic sodium chloride solution: the solution is clear and colorless or pale yellow.

**Loss on drying** <2.41> Not more than 3.0% (1 g, in vacuum, 60°C, 3 hours).

**Bacterial endotoxins** <4.01> Less than 0.25 EU/mg (potency).

**Uniformity of dosage units** <6.02> Perform the test according to the following method: it meets the requirement of the Content uniformity test. Calculate the acceptance value by using the average of the limits specified in the potency definition for *T*.

Dissolve the total amount of the content of one Imipenem and Cilastatin Sodium for Injection in isotonic sodium chloride solution to make 100 mL. Measure exactly *V* mL of this solution, equivalent to about 25 mg (potency) of Imipenem Hydrate according to the labeled amount, add 0.1 mol/L 3-(*N*-morpholino)propanesulfonic acid buffer solution, pH 7.0 to make exactly 50 mL, and use this solution as the sample solution. Proceed hereafter as directed in the Assay.

$$\begin{aligned} &\text{Amount [mg (potency)] of imipenem (C}_{12}\text{H}_{17}\text{N}_3\text{O}_4\text{S)} \\ &= W_S \times (A_{\text{TI}}/A_{\text{SI}}) \times (100/V) \end{aligned}$$

$W_S$ : Amount [mg (potency)] of Imipenem Reference Standard

$$\begin{aligned} &\text{Amount (mg) of cilastatin (C}_{16}\text{H}_{26}\text{N}_2\text{O}_5\text{S)} \\ &= W_S \times (A_{\text{TC}}/A_{\text{SC}}) \times (100/V) \times 0.955 \end{aligned}$$

$W_S$ : Amount (mg) of cilastatin ammonium for assay, calculated on the anhydrous basis and corrected on the amount of the residual solvent

**Foreign insoluble matter** <6.06> Perform the test according to the Method 2: the Injection which is dissolved before use meets the requirement.

**Insoluble particulate matter** <6.07> Perform the test according to the Method 1: the Injection which is dissolved before use meets the requirement.

**Sterility** <4.06> Perform the test according to the Membrane filtration method: it meets the requirement.

**Assay** Weigh accurately the mass of the contents of not less than 10 Imipenem and Cilastatin Sodium for Injections. Weigh accurately an amount of the content, equivalent to one Imipenem and Cilastatin Sodium for Injection, dissolve in isotonic sodium chloride solution to make exactly 100 mL. Measure exactly an amount of this solution, equivalent to about 25 mg (potency) of imipenem, add 0.1 mol/L 3-(*N*-morpholino)propanesulfonic acid buffer solution, pH 7.0 to make exactly 50 mL, and use this as the sample solution. Separately, weigh accurately an amount of Imipenem Reference Standard, equivalent to about 25 mg (potency), and weigh accurately about 25 mg of cilastatin ammonium for assay, separately determined the water content, dissolve in 10

mL of isotonic sodium chloride solution, add 0.1 mol/L 3-(*N*-morpholino)propanesulfonic acid buffer solution, pH 7.0 to make exactly 50 mL, and use this solution as the standard solution. Perform the test with exactly 10  $\mu$ L each of the sample solution and standard solution as directed under Liquid Chromatography <2.01> according to the following conditions, and determine the peak areas of imipenem,  $A_{\text{TI}}$  and  $A_{\text{SI}}$ , and that of cilastatin,  $A_{\text{TC}}$  and  $A_{\text{SC}}$ .

$$\begin{aligned} &\text{Amount [mg (potency)] of imipenem (C}_{12}\text{H}_{17}\text{N}_3\text{O}_4\text{S)} \\ &= W_S \times (A_{\text{TI}}/A_{\text{SI}}) \end{aligned}$$

$W_S$ : Amount [mg (potency)] of Imipenem Reference Standard

$$\begin{aligned} &\text{Amount (mg) of cilastatin (C}_{16}\text{H}_{26}\text{N}_2\text{O}_5\text{S)} \\ &= W_S \times (A_{\text{TC}}/A_{\text{SC}}) \times 0.955 \end{aligned}$$

$W_S$ : Amount (mg) of cilastatin ammonium for assay, calculated on the anhydrous basis and corrected on the amount of ethanol

**Operating conditions—**

**Detector:** An ultraviolet absorption photometer (wavelength: 250 nm).

**Column:** A stainless steel column 4.6 mm in inside diameter and 20 cm in length, packed with octylsilanized silica gel for liquid chromatography (10  $\mu$ m in particle diameter).

**Column temperature:** A constant temperature of about 50°C.

**Mobile phase:** Dissolve 0.836 g of 3-(*N*-morpholino)propanesulfonic acid, 1.0 g of sodium 1-hexane sulfonate and 50 mg of disodium dihydrogen ethylenediamine tetraacetate dihydrate in 800 mL of water, adjust to pH 7.0 with 0.1 mol/L sodium hydrate TS, and add water to make 1000 mL.

**Flow rate:** Adjust the flow rate so that the retention time of imipenem is about 3 minutes.

**System suitability—**


**System performance:** When the procedure is run with 10  $\mu$ L of the standard solution under the above operating conditions, imipenem and cilastatin are eluted in this order with the resolution between these peaks being not less than 2.0, and the symmetry factors of the peak of imipenem and cilastatin are not more than 2.0, respectively.

**System repeatability:** When the test is repeated 6 times with 10  $\mu$ L of the standard solution under the above operating conditions, the relative standard deviations of the peak area of imipenem and cilastatin are not more than 2.0%, respectively.

**Containers and storage** Containers—Hermetic containers.

## Imipramine Hydrochloride

イミプラミン塩酸塩



$\text{C}_{19}\text{H}_{24}\text{N}_2 \cdot \text{HCl}$ : 316.87  
3-(10,11-Dihydro-5*H*-dibenzo[*b,f*]azepin-5-yl)-



*N,N*-dimethylpropylamine monohydrochloride [113-52-0]

Imipramine Hydrochloride, when dried, contains not less than 98.5% of  $C_{19}H_{24}N_2 \cdot HCl$ .

**Description** Imipramine Hydrochloride occurs as a white to pale yellowish white, crystalline powder. It is odorless.

It is freely soluble in water and in ethanol (95), and practically insoluble in diethyl ether.

The pH of the aqueous solution (1 in 10) is between 4.2 and 5.2.

It is gradually colored by light.

**Identification (1)** Dissolve 5 mg of Imipramine Hydrochloride in 2 mL of nitric acid: a deep blue color develops.

(2) Dissolve 5 mg of Imipramine Hydrochloride in 250 mL of 0.01 mol/L hydrochloric acid TS. Determine the absorption spectrum of the solution as directed under Ultraviolet-visible Spectrophotometry <2.24>, and compare the spectrum with the Reference Spectrum or the spectrum of a solution of Imipramine Hydrochloride Reference Standard prepared in the same manner as the sample solution: both spectra exhibit similar intensities of absorption at the same wavelengths.

(3) Dissolve 0.05 g of Imipramine Hydrochloride in 5 mL of water, add 1 mL of ammonia TS, allow to stand for 5 minutes, filter, and acidify the filtrate with dilute nitric acid: it responds to the Qualitative Tests <1.09> (2) for chloride.

**Melting point** <2.60> 170 – 174°C (with decomposition).

**Purity (1)** Clarity and color of solution—Dissolve 1.0 g of Imipramine Hydrochloride in 10 mL of water: the solution is clear, and has no more color than the following control solution.

Control solution: Take exactly 1.0 mL of Cobaltous Chloride Colorimetric Stock Solution, 2.4 mL of Ferric Chloride Colorimetric Solution, 0.4 mL of Cupric Sulfate Colorimetric Stock Solution and 6.2 mL of diluted hydrochloric acid (1 in 40), and mix them. Pipet 0.5 mL of this solution, and add exactly 9.5 mL of water.

(2) Iminodibenzyl—Dissolve 50 mg of Imipramine Hydrochloride in 10 mL of a mixture of hydrochloric acid and ethanol (95) (1:1) in a 25-mL brown volumetric flask. Cool the flask in ice water, add 5 mL of an ethanol (95) solution of furfural (1 in 250) and 5 mL of hydrochloric acid, and allow to stand at 25°C for 3 hours. Add a mixture of hydrochloric acid and ethanol (95) (1:1) to make 25 mL, and determine the absorbance of this solution at 565 nm as directed under Ultraviolet-visible Spectrophotometry <2.24>: it is not more than 0.16.

(3) Related substances—Dissolve 0.20 g of Imipramine Hydrochloride in 10 mL of ethanol (95), and use this solution as the sample solution. Pipet 1 mL of this solution, and add ethanol (95) to make exactly 50 mL. Pipet 5 mL of this solution, add ethanol (95) to make exactly 50 mL, and use this solution as the standard solution. Perform the test with these solutions as directed under Thin-layer Chromatography <2.03>. Spot 5  $\mu$ L each of the sample solution and standard solution on a plate of silica gel for thin-layer chromatography. Develop the plate with a mixture of ethyl acetate, acetic acid (100), hydrochloric acid and water (11:7:1:1) to a distance of about 12 cm, and air-dry the plate. Spray evenly potassium dichromate-sulfuric acid TS on the

plate: the spots other than the principal spot from the sample solution are not more intense than the spot from the standard solution.

**Loss on drying** <2.41> Not more than 0.5% (1 g, 105°C, 2 hours).

**Residue on ignition** <2.44> Not more than 0.1% (1 g).

**Assay** Weigh accurately about 0.3 g of Imipramine Hydrochloride, previously dried, and dissolve in 20 mL of water. Add 5 mL of sodium hydroxide TS, and extract with three 20-mL portions of chloroform. Filter each extract through a pledget of absorbent cotton on which a small quantity of anhydrous sodium sulfate is placed. Combine the chloroform extracts, and titrate <2.50> with 0.1 mol/L perchloric acid VS until the yellow solution changes to red-purple (indicator: 10 drops of metanil yellow TS). Perform a blank determination.

Each mL of 0.1 mol/L perchloric acid VS  
= 31.69 mg of  $C_{19}H_{24}N_2 \cdot HCl$

**Containers and storage** Containers—Tight containers.  
Storage—Light-resistant.

## Imipramine Hydrochloride Tablets

イミプラミン塩酸塩錠

Imipramine Hydrochloride Tablets contain not less than 93% and not more than 107% of the labeled amount of imipramine hydrochloride ( $C_{19}H_{24}N_2 \cdot HCl$ : 316.87).

**Method of preparation** Prepare as directed under Tablets, with Imipramine Hydrochloride.

**Identification (1)** Weigh a quantity of powdered Imipramine Hydrochloride Tablets, equivalent to 0.25 g of Imipramine Hydrochloride according to the labeled amount, add 25 mL of chloroform, shake thoroughly, and filter. Evaporate the filtrate on a water bath, and proceed with the residue as directed in the Identification (1) under Imipramine Hydrochloride.

(2) Dissolve an amount of the residue obtained in (1), equivalent to 5 mg of Imipramine Hydrochloride, in 250 mL of 0.01 mol/L hydrochloric acid TS, and determine the absorption spectrum as directed under Ultraviolet-visible Spectrophotometry <2.24>: it exhibits a maximum between 249 nm and 253 nm, and a shoulder between 270 nm and 280 nm.

(3) Dry the residue obtained in (1) at 105°C for 2 hours: the residue melts between 170°C and 174°C (with decomposition).

**Dissolution** <6.10> Perform the test according to the following method: it meets the requirement.

Perform the test with 1 tablet of Imipramine Hydrochloride Tablet at 75 revolutions per minute according to the Paddle method, using 900 mL of 2nd fluid for dissolution test as the test solution. Take 20 mL or more of the dissolved solution after 60 minutes from the start of the dissolution test, and filter through a membrane filter with pore size of not more than 0.8  $\mu$ m. Discard the first 10 mL of the filtrate, pipet the subsequent *V* mL, add 2nd fluid for dissolution test

to make exactly  $V'$  mL so that each mL of the filtrate contains about  $10\text{ }\mu\text{g}$  of imipramine hydrochloride ( $\text{C}_{19}\text{H}_{24}\text{N}_2\cdot\text{HCl}$ ) according to the labeled amount, and use this solution as the sample solution. Separately, weigh accurately about  $0.025\text{ g}$  of Imipramine Hydrochloride Reference Standard, previously dried at  $105^\circ\text{C}$  for 2 hours, dissolve in 2nd fluid for dissolution test to make exactly  $100\text{ mL}$ . Pipet  $4\text{ mL}$  of this solution, add 2nd fluid for dissolution test to make exactly  $100\text{ mL}$ , and use this solution as the standard solution. Determine the absorbances,  $A_T$  and  $A_S$ , of the sample solution and the standard solution at  $250\text{ nm}$  as directed under Ultraviolet-visible Spectrophotometry <2.24>. The dissolution rate of Imipramine Hydrochloride Tablets in 60 minutes should be not less than 75%.

Dissolution rate (%) with respect to the labeled amount of imipramine hydrochloride ( $\text{C}_{19}\text{H}_{24}\text{N}_2\cdot\text{HCl}$ )

$$= W_S \times (A_T/A_S) \times (V'/V) \times (1/C) \times 36$$

$W_S$ : Amount (mg) of Imipramine Hydrochloride Reference Standard.

$C$ : Labeled amount (mg) of imipramine hydrochloride ( $\text{C}_{19}\text{H}_{24}\text{N}_2\cdot\text{HCl}$ ) in 1 tablet.

**Assay** Take 20 Imipramine Hydrochloride Tablets, add exactly  $200\text{ mL}$  of  $0.01\text{ mol/L}$  hydrochloric acid TS, and shake well until the tablets are completely disintegrated. After centrifuging the solution, pipet a volume of the supernatant liquid, equivalent to about  $25\text{ mg}$  of imipramine hydrochloride ( $\text{C}_{19}\text{H}_{24}\text{N}_2\cdot\text{HCl}$ ) according to the labeled amount, add  $0.01\text{ mol/L}$  hydrochloric acid TS to make exactly  $100\text{ mL}$ , and use this solution as the sample solution. Separately, weigh accurately about  $25\text{ mg}$  of Imipramine Hydrochloride for Assay, previously dried at  $105^\circ\text{C}$  for 2 hours, dissolve in  $0.01\text{ mol/L}$  hydrochloric acid TS to make exactly  $100\text{ mL}$ , and use this solution as the standard solution. Pipet  $3\text{ mL}$  each of these solutions into separators which contain  $15\text{ mL}$  of potassium hydrogen phthalate buffer solution, pH 5.6,  $8\text{ mL}$  of bromocresol green-sodium hydroxide TS and  $30\text{ mL}$  of chloroform, and shake. Filter the chloroform layer through a pledget of absorbent cotton into a  $100\text{-mL}$  volumetric flask. Repeat the extraction with two  $30\text{-mL}$  portions of chloroform, combine the chloroform layers in the  $100\text{-mL}$  volumetric flask, and add chloroform to make exactly  $100\text{ mL}$ . Perform the test with these solutions as directed under Ultraviolet-visible Spectrophotometry <2.24>, using a solution obtained by proceeding with  $3\text{ mL}$  of  $0.01\text{ mol/L}$  hydrochloric acid TS in the same manner as the blank. Determine the absorbances,  $A_T$  and  $A_S$ , of these solutions at  $416\text{ nm}$ .

Amount (mg) of imipramine hydrochloride ( $\text{C}_{19}\text{H}_{24}\text{N}_2\cdot\text{HCl}$ )


$$= W_S \times (A_T/A_S)$$

$W_S$ : Amount (mg) of Imipramine Hydrochloride for Assay

**Containers and storage** Containers—Tight containers.

## Indenolol Hydrochloride

インデノロール塩酸塩



$\text{C}_{15}\text{H}_{21}\text{NO}_2\cdot\text{HCl}$ : 283.79

(2*RS*)-1-(3*H*-Inden-4-yloxy)-

3-(1-methylethyl)aminopropan-2-ol monohydrochloride

(2*RS*)-1-(3*H*-Inden-7-yloxy)-

3-(1-methylethyl)aminopropan-2-ol monohydrochloride  
[68906-88-7]

Indenolol Hydrochloride is a mixture of (2*RS*)-1-(3*H*-Inden-4-yloxy)-3-(1-methylethyl)aminopropan-2-ol monohydrochloride and (2*RS*)-1-(3*H*-Inden-7-yloxy)-3-(1-methylethyl)aminopropan-2-ol monohydrochloride.

When dried, it contains not less than 98.5% of  $\text{C}_{15}\text{H}_{21}\text{NO}_2\cdot\text{HCl}$ .

**Description** Indenolol Hydrochloride occurs as white to pale yellow crystals or crystalline powder.

It is freely soluble in water and in acetic acid (100), soluble in ethanol (95) and in chloroform, slightly soluble in acetic anhydride, very slightly soluble in ethyl acetate, and practically insoluble in diethyl ether.

The pH of a solution of Indenolol Hydrochloride (1 in 10) is between 3.5 and 5.5.

It is colored by light.

**Identification** (1) Dissolve  $0.1\text{ g}$  of Indenolol Hydrochloride in 1 to 2 drops of dilute hydrochloric acid and  $5\text{ mL}$  of water, and add  $1\text{ mL}$  of Reinecke salt TS: a red-purple precipitate is formed.

(2) Determine the absorption spectrum of a solution of Indenolol Hydrochloride (1 in 50,000) as directed under Ultraviolet-visible Spectrophotometry <2.24>, and compare the spectrum with the Reference Spectrum 1: both spectra exhibit similar intensities of absorption at the same wavelengths. Separately, determine the absorption spectrum of a solution of Indenolol Hydrochloride (1 in 10,000) as directed under Ultraviolet-visible Spectrophotometry <2.24>, and compare the spectrum with the Reference Spectrum 2: both spectra exhibit similar intensities of absorption at the same wavelengths.

(3) Determine the infrared absorption spectrum of Indenolol Hydrochloride, previously dried, as directed in the potassium chloride disk method under Infrared Spectrophotometry <2.25>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wave numbers.

(4) A solution of Indenolol Hydrochloride (1 in 10) responds to the Qualitative Tests <1.09> for chloride.

**Absorbance** <2.24>  $E_{1\text{cm}}^{1\%}$  (250 nm): 330 – 340 (after drying,  $10\text{ mg}$ , water,  $1000\text{ mL}$ ).

**Melting point** <2.60>  $140 - 143^\circ\text{C}$

**Purity (1)** Clarity and color of solution—Dissolve 1.0 g of Indenolol Hydrochloride in 10 mL of water: the solution is clear and colorless to pale yellow.

(2) Heavy metals <1.07>—Proceed with 1.0 g of Indenolol Hydrochloride according to Method 2, and perform the test. Prepare the control solution with 2.0 mL of Standard Lead Solution (not more than 20 ppm).

(3) Arsenic <1.11>—Prepare the test solution with 1.0 g of Indenolol Hydrochloride according to Method 1, and perform the test (not more than 2 ppm).

(4) Related substances—Dissolve 0.20 g of Indenolol Hydrochloride in 10 mL of chloroform, and use this solution as the sample solution. Pipet 1 mL of the sample solution, add chloroform to make exactly 200 mL, and use this solution as the standard solution. Perform the test with these solutions as directed under Thin-layer Chromatography <2.03>. Spot 10  $\mu$ L each of the sample solution and standard solution on a plate of silica gel with fluorescent indicator for thin-layer chromatography. Develop the plate with a mixture of 1,2-dichloroethane, ethanol (99.5) and ammonia solution (28) (70:15:2) to a distance of about 12 cm, and air-dry the plate. Examine under ultraviolet light (main wavelength: 254 nm): the spots other than the principal spot from the sample solution are not more intense than the spot from the standard solution.

**Loss on drying** <2.41> Not more than 0.5% (1 g, in vacuum, phosphorus (V) oxide, 4 hours).

**Residue on ignition** <2.44> Not more than 0.1% (1 g).

**Isomer ratio** Dissolve 5 mg of Indenolol Hydrochloride in 1.0 mL of a mixture of ethyl acetate and dehydrated trifluoroacetic acid (9:1), and use this solution as the sample solution. Perform the test with 2  $\mu$ L of the sample solution as directed under Gas Chromatography <2.02> according to the following conditions. Determine the areas of two adjacent peaks,  $A_a$  and  $A_b$ , having the retention times of about 16 minutes, where  $A_a$  is the peak area of shorter retention time and  $A_b$  is the peak area of longer retention time: the ratio  $A_a/(A_a + A_b)$  is between 0.6 and 0.7.

**Operating conditions**—

Detector: A hydrogen flame-ionization detector.

Column: A glass column about 2 mm in inside diameter and about 2 m in length, packed with siliceous earth for gas chromatography (150 to 180  $\mu$ m in particle diameter) coated with 65% phenyl-methyl silicon polymer for gas chromatography at the ratio of 2%.

Column temperature: A constant temperature between 150°C and 170°C.

Carrier gas: Helium.

Flow rate: Adjust the flow rate so that the retention time of the peak showing earlier elution of the two peaks of indenolol hydrochloride is about 16 minutes.

Selection of column: Proceed with 2  $\mu$ L of the sample solution under the above operating conditions, and calculate the resolution. Use a column with the resolution between the two peaks being not less than 1.1.

**Assay** Weigh accurately about 0.5 g of Indenolol Hydrochloride, previously dried, dissolve in 50 mL of a mixture of acetic anhydride and acetic acid (100) (4:1), and titrate <2.50> with 0.1 mol/L perchloric acid VS until the color of the solution changes from purple through blue to green (indi-

cator: 3 drops of crystal violet TS). Perform a blank determination, and make any necessary correction.


Each mL of 0.1 mol/L perchloric acid VS  
= 28.38 mg of  $C_{15}H_{21}NO_2 \cdot HCl$

**Containers and storage** Containers—Well-closed containers.

Storage—Light-resistant.

## Indigocarmine

インジゴカルミン



$C_{16}H_8N_2Na_2O_8S_2$ ; 466.35

Disodium 3,3'-dioxo-[4<sup>2,2'</sup>-biindoline]-5,5'-disulfonate  
[860-22-0]

Indigocarmine, when dried, contains not less than 95.0% of  $C_{16}H_8N_2Na_2O_8S_2$ .

**Description** Indigocarmine occurs as blue to dark blue powder or granules. It is odorless.

It is sparingly soluble in water, and practically insoluble in ethanol (95) and in diethyl ether.

It is hygroscopic.

When compressed, it has a coppery luster.

**Identification (1)** A solution of Indigocarmine (1 in 100) is dark blue in color. Perform the following tests with this solution as the sample solution: the dark blue color of each solution disappears.

(i) Add 1 mL of nitric acid to 2 mL of the sample solution;

(ii) Add 1 mL of bromine TS to 2 mL of the sample solution;

(iii) Add 1 mL of chlorine TS to 2 mL of the sample solution;

(iv) Add 2 mL of sodium hydroxide TS and 0.2 g of zinc powder to 2 mL of the sample solution, and warm.

(2) Dissolve 0.1 g of Indigocarmine in 100 mL of a solution of ammonium acetate (1 in 650). To 1 mL of the solution add a solution of ammonium acetate (1 in 650) to make 100 mL. Determine the absorption spectrum of the solution as directed under Ultraviolet-visible Spectrophotometry <2.24>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wavelengths.

(3) Ignite 1 g of Indigocarmine to carbonize. After cooling, add 20 mL of water to the residue, shake, and filter the mixture: the filtrate responds to the Qualitative Tests <1.09> for sodium salt and for sulfate.

**pH** <2.54> Dissolve 0.10 g of Indigocarmine in 20 mL of water: the pH of the solution is between 5.0 and 6.0.

**Purity (1)** Water-insoluble substances—To 1.00 g of Indigocarmine add 200 mL of water, shake, and filter through a tared glass filter (G4). Wash the residue with water until the blue color of the filtrate becomes practically colorless, and

dry the residue at 105°C for 4 hours: the mass of the residue does not exceed 5.0 mg.

(2) **Arsenic** <1.11>—Place 0.8 g of Indigocarmine in a Kjeldahl flask, add 5 mL of sulfuric acid and 5 mL of nitric acid, and ignite gently. Repeat the addition of 2 to 3 mL of nitric acid occasionally, and continue to heat until a colorless to light yellow solution is obtained. After cooling, add 15 mL of a saturated ammonium oxalate solution, heat the solution until dense white fumes are evolved, and concentrate to 2 to 3 mL. After cooling, dilute with water to 10 mL, and perform the test with 5 mL of this solution as the test solution (not more than 5 ppm).

**Loss on drying** <2.41> Not more than 10.0% (1 g, 105°C, 2 hours).

**Residue on ignition** <2.44> Not less than 28.0% and not more than 38.0% (after drying, 1 g).

**Assay** Weigh accurately about 0.5 g of Indigocarmine, previously dried, add 15 g of sodium hydrogen tartrate monohydrate, and dissolve in 200 mL of water, boil with bubbling of a stream of carbon dioxide, and titrate <2.50>, while being hot, with 0.1 mol/L titanium (III) chloride VS until the color of the solution changes from blue through yellow to orange.

Each mL of 0.1 mol/L titanium (III) chloride VS  
= 23.32 mg of  $C_{16}H_8N_2Na_2O_8S_2$

**Containers and storage** Containers—Tight containers.  
Storage—Light-resistant.

## Indigocarmine Injection

インジゴカルミン注射液

Indigocarmine Injection is an aqueous solution for injection. It contains not less than 95% and not more than 105% of the labeled amount of indigocarmine ( $C_{16}H_8N_2Na_2O_8S_2$ ; 466.35).

**Method of preparation** Prepare as directed under Injection, with Indigocarmine.

**Description** Indigocarmine Injection is a dark blue liquid.  
pH: 3.0 – 5.0

**Identification** (1) To a volume of Indigocarmine Injection, equivalent to 0.02 g of Indigocarmine according to the labeled amount, add 1 mL of nitric acid: the dark blue color of the liquid disappears, and a yellow-brown color develops.

(2) To a volume of Indigocarmine Injection, equivalent to 0.02 g of Indigocarmine according to the labeled amount, add 1 mL of bromine TS: the dark blue color disappears, and a yellow-brown color develops.

(3) To a volume of Indigocarmine Injection, equivalent to 0.02 g of Indigocarmine according to the labeled amount, add 1 mL of chlorine TS: the dark blue color disappears, and a yellow-brown color develops.

(4) To a volume of Indigocarmine Injection, equivalent to 0.01 g of Indigocarmine according to the labeled amount, add ammonium acetate solution (1 in 650) to make 1000 mL, and determine the absorbance of the solution as directed under Ultraviolet-visible Spectrophotometry <2.24>: it exhibits a

maximum between 610 nm and 614 nm.

**Assay** Measure exactly a volume of Indigocarmine Injection, equivalent to about 0.2 g of indigocarmine ( $C_{16}H_8N_2Na_2O_8S_2$ ), add 6 g of sodium hydrogen tartrate monohydrate, and dissolve in water to make 200 mL. Then boil under a carbon dioxide stream, and proceed as directed in the Assay under Indigocarmine.

Each mL of 0.1 mol/L titanium (III) chloride VS  
= 23.32 mg of  $C_{16}H_8N_2Na_2O_8S_2$

**Containers and storage** Containers—Hermetic containers, and colored containers may be used.

Storage—Light-resistant.

## Indium ( $^{111}\text{In}$ ) Chloride Injection

塩化インジウム ( $^{111}\text{In}$ ) 注射液

Indium ( $^{111}\text{In}$ ) Chloride Injection is an aqueous solution for injection

It contains indium-111 ( $^{111}\text{In}$ ) in the form of indium chloride.


It conforms to the requirements of Indium ( $^{111}\text{In}$ ) Chloride Injection in the Minimum Requirements for Radiopharmaceuticals.

Test for Extractable Volume of Parenteral Preparations and Insoluble Particulate Matter Test for Injections are not applied to this injection.

**Description** Indium ( $^{111}\text{In}$ ) Chloride Injection is a clear, colorless liquid.

## Indometacin

インドメタシン



$C_{19}H_{16}ClNO_4$ ; 357.79

[1-(4-Chlorobenzoyl)-5-methoxy-2-methyl-1H-indol-3-yl]acetic acid [53-86-1]

Indometacin, when dried, contains not less than 98.0% of  $C_{19}H_{16}ClNO_4$ .

**Description** Indometacin occurs as a white to light yellowish white, very fine crystalline powder.

It is sparingly soluble in methanol, in ethanol (95) and in diethyl ether, and practically insoluble in water.

It dissolves in sodium hydroxide TS.

It is colored by light.

Melting point: 155 – 162°C

**Identification** (1) Dissolve 2 mg of Indometacin in 100 mL of methanol. Determine the absorption spectrum of the solution as directed under Ultraviolet-visible Spectrophotometry

<2.24>, and compare the spectrum with the Reference Spectrum or the spectrum of a solution of Indometacin Reference Standard prepared in the same manner as the sample solution: both spectra exhibit similar intensities of absorption at the same wavelengths.

(2) Determine the infrared absorption spectrum of Indometacin, previously dried, as directed in the potassium bromide disk method under Infrared Spectrophotometry <2.25>, and compare the spectrum with the Reference Spectrum or the spectrum of dried Indometacin Reference Standard: both spectra exhibit similar intensities of absorption at the same wave numbers. If any difference appears between the spectra, recrystallize the sample and the Reference Standard with diethyl ether, filter and dry the crystals, and perform the test with the crystals.

(3) Perform the test with Indometacin as directed under Flame Coloration Test <1.04> (2): a green color appears.

**Purity** (1) Acidity—To 1.0 g of Indometacin add 50 mL of water, shake for 5 minutes, and filter. To the filtrate add 0.20 mL of 0.1 mol/L sodium hydroxide VS and 1 drop of phenolphthalein TS: a red color develops.

(2) Heavy metals <1.07>—Proceed with 1.0 g of Indometacin according to Method 2, and perform the test. Prepare the control solution with 2.0 mL of Standard Lead Solution (not more than 20 ppm).

(3) Arsenic <1.11>—Prepare the test solution with 1.0 g of Indometacin according to Method 3, and perform the test (not more than 2 ppm).

(4) Related substances—Dissolve 0.10 g of Indometacin in 10 mL of methanol, and use this solution as the sample solution. Pipet 1 mL of the sample solution, and add methanol to make exactly 50 mL. Pipet 5 mL of this solution, add methanol to make exactly 20 mL, and use this solution as the standard solution. Perform the test with these solutions as directed under Thin-layer Chromatography <2.03>. Spot 25  $\mu$ L each of the sample solution and standard solution on a plate of silica gel with fluorescent indicator for thin-layer chromatography. Develop the plate with a mixture of dehydrated diethyl ether and acetic acid (100) (100:3) to a distance of about 10 cm, and air-dry the plate. Examine under ultraviolet light (main wavelength: 254 nm): the spots other than the principal spot from the sample solution are not more intense than the spot from the standard solution.

**Loss on drying** <2.41> Not more than 0.5% (1 g, 105°C, 4 hours).

**Residue on ignition** <2.44> Not more than 0.1% (1 g).

**Assay** Weigh accurately about 0.7 g of Indometacin, previously dried, dissolve in 60 mL of methanol, add 30 mL of water, and titrate <2.50> with 0.1 mol/L sodium hydroxide VS (indicator: 3 drops of phenolphthalein TS). Perform a blank determination.

Each mL of 0.1 mol/L sodium hydroxide VS  
= 35.78 mg of  $C_{19}H_{16}ClNO_4$

**Containers and storage** Containers—Tight containers.  
Storage—Light-resistant.

## Indometacin Capsules

インドメタシンカプセル

Indometacin Capsules contain not less than 90% and not more than 110% of the labeled amount of indometacin ( $C_{19}H_{16}ClNO_4$ ; 357.79).

**Method of preparation** Prepare as directed under Capsules, with Indometacin.

**Identification** Powder the contents of Indometacin Capsules. To a quantity of the powder, equivalent to 0.1 g of Indometacin according to the labeled amount, add 20 mL of chloroform, shake well, and centrifuge. Filter the supernatant liquid, and evaporate the filtrate to dryness. After cooling, dissolve the residue in 20 mL of methanol. To 10 mL of this solution add methanol to make 50 mL, then to 2 mL of this solution add methanol to make 100 mL, and use this solution as the sample solution. Determine the absorption spectrum of the sample solution as directed under Ultraviolet-visible Spectrophotometry <2.24>: it exhibits a maximum between 317 nm and 321 nm.

**Purity** Related substances—Powder the content of Indometacin Capsules. To a quantity of the powder, equivalent to 0.10 g of Indometacin according to the labeled amount, add exactly 10 mL of methanol, shake well, filter, and use the filtrate as the sample solution. Dissolve 25 mg of Indometacin Reference Standard in methanol to make exactly 50 mL. Pipet 1 mL of the solution, add methanol to make exactly 10 mL, and use this solution as the standard solution. Proceed as directed in the Purity (4) under Indometacin.

**Dissolution** <6.10> Perform the test according to the following method: it meets the requirement.

Take 1 capsule of Indometacin Capsules, and perform the test using 900 mL of a mixture of water and phosphate buffer solution, pH 7.2, (4:1) as the test solution at 100 revolutions per minute as directed in the Basket method. Take 20 mL or more of the dissolved solution at 20 minutes after starting the test, and filter through a membrane filter (less than 0.8  $\mu$ m in pore size). Discard the first 10 mL of the filtrate, and use the subsequent filtrate as the sample solution. Separately, weigh accurately about 30 mg of Indometacin Reference Standard, previously dried at 105°C for 4 hours, dissolve in a mixture of water and phosphate buffer solution, pH 7.2, (4:1) to make exactly 1000 mL, and use this as the standard solution. Determine the absorbances,  $A_T$  and  $A_S$ , of the sample solution and the standard solution at 320 nm as directed under Ultraviolet-visible Spectrophotometry <2.24>. The dissolution rate of Indometacin Capsules in 20 minutes should be not less than 75%.

Dissolution rate (%) with respect to the labeled amount of indometacin ( $C_{19}H_{16}ClNO_4$ )  
=  $W_S \times (A_T/A_S) \times (90/C)$

$W_S$ : Amount (mg) of Indometacin Reference Standard.

$C$ : Labeled amount (mg) of indometacin ( $C_{19}H_{16}ClNO_4$ ) in 1 capsule.

**Assay** Weigh accurately the contents of not less than 20 In-

dometacin Capsules. Powder the combined contents, and weigh accurately a portion of the powder, equivalent to about 50 mg of indometacin ( $C_{19}H_{16}ClNO_4$ ). Dissolve in 40 mL of methanol, and add methanol to make exactly 50 mL. Filter this solution, discarding the first 10-mL portion of the filtrate. Pipet the subsequent 5 mL of the filtrate, add exactly 3 mL of the internal standard solution, add the mobile phase to make 100 mL, and use this solution as the sample solution. Separately, weigh accurately about 50 mg of Indometacin Reference Standard, previously dried at 105°C for 4 hours, and dissolve in methanol to make exactly 50 mL. Pipet 5 mL of the solution, add exactly 3 mL of the internal standard solution, add the mobile phase to make 100 mL, and use this solution as the standard solution. Perform the test with 20  $\mu$ L each of the sample solution and standard solution as directed under Liquid Chromatography <2.01> according to the following conditions, and calculate the ratios,  $Q_T$  and  $Q_S$ , of the peak area of indometacin to that of the internal standard, respectively.

$$\begin{aligned} &\text{Amount (mg) of indometacin (C}_{19}\text{H}_{16}\text{ClNO}_4\text{)} \\ &= W_S \times (Q_T/Q_S) \end{aligned}$$

$W_S$ : Amount (mg) of Indometacin Reference Standard

**Internal standard solution**—A solution of butyl parahydroxybenzoate in methanol (1 in 1000).

**Operating conditions**—

**Detector**: An ultraviolet absorption photometer (wavelength: 254 nm).

**Column**: A stainless steel column 4.6 mm in inside diameter and 25 cm in length, packed with octadecylsilanized silica gel for liquid chromatography (7  $\mu$ m in particle diameter).

**Column temperature**: A constant temperature of about 25°C.

**Mobile phase**: A mixture of methanol and diluted phosphoric acid (1 in 1000) (7:3).

**Flow rate**: Adjust the flow rate so that the retention time of indometacin is about 8 minutes.

**System suitability**—

**System performance**: Dissolve 50 mg of 4-chlorobenzoic acid, 30 mg of butyl parahydroxybenzoate and 50 mg of indometacin in 50 mL of methanol. To 5 mL of this solution add the mobile phase to make 100 mL. When the procedure is run with 20  $\mu$ L of this solution under the above operating conditions, 4-chlorobenzoic acid, butyl parahydroxybenzoate and indometacin are eluted in this order, with the resolution between the peaks of 4-chlorobenzoic acid and butyl parahydroxybenzoate being not less than 2.0, and between the peaks of butyl parahydroxybenzoate and indometacin being not less than 5.

**System repeatability**: When the test is repeated 6 times with 20  $\mu$ L of the standard solution under the above operating conditions, the relative standard deviation of the ratios of the peak area of indometacin to that of the internal standard is not more than 1.0%.

**Containers and storage** Containers—Tight containers.

## Indometacin Suppositories

インドメタシン坐剤

Indometacin Suppositories contain not less than 90% and not more than 110% of the labeled amount of indometacin ( $C_{19}H_{16}ClNO_4$ ; 357.79).

**Method of preparation** Prepare as directed under Suppositories, with Indometacin.

**Identification** Dissolve a quantity of Indometacin Suppositories, equivalent to 0.05 g of Indometacin according to the labeled amount, in 20 mL of methanol by warming, add methanol to make 50 mL, and filter if necessary. To 2 mL of this solution add methanol to make 100 mL, and use this solution as the sample solution. Determine the absorption spectrum of the sample solution as directed under Ultraviolet-visible Spectrophotometry <2.24>: it exhibits a maximum between 317 nm and 321 nm.

**Assay** Weigh accurately not less than 20 Indometacin Suppositories, cut into small pieces carefully, and mix well. Weigh accurately a portion of the mass, equivalent to about 50 mg of indometacin ( $C_{19}H_{16}ClNO_4$ ), add 40 mL of tetrahydrofuran, warm at 40°C, dissolve by shaking, cool, and add tetrahydrofuran to make exactly 50 mL. Filter the solution, discard the first 10 mL of the filtrate, pipet the subsequent 5 mL of the filtrate, add exactly 3 mL of the internal standard solution, and add the mobile phase to make 100 mL. Allow the solution to stand for 30 minutes, filter through a membrane filter (0.5  $\mu$ m pore size), discard the first 10 mL of the filtrate, and use the subsequent filtrate as the sample solution. Separately, weigh accurately about 50 mg of Indometacin Reference Standard, previously dried at 105°C for 4 hours, and dissolve in tetrahydrofuran to make exactly 50 mL. Pipet 5 mL of the solution, proceed in the same manner as the sample solution, and use as the standard solution. Perform the test with 20  $\mu$ L each of the sample solution and standard solution as directed under Liquid Chromatography <2.01> according to the following conditions, and calculate the ratios,  $Q_T$  and  $Q_S$ , of the peak area of indometacin to that of the internal standard, respectively.

$$\begin{aligned} &\text{Amount (mg) of indometacin (C}_{19}\text{H}_{16}\text{ClNO}_4\text{)} \\ &= W_S \times (Q_T/Q_S) \end{aligned}$$

$W_S$ : Amount (mg) of Indometacin Reference Standard

**Internal standard solution**—A solution of butyl parahydroxybenzoate in methanol (1 in 1000).

**Operating conditions**—

**Detector**: An ultraviolet absorption photometer (wavelength: 254 nm).

**Column**: A stainless steel column 4.0 mm in inside diameter and 25 cm in length, packed with octadecylsilanized silica gel for liquid chromatography (7  $\mu$ m in particle diameter).

**Column temperature**: A constant temperature of about 25°C.

**Mobile phase**: A mixture of methanol and diluted phosphoric acid (1 in 1000) (7:3).

**Flow rate**: Adjust the flow rate so that the retention time of

indometacin is about 8 minutes.

*System suitability*—

**System performance:** Dissolve 50 mg of 4-chlorobenzoic acid, 30 mg of butyl parahydroxybenzoate and 50 mg of indometacin in 50 mL of methanol. To 5 mL of this solution add the mobile phase to make 100 mL. When the procedure is run with 20  $\mu$ L of this solution under the above operating conditions, 4-chlorobenzoic acid, butyl parahydroxybenzoate and indometacin are eluted in this order with the resolution between the peaks of 4-chlorobenzoic acid and butyl parahydroxybenzoate being not less than 2.0 and between the peaks of parahydroxybenzoate and indometacin being not less than 5.

**System repeatability:** When the test is repeated 6 times with 20  $\mu$ L of the standard solution under the above operating conditions, the relative standard deviation of the ratios of the peak area of indometacin to that of the internal standard is not more than 1.0%.

**Containers and storage** Containers—Well-closed containers.

Storage—Light-resistant, and in a cold place.

## Influenza HA Vaccine

インフルエンザ HA ワクチン

Influenza HA Vaccine is a liquid for injection containing hemagglutinin of influenza virus.

It conforms to the requirements of Influenza HA Vaccine in the Minimum Requirements for Biological Products.

**Description** Influenza HA Vaccine is a clear liquid or a slightly whitish turbid liquid.

## Insulin

インスリン

Insulin is obtained from the pancreas of healthy bovine or porcine, that has blood sugar-decreasing activity.

Its potency, calculated on the dried basis, is not less than 26 Insulin Units in each mg.

It is labeled to indicate the animal species from which it is derived.

**Description** Insulin occurs as a white, crystalline powder. It is odorless.

It is practically insoluble in water, in ethanol (95) and in diethyl ether.

It dissolves in diluted hydrochloric acid (1 in 360) and in dilute sodium hydroxide TS.

It is hygroscopic.

**Identification** Dissolve 0.01 g of Insulin in 10 mL of 0.1 mol/L hydrochloric acid TS, and use this solution as the sample solution. Adjust the pH of the sample solution to between 5.1 and 5.3 with a solution of sodium hydroxide (1 in 100): a precipitate is produced. Adjust the solution to a pH between

2.5 and 3.5 with dilute hydrochloric acid: the precipitate dissolves.

**Purity** Clarity and color of solution—Dissolve 0.10 g of Insulin in 10 mL of diluted hydrochloric acid (1 in 360): the solution is clear and colorless to light yellow.

**Zinc content** Weigh accurately about 10 mg of Insulin, dissolve in 5 mL of 0.1 mol/L hydrochloric acid TS and water to make exactly 50 mL. If necessary, dilute the solution with water so as to contain 0.4 to 1.0  $\mu$ g of zinc (Zn: 65.41) per mL, and use as the sample solution. Add water to an accurately measured volume of Standard Zinc Solution for atomic absorption spectrophotometry to make a solution containing 0.3 to 1.2  $\mu$ g of zinc (Zn: 65.41) per mL, and use as the standard solution. Perform the test with the sample solution and standard solution as directed under Atomic Absorption Spectrophotometry <2.23> according to the following conditions, and determine the amount of zinc in the sample solution using the calibration curve obtained from the absorbance of the standard solution: the amount of zinc is not less than 0.27% and not more than 1.08%, calculated on the dried basis.

Gas: Combustible gas—Acetylene gas

Supporting gas—Air

Lamp: Zinc hollow-cathode lamp

Wavelength: 213.9 nm

**Loss on drying** <2.41> Not more than 10.0% (0.2 g, 105°C, 16 hours).

**Residue on ignition** <2.44> Weigh accurately 0.02 to 0.04 g of Insulin in a tared platinum dish, add 2 drops of nitric acid, and heat the dish at first very gently and then strongly to incinerate. Place the dish in a muffle furnace, heat at 600°C for 15 minutes, cool in a desiccator (silica gel), and weigh: the mass of the residue is not more than 2.5%.

**Nitrogen content** Weigh accurately about 20 mg of Insulin, dissolve in 10 mL of 0.1 mol/L hydrochloric acid TS, and perform the test as directed under Nitrogen Determination <1.08>: not less than 14.5% and not more than 16.5% of nitrogen (N: 14.01) is found, calculated on the dried basis.

### Assay

(i) Animals: Select healthy rabbits each weighing not less than 1.8 kg. Keep the rabbits in the laboratory not less than 1 week before use in the assay by feeding them with an appropriate uniform diet and water.

(ii) Diluent for insulin: Dissolve 1.0 to 2.5 g of phenol or *m*-cresol in 500 mL of 0.01 mol/L hydrochloric acid VS, and add 14 to 18 g of glycerin and 0.01 mol/L hydrochloric acid VS to make 1000 mL.

(iii) Standard stock solution: Weigh accurately about 20 mg of Insulin Reference Standard, and dissolve it in the diluent for insulin to make a standard stock solution containing exactly 20.0 Units in each mL. Preserve this solution between 1°C and 15°C, and use within 6 months.

(iv) Standard solution: Dilute two portions of the standard stock solution to make two standard solutions with the diluent for insulin, one to contain exactly 2.0 Units in each mL which is designated as the high-dose standard solution  $S_H$ , and the other to contain exactly 1.0 Unit in each mL which is designated as the low-dose standard solution  $S_L$ .

(v) Sample solution: Weigh accurately about 20 mg of Insulin according to the labeled Units, dissolve with the diluent

for insulin to make two different sample solutions, one to contain exactly 2.0 Units in each mL which is designated as the high-dose sample solution,  $T_H$ , and the other to contain exactly 1.0 Unit in each mL which is designated as the low-dose sample solution,  $T_L$ .

(vi) Dose for injection: Select the dose for injection on the basis of trial or experience. Inject a fixed identical volume, usually 0.3 to 0.5 mL, of the standard solutions and the sample solutions throughout the whole run.

(vii) Procedure: Divide the animals into 4 equal groups of not less than 6 animals each, with least difference in body mass. Withhold all food, except water, for not less than 14 hours before the injections, and withhold water during the assay until the final blood sample is taken. Handle the animals with care in order to avoid undue excitement.

Inject into each of the animals subcutaneously the dose of the standard solutions and the sample solutions indicated in the following design.

|              |       |              |       |
|--------------|-------|--------------|-------|
| First group  | $S_H$ | Third group  | $T_H$ |
| Second group | $S_L$ | Fourth group | $T_L$ |

The second injection should be made on the day after the first injection or within 1 week, using the dose of the standard solutions and the sample solutions indicated in the following design.

|              |       |              |       |
|--------------|-------|--------------|-------|
| First group  | $T_L$ | Third group  | $S_L$ |
| Second group | $T_H$ | Fourth group | $S_H$ |

At 1 hour and 2.5 hours after the time of injection, obtain a sufficient blood sample to perform the test from a marginal ear vein of each animal, and determine the blood sugar content of the blood samples according to (viii).

(viii) Blood sugar determination: Place 5.0 mL of a solution of zinc sulfate heptahydrate (9 in 2000) in a test tube 18 mm in outside diameter and 165 mm in length, add 1.0 mL of a solution of sodium hydroxide (1 in 250), and add gently 0.10 mL of the blood sample to the mixture in the test tube using a blood sugar pipet. Suck up the supernatant liquid into the pipet, wash out the remaining blood in the inner wall of the pipet, and repeat this procedure. Shake thoroughly the contents in the test tube, and heat the test tube in a water bath for 3 minutes. Filter the mixture through a funnel 30 to 40 mm in diameter in which a pledget of absorbent cotton, previously washed with two 3-mL portions of warm water, has been placed, receive the filtrate into a test tube 30 mm in inside diameter and 90 mm in length, wash the test tube and the funnel with two 3-mL portions of water, and combine the washings with the filtrate. Add 2.0 mL of alkaline potassium hexacyanoferrate (III) TS, heat in a water bath for 15 minutes, cool immediately, add 3.0 mL of potassium iodide-zinc sulfate TS and 2.0 mL of diluted acetic acid (100) (3 in 100), and titrate <2.50> the liberated iodine with 0.005 mol/L sodium thiosulfate VS (indicator: 2 to 4 drops of starch-sodium chloride TS). Perform a blank determination. From the consumed volume (mL) of 0.005 mol/L sodium thiosulfate VS, obtain the blood sugar content (%) according to the following table.

Conversion Table for Blood sugar content (%)

| mL* | 0     | 1     | 2     | 3     | 4     | 5     | 6     | 7     | 8     | 9     |
|-----|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|
| 0.0 | 0.385 | 0.382 | 0.379 | 0.376 | 0.373 | 0.370 | 0.367 | 0.364 | 0.361 | 0.358 |
| 0.1 | 0.355 | 0.352 | 0.350 | 0.348 | 0.345 | 0.343 | 0.341 | 0.338 | 0.336 | 0.333 |
| 0.2 | 0.331 | 0.329 | 0.327 | 0.325 | 0.323 | 0.321 | 0.318 | 0.316 | 0.314 | 0.312 |
| 0.3 | 0.310 | 0.308 | 0.306 | 0.304 | 0.302 | 0.300 | 0.298 | 0.296 | 0.294 | 0.292 |
| 0.4 | 0.290 | 0.288 | 0.286 | 0.284 | 0.282 | 0.280 | 0.278 | 0.276 | 0.274 | 0.272 |
| 0.5 | 0.270 | 0.268 | 0.266 | 0.264 | 0.262 | 0.260 | 0.259 | 0.257 | 0.255 | 0.253 |
| 0.6 | 0.251 | 0.249 | 0.247 | 0.245 | 0.243 | 0.241 | 0.240 | 0.238 | 0.236 | 0.234 |
| 0.7 | 0.232 | 0.230 | 0.228 | 0.226 | 0.224 | 0.222 | 0.221 | 0.219 | 0.217 | 0.215 |
| 0.8 | 0.213 | 0.211 | 0.209 | 0.208 | 0.206 | 0.204 | 0.202 | 0.200 | 0.199 | 0.197 |
| 0.9 | 0.195 | 0.193 | 0.191 | 0.190 | 0.188 | 0.186 | 0.184 | 0.182 | 0.181 | 0.179 |
| 1.0 | 0.177 | 0.175 | 0.173 | 0.172 | 0.170 | 0.168 | 0.166 | 0.164 | 0.163 | 0.161 |
| 1.1 | 0.159 | 0.157 | 0.155 | 0.154 | 0.152 | 0.150 | 0.148 | 0.146 | 0.145 | 0.143 |
| 1.2 | 0.141 | 0.139 | 0.138 | 0.136 | 0.134 | 0.132 | 0.131 | 0.129 | 0.127 | 0.125 |
| 1.3 | 0.124 | 0.122 | 0.120 | 0.119 | 0.117 | 0.115 | 0.113 | 0.111 | 0.110 | 0.108 |
| 1.4 | 0.108 | 0.104 | 0.102 | 0.101 | 0.099 | 0.097 | 0.095 | 0.093 | 0.092 | 0.090 |
| 1.5 | 0.088 | 0.086 | 0.084 | 0.083 | 0.081 | 0.079 | 0.077 | 0.075 | 0.074 | 0.072 |
| 1.6 | 0.070 | 0.068 | 0.066 | 0.065 | 0.063 | 0.061 | 0.059 | 0.057 | 0.056 | 0.054 |
| 1.7 | 0.052 | 0.050 | 0.048 | 0.047 | 0.045 | 0.043 | 0.041 | 0.039 | 0.038 | 0.036 |
| 1.8 | 0.034 | 0.032 | 0.031 | 0.029 | 0.027 | 0.025 | 0.024 | 0.022 | 0.020 | 0.019 |
| 1.9 | 0.017 | 0.015 | 0.014 | 0.012 | 0.010 | 0.008 | 0.007 | 0.005 | 0.003 | 0.002 |

\*Indicates the volume of 0.005 mol/L sodium thiosulfate VS required in titration. For example, if the amount was 1.28 mL, the blood sugar content would be 0.127% from the above table.

(ix) Calculation: Sum up the two blood sugar values of each animal after each injection. Subtract the blood sugar value effected by the first injection from that effected by the second injection of each animal in the first group and the third group. The differences are symbolized as  $y_1$  and  $y_3$ , respectively. Subtract the blood sugar value effected by the second injection from that effected by the first injection of each animal in the second group and the fourth group. The differences are symbolized as  $y_2$  and  $y_4$ , respectively. Sum up not less than 6 values of individual differences in the blood sugar values  $y_1$ ,  $y_2$ ,  $y_3$ , and  $y_4$  to obtain  $Y_1$ ,  $Y_2$ ,  $Y_3$ , and  $Y_4$ , respectively.

Units in each mg of Insulin

= antilog  $M \times (\text{Units in each mL of } S_H) \times (b/a)$

$M = 0.301 \times (Y_a/Y_b)$

$Y_a = -Y_1 + Y_2 + Y_3 - Y_4$

$Y_b = Y_1 + Y_2 + Y_3 + Y_4$

$a$ : Amount (mg) of the sample.

$b$ : Total volume (mL) of the high-dose sample solution prepared by dissolving the sample with the diluent for insulin.

Compute  $L$  ( $P=0.95$ ) by using the following equation:  $L$  should be not more than 0.1212. If  $L$  exceeds 0.1212, repeat the assay by increasing the number of animals or improving the assay conditions in a better way until  $L$  becomes not more than 0.1212.

$$L = 2\sqrt{(C-1)(CM^2 + 0.09062)}$$

$$C = Y_b^2 / (Y_b^2 - 4fs^2t^2)$$

$f$ : Number of the animals of each group.

$$s^2 = \{ \sum y^2 - (Y/f) \} / n$$

$\sum y^2$ : The sum of squares of  $y_1$ ,  $y_2$ ,  $y_3$ , and  $y_4$  in each group.



$$Y = Y_1^2 + Y_2^2 + Y_3^2 + Y_4^2$$

$$n = 4(f - 1)$$


$t^2$ : Value shown in the following table against  $n$  for which  $s^2$  is calculated.

| $n$ | $t^2 = F_1$ | $n$ | $t^2 = F_1$ | $n$      | $t^2 = F_1$ |
|-----|-------------|-----|-------------|----------|-------------|
| 1   | 161.45      | 13  | 4.667       | 25       | 4.242       |
| 2   | 18.51       | 14  | 4.600       | 26       | 4.225       |
| 3   | 10.129      | 15  | 4.543       | 27       | 4.210       |
| 4   | 7.709       | 16  | 4.494       | 28       | 4.196       |
| 5   | 6.608       | 17  | 4.451       | 29       | 4.183       |
| 6   | 5.987       | 18  | 4.414       | 30       | 4.171       |
| 7   | 5.591       | 19  | 4.381       | 40       | 4.085       |
| 8   | 5.318       | 20  | 4.351       | 60       | 4.001       |
| 9   | 5.117       | 21  | 4.325       | 120      | 3.920       |
| 10  | 4.965       | 22  | 4.301       | $\infty$ | 3.841       |
| 11  | 4.844       | 23  | 4.279       |          |             |
| 12  | 4.747       | 24  | 4.260       |          |             |

**Containers and storage** Containers—Tight containers.  
Storage—Not exceeding 8°C.

## Insulin Human (Genetical Recombination)

ヒトインスリン(遺伝子組換え)



$C_{257}H_{383}N_{65}O_{77}S_6$ : 5807.57  
[11061-68-0]

Insulin Human (Genetical Recombination) is a human insulin prepared by genetical recombinant technology.

It has an activity to reduce the blood sugar concentration. It contains not less than 27.5 Insulin Units per mg, calculated on the dried basis.

**Description** Insulin Human (Genetical Recombination) occurs as a white powder.

It is practically insoluble in water and in ethanol (95).

It dissolves in 0.01 mol/L hydrochloric acid TS and in sodium hydroxide TS with decomposition.

It is hygroscopic.

**Identification** Weigh accurately a suitable amount of Insulin Human (Genetical Recombination), and dissolve in 0.01 mol/L hydrochloric acid TS to make a solution so that each mL contains 2.0 mg. Transfer 500  $\mu$ L of this solution into a clean test tube, add 2.0 mL of HEPES buffer solution, pH 7.5 and 400  $\mu$ L of V8-protease TS, react at 25°C for 6 hours, then add 2.9 mL of ammonium sulfate buffer solution to stop the reaction, and use this solution as the sample solution.

Separately, proceed with Human Insulin Reference Standard in the same manner as above, and use this solution as the standard solution. Perform the test with exactly 50  $\mu$ L each of the sample solution and standard solution as directed under Liquid Chromatography <2.01> according to the following conditions, and compare the chromatograms obtained from these solutions: the peak appears just after the peak of the solvent and the succeeding three peaks with apparently higher peak height show the same retention time and similar peak height each other on both chromatograms.

**Operating conditions—**

Detector: An ultraviolet absorption photometer (wavelength: 214 nm).

Column: A stainless steel column 4.6 mm in inside diameter and 10 cm in length, packed with octadecylsilanized silica gel for liquid chromatography (3  $\mu$ m in particle diameter).

Column temperature: A constant temperature of about 40°C.

Mobile phase: Solution A—A mixture of water, ammonium sulfate buffer solution and acetonitrile (7:2:1). Solution B—A mixture of water, acetonitrile and ammonium sulfate buffer solution (2:2:1).

Change the mixing ratio of the solutions A and B linearly from 9:1 to 3:7 in 60 minutes after sample injection, further change to 0:10 linearly in 5 minutes, and then flow the solution B only for 5 minutes.

Flow rate: 1.0 mL per minute.

**System suitability—**

System performance: When the procedure is run with 50  $\mu$ L of the standard solution under the above operating conditions, the symmetry factor of the two larger peaks which appear next to the first peak just after the solvent peak are not more than 1.5, and the resolution between these peaks is not less than 3.4.

**Purity (1)** Related substances—Perform this procedure rapidly. Dissolve 7.5 mg of Insulin Human (Genetical Recombination) in 2 mL of 0.01 mol/L hydrochloric acid TS, and use this solution as the sample solution. Perform the test with 20  $\mu$ L of the sample solution as directed under Liquid Chromatography <2.01> according to the following conditions, and determine the peak area of human insulin,  $A_1$ , the peak area of the desamide substance at the relative retention time of 1.3 to the human insulin,  $A_D$ , and the total area of the peaks other than the solvent peak,  $A_T$ : the amounts of the desamide substance and related substances other than the desamide substance are each not more than 2.0%. Previously, perform the test with 0.01 mol/L hydrochloric acid TS in the same manner to confirm the solvent peak.

Amount (%) of the desamide substance =  $(A_D/A_T) \times 100$

Amount (%) of related substances other than the desamide substance =  $\{[A_T - (A_1 + A_D)]/A_T\} \times 100$

**Operating conditions—**

Detector: An ultraviolet absorption photometer (wavelength: 214 nm).

Column: A stainless steel column 4.6 mm in inside diameter and 25 cm in length, packed with octadecylsilanized silica gel for liquid chromatography (5  $\mu$ m in particle diameter).

Column temperature: A constant temperature of about 40°C.

Mobile phase: Solution A—A mixture of phosphoric acid-sodium sulfate buffer solution, pH 2.3 and acetonitrile for liquid chromatography (41:9). Solution B—A mixture of phosphoric acid-sodium sulfate buffer solution, pH 2.3 and acetonitrile for liquid chromatography (1:1).

Flow a mixture of the solution A and the solution B (78:22) for 36 minutes before and after the sample injection, then change the mixing ratio to 33:67 linearly in 25 minutes, and maintain this ratio for 6 minutes. Then flow the first mixture (78:22) for the next 15 minutes. Adjust the mixing ratio of the first mixture so that the retention time of human insulin is about 25 minutes.

Flow rate: 1.0 mL per minute.

Time span of measurement: For about 75 minutes after the sample is injected.

*System suitability—*

Test for required detection: Confirm that the peak height of the desamide substance obtained from 20  $\mu$ L of human insulin desamide substance-containing TS is between 30% and 70% of the full scale.

System performance: When the procedure is run with 20  $\mu$ L of human insulin desamide substance-containing TS under the above operating conditions, human insulin and human insulin desamide substance are eluted in this order with the resolution between these peaks being not less than 2.0, and the symmetry factor of the peak of human insulin is not more than 1.8.

(2) High-molecular proteins—Dissolve 4 mg of Insulin Human (Genetical Recombination) in 1 mL of 0.01 mol/L hydrochloric acid TS. Perform the test with 100  $\mu$ L of this solution as directed under Liquid Chromatography <2.01> according to the following conditions, and calculate each peak area: the total of areas of the peaks having smaller retention time than human insulin is not more than 1.0% of the total area of all peaks.

*Operating conditions—*

Detector: An ultraviolet absorption photometer (wavelength: 276 nm).

Column: A stainless steel column 7.5 mm in inside diameter and 30 cm in length, packed with hydrophilic silica gel for liquid chromatography.

Column temperature: A constant temperature of about 25°C.

Mobile phase: A mixture of a solution of L-arginine (1 in 1000), acetonitrile and acetic acid (100) (13:4:3).

Flow rate: Adjust the flow rate so that the retention time of human insulin is about 20 minutes.

Time span of measurement: Until the peak of human insulin monomer has appeared.

*System suitability—*

Test for required detection: Confirm that the peak height of the dimer obtained from 100  $\mu$ L of human insulin dimer containing TS is between 10% and 50% of the full scale.

System performance: When the procedure is run with 100  $\mu$ L of human insulin dimer containing TS under the above operating conditions, polymer, dimer and monomer are eluted in this order, and the ratio,  $H_1/H_2$ , of the peak height of the dimer  $H_1$  to the height of the bottom between the peaks of the dimer and the monomer  $H_2$  is not less than 2.0.

(3) Product related impurities—Within the limits specified in each application dossier.

(4) Process related impurities—Within the limits specified in each application dossier.

**Zinc content** Weigh accurately about 50 mg of Insulin Human (Genetical Recombination), and dissolve in 0.01 mol/L hydrochloric acid TS to make exactly 25 mL. If necessary, dilute with 0.01 mol/L hydrochloric acid TS to make a solution so that each mL contains between 0.4  $\mu$ g and 1.6  $\mu$ g of zinc (Zn: 65.41), and use this solution as the sample solution. Separately, take exactly a suitable amount of Standard Zinc Solution for Atomic Absorption Spectrophotometry, dilute with 0.01 mol/L hydrochloric acid TS to make solutions containing 0.40  $\mu$ g, 0.80  $\mu$ g, 1.20  $\mu$ g and 1.60  $\mu$ g of zinc (Zn: 65.41) in each mL, respectively, and use these solutions as the standard solutions. Perform the test with the sample solution and standard solutions as directed under Atomic Absorption Spectrophotometry <2.23>, and determine the amount of zinc (Zn: 65.41) in the sample solution by using a calibration curve obtained from the absorbances of the standard solutions: not more than 1.0%, calculated on the dried basis.

Gas: Combustible gas—Acetylene

Supporting gas—Air

Lamp: Zinc hollow cathode lamp

Wavelength: 213.9 nm

**Loss on drying** <2.41> Not more than 10.0% (0.2 g, 105°C, 24 hours).

**Bacterial endotoxins** <4.01> Less than 10 EU/mg.

**Assay** Perform this procedure quickly. Weigh accurately about 7.5 mg of Insulin Human (Genetical Recombination), dissolve in 0.01 mol/L hydrochloric acid TS to make exactly 5 mL, and use this solution as the sample solution. Separately, weigh accurately a suitable amount of Human Insulin Reference Standard, dissolve exactly in 0.01 mol/L hydrochloric acid TS to make a solution so that each mL contains about 40 Insulin Units, and use this solution as the standard solution. Perform the test with exactly 20  $\mu$ L each of the sample solution and standard solution as directed under Liquid Chromatography <2.01> according to the following conditions, and determine the peak areas of human insulin,  $A_{TI}$  and  $A_{SI}$ , and the peak areas of the desamide substance at the relative retention time of 1.3 to the human insulin,  $A_{TD}$  and  $A_{SD}$ , respectively, of these solutions.

Amount (Insulin Unit/mg) of human insulin

$$\begin{aligned} & (C_{257}H_{383}N_{65}O_{77}S_6) \\ &= \{(W_S \times F)/D\} \times \{(A_{TI} + A_{TD})/(A_{SI} + A_{SD})\} \\ & \times (5/W_T) \end{aligned}$$

$F$ : Label unit (Insulin Unit/mg) of Human Insulin Reference Standard.

$D$ : Volume (mL) of 0.01 mol/L hydrochloric acid TS used to dissolve the reference standard.

$W_T$ : Amount (mg) of the sample calculated on the dried basis.

$W_S$ : Amount (mg) of Human Insulin Reference Standard.

*Operating conditions—*

Detector: An ultraviolet absorption photometer (wavelength: 214 nm).

Column: A stainless steel column 4.6 mm in inside diameter and 15 cm in length, packed with octadecylsilanized silica gel for liquid chromatography (5  $\mu$ m in particle diameter).

Column temperature: A constant temperature of about 40°C.

Mobile phase: A mixture of phosphoric acid-sodium sul-

fate buffer solution, pH 2.3 and acetonitrile for liquid chromatography (3:1). Adjust the mixing ratio of the component of the mobile phase so that the retention time of human insulin is between 10 minutes and 17 minutes.

Flow rate: 1.0 mL per minute.

**System suitability—**

**System performance:** When the procedure is run with 20  $\mu$ L of human insulin desamide substance-containing TS under the above operating conditions, human insulin and human insulin desamide substance are eluted in this order with the resolution between these peaks being not less than 2.0, and the symmetry factor of the peak of human insulin is not more than 1.8.

**System repeatability:** When the test is repeated 6 times with 20  $\mu$ L of the standard solution under the above operating conditions, the relative standard deviation of the peak areas of human insulin is not more than 1.6%.

**Containers and storage** Containers—Tight containers.

Storage—Light-resistant, and at  $-20^{\circ}\text{C}$  or below.

## Insulin Injection

### インスリン注射液

Insulin Injection is an aqueous solution for injection.

It contains not less than 95% and not more than 105% of the labeled Insulin Units.

**Method of preparation** Suspend Insulin in Water for Injection, dissolve by adding Hydrochloric Acid, and prepare as directed under Injections. It contains 0.10 to 0.25 g of Phenol or Cresol and 1.4 to 1.8 g of Concentrated Glycerin for each 100 mL of Insulin Injection. It should not contain sodium chloride.

**Description** Insulin Injection is a clear, colorless or pale yellow liquid.

**Identification** Adjust Insulin Injection to pH between 5.1 and 5.3 with a solution of sodium hydroxide (1 in 100): a precipitate is produced. Adjust the solution to a pH between 2.5 and 3.5 with dilute hydrochloric acid: the precipitate dissolves.

**pH** <2.54> 2.5 – 3.5

**Residue on ignition** <2.44> Measure exactly a volume of Insulin Injection, equivalent to 500 to 1000 Units according to the labeled Units, in a tared platinum dish, and evaporate slowly by heating on a water bath to dryness. Add 2 drops of nitric acid to the residue, and heat at first very gently, then strongly to incinerate. Place in a muffle furnace, and heat at  $600^{\circ}\text{C}$  for 15 minutes, cool in a desiccator (silica gel), and weigh: the mass of the residue is not more than 1 mg for each labeled 1000 Units.

**Extractable volume** <6.05> It meets the requirement.

**Nitrogen content** Perform the test as directed under Nitrogen Determination <1.08>: not less than 0.50 mg and not more than 0.64 mg of nitrogen (N: 14.01) is found for each labeled 100 Units.

**Assay** Proceed with Insulin Injection as directed in the As-

say under Insulin with alterations in (v) Sample solution and (ix) Calculation as follows.

(v) Sample solution: According to the labeled Units, dilute two portions of Insulin Injection to make two sample solutions with the diluent for insulin, one to contain exactly 2.0 Units in each mL which is designated as the high-dose sample solution  $T_H$ , and the other to contain exactly 1.0 Unit in each mL which is designated as the low-dose sample solution  $T_L$ .

(ix) Calculation: Proceed as directed in the Assay (ix) Calculation under Insulin, replacing the equation as follows.

Units in each mL of Insulin Injection

$$= \text{antilog } M \times (\text{Units in each mL of } S_H) \times (b/a)$$

$a$ : Volume (mL) of the sample.

$b$ : Total volume (mL) of the high-dose sample solution prepared by diluting the volume of the sample with diluent for insulin.

**Containers and storage** Containers—Hermetic containers.

Storage—In a cold place, and avoid freezing.

**Expiration date** 24 months after preparation.

## Isophane Insulin Injection (Aqueous Suspension)

### イソフェンインスリン水性懸濁注射液

Isophane Insulin Injection (Aqueous Suspension) is an aqueous suspension for injection.

It contains not less than 90% and not more than 110% of the labeled Insulin Units, and not less than 0.01 mg and not more than 0.04 mg of zinc (Zn: 65.41) for each labeled 100 Units.

When Sodium Chloride is used in the preparation of Isophane Insulin Injection (Aqueous Suspension), this should be stated on the label.

**Method of preparation** Prepare as directed under Injections, with Insulin and Protamine Sulfate. To each 100 mL of Isophane Insulin Injection (Aqueous Suspension) add either 0.38 to 0.63 g of Dibasic Sodium Phosphate, 1.4 to 1.8 g of Concentrated Glycerin, 0.15 to 0.17 g of Cresol, and 0.06 to 0.07 g of Phenol, or 0.38 to 0.63 g of Dibasic Sodium Phosphate, 0.42 to 0.45 g of Sodium Chloride, 0.7 to 0.9 g of Concentrated Glycerin, and 0.18 to 0.22 g of Cresol.

**Description** Isophane Insulin Injection (Aqueous Suspension) is a white aqueous suspension. When allowed to stand, it separates into a white precipitate and colorless supernatant liquid, and the precipitate returns easily to the suspension state on gentle shaking.

When examined microscopically, the precipitate mostly consists of fine, oblong crystals of 5 to 30  $\mu\text{m}$  in major axis, and does not contain amorphous substances or large aggregates.

**Identification** Proceed as directed in the Identification under Insulin Zinc Protamine Injection (Aqueous Suspension).

**pH** <2.54> 7.0 – 7.4

**Purity** (1) Protein—Perform the test as directed under Nitrogen Determination <1.08>: not exceeding 0.85 mg of

nitrogen (N: 14.01) is found for each labeled 100 Units.

(2) Isophane ratio—(i) Buffer solution A: Dissolve 2.0 g of anhydrous disodium hydrogenphosphate, 16 g of glycerin, 1.6 g of *m*-cresol, and 0.65 g of phenol in water to make exactly 200 mL.

(ii) Buffer solution B: Dissolve 2.0 g of anhydrous disodium hydrogenphosphate, 4.35 g of sodium chloride, 8.0 g of glycerin, and 2.0 g of *m*-cresol in water to make exactly 200 mL.

(iii) Insulin solution: Weigh accurately 1000 Units of Insulin Reference Standard, dissolve in 1.5 mL of diluted hydrochloric acid (1 in 360), and add 5.0 mL of buffer solution A and water to make 20 mL. Adjust the pH to 7.2 with dilute hydrochloric acid or sodium hydroxide TS. The solution is clear. Dilute with water to make exactly 25 mL. The solution is clear, and the pH is between 7.1 and 7.4. When it is stated on the label that Sodium Chloride is used in the preparation, use 5.0 mL of buffer solution B instead of buffer solution A in the above procedure.

(iv) Protamine solution: Weigh accurately 50 mg of Protamine Sulfate Reference Standard, and dissolve in 2 mL of buffer solution A and water to make 8 mL. Adjust the pH to 7.2 with dilute hydrochloric acid or sodium hydroxide TS, and dilute with water to exactly 10 mL. The solution is clear, and the pH is between 7.1 and 7.4. When it is stated on the label that Sodium Chloride is used in the preparation, use 2 mL of buffer solution B instead of buffer solution A in the above procedure.

(v) Procedure: When Isophane Insulin Injection (Aqueous Suspension) contains 40 Units per mL, centrifuge a portion of the suspension, measure exactly two 10-mL portions of the supernatant liquid in two tubes A and B, respectively, add exactly 1 mL of the insulin solution to tube A, and 1 mL of the protamine solution to tube B, mix the contents of each tube, allow to stand for 10 minutes, and determine the turbidity of each mixture by using a photometer or a nephelometer: the turbidity of the mixture in tube B is not greater than that in tube A. When Isophane Insulin Injection (Aqueous Suspension) contains 80 Units per mL, measure exactly 5 mL of the supernatant liquid, and proceed in the same manner.

**Extractable volume** <6.05> It meets the requirement.

**Assay** (1) Insulin—To Isophane Insulin Injection (Aqueous Suspension) add diluted hydrochloric acid (1 in 1000) to adjust pH to about 2.5, and proceed with the clear solution as directed in the Assay under Insulin, with alterations in (v) Sample solution and (ix) Calculation as follows.

(v) Sample solution: According to the labeled Units, dilute Insulin Injection (Aqueous Suspension) to make two sample solutions with the diluent for insulin, one to contain exactly 2.0 Units in each mL which is designated as the high-dose sample solution  $T_H$ , and the other to contain exactly 1.0 Unit in each mL which is designated as the low-dose sample solution  $T_L$ .

(ix) Calculation: Proceed as directed in the Assay (ix) Calculation under Insulin, using the following equation,

Units in each mL of Isophane Insulin Injection (Aqueous Suspension)

$$= \text{antilog } M \times (\text{Units in each mL of } S_H) \times (b/a)$$

$a$ : Volume (mL) of the sample,

instead of the following equation,

Units in each mg of Insulin

$$= \text{antilog } M \times (\text{Units in each mL of } S_H) \times (b/a)$$

$a$ : Mass (mg) of the sample.

(2) Zinc—Pipet a volume of Isophane Insulin Injection (Aqueous Suspension), equivalent to about 400 Units according to the labeled Units, add 1 mL of 0.1 mol/L hydrochloric acid TS and water to make exactly 100 mL, dilute, if necessary, with water to contain 0.6 to 1.0  $\mu\text{g}$  of zinc (Zn: 65.41) per mL, and use this solution as the sample solution. Separately, pipet a volume of Standard Zinc Solution for Atomic Absorption Spectrophotometry, dilute with water to contain 0.4 to 1.2  $\mu\text{g}$  of zinc (Zn: 65.41) per mL, and use this solution as the standard solution. Perform the test with the sample solution and the standard solution according to Atomic Absorption Spectrophotometry <2.23> under the following conditions, and determine the amount of zinc in the sample solution using the analytical curve obtained from the absorbance of the standard solution.

Gas: Combustible gas—Acetylene gas

Supporting gas—Air

Lamp: Zinc hollow – cathode lamp

Wavelength: 213.9 nm

**Containers and storage** Containers—Hermetic containers.

Storage—In a cold place, and avoid freezing.

**Expiration date** 24 months after preparation.

## Insulin Zinc Injection (Aqueous Suspension)

インスリン亜鉛水性懸濁注射液

Insulin Zinc Injection (Aqueous Suspension) is an aqueous suspension for injection.

It contains not less than 90% and not more than 110 % of the labeled Insulin Units, and not less than 0.12 mg and not more than 0.30 mg of zinc (Zn: 65.41) for each labeled 100 Units.

**Method of preparation** Prepare as directed under Injections, with Insulin and Zinc Chloride. It contains 0.15 to 0.17 g of Sodium Acetate Hydrate, 0.65 to 0.75 g of Sodium Chloride and 0.09 to 0.11 g of Methyl Parahydroxybenzoate for each 100 mL of Insulin Zinc Injection (Aqueous Suspension).

**Description** Insulin Zinc Injection (Aqueous Suspension) is a white suspension. When allowed to stand, it separates into a white precipitate and a colorless supernatant liquid, and it readily becomes a suspension again on gentle shaking.

When it is examined microscopically, the majority of the particles in the suspension are crystals, the dimension of which is 10 to 40  $\mu\text{m}$ . The rest is amorphous and does not exceed 2  $\mu\text{m}$  in dimension.

**Identification** Adjust the pH of Amorphous Insulin Zinc Injection (Aqueous Suspension) to between 2.5 and 3.5 with dilute hydrochloric acid: the particles dissolve, and the solution is clear and colorless.

**pH** <2.54> 7.1 – 7.5

**Purity** Dissolved insulin—Perform the following test with a

clear liquid obtained by centrifuging Insulin Zinc Injection (Aqueous Suspension): not more than 2.5% of the labeled units is found.

Use the clear liquid of Insulin Zinc Injection (Aqueous Suspension) as the sample solution. Prepare the standard solution having a concentration of 2.5% of the labeled units of Insulin Zinc Injection (Aqueous Suspension) by proceeding as directed in the Assay (iv) under Insulin. Divide the healthy rabbits weighing not less than 1.8 kg, fasted for not less than 14 hours before injection, into 2 equal groups of not less than 3. Inject subcutaneously an amount of the standard solution or the sample solution equivalent to 0.3 units per kg of body mass to the animals of each group. Collect blood before and 1 hour and 2.5 hours after injection, then proceed as directed in the Assay (viii) under Insulin, and calculate the ratio of the average blood sugar level of 1 hour and 2.5 hours after to that of before injection of each animal: the mean value for the group injected the sample solution is not less than that for the group injected the standard solution.

**Extractable volume** <6.05> It meets the requirement.

**Nitrogen content** Perform the test as directed under Nitrogen Determination <1.08>: the amount of nitrogen (N: 14.01) is not less than 0.50 mg and not more than 0.64 mg for each labeled 100 Units.

**Assay (1)** Insulin—Proceed as directed in the Assay under Insulin with the clear liquid obtained from Insulin Zinc Injection (Aqueous Suspension) by adjusting the pH to about 2.5 with diluted hydrochloric acid (1 in 1000), with alterations in (v) Sample solution and (ix) Calculation as follows.

(v) Sample solution: According to the labeled Units, dilute two portions of Insulin Zinc Injection (Aqueous Suspension) to make two sample solutions with the diluent for insulin, one to contain exactly 2.0 Units in each mL which is designated as the high-dose sample solution  $T_H$ , and the other to contain exactly 1.0 Unit in each mL which is designated as the low-dose sample solution  $T_L$ .

(ix) Calculation: Proceed as directed in the Assay (ix) Calculation under Insulin, using the following equation,

Units in each mL of Insulin Zinc Injection (Aqueous Suspension)

$$= \text{antilog } M \times (\text{Units in each mL of } S_H) \times (b/a)$$

$a$ : Volume (mL) of the sample,

instead of the following equation,

Units in each mg of Insulin

$$= \text{antilog } M \times (\text{Units in each mL of } S_H) \times (b/a)$$

$a$ : Mass (mg) of the sample.

(2) Zinc—Measure exactly a volume of Insulin Zinc Injection (Aqueous Suspension), equivalent to about 200 Units according to the labeled units, add 1 mL of 0.1 mol/L hydrochloric acid TS and sufficient water to make exactly 200 mL, then dilute with water to contain 0.6 to 1.0  $\mu\text{g}$  of zinc (Zn: 65.41) per mL, and use this solution as the sample solution. Separately, pipet a volume of Standard Zinc Solution for atomic absorption spectrophotometry, dilute with water to contain 0.4 to 1.2  $\mu\text{g}$  of zinc (Zn: 65.41) per mL, and use this solution as the standard solution. Perform the test with the sample solution and standard solution as directed under Atomic Absorption Spectrophotometry <2.23> according to

the following conditions, and determine the amount of zinc in the sample solution using the calibration curve obtained from the absorbance of the standard solution.

Gas: Combustible gas—Acetylene gas

Supporting gas—Air

Lamp: Zinc hollow-cathode lamp

Wavelength: 213.9 nm

(3) Crystalline insulin—Measure exactly a volume of Amorphous Insulin Zinc Injection (Aqueous Suspension), equivalent to about 600 Units according to the labeled units, centrifuge, discard the supernatant liquid, suspend the residue in 5 mL of water, add 10 mL of sodium acetate-acetone TS, shake for 3 minutes, and centrifuge. Discard the supernatant liquid, and repeat the above treatment on the residue. Wash down the residue into a Kjeldahl flask with 15 mL of sulfuric acid, and perform the test as directed under the Nitrogen Determination <1.08>: the amount of nitrogen (N: 14.01) is not less than 55% and not more than 70% of the total nitrogen content. Calculate the total nitrogen content for insulin Units of the sample taken from the values of nitrogen obtained in the Nitrogen content.

**Containers and storage** Containers—Hermetic containers.

Storage—In a cold place, and avoid freezing.

**Expiration date** 24 months after preparation.

## Amorphous Insulin Zinc Injection (Aqueous Suspension)

無晶性インスリン亜鉛水性懸濁注射液

Amorphous Insulin Zinc Injection (Aqueous Suspension) is an aqueous suspension for injection.

It contains not less than 90% and not more than 110 % of the labeled Insulin Units, and not less than 0.12 mg and not more than 0.30 mg of zinc (Zn: 65.41) for each labeled 100 Units.

**Method of preparation** Prepare as directed under Injections, with Insulin and Zinc Chloride. Each 100 mL of Amorphous Insulin Zinc Injection (Aqueous Suspension) contains 0.15 to 0.17 g of Sodium Acetate Hydrate, 0.65 to 0.75 g of Sodium Chloride, and 0.09 to 0.11 g of Methyl Parahydroxybenzoate.

**Description** Amorphous Insulin Zinc Injection (Aqueous Suspension) is a white suspension. When allowed to stand, it separates into a white precipitate and a colorless supernatant liquid, and it readily becomes a suspension again on gentle shaking.

When examined microscopically, most of the particles in the suspension are amorphous and have no uniform shape, and most of the dimension does not exceed 2  $\mu\text{m}$ .

**Identification** Adjust the pH of Amorphous Insulin Zinc Injection (Aqueous Suspension) to between 2.5 and 3.5 with dilute hydrochloric acid: the particles dissolve, and the solution is clear and colorless.

**pH** <2.54> 7.1 – 7.5

**Purity** Dissolved insulin—Perform the following test with a clear liquid obtained by centrifuging Amorphous Insulin

Zinc Injection (Aqueous Suspension): not more than 2.5% of the labeled units is found.

Use the clear liquid of Amorphous Insulin Zinc Injection (Aqueous Suspension) as the sample solution. Prepare the standard solution having a concentration of 2.5% of the labeled units of Amorphous Insulin Zinc Injection (Aqueous Suspension) by proceeding as directed in the Assay (iv) under Insulin. Divide healthy rabbits weighing more than 1.8 kg, fasted for not less than 14 hours before injection, into 2 equal groups of not less than 3. Inject subcutaneously an amount of the standard solution or the sample solution equivalent to 0.3 units per kg of body mass to the animals of each group. Collect blood before and 1 hour and 2.5 hours after injection, then proceed as directed in the Assay (viii) under Insulin, and calculate the ratio of the average blood sugar level of 1 hour and 2.5 hours after to that of before injection of each animal: the mean value for the group injected the sample solution is not less than that for the group injected the standard solution.

**Extractable volume** <6.05> It meets the requirement.

**Nitrogen content** Perform the test as directed under Nitrogen Determination <1.08>: the amount of nitrogen (N: 14.01) is not less than 0.50 mg and not more than 0.64 mg for each labeled 100 Units.

**Assay (1)** Insulin—Proceed as directed in the Assay under Insulin with the clear liquid obtained from Amorphous Insulin Zinc Injection (Aqueous Suspension) by adjusting the pH to about 2.5 with diluted hydrochloric acid (1 in 1000), with alteration in (v) Sample solution and (ix) Calculation as follows.

(v) Sample solution: According to the labeled Units, dilute two portions of Amorphous Insulin Zinc Injection (Aqueous Suspension) to make two sample solutions with the diluent for insulin, one to contain exactly 2.0 Units in each mL which is designated as the high-dose sample solution  $T_H$ , and the other to contain exactly 1.0 Unit in each mL which is designated as the low-dose sample solution  $T_L$ .

(ix) Calculation: Proceed as directed in the Assay (ix) Calculation under Insulin, using the following equation,

Units in each mL of Amorphous Insulin Zinc Injection (Aqueous Suspension)

$$= \text{antilog } M \times (\text{Units in each mL of } S_H) \times (b/a)$$

$a$ : Volume (mL) of the sample,

instead of the following equation,

Units in each mg of Insulin

$$= \text{antilog } M \times (\text{Units in each mL of } S_H) \times (b/a)$$

$a$ : Mass (mg) of the sample.

(2) Zinc—Measure exactly a volume of Amorphous Insulin Zinc Injection (Aqueous Suspension), equivalent to about 200 Units according to the labeled units, add 1 mL of 0.1 mol/L hydrochloric acid TS and sufficient water to make exactly 200 mL, then dilute with water to contain 0.6 to 1.0  $\mu\text{g}$  of zinc (Zn: 65.41) per mL, and use this solution as the sample solution. Separately, pipet a volume of Standard Zinc Solution for atomic absorption spectrophotometry, dilute with water to contain 0.4 to 1.2  $\mu\text{g}$  of zinc (Zn: 65.41) per mL, and use this solution as the standard solution. Perform the test with the sample solution and the standard solution as directed

under Atomic Absorption Spectrophotometry <2.23> according to the following conditions, and determine the amount of zinc in the sample solution using the calibration curve obtained from the absorbance of the standard solution.

Gas: Combustible gas—Acetylene gas

Supporting gas—Air

Lamp: Zinc hollow-cathode lamp

Wavelength: 213.9 nm

(3) Crystalline insulin—Measure exactly a volume of Amorphous Insulin Zinc Injection (Aqueous Suspension), equivalent to about 1000 Units according to the labeled units, centrifuge, discard the supernatant liquid, suspend the residue in 5 mL of water, add 10 mL of sodium acetate-acetone TS, shake for 3 minutes, and centrifuge. Discard the supernatant liquid, and repeat the above treatment on the residue. Wash down the residue into a Kjeldahl flask with 15 mL of sulfuric acid, and perform the test as directed under the Nitrogen Determination <1.08>: the amount of nitrogen (N: 14.01) is not more than 10% of the total nitrogen content. Calculate the total nitrogen content for insulin Units of the sample taken from the values of nitrogen obtained in the Nitrogen content.

**Containers and storage** Containers—Hermetic containers.

Storage—In a cold place, and avoid freezing.

**Expiration date** 24 months after preparation.

## Crystalline Insulin Zinc Injection (Aqueous Suspension)

結晶性インスリン亜鉛水性懸濁注射液

Crystalline Insulin Zinc Injection (Aqueous Suspension) is an aqueous suspension for injection.

It contains not less than 90% and not more than 110 % of the labeled Insulin Units, and not less than 0.12 mg and not more than 0.30 mg of zinc (Zn: 65.41) for each labeled 100 Units.

**Method of preparation** Prepare as directed under Injections, with Insulin and Zinc Chloride. Each 100 mL of Crystalline Insulin Zinc Injection (Aqueous Suspension) contains 0.15 to 0.17 g of Sodium Acetate Hydrate, 0.65 to 0.75 g of Sodium Chloride, and 0.09 to 0.11 g of Methyl Parahydroxybenzoate.

**Description** Crystalline Insulin Zinc Injection (Aqueous Suspension) is a white suspension. When allowed to stand, it separates into a white precipitate and a colorless supernatant liquid, and it readily becomes a suspension again on gentle shaking.

When it is examined microscopically, most part of the particles in the suspension are crystals, the dimension of which is mostly 10 to 40  $\mu\text{m}$ .

**Identification** Adjust the pH of Crystalline Insulin Zinc Injection (Aqueous Suspension) to between 2.5 and 3.5 with dilute hydrochloric acid: the particles dissolve, and the solution is clear and colorless.

**pH** <2.54> 7.1 – 7.5

**Purity** Dissolved insulin—Perform the following test with a

clear liquid obtained by centrifuging Crystalline Insulin Zinc Injection (Aqueous Suspension): not more than 2.5% of the labeled units is found.

Use the clear liquid of Crystalline Insulin Zinc Injection (Aqueous Suspension) as the sample solution. Prepare the standard solution having a concentration of 2.5% of the labeled units of Crystalline Insulin Zinc Injection (Aqueous Suspension) by proceeding as directed in the Assay (iv) under Insulin. Divide healthy rabbits weighing more than 1.8 kg, fasted for not less than 14 hours before injection, into 2 equal groups of not less than 3. Inject subcutaneously an amount of the standard solution or the sample solution equivalent to 0.3 units per kg of body mass to the animals of each group. Collect blood before and 1 hour and 2.5 hours after injection, then proceed as directed in the Assay (viii) under Insulin, and calculate the ratio of the average blood sugar level of 1 hour and 2.5 hours after to that of before injection of each animal: the mean value for the group injected the sample solution is not less than that for the group injected the standard solution.

**Extractable volume** <6.05> It meets the requirement.

**Nitrogen content** Perform the test as directed under Nitrogen Determination <1.08>: not less than 0.50 mg and not more than 0.64 mg of nitrogen (N: 14.01) is found for each labeled 100 Units.

**Assay (1)** Insulin—Proceed as directed in the Assay under Insulin with the clear liquid obtained from Crystalline Insulin Zinc Injection (Aqueous Suspension) by adjusting the pH to about 2.5 with diluted hydrochloric acid (1 in 1000), with alterations in (v) Sample solution and (ix) Calculation as follows.

(v) Sample solution: According to the labeled Units, dilute two portions of Crystalline Insulin Zinc Injection (Aqueous Suspension) to make two sample solutions with the diluent for insulin, one to contain exactly 2.0 Units in each mL which is designated as the high-dose sample solution  $T_H$ , and the other to contain exactly 1.0 Unit in each mL which is designated as the low-dose sample solution  $T_L$ .

(ix) Calculation: Proceed as directed in the Assay (ix) Calculation under Insulin, using the following equation,

Units in each mL of Crystalline Insulin Zinc Injection (Aqueous Suspension)

$$= \text{antilog } M \times (\text{Units in each mL of } S_H) \times (b/a)$$

$a$ : Volume (mL) of the sample,

instead of the following equation,

Units in each mg of Insulin

$$= \text{antilog } M \times (\text{Units in each mL of } S_H) \times (b/a)$$

$a$ : Mass (mg) of the sample.

(2) Zinc—Measure exactly a volume of Crystalline Insulin Zinc Injection (Aqueous Suspension), equivalent to about 200 Units according to the labeled units, add 1 mL of 0.1 mol/L hydrochloric acid TS and sufficient water to make exactly 200 mL, then dilute with water to contain 0.6 to 1.0  $\mu\text{g}$  of zinc (Zn: 65.41) per mL, and use this solution as the sample solution. Separately, pipet a volume of Standard Zinc Solution for atomic absorption spectrophotometry, dilute with water to contain 0.4 to 1.2  $\mu\text{g}$  of zinc (Zn: 65.41) per mL, and use this solution as the standard solution. Perform the test

with the sample solution and the standard solution as directed under Atomic Absorption Spectrophotometry <2.23> according to the following conditions, and determine the amount of zinc in the sample solution using the calibration curve obtained from the absorbance of the standard solution.

Gas: Combustible gas—Acetylene gas

Supporting gas—Air

Lamp: Zinc hollow-cathode lamp

Wavelength: 213.9 nm

(3) Crystalline insulin—Measure accurately a volume of Crystalline Insulin Zinc Injection (Aqueous Suspension), equivalent to about 400 Units according to the labeled Units, centrifuge, discard the supernatant liquid, suspend the residue in 5 mL of water, add 10 mL of sodium acetate-acetone TS, shake for 3 minutes, and centrifuge. Discard the supernatant liquid, and repeat the above treatment on the residue. Wash down the residue into a Kjeldahl flask with 15 mL of sulfuric acid, and perform the test as directed under the Nitrogen Determination <1.08>: the amount of nitrogen (N: 14.01) is not less than 85% of the total nitrogen content. Calculate the total nitrogen content for insulin Units of a sample from the values of nitrogen obtained in the Nitrogen content.

**Containers and storage** Containers—Hermetic containers.

Storage—In a cold place, and avoid freezing.

**Expiration date** 24 months after preparation.

## Insulin Zinc Protamine Injection (Aqueous Suspension)

プロタミンインスリン亜鉛水性懸濁注射液

Insulin Zinc Protamine Injection (Aqueous Suspension) is an aqueous suspension for injection.

It contains not less than 90% and not more than 110 % of the labeled Insulin Units, and not less than 0.20 mg and not more than 0.30 mg of zinc (Zn: 65.41) for each labeled 100 Units.

**Method of preparation** Prepare as directed under Injections, with Insulin, Protamine Sulfate and Zinc Chloride. It contains 0.38 to 0.63 g of Dibasic Sodium Phosphate Hydrate, 1.4 to 1.8 g of Concentrated Glycerin, and 0.18 to 0.22 g of Cresol or 0.22 to 0.28 g of Phenol for each 100 mL of Insulin Zinc Protamine Injection (Aqueous Suspension).

**Description** Insulin Zinc Protamine Injection (Aqueous Suspension) is a white suspension. When allowed to stand, it separates into a white precipitate and a colorless, supernatant liquid, and it readily becomes suspension again on gentle shaking.

When it is examined microscopically, no large particles are seen.

**Identification** Adjust the pH of Insulin Zinc Protamine Injection (Aqueous Suspension) to between 2.5 and 3.5 with dilute hydrochloric acid: the particles dissolve, and the solution is clear and colorless.

**pH** <2.54> 7.0 – 7.4

**Purity (1)** Protein—Perform the test as directed under

Nitrogen Determination <1.08>: not exceeding 1.25 mg of nitrogen (N: 14.01) is found for each labeled 100 Units.

(2) Dissolved insulin—Perform the following test with the clear liquid obtained by centrifuging Insulin Zinc Protamine Injection (Aqueous Suspension): not more than 2.5% of the labeled Units is found.

Use a clear liquid of Insulin Zinc Protamine Injection (Aqueous Suspension) as the sample solution, and prepare the standard solution by proceeding as directed in the Assay (iv) under Insulin to adjust its concentration to 2.5% of the labeled units of Insulin Zinc Protamine Injection (Aqueous Suspension). Divide the healthy rabbits weighing not less than 1.8 kg, fasted for not less than 14 hours before injection, into 2 equal groups of not less than 3. Inject subcutaneously an amount of the standard solution or the sample solution equivalent to 0.3 units per kg of body mass. Collect blood before and 1 hour and 2.5 hours after injection, proceed as directed in the Assay (viii) under Insulin, and calculate the ratios of the average blood sugar content in each rabbit measured 1 hour and 2.5 hours after injection to the content before injection: the mean value for the group injected with the sample solution is not less than the mean value for the group injected with the standard solution.

**Extractable volume** <6.05> It meets the requirement.

**Assay (1) Insulin**—Proceed as directed in the Assay under Insulin with the clear liquid obtained by adjusting the pH to 2.5 with diluted hydrochloric acid (1 in 1000), with alterations in (v) Sample solution and (ix) Calculation as follows.

(v) Sample solution: According to the labeled Units, dilute Insulin Zinc Protamine Injection (Aqueous Suspension) to make two sample solutions with the diluent for insulin, one to contain exactly 2.0 Units in each mL which is designated as the high-dose sample solution  $T_H$ , and the other to contain exactly 1.0 Unit in each mL which is designated as the low-dose sample solution  $T_L$ .

(ix) Calculation: Proceed as directed in the Assay (ix) Calculation under Insulin, using the following equation,

Units in each mL of Insulin Zinc Protamine Injection (Aqueous Suspension)

$$= \text{antilog } M \times (\text{Units in each mL of } S_H) \times (b/a)$$

$a$ : Volume (mL) of the sample,

instead of the following equation,

Units in each mg of Insulin

$$= \text{antilog } M \times (\text{Units in each mL of } S_H) \times (b/a)$$

$a$ : Mass (mg) of the sample.

(2) Zinc—Measure accurately a volume of Insulin Zinc Protamine Injection (Aqueous Suspension), equivalent to about 200 Units according to the labeled units, add 1 mL of 0.1 mol/L hydrochloric acid TS and sufficient water to make exactly 200 mL, dilute with water to contain 0.6 to 1.0  $\mu\text{g}$  of zinc (Zn: 65.41) in 1 mL, and use this solution as the sample solution. Separately, pipet a volume of Standard Zinc Solution for Atomic Absorption Spectrophotometry, dilute with water to contain 0.4 to 1.2  $\mu\text{g}$  of zinc (Zn: 65.41) per mL, and use this solution as the standard solution. Perform the test with the sample solution and standard solution according to the Atomic Absorption Spectrophotometry <2.23> under the following conditions, and determine the amount of zinc in the sample solution using the analytical curve obtained from

the absorbance of the standard solution.

Gas: Combustible gas—Acetylene gas

Supporting gas—Air

Lamp: Zinc hollow-cathode lamp

Wavelength: 213.9 nm


**Containers and storage** Containers—Hermetic containers.

Storage—In a cold place, and avoid freezing.

**Expiration date** 24 months after preparation.

## Iodamide

ヨードミド



$\text{C}_{12}\text{H}_{11}\text{I}_3\text{N}_2\text{O}_4$ : 627.94

3-Acetylaminomethyl-2,4,6-triiodobenzoic acid [440-58-4]

Iodamide, calculated on the dried basis, contains not less than 98.5% of  $\text{C}_{12}\text{H}_{11}\text{I}_3\text{N}_2\text{O}_4$ .

**Description** Iodamide occurs as a white, crystalline powder. It is odorless.

It is slightly soluble in water and in ethanol (95), and practically insoluble in diethyl ether.

It dissolves in sodium hydroxide TS and in sodium carbonate TS.

It gradually changes in color by light.

**Identification (1)** To 0.01 g of Iodamide add 5 mL of hydrochloric acid, and heat in a water bath for 5 minutes: the solution responds to the Qualitative Tests <1.09> for primary aromatic amines.

(2) Heat 0.1 g of Iodamide over a flame: a purple gas evolves.

(3) Determine the infrared absorption spectrum of Iodamide, previously dried, as directed in the potassium bromide disk method under Infrared Spectrophotometry <2.25>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wave numbers. If any difference appears between the spectra, dissolve 1 g of Iodamide in 100 mL of water by heating, and concentrate the solution to about 30 mL by gentle boiling. After cooling, collect the formed crystals by filtration, dry, and repeat the test on the dried crystals.

**Purity (1)** Clarity and color of solution—Dissolve 1.0 g of Iodamide in 10 mL of diluted sodium hydroxide TS (1 in 5): the solution is clear and colorless.

(2) Primary aromatic amines—Dissolve 0.20 g of Iodamide in 5 mL of water and 1 mL of sodium hydroxide TS, add 4 mL of a solution of sodium nitrite (1 in 100) and 10 mL of 1 mol/L hydrochloric acid TS, shake well, and allow to stand for 2 minutes. To this solution add 5 mL of ammonium amidosulfate TS, shake thoroughly, allow to stand for 1 minute, add 0.4 mL of a solution of 1-naphthol in ethanol



(95) (1 in 10), 15 mL of sodium hydroxide TS and water to make exactly 50 mL, and determine the absorbance at 485 nm as directed under Ultraviolet-visible Spectrophotometry <2.24>, using a solution, prepared in the same manner, as the blank: the absorbance of the solution is not more than 0.12.

(3) Soluble halide—Dissolve 2.5 g of Iodamide in 20 mL of water and 2.5 mL of ammonia TS, then add 20 mL of dilute nitric acid and water to make 100 mL. Allow to stand for 15 minutes with occasional shaking, and filter. Discard the first 10 mL of the filtrate, transfer 25 mL of the subsequent filtrate to a Nessler tube, and add ethanol (95) to make 50 mL. Use this solution as the test solution, and proceed as directed under Chloride Limit Test <1.03>. Prepare the control solution with 0.10 mL of 0.01 mol/L hydrochloric acid VS and 6 mL of dilute nitric acid, and dilute with water to 25 mL, then with ethanol (95) to 50 mL.

(4) Iodine—Dissolve 0.20 g of Iodamide in 2 mL of sodium hydroxide TS, add 2.5 mL of 0.5 mol/L sulfuric acid TS, allow to stand for 10 minutes with occasional shaking, then add 5 mL of chloroform, shake vigorously and allow to stand: the chloroform layer remains colorless.

(5) Heavy metals <1.07>—Proceed with 2.0 g of Iodamide according to Method 4, and perform the test. Prepare the control solution with 2.0 mL of Standard Lead Solution (not more than 10 ppm).

(6) Arsenic <1.11>—Prepare the test solution with 1.0 g of Iodamide according to Method 3, and perform the test (not more than 2 ppm).

**Loss on drying** <2.41> Not more than 3.0% (1 g, 105°C, 4 hours).

**Residue on ignition** <2.44> Not more than 0.1% (1 g).

**Assay** Weigh accurately about 0.5 g of Iodamide in a saponification flask, dissolve in 40 mL of sodium hydroxide TS, add 1 g of zinc powder, connect the flask with a reflux condenser, boil for 30 minutes, cool, and filter. Wash the flask and filter paper with 50 mL of water, and combine the washings with the filtrate. Add 5 mL of acetic acid (100), and titrate <2.50> with 0.1 mol/L silver nitrate VS until the color of the precipitate changes from yellow to green (indicator: 1 mL of tetrabromophenolphthalein ethyl ester TS).

Each mL of 0.1 mol/L silver nitrate VS  
= 20.93 mg of  $\text{C}_{12}\text{H}_{11}\text{I}_3\text{N}_2\text{O}_4$

**Containers and storage** Containers—Tight containers.

Storage—Light-resistant.

## Iodinated ( $^{131}\text{I}$ ) Human Serum Albumin Injection

ヨウ化人血清アルブミン ( $^{131}\text{I}$ ) 注射液

Iodinated ( $^{131}\text{I}$ ) Human Serum Albumin Injection is an aqueous solution for injection containing normal human serum albumin iodinated by iodine-131 ( $^{131}\text{I}$ ).

It conforms to the requirements of Iodinated ( $^{131}\text{I}$ ) Human Serum Albumin Injection in the Minimum Requirements for Radiopharmaceuticals.

Test for Extractable Volume of Parenteral Preparations and Insoluble Particulate Matter Test for Injec-

tions are not applied to this injection.

**Description** Iodinated ( $^{131}\text{I}$ ) Human Serum Albumin Injection is a clear, colorless or light yellow liquid.

## Iodine

ヨウ素

I: 126.90

Iodine contains not less than 99.5% of I.

**Description** Iodine occurs as grayish black plates or heavy, granular crystals, having a metallic luster and a characteristic odor.

It is freely soluble in diethyl ether, soluble in ethanol (95), sparingly soluble in chloroform, and very slightly soluble in water.

It dissolves in potassium iodide TS.

Iodine sublimes at room temperature.

**Identification** (1) A solution of Iodine in ethanol (95) (1 in 50) shows a red-brown color.

(2) A solution of Iodine in chloroform (1 in 1000) shows a red-purple to purple color.

(3) Add 0.5 mL of starch TS to 10 mL of a saturated solution of Iodine: a dark blue color is produced. When the mixture is boiled, the color disappears, and it reappears on cooling.

**Purity** (1) Non-volatile residue—Sublime 2.0 g of Iodine on a water bath, and dry the residue at 105°C for 1 hour: the mass of the residue is not more than 1.0 mg.

(2) Chloride or bromide—Mix 1.0 g of finely powdered Iodine with 20 mL of water, and filter the mixture. To 10 mL of the filtrate add dropwise diluted sulfurous acid solution (1 in 5) until the yellow color disappears. Add 1 mL of ammonia TS, followed by 1 mL of silver nitrate TS in small portions, and add water to make 20 mL. Shake well, filter, and after discarding the first 2 mL of the filtrate, take 10 mL of the subsequent filtrate. To the filtrate add 2.0 mL of nitric acid and water to make 20 mL: the solution so obtained has no more turbidity than the following control solution.

Control solution: To 0.20 mL of 0.01 mol/L hydrochloric acid VS add 5 mL of water, 2.5 mL of ammonia TS, 1 mL of silver nitrate TS, 2.0 mL of nitric acid and water to make 20 mL.

**Assay** Place 1 g of potassium iodide and 1 mL of water in a glass-stoppered flask, weigh accurately, add about 0.3 g of Iodine to the flask, and weigh accurately again. Dissolve the iodine by gentle shaking, add 20 mL of water and 1 mL of dilute hydrochloric acid, and titrate <2.50> with 0.1 mol/L sodium thiosulfate VS (indicator: 1 mL of starch TS).

Each mL of 0.1 mol/L sodium thiosulfate VS  
= 12.69 mg of I

**Containers and storage** Containers—Tight containers.

## Iodine Tincture

ヨードチンキ

Iodine Tincture contains not less than 5.7 w/v% and not more than 6.3 w/v% of iodine (I: 126.90), and not less than 3.8 w/v% and not more than 4.2 w/v% of potassium iodide (KI: 166.00).

### Method of preparation

|                  |                       |
|------------------|-----------------------|
| Iodine           | 60 g                  |
| Potassium Iodide | 40 g                  |
| 70 vol% Ethanol  | a sufficient quantity |
| To make 1000 mL  |                       |

Prepare as directed under Tinctures, with the above ingredients. It may be prepared with an appropriate quantity of Ethanol or Ethanol for Disinfectant and Purified Water in place of 70 vol% Ethanol.

**Description** Iodine Tincture is a dark red-brown liquid, and has a characteristic odor.

Specific gravity  $d_{20}^{20}$ : about 0.97

**Identification (1)** To a mixture of 1 mL of starch TS and 9 mL of water add 1 drop of Iodine Tincture: a dark blue-purple color develops.

**(2)** Evaporate 3 mL of Iodine Tincture to dryness on a water bath, and heat gently over a free flame: a white residue is formed which responds to the Qualitative Tests <1.09> for potassium salt and iodide.

**Alcohol number** <1.01> Not less than 6.6 (Method 2). Perform the pretreatment (ii) in the Method 1.

**Assay (1)** Iodine—Pipet 5 mL of Iodine Tincture, add 0.5 g of potassium iodide, 20 mL of water and 1 mL of dilute hydrochloric acid, and titrate <2.50> with 0.1 mol/L sodium thiosulfate VS (indicator: 2 mL of starch TS).

Each mL of 0.1 mol/L sodium thiosulfate VS  
= 12.69 mg of I

**(2)** Potassium iodide—Pipet 5 mL of Iodine Tincture into an iodine flask, add 20 mL of water, 50 mL of hydrochloric acid and 5 mL of chloroform. Cool to room temperature, and titrate <2.50> with 0.05 mol/L potassium iodate VS until the red-purple color disappears from the chloroform layer, with agitating the mixture vigorously and continuously. After the chloroform layer has been decolorized, allow the mixture to stand for 5 minutes. If the color reappears, the mixture should be titrated <2.50> further with 0.05 mol/L potassium iodate VS. Calculate the amount (mg) of potassium iodide from the number of mL ( $a$ ) of 0.05 mol/L potassium iodate VS used as above and the number of mL ( $b$ ) of 0.1 mol/L sodium thiosulfate VS used in the titration under the Assay (1).

Amount (mg) of potassium iodide (KI)  
=  $16.600 \times \{a - (b/2)\}$

**Containers and storage** Containers—Tight containers.

## Dilute Iodine Tincture

希ヨードチンキ

Dilute Iodine Tincture contains not less than 2.8 w/v% and not more than 3.2 w/v% of iodine (I: 126.90), and not less than 1.9 w/v% and not more than 2.1 w/v% of potassium iodide (KI: 166.00).

### Method of preparation

|                  |                       |
|------------------|-----------------------|
| Iodine           | 30 g                  |
| Potassium Iodide | 20 g                  |
| 70 vol% Ethanol  | a sufficient quantity |
| To make 1000 mL  |                       |

Prepare as directed under Medicated Spirits, with the above ingredients. It may be prepared with an appropriate quantity of Ethanol or Ethanol for Disinfection and Purified Water in place of 70 vol% Ethanol. It may also be prepared by adding 70 vol% Ethanol to 500 mL of Iodine Tincture to make 1000 mL.

**Description** Dilute Iodine Tincture is a dark red-brown liquid, and has a characteristic odor.

Specific gravity  $d_{20}^{20}$ : about 0.93

**Identification (1)** To a mixture of 1 mL of starch TS and 9 mL of water add 1 drop of Dilute Iodine Tincture: a dark blue-purple color develops.

**(2)** Evaporate 3 mL of Diluted Iodine Tincture to dryness on a water bath, and heat gently over a free flame: a white residue is formed which responds to the Qualitative Tests <1.09> for potassium salt and iodide.

**Alcohol number** <1.01> Not less than 6.7 (Method 2). Perform the pretreatment (ii) in the Method 1.

**Assay (1)** Iodine—Pipet exactly 10 mL of Dilute Iodine Tincture, add 0.5 g of potassium iodide, 20 mL of water and 1 mL of dilute hydrochloric acid, and titrate <2.50> with 0.1 mol/L sodium thiosulfate VS (indicator: 2 mL of starch TS).

Each mL of 0.1 mol/L sodium thiosulfate VS  
= 12.69 mg of I

**(2)** Potassium iodide—Pipet exactly 10 mL of Dilute Iodine Tincture into an iodine flask, add 20 mL of water, 50 mL of hydrochloric acid and 5 mL of chloroform. Cool to room temperature, and titrate <2.50> with 0.05 mol/L potassium iodate VS until the red-purple color in the chloroform layer disappears while agitating vigorously and continuously. After the chloroform layer has been decolorized, allow the mixture to stand for 5 minutes. If the color reappears, the mixture should be titrated <2.50> further with 0.05 mol/L potassium iodate VS. Calculate the amount (mg) of potassium iodide from the volume ( $a$  mL) of 0.05 mol/L potassium iodate VS consumed as above and the volume ( $b$  mL) of 0.1 mol/L sodium thiosulfate VS consumed in the titration under Assay (1).

Amount (mg) of potassium iodide (KI)  
=  $16.600 \times \{a - (b/2)\}$

**Containers and storage** Containers—Tight containers.

## Compound Iodine Glycerin

複方ヨード・グリセリン

Compound Iodine Glycerin contains not less than 1.1 w/v% and not more than 1.3 w/v% of iodine (I: 126.90), not less than 2.2 w/v% and not more than 2.6 w/v% of potassium iodide (KI: 166.00), not less than 2.7 w/v% and not more than 3.3 w/v% of total iodine (as I), and not less than 0.43 w/v% and not more than 0.53 w/v% of phenol (C<sub>6</sub>H<sub>6</sub>O: 94.11).

### Method of preparation

|                  |                       |
|------------------|-----------------------|
| Iodine           | 12 g                  |
| Potassium Iodide | 24 g                  |
| Glycerin         | 900 mL                |
| Mentha Water     | 45 mL                 |
| Liquefied Phenol | 5 mL                  |
| Purified Water   | a sufficient quantity |
| To make 1000 mL  |                       |

Dissolve Potassium Iodide and Iodine in about 25 mL of Purified Water. After adding Glycerin, add Mentha Water, Liquefied Phenol and sufficient Purified Water to make 1000 mL, mixing thoroughly. It may be prepared with an appropriate quantity of Concentrated Glycerin and Purified Water in place of Glycerin.

**Description** Compound Iodine Glycerin is a red-brown, viscous liquid. It has a characteristic odor.

Specific gravity  $d_{20}^{20}$ : about 1.23

**Identification (1)** The colored solution obtained in the Assay (1) acquires a red color. Determine the absorption spectrum of this solution as directed under Ultraviolet-visible Spectrophotometry <2.24>: it exhibits a maximum between 510 nm and 514 nm (iodine).

**(2)** The colored solution obtained in the Assay (2) acquires a red color. Determine the absorption spectrum of this solution as directed under Ultraviolet-visible Spectrophotometry <2.24>: it exhibits a maximum between 510 nm and 514 nm (potassium iodide).

**(3)** The colored solution obtained in the Assay (4) has a yellow color. Determine the absorption spectrum of this solution as directed under Ultraviolet-visible Spectrophotometry <2.24>: it exhibits a maximum between 401 nm and 405 nm (phenol).

**(4)** Take 1 mL of Compound Iodine Glycerin in a glass-stoppered test tube, add 10 mL of ethanol (95), and mix. Then add 2 mL of sodium hydroxide TS, add 1 mL of a solution of copper (II) chloride dihydrate in ethanol (95) (1 in 10), and shake: a blue color develops (glycerin).

**Assay (1)** Iodine—Measure the specific gravity <2.56> of Compound Iodine Glycerin according to Method 2. Weigh exactly about 7 mL of it, add ethanol (95) to make exactly 200 mL, and use this solution as the sample solution. On the other hand, weigh accurately about 80 mg of iodine for assay and about 0.17 g of potassium iodide for assay, previously dried at 105°C for 4 hours, dissolve in ethanol (95) to make exactly 200 mL, and use this solution as the standard solution.

Pipet 3 mL each of the sample solution and the standard solution into 50-mL separators, to each add exactly 10 mL of a mixture of chloroform and hexane (2:1) and 15 mL of water successively, and shake immediately and vigorously. Separate the chloroform-hexane layers [use the water layers in (2)], and filter through a pledget of cotton. Determine the absorbances of the filtrates,  $A_T$  and  $A_S$ , at 512 nm as directed under Ultraviolet-visible Spectrophotometry <2.24>, using a mixture of chloroform and hexane (2:1) as the blank.

$$\text{Amount (mg) of iodine (I)} = W_S \times (A_T/A_S)$$

$W_S$ : Amount (mg) of iodine for assay

**(2)** Potassium iodide—Separate the water layers of the sample solution and the standard solution obtained in (1), pipet 10 mL of each of the water layers, and to each add 1 mL of diluted dilute hydrochloric acid (1 in 2), 1 mL of sodium nitrite TS and exactly 10 mL of a mixture of chloroform and hexane (2:1). Shake immediately and vigorously, separate the chloroform-hexane layers, and filter through a pledget of cotton. Determine the absorbances,  $A_T$  and  $A_S$ , of both solutions at 512 nm as directed under Ultraviolet-visible Spectrophotometry <2.24>, using a mixture of chloroform and hexane (2:1) as the blank.

$$\text{Amount (mg) of potassium iodide (KI)} = W_S \times (A_T/A_S)$$

$W_S$ : Amount (mg) of potassium iodide for assay

**(3)** Total iodine—Measure the specific gravity <2.56> of Compound Iodine Glycerin according to Method 2. Weigh exactly about 5 mL of it, and add water to make exactly 50 mL. Pipet 5 mL of this solution into a 50-mL flask, and add 0.5 g of zinc powder and 5 mL of acetic acid (100). Shake until the color of iodine disappears, and heat under a reflux condenser on a water bath for 30 minutes. Wash the condenser with 10 mL of hot water, and filter through a glass filter (G3). Wash the flask with two 10-mL portions of warm water, and combine the filtrate and the washings. After cooling, add water to make exactly 50 mL, and use this solution as the sample solution. On the other hand, dissolve about 0.2 g of potassium iodide for assay, previously dried at 105°C for 4 hours and accurately weighed, in water to make exactly 50 mL. Pipet 5 mL of this solution, add 5 mL of acetic acid (100) and water to make exactly 50 mL, and use this solution as the standard solution. Pipet 4 mL each of the sample solution and standard solution into 30-mL separators, and to each add 5 mL of water, 1 mL of diluted dilute hydrochloric acid (1 in 2), 1 mL of sodium nitrite TS and 10 mL of a mixture of chloroform and hexane (2:1). Shake well immediately, and proceed as directed in (2).

$$\text{Amount (mg) of total iodine (I)} = W_S \times (A_T/A_S) \times 0.7644$$

$W_S$ : Amount (mg) of potassium iodide for assay

**(4)** Phenol—Measure the specific gravity <2.56> of Compound Iodine Glycerin according to Method 2. Weigh exactly about 2 mL of it, add 3 mL of 0.1 mol/L sodium thiosulfate VS, and shake. Add 2 mL of dilute hydrochloric acid, and shake with two 10-mL portions of chloroform. Separate the chloroform layer, and shake with two 10-mL portions of 0.5 mol/L sodium hydroxide TS. Separate the water layer, add water to make exactly 500 mL, and use this solution as the sample solution. Dissolve about 0.5 g of phenol for assay, accurately weighed, in ethanol (95) to make exactly 100 mL,

pipet 2 mL of this solution, proceed in the same manner as the sample solution, and use so obtained solution as the standard solution. Pipet 3 mL each of the sample solution and standard solution, to each add 2 mL of dilute hydrochloric acid, and place in a water bath at 30°C. Allow to stand for 10 minutes, and add exactly 2 mL of a solution of sodium nitrite (1 in 100), shake, and allow to stand at 30°C for 60 minutes. Add dilute potassium hydroxide-ethanol TS to make exactly 25 mL, and determine the absorbances of these solutions,  $A_T$  and  $A_S$ , at 403 nm as directed under Ultraviolet-visible Spectrophotometry <2.24>, using the solution prepared in the same manner with 3 mL of water instead of the sample solution as the blank.

$$\begin{aligned} &\text{Amount (mg) of phenol (C}_6\text{H}_6\text{O)} \\ &= W_S \times (A_T/A_S) \times (1/50) \end{aligned}$$

$W_S$ : Amount (mg) of phenol for assay

**Containers and storage** Containers—Tight containers.  
Storage—Light-resistant.

## Dental Iodine Glycerin

歯科用ヨード・グリセリン

Dental Iodine Glycerin contains not less than 9.0 w/v% and not more than 11.0 w/v% of iodine (I: 126.90), not less than 7.2 w/v% and not more than 8.8 w/v% of potassium iodide (KI: 166.00), and not less than 0.9 w/v% and not more than 1.1 w/v% of zinc sulfate hydrate ( $\text{ZnSO}_4 \cdot 7\text{H}_2\text{O}$ : 287.58).

### Method of preparation

|                      |                       |
|----------------------|-----------------------|
| Iodine               | 10 g                  |
| Potassium Iodide     | 8 g                   |
| Zinc Sulfate Hydrate | 1 g                   |
| Glycerin             | 35 mL                 |
| Purified Water       | a sufficient quantity |
| To make 100 mL       |                       |

Dissolve and mix the above ingredients.

**Description** Dental Iodine Glycerin is a dark red-brown liquid, having the odor of iodine.

**Identification** (1) The colored solution obtained in the Assay (1) acquires a red color. Determine the absorption spectrum of this solution as directed under Ultraviolet-visible Spectrophotometry <2.24>: it exhibits a maximum between 510 nm and 514 nm (iodine).

(2) The colored solution obtained in the Assay (2) acquires a red color. Determine the absorption spectrum of this solution as directed under Ultraviolet-visible Spectrophotometry <2.24>: it exhibits a maximum between 510 nm and 514 nm (potassium iodide).

(3) Put 1 mL of Dental Iodine Glycerin in a glass-stoppered, test tube, add 10 mL of ethanol (95), and mix. Then add 2 mL of sodium hydroxide TS, add 1 mL of a solution of copper (II) chloride dihydrate in ethanol (95) (1 in 10), and shake: a blue color develops (glycerin).

(4) The colored solution obtained in the Assay (3) acquires a red-purple to purple color. Determine the absorption

spectrum of this solution as directed under Ultraviolet-visible Spectrophotometry <2.24>: it exhibits a maximum between 618 nm and 622 nm (zinc sulfate hydrate).

**Assay** (1) Iodine—Pipet 5 mL of Dental Iodine Glycerin, and add diluted ethanol (3 in 10) to make exactly 50 mL. Pipet 5 mL of this solution, add water to make exactly 200 mL, and use this solution as the sample solution. On the other hand, weigh accurately about 0.5 g of iodine for assay and about 0.4 g of potassium iodide for assay, previously dried at 105°C for 4 hours, and dissolve in diluted ethanol (3 in 10) to make exactly 50 mL. Pipet 5 mL of this solution, add water to make exactly 200 mL, and use this solution as the standard solution. Pipet 10 mL each of the sample solution and standard solution, to each add exactly 20 mL of a mixture of chloroform and hexane (2:1), shake immediately, and separate the chloroform-hexane layer [use the water layer in (2)]. Filter through a pledget of cotton. Determine the absorbances,  $A_T$  and  $A_S$ , of the filtrates obtained from the sample solution and standard solution, respectively, at 512 nm as directed under Ultraviolet-visible Spectrophotometry <2.24>, using a mixture of chloroform and hexane (2:1) as the blank.

$$\text{Amount (mg) of iodine (I)} = W_S \times (A_T/A_S)$$

$W_S$ : Amount (mg) of iodine for assay

(2) Potassium iodide—Separate the water layers of the sample solution and standard solution obtained in (1), pipet 7 mL each of the water layers, and to each add exactly 1 mL of diluted hydrochloric acid (1 in 2), 1 mL of sodium nitrite TS and 10 mL of a mixture of chloroform and hexane (2:1), and shake immediately. Separate the chloroform-hexane layer, and filter through a pledget of cotton. Determine the absorbances,  $A_T$  and  $A_S$ , of the filtrates obtained from the sample solution and standard solution, respectively, at 512 nm as directed under Ultraviolet-visible Spectrophotometry, using a mixture of chloroform and hexane (2:1) as the blank.

$$\text{Amount (mg) of potassium iodide (KI)} = W_S \times (A_T/A_S)$$

$W_S$ : Amount (mg) of potassium iodide for assay

(3) Zinc sulfate Hydrate—Pipet 5 mL of Dental Iodine Glycerin, and add diluted ethanol (3 in 10) to make exactly 50 mL. Pipet 5 mL of this solution, add water to make exactly 100 mL, and use this solution as the sample solution. On the other hand, pipet 10 mL of Standard Zinc Stock Solution, add diluted ethanol (3 in 200) to make exactly 1000 mL, and use this solution as the standard solution. Pipet 10 mL each of the sample solution and standard solution, to each add 10 mL of a mixture of chloroform and hexane (2:1), shake, and allow to stand. Pipet 3 mL each of the water layers, and to each add 2 mL of boric acid-potassium chloride-sodium hydroxide buffer solution, pH 10.0, 2 mL of zincon TS and water to make exactly 25 mL. Determine the absorbances,  $A_T$  and  $A_S$ , obtained from the sample solution and standard solution, respectively, at 620 nm as directed under Ultraviolet-visible Spectrophotometry <2.24>, using the solution prepared in the same manner with 3 mL of water as the blank.

$$\begin{aligned} &\text{Amount (mg) of zinc sulfate Hydrate (ZnSO}_4 \cdot 7\text{H}_2\text{O)} \\ &= W_S \times (A_T/A_S) \times 4.397 \end{aligned}$$

$W_S$ : Amount (mg) of zinc in 10 mL of Standard Zinc Stock Solution

**Containers and storage** Containers—Tight containers.

Storage—Light-resistant.

## Iodine, Salicylic Acid and Phenol Spirit

コード・サリチル酸・フェノール精

Iodine, Salicylic Acid and Phenol Spirit contains not less than 1.08 w/v% and not more than 1.32 w/v% of iodine (I: 126.90), not less than 0.72 w/v% and not more than 0.88 w/v% of potassium iodide (KI: 166.00), not less than 4.5 w/v% and not more than 5.5 w/v% of salicylic acid ( $C_7H_6O_3$ : 138.12), not less than 1.8 w/v% and not more than 2.2 w/v% of phenol ( $C_6H_6O$ : 94.11), and not less than 7.2 w/v% and not more than 8.8 w/v% of benzoic acid ( $C_7H_6O_2$ : 122.12).

### Method of preparation

|                          |                       |
|--------------------------|-----------------------|
| Iodine Tincture          | 200 mL                |
| Salicylic Acid           | 50 g                  |
| Phenol                   | 20 g                  |
| Benzoid Acid             | 80 g                  |
| Ethanol for Disinfection | a sufficient quantity |
| To make 1000 mL          |                       |

Prepare as directed under Medicated Spirits, with the above ingredients. It may be prepared with an appropriate quantity of Ethanol and Purified Water in place of Ethanol for Disinfection.

**Description** Iodine, Salicylic Acid and Phenol Spirit is a dark red-brown liquid, having the odor of phenol.

**Identification (1)** To a mixture of 1 mL of starch TS and 9 mL of water add 1 drop of Iodine, Salicylic Acid and Phenol Spirit: a dark blue-purple color develops (iodine).

**(2)** To 1 mL of Iodine, Salicylic Acid and Phenol Spirit add 5 mL of ethanol (95) and water to make 50 mL. To 1 mL of this solution add hydrochloric acid-potassium chloride buffer solution, pH 2.0, to make 50 mL, and to 15 mL of this solution add 5 mL of a solution of iron (III) nitrate enneahydrate (1 in 200): a red-purple color is produced (salicylic acid).

**(3)** Shake 1 mL of Iodine, Salicylic Acid and Phenol Spirit with 1 mL of sodium thiosulfate TS, add 20 mL of water and 5 mL of dilute hydrochloric acid, and extract with 25 mL of diethyl ether. Wash the diethyl ether extract with two 25-mL portions of sodium hydrogen carbonate TS, and extract with 10 mL of dilute sodium hydroxide TS. Shake 1 mL of the extract with 1 mL of sodium nitrite TS and 1 mL of dilute hydrochloric acid, and add 3 mL of sodium hydroxide TS: a yellow color is developed (phenol).

**(4)** Shake 1 mL of Iodine, Salicylic Acid and Phenol Spirit with 1 mL of sodium thiosulfate TS, add 20 mL of water and 5 mL of dilute hydrochloric acid, extract with 10 mL of diethyl ether, and use the diethyl ether extract as the sample solution. Dissolve 25 mg of salicylic acid, 0.01 g of phenol and 0.04 g of benzoic acid in 5 mL each of diethyl ether, respectively, and use these solutions as the standard solutions (1), (2) and (3). Perform the test with the sample solution and standard solutions (1), (2) and (3) as directed under

Thin-layer Chromatography <2.03>. Spot 5  $\mu$ L of each solution on a plate of silica gel with fluorescent indicator for thin-layer chromatography. Develop the plate with a mixture of chloroform, acetone and acetic acid (100) (45:5:1) to a distance of about 10 cm, and air-dry the plate. Examine under ultraviolet light (main wavelength: 254 nm): the 3 spots from the sample solution show the same  $R_f$  value as the corresponding spots of the standard solutions (1), (2) and (3). Spray evenly iron (III) chloride TS on the plate: the spot from standard solution (1) and the corresponding spot from the sample solution acquires a purple color.

**Assay (1)** Iodine—Pipet 4 mL of Iodine, Salicylic Acid and Phenol Spirit, add ethanol (95) to make exactly 50 mL, and use this solution as the sample solution. On the other hand, weigh accurately about 1.2 g of iodine for assay and about 0.8 g of potassium iodide for assay, previously dried at 105°C for 4 hours, and dissolve in ethanol (95) to make exactly 100 mL. Pipet 4 mL of this solution, add ethanol (95) to make exactly 50 mL, and use this solution as the standard solution. Pipet 3 mL each of the sample solution and standard solution, to each add exactly 25 mL of a mixture of chloroform and hexane (2:1), and shake. Further add exactly 10 mL of water, shake and separate the chloroform-hexane layers [use the water layers in (2)]. Filter through a pledget of absorbent cotton, and determine the absorbances of the filtrates from the sample solution and standard solution, respectively,  $A_T$  and  $A_S$ , at 512 nm as directed under Ultraviolet-visible Spectrophotometry <2.24>, using a mixture of chloroform and hexane (2:1) as the blank.

$$\text{Amount (mg) of iodine (I)} = W_S \times (A_T/A_S) \times (1/25)$$

$W_S$ : Amount (mg) of iodine for assay

**(2)** Potassium iodide—Separate the water layers of the sample solution and standard solution obtained in the Assay (1), pipet 8 mL each of the water layers, and add 1 mL of diluted dilute hydrochloric acid (1 in 2) and 1 mL of sodium nitrite TS. Immediately after shaking, add exactly 10 mL of a mixture of chloroform and hexane (2:1), shake, and proceed in the same manner as for the Assay (1).

$$\begin{aligned} \text{Amount (mg) of potassium iodide (KI)} \\ = W_S \times (A_T/A_S) \times (1/25) \end{aligned}$$

$W_S$ : Amount (mg) of potassium iodide for assay

**(3)** Salicylic acid, phenol and benzoic acid—Pipet 2 mL of Iodine, Salicylic Acid and Phenol Spirit, add 20 mL of diluted methanol (1 in 2) and 0.1 mol/L sodium thiosulfate VS until the color of iodine disappears, add exactly 20 mL of the internal standard solution, then add diluted methanol (1 in 2) to make 200 mL, and use this solution as the sample solution. Weigh accurately about 0.2 g of salicylic acid for assay, previously dried in a desiccator (silica gel) for 3 hours, about 80 mg of phenol for assay, and 0.32 g of benzoic acid, previously dried in a desiccator (silica gel) for 3 hours, dissolve in diluted methanol (1 in 2) to make exactly 50 mL. Pipet 25 mL of this solution, add exactly 20 mL of the internal standard solution and diluted methanol (1 in 2) to make 200 mL, and use this solution as the standard solution. Perform the test with 3  $\mu$ L of the sample solution and standard solution as directed under Liquid Chromatography <2.01> according to the operating conditions in the Assay under Compound Salicylic Acid Spirit. Calculate the ratios,  $Q_{Ta}$ ,  $Q_{Tb}$  and  $Q_{Tc}$ , of

the peak areas of salicylic acid, phenol and benzoic acid to those of the internal standard of the sample solution, and the ratios,  $Q_{Sa}$ ,  $Q_{Sb}$  and  $Q_{Sc}$ , of the peak areas of salicylic acid, phenol and benzoic acid to those of the internal standard of the standard solution.

$$\begin{aligned} \text{Amount (mg) of salicylic acid (C}_7\text{H}_6\text{O}_3\text{)} \\ = W_{Sa} \times (Q_{Ta}/Q_{Sa}) \times (1/2) \end{aligned}$$

$$\begin{aligned} \text{Amount (mg) of phenol (C}_6\text{H}_6\text{O)} \\ = W_{Sb} \times (Q_{Tb}/Q_{Sb}) \times (1/2) \end{aligned}$$

$$\begin{aligned} \text{Amount (mg) of benzoic acid (C}_7\text{H}_6\text{O}_2\text{)} \\ = W_{Sc} \times (Q_{Tc}/Q_{Sc}) \times (1/2) \end{aligned}$$

$W_{Sa}$ : Amount (mg) of salicylic acid for assay

$W_{Sb}$ : Amount (mg) of phenol for assay

$W_{Sc}$ : Amount (mg) of benzoic acid

**Internal standard solution**—A solution of theophylline in methanol (1 in 1000).

**Containers and storage** Containers—Tight containers.

Storage—Light-resistant.

## Iodoform

ヨードホルム



$\text{CHI}_3$ : 393.73

Triiodomethane [75-47-8]

Iodoform, when dried, contains not less than 99.0% of  $\text{CHI}_3$ .

**Description** Iodoform occurs as lustrous, yellow crystals or crystalline powder. It has a characteristic odor.

It is freely soluble in diethyl ether, sparingly soluble in ethanol (95), and practically insoluble in water.

It is slightly volatile at ordinary temperature.

Melting point: about 120°C (with decomposition).

**Identification** Heat 0.1 g of Iodoform: a purple gas is evolved.

**Purity (1)** Water-soluble colored substances and acidity or alkalinity—Shake well 2.0 g of Iodoform, previously powdered, with 5 mL of water for 1 minute, allow to stand, and filter the supernatant liquid: the filtrate is colorless and neutral.

**(2) Chloride** <1.03>—Shake well 3.0 g of Iodoform, previously powdered, with 75 mL of water for 1 minute, allow to stand, and filter the supernatant liquid. To 25 mL of the filtrate add 6 mL of dilute nitric acid and water to make 50 mL, and perform the test using this solution as the test solution. Prepare the control solution with 0.30 mL of 0.01 mol/L hydrochloric acid VS (not more than 0.011%).

**(3) Sulfate** <1.14>—To 25 mL of the filtrate obtained in (2) add 1 mL of dilute hydrochloric acid and water to make 50 mL, and perform the test using this solution as the test solution. Prepare the control solution with 0.35 mL of 0.005 mol/L sulfuric acid VS (not more than 0.017%).

**Loss on drying** <2.41> Not more than 0.5% (1 g, silica gel,

24 hours).

**Residue on ignition** <2.44> Not more than 0.1% (1 g).

**Assay** Weigh accurately about 0.2 g of Iodoform, previously dried, in a 500-mL glass-stoppered flask, and dissolve it in 20 mL of ethanol (95). Add exactly 30 mL of 0.1 mol/L silver nitrate VS and 10 mL of nitric acid, stopper the flask, shake well, and allow to stand in a dark place over 16 hours. Add 150 mL of water, and titrate <2.50> the excess silver nitrate with 0.1 mol/L ammonium thiocyanate VS (indicator: 5 mL of ammonium iron (III) sulfate TS). Perform a blank determination.

Each mL of 0.1 mol/L silver nitrate VS


= 13.12 mg of  $\text{CHI}_3$

**Containers and storage** Containers—Tight containers.

Storage—Light-resistant.

## Iopamidol

イオパミドール



$\text{C}_{17}\text{H}_{22}\text{I}_3\text{N}_3\text{O}_8$ : 777.09

*N,N'*-Bis[2-hydroxy-1-(hydroxymethyl)ethyl]-5-[(2*S*)-2-hydroxypropanoylamino]-2,4,6-triiodoisophthalamide [62883-00-5]

Iopamidol, when dried, contains not less than 99.0% of  $\text{C}_{17}\text{H}_{22}\text{I}_3\text{N}_3\text{O}_8$ .

**Description** Iopamidol occurs as a white crystalline powder.

It is very soluble in water, sparingly soluble in methanol, and very slightly soluble in ethanol (99.5).

**Identification (1)** To 0.05 g of Iopamidol add 5 mL of hydrochloric acid, heat for 10 minutes in a water bath: the test solution responds to the Qualitative Tests <1.09> for primary aromatic amines.

**(2)** Heat 0.1 g of Iopamidol over a flame: a purple gas is evolved.

**(3)** Determine the infrared absorption spectrum of Iopamidol, previously dried, as directed in the potassium bromide disk method under Infrared Spectrophotometry <2.25>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wave numbers.

**Optical rotation** <2.49>  $[\alpha]_{436}^{20}$ :  $-4.6 - -5.2^\circ$  (after drying, 4 g, water, warm, after cooling, 10 mL, 100 mm).

**Purity (1)** Clarity and color of solution—Dissolve 1.0 g of Iopamidol in 10 mL of water: the solution is clear and colorless.

**(2)** Primary aromatic amines—Dissolve 0.60 g of

Iopamidol in 8 mL of water, add 1 mL of a solution of sodium nitrite (1 in 50) and 12 mL of 2 mol/L hydrochloric acid TS, shake, and allow to stand for 2 minutes. Add 1 mL of a solution of ammonium amidosulfate (1 in 10), shake well, allow to stand for 1 minute, and add 1 mL of naphthylethylenediamine TS and water to make exactly 50 mL. Determine the absorbance of this solution at 495 nm as directed under Ultraviolet-visible Spectrophotometry <2.24> using a solution, prepared in the same manner, as the blank: the absorbance is not more than 0.12 (not more than 0.020%).

(3) Iodine—Dissolve 2.0 g of Iopamidol in 25 mL of water, add 5 mL of 1 mol/L sulfuric acid and 5 mL of toluene, shake well, and allow to stand: the toluene layer is colorless.

(4) Free iodine ion—Weigh accurately about 5.0 g of Iopamidol, dissolve in 70 mL of water, and adjust the pH to about 4.5 with dilute acetic acid. To this solution add 2 mL of 0.1 mol/L sodium chloride TS, and titrate <2.50> with 0.001 mol/L silver nitrate VS (potentiometric titration).

Each mL of 0.001 mol/L silver nitrate VS  
= 0.1269 mg of I

Content of iodine ion in Iopamidol is not more than 0.001%.

(5) Heavy metals <1.07>—Moisten 1.0 g of Iopamidol with a small quantity of sulfuric acid, heat gradually to almost incinerate by a possibly lower temperature. After cooling, moisten again with a small quantity of sulfuric acid, heat gradually until white fumes no longer are evolved, and incinerate by ignition between 450 to 550°C. Proceed as directed in Method 2, and perform the test. Prepare the control solution with 1.0 mL of Standard Lead Solution (not more than 10 ppm).

(6) Related substances—Dissolve 0.10 g of Iopamidol in water to make exactly 10 mL, and use this solution as the sample solution. Separately, dissolve 10 mg of *N,N'*-bis[2-hydroxy-1-(hydroxymethyl)ethyl]-5-hydroxyacetylamin-2,4,6-triiodoisophthalamide in water to make exactly 100 mL. Pipet 5 mL of this solution, add water to make exactly 50 mL, and use this solution as the standard solution. Perform the test with exactly 20 µL each of the sample solution and standard solution as directed under Liquid Chromatography <2.01> according to the following conditions, and determine each peak area of the both solutions by the automatic integration method: each area of the peaks other than the peak of iopamidol from the sample solution is not larger than the peak area of the standard solution, and the total of these areas is not larger than 2.5 times of the peak area of the standard solution.

#### Operating conditions—

Detector: An ultraviolet absorption photometer (wavelength: 240 nm).

Column: A stainless steel column 4.6 mm in inside diameter and 25 cm in length, packed with octadecylsilanized silica gel for liquid chromatography (5 µm in particle diameter).

Column temperature: A constant temperature of about 35°C.

Mobile phase: Use water as the mobile phase A, and a mixture of water and methanol (3:1) as the mobile phase B. Change the mixed ratios of the mobile phase A and the mobile phase B stepwise as follows:

| Time after injection<br>of sample (min) | Mobile phase<br>A (vol%) | Mobile phase<br>B (vol%) |
|---|--------------------------|--------------------------|
| 0 – 6                                   | 92                       | 8                        |
| 6 – 18                                  | 92 → 65                  | 8 → 35                   |
| 18 – 30                                 | 65 → 8                   | 35 → 92                  |
| 30 – 34                                 | 8                        | 92                       |

Flow rate: Adjust the flow rate to 1.5 mL per minute.

Time span of measurement: About 4.3 times as long as the retention time of iopamidol.

#### System suitability—

System performance: Dissolve 1 mL of the sample solution and 10 mg of *N,N'*-bis[2-hydroxy-1-(hydroxymethyl)ethyl]-5-hydroxyacetylamin-2,4,6-triiodoisophthalamide in water to make 100 mL. When the procedure is run with 20 µL of this solution under the above operating conditions, *N,N'*-bis[2-hydroxy-1-(hydroxymethyl)ethyl]-5-hydroxyacetylamin-2,4,6-triiodoisophthalamide and iopamidol are eluted in this order with the resolution between these peaks being not less than 7.

System repeatability: When the test is repeated 6 times with 20 µL of the standard solution under the above operating conditions, the relative standard deviation of the peak areas of *N,N'*-bis[2-hydroxy-1-(hydroxymethyl)ethyl]-5-hydroxyacetylamin-2,4,6-triiodoisophthalamide is not more than 1.0%.

**Loss on drying** <2.41> Not more than 0.30% (1 g, 105°C, 3 hours).

**Residue on ignition** <2.44> Not more than 0.1% (1 g).

**Assay** Weigh accurately about 0.5 g of Iopamidol, previously dried, transfer to a saponification flask, dissolve in 40 mL of sodium hydroxide TS, add 1 g of zinc powder, boil for 30 minutes under a reflux condenser, cool, and filter. Wash the flask and the filter paper with 50 mL of water, and combine the washing with the filtrate. Add 5 mL of acetic acid (100) to this solution, and titrate <2.50> with 0.1 mol/L silver nitrate VS (potentiometric titration).


Each mL of 0.1 mol/L silver nitrate VS  
= 25.90 mg of C<sub>17</sub>H<sub>22</sub>I<sub>3</sub>N<sub>3</sub>O<sub>8</sub>

**Containers and storage** Containers—Well-closed containers.

Storage—Light-resistant.

## Iotalamic Acid

イオタラム酸



C<sub>11</sub>H<sub>9</sub>I<sub>3</sub>N<sub>2</sub>O<sub>4</sub>: 613.91  
3-Acetylamin-2,4,6-triiodo-5-(methylaminocarbonyl)benzoic acid [2276-90-6]

Iotalamic Acid, when dried, contains not less than 99.0% of  $C_{11}H_9I_3N_2O_4$ .

**Description** Iotalamic Acid occurs as a white powder. It is odorless.

It is sparingly soluble in ethanol (95), very slightly soluble in water, and practically insoluble in diethyl ether.

It dissolves in sodium hydroxide TS.

It gradually colored by light.

**Identification (1)** Heat 0.1 g of Iotalamic Acid over a flame: a purple gas is evolved.

(2) Determine the infrared spectrum of Iotalamic Acid, previously dried, as directed in the potassium bromide disk method under Infrared Spectrophotometry <2.25>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wave numbers.

**Purity (1)** Clarity and color of solution—Dissolve 2.0 g of Iotalamic Acid in 10 mL of sodium hydroxide TS: the solution is clear and colorless.

(2) Primary aromatic amines—To 0.50 g of Iotalamic Acid add 15 mL of water, and dissolve it in 1 mL of sodium hydroxide TS while ice-cooling. Add 4 mL of a solution of sodium nitrite (1 in 100) to the solution, immediately add 12 mL of 1 mol/L hydrochloric acid TS, and shake gently. Then allow the mixture to stand for exactly 2 minutes, add 8 mL of ammonium amidosulfate TS, and shake occasionally for 5 minutes. Add 3 drops of a solution of 1-naphthol in ethanol (95) (1 in 10), allow to stand for 1 minute, add 3.5 mL of ammonia-ammonium chloride buffer solution, pH 10.7, mix, and immediately add water to make 50 mL. Determine within 20 minutes the absorbance of this solution at 485 nm as directed under Ultraviolet-visible Spectrophotometry <2.24>, using a solution, prepared in the same manner, as the blank: the absorbance is not more than 0.25.

(3) Soluble halides—Dissolve 0.5 g of Iotalamic Acid in 20 mL of diluted ammonia TS (1 in 40), add 6 mL of dilute nitric acid, shake, allow to stand for 5 minutes, and filter. Transfer the filtrate to a Nessler tube, wash the residue with 20 mL of water, combine the filtrate and the washings, and add water to make 50 mL. Proceed as directed for the Chloride Limit Test <1.03> using this solution as the test solution. Prepare the control solution as follows: to 0.10 mL of 0.01 mol/L hydrochloric acid VS and add 20 mL of diluted ammonia TS (1 in 40), 6 mL of dilute nitric acid and water to make 50 mL.

(4) Iodine—Dissolve 0.20 g of Iotalamic Acid in 2.0 mL of sodium hydroxide TS, add 2.5 mL of 0.5 mol/L sulfuric acid TS, and allow to stand for 10 minutes with occasional shaking. Add 5 mL of chloroform, shake well, and allow to stand: the chloroform layer remains colorless.

(5) Heavy metals <1.07>—Proceed with 1.0 g of Iotalamic Acid according to Method 2, and perform the test. Prepare the control solution with 2.0 mL of Standard Lead Solution (not more than 20 ppm).

(6) Arsenic <1.11>—Prepare the test solution with 0.6 g of Iotalamic Acid according to Method 3, and perform the test (not more than 3.3 ppm).

**Loss on drying** <2.41> Not more than 0.5% (1 g, 105°C, 4 hours).

**Residue on ignition** <2.44> Not more than 0.1% (1 g).


**Assay** Weigh accurately about 0.4 g of Iotalamic Acid, previously dried, place it in a saponification flask, dissolve in 40 mL of sodium hydroxide TS, add 1 g of zinc powder, and heat for 30 minutes under a reflux condenser. Cool, filter, wash the flask and the filter paper with 50 mL of water, and combine the washings and the filtrate. Add 5 mL of acetic acid (100) to this solution, and titrate <2.50> with 0.1 mol/L silver nitrate VS, until the color of the precipitate changes from yellow to green (indicator: 1 mL of tetrabromophenolphthalein ethyl ester TS).

Each mL of 0.1 mol/L silver nitrate VS  
= 20.46 mg of  $C_{11}H_9I_3N_2O_4$

**Containers and storage** Containers—Tight containers.  
Storage—Light-resistant.

## Iotroxic Acid

イオトロクス酸



$C_{22}H_{18}I_6N_2O_9$ : 1215.81

3,3'-(3,6,9-Trioxadecanedioyl)diiminobis-(2,4,6-triiodobenzoic acid) [51022-74-3]

Iotroxic Acid contains not less than 98.5% of  $C_{22}H_{18}I_6N_2O_9$ , calculated on the anhydrous basis.

**Description** Iotroxic Acid occurs as a white crystalline powder.

It is soluble in methanol, slightly soluble in ethanol (95), and practically insoluble in water and in diethyl ether.

It is gradually colored by light.

**Identification (1)** Heat 0.1 g of Iotroxic Acid over a flame: a purple gas evolves.

(2) Dissolve a suitable amount of Iotroxic Acid in a suitable amount of methanol, evaporate the methanol under reduced pressure, and determine the infrared absorption spectrum of the residue so obtained as directed in the potassium bromide disk method under Infrared Spectrophotometry <2.25>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wave numbers.

**Purity (1)** Clarity and color of solution—Dissolve 1.0 g of Iotroxic Acid in 10 mL of diluted sodium hydroxide TS (1 in 5): the solution is clear and colorless.

(2) Primary aromatic amines—Dissolve 0.20 g of Iotroxic Acid in 5 mL of water and 1 mL of sodium hydroxide TS, add 4 mL of a solution of sodium nitrite (1 in 100) and 10 mL of 1 mol/L hydrochloric acid TS, mix, and allow to stand for 2 minutes. Add 5 mL of ammonium amidosulfate TS, shake well, allow to stand for 1 minute, then add 0.4 mL of a solution of  $\alpha$ -naphthol in ethanol (95) (1 in 10), 15 mL of sodium hydroxide TS and water to make exactly 50 mL. Read the absorbance of this solution at 485 nm as directed under Ultraviolet-visible Spectrophotometry <2.24>.



iolet-visible Spectrophotometry <2.24>, using a blank solution obtained in the same manner as above: the absorbance is not more than 0.22.

(3) Iodine—Dissolve 0.20 g of Iotrox Acid in 2.0 mL of sodium hydrogen carbonate TS, add 5 mL of toluene, mix well, and allow to stand: the toluene layer is colorless.

(4) Free iodine ion—Weigh accurately about 5.0 g of Iotrox Acid, dissolve in 12 mL of a solution of meglumine (3 in 20), add water to make 70 mL, and adjust the pH to about 4.5 with acetic acid (100). To this solution add 2 mL of 0.1 mol/L sodium chloride TS, and titrate <2.50> with 0.001 mol/L silver nitrate VS (potentiometric titration).

Each mL of 0.001 mol/L silver nitrate  
= 0.1269 mg of I

Content of iodine ion in Iotrox Acid, calculated on the anhydrous basis, is not more than 0.004%.

(5) Heavy metals <1.07>—Heat strongly 1.0 g of Iotrox Acid as directed under Residue on Ignition Test, then proceed according to Method 2, and perform the test. Prepare the control solution with 1.0 mL of Standard Lead Solution (not more than 10 ppm).

(6) Related substances—Dissolve 0.15 g of Iotrox Acid in 10 mL of methanol, and use this solution as the sample solution. Pipet 1 mL of the sample solution, add methanol to make exactly 200 mL, and use this solution as the standard solution. Perform the test with these solutions as directed under Thin-layer Chromatography <2.03>. Spot 10  $\mu$ L each of the sample solution and standard solution on a plate of silica gel with fluorescent indicator for thin-layer chromatography. Develop the plate with a mixture of toluene, acetone and formic acid (6:4:1) to a distance of about 15 cm, and air-dry the plate. Examine under ultraviolet light (main wavelength: 254 nm): the spots other than the principal spot from the sample solution are not more intense than the spot from the standard solution.

**Water** <2.48> 1.0 – 2.0% (0.5 g, direct titration).

**Residue on ignition** <2.44> Not more than 0.1% (1 g).


**Assay** Weigh accurately about 0.5 g of Iotrox Acid, dissolve in 40 mL of sodium hydroxide TS in a saponification flask, add 1 g of zinc powder, and boil for 30 minutes under a reflux condenser. After cooling, filter, wash the flask and the filter paper with 50 mL of water, and combine the washings to the filtrate. To this solution add 5 mL of acetic acid (100), and titrate <2.50> with 0.1 mol/L silver nitrate VS (potentiometric titration).

Each mL of 0.1 mol/L silver nitrate VS  
= 20.26 mg of  $C_{22}H_{18}I_6N_2O_9$

**Containers and storage** Containers—Tight containers.  
Storage—Light-resistant.

## Ipratropium Bromide Hydrate

イプラトロピウム臭化物水和物



$C_{20}H_{30}BrNO_3 \cdot H_2O$ : 430.38  
(1*R*,3*r*,5*S*)-3-[(2*RS*)-3-Hydroxy-2-phenylpropanoyloxy]-8-methyl-8-(1-methylethyl)-8-azoniabicyclo[3.2.1]octane bromide monohydrate [66985-17-9]

Ipratropium Bromide Hydrate, when dried, contains not less than 99.0% of ipratropium bromide ( $C_{20}H_{30}BrNO_3$ : 412.36).

**Description** Ipratropium Bromide Hydrate occurs as a white, crystalline powder.

It is freely soluble in water, soluble in ethanol (99.5), slightly soluble in acetonitrile and in acetic acid (100), and practically insoluble in diethyl ether.

The pH of a solution of Ipratropium Bromide Hydrate (1 in 20) is between 5.0 and 7.5.

Melting point: about 223°C (with decomposition, after drying).

**Identification** (1) To 5 mg of Ipratropium Bromide Hydrate add 0.5 mL of fuming nitric acid, and evaporate on a water bath to dryness. After cooling, dissolve the residue in 5 mL of acetone, and add 2 drops of potassium hydroxide-ethanol TS: a purple color develops.

(2) Determine the absorption spectrum of a solution of Ipratropium Bromide Hydrate in 0.01 mol/L hydrochloric acid TS (3 in 2000) as directed under Ultraviolet-visible Spectrophotometry <2.24>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wavelengths.

(3) Determine the infrared absorption spectrum of Ipratropium Bromide Hydrate as directed in the potassium bromide disk method under Infrared Spectrophotometry <2.25>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wave numbers.

(4) The solution of Ipratropium Bromide Hydrate (1 in 100) responds to the Qualitative Tests <1.09> for bromide.

**Purity** (1) Clarity and color of solution—Dissolve 1.0 g of Ipratropium Bromide Hydrate in 20 mL of water: the solution is clear and colorless.

(2) Sulfate <1.14>—Perform the test with 1.0 g of Ipratropium Bromide Hydrate. Prepare the control solution with 0.50 mL of 0.005 mol/L sulfuric acid VS (not more than 0.024%).

(3) Heavy metals <1.07>—Proceed with 2.0 g of Ipratropium Bromide Hydrate according to Method 4, and perform the test. Prepare the control solution with 2.0 mL of Standard Lead Solution (not more than 10 ppm).

(4) Arsenic <1.11>—Prepare the test solution with 2.0 g of Ipratropium Bromide Hydrate according to Method 3,

and perform the test. Use a solution of magnesium nitrate hexahydrate in ethanol (95) (1 in 10) (not more than 1 ppm).

(5) Isopropylatropine bromide—Dissolve 25 mg of Ipratropium Bromide Hydrate in the mobile phase to make exactly 100 mL, and use this solution as the sample solution. Perform the test with 25  $\mu$ L of the sample solution as directed under Liquid Chromatography <2.01> according to the following conditions. Determine the peak area,  $A_a$ , of ipratropium and the peak area,  $A_b$ , having a relative retention time to ipratropium about 1.3 by the automatic integration method:  $A_b/(A_a + A_b)$  is not more than 0.01, and no peak other than the peak of ipratropium and the peak having a relative retention time to ipratropium about 1.3 appears within about 14 minutes of the retention time after the solvent peak.

#### Operating conditions—

**Detector:** An ultraviolet absorption photometer (wavelength: 210 nm).

**Column:** A stainless steel column about 4 mm in inside diameter and 10 to 15 cm in length, packed with octylsilanized silica gel for liquid chromatography (5  $\mu$ m in particle diameter).

**Column temperature:** Room temperature.

**Mobile phase:** A mixture of diluted phosphoric acid (1 in 200), acetonitrile and methanesulfonic acid (1000:120:1).

**Flow rate:** Adjust the flow rate so that the retention time of ipratropium is about 7 minutes.

**Selection of column:** Heat a solution of Ipratropium Bromide in 1 mol/L hydrochloric acid TS (1 in 100) at 100°C for 1 hour, and cool. To 2.5 mL of this solution add the mobile phase to make 100 mL. Proceed with 25  $\mu$ L of this solution under the above operating conditions, and calculate the resolution. Use a column showing a resolution not less than 3 between the peak of ipratropium and the peak having a relative retention time to ipratropium about 0.6.

**Detection sensitivity:** Adjust the detection sensitivity so that the peak height of ipratropium obtained from 25  $\mu$ L of the sample solution composes 50 to 80% of the full scale.

(6) Apo-compounds—Dissolve 0.14 g of Ipratropium Bromide in 0.01 mol/L hydrochloric acid TS to make 100 mL. Perform the test with this solution as directed under Ultraviolet-visible Spectrophotometry <2.24>, and determine the absorbances,  $A_1$  and  $A_2$ , at 246 nm and 263 nm, respectively:  $A_1/A_2$  is not more than 0.91.

**Loss on drying** <2.41> 3.9 – 4.4% (1 g, 105°C, 4 hours).

**Residue on ignition** <2.44> Not more than 0.1% (1 g).


**Assay** Weigh accurately about 0.3 g of Ipratropium Bromide Hydrate, previously dried, dissolve in 40 mL of acetic acid (100), add 40 mL of 1,4-dioxane and 2.5 mL of bismuth nitrate TS, and titrate <2.50> with 0.1 mol/L perchloric acid VS (potentiometric titration). Perform a blank determination, and make any necessary correction.

Each mL of 0.1 mol/L perchloric acid VS  
= 41.24 mg of  $C_{20}H_{30}BrNO_3$

**Containers and storage** Containers—Tight containers.

## Isepamicin Sulfate

イセパマイシン硫酸塩



$C_{22}H_{43}N_5O_{12} \cdot xH_2SO_4$

6-Amino-6-deoxy- $\alpha$ -D-glucopyranosyl-(1 $\rightarrow$ 4)-

[3-deoxy-4-C-methyl-3-methylamino- $\beta$ -L-arabinopyranosyl-(1 $\rightarrow$ 6)]-2-deoxy-1-N-[(2S)-3-amino-2-hydroxypropanoyl]-D-streptamine sulfate [67814-76-0]

Isepamicin Sulfate is the sulfate of a derivative of gentamycin B, an aminoglycoside substance, having antibacterial activity produced by the growth of *Micromonospora purpurea*.

It contains not less than 680  $\mu$ g (potency) and not more than 780  $\mu$ g (potency) per mg, calculated on the anhydrous basis. The potency of Isepamicin Sulfate is expressed as mass (potency) of isepamicin ( $C_{22}H_{43}N_5O_{12}$ : 569.60).

**Description** Isepamicin Sulfate occurs as a white to pale yellowish white powder.

It is very soluble in water, and practically insoluble in methanol and in ethanol (95).

It is hygroscopic.

**Identification (1)** Dissolve 0.02 g of Isepamicin Sulfate in 1 mL of water, add 3 mL of anthrone TS, shake, and allow to stand: a blue-purple color develops.

(2) Dissolve 10 mg each of Isepamicin Sulfate and Isepamicin Sulfate Reference Standard in 5 mL of water, and use these solutions as the sample solution and the standard solution. Perform the test with these solutions as directed under Thin-layer chromatography <2.03>. Spot 5  $\mu$ L of the sample solution and standard solution on a plate of silica gel for thin-layer chromatography. Develop the plate with a mixture of ammonia water (28), ethanol (99.5), 1-butanol and chloroform (5:5:4:2) to a distance of about 15 cm, and air-dry the plate. Spray evenly 0.2% ninhydrin-water saturated 1-butanol TS on the plate, and heat at about 100°C for about 10 minutes: the principal spots from the sample solution and the standard solution exhibit a red-brown color and show the same  $R_f$  value.

(3) Dissolve 0.01 g of Isepamicin Sulfate in 1 mL of water, and add 1 drop of barium chloride TS: a white precipitate is produced.

**Optical rotation** <2.49>  $[\alpha]_D^{20}$ : +100 – +120° (0.25 g calculated on the anhydrous bases, water, 25 mL, 100 mm).

**pH** <2.54> Dissolve 0.5 g of Isepamicin Sulfate in 5 mL of water: the pH of the solution is between 5.5 and 7.5.

**Purity** (1) Clarity and color of solution—Dissolve 1.0 g of Isepamicin Sulfate in 10 mL of water: the solution is clear and colorless.

(2) Heavy metals <1.07>—Proceed with 1.0 g of Isepamicin Sulfate according to Method 4, and perform the test. Prepare the control solution with 1.0 mL Standard Lead Solution (not more than 10 ppm).

(3) Related substances—Perform the test with 5  $\mu$ L of the sample solution obtained in the Assay as directed under Assay. Determine each peak area of the sample solution by the automatic integration method, and calculate the amounts of their peaks by the area percentage method: the amount of HAPA-gentamicin-B equivalent to about 0.4 of the relative retention time to isepamicin is not more than 5.0%, and gentamicin B equivalent to about 1.3 of that is not more than 3.0%. Correct the peak area of gentamicin B by multiplying the sensitivity coefficient, 1.11.

**Operating conditions—**

Apparatus, detector, column, column temperature, reaction coil, mobile phase, reagent, reaction temperature, flow rate of the mobile phase, and flow rate of the reagent: Proceed as directed in the operating conditions in the Assay.

Time span of measurement: About 2 times as long as the retention time of isepamicin.

**System suitability—**

Test for required detection: Pipet 1 mL of the standard solution, add water to make exactly 10 mL, and use this solution as the solution for the test for required detection. Pipet 1 mL of the solution, and add water to make exactly 10 mL. Confirm that the peak area of isepamicin obtained from 5  $\mu$ L of this solution is equivalent to 7 to 13% of that obtained from 5  $\mu$ L of the solution for the test for required detection.

System performance and system repeatability: Proceed as directed in the system suitability in the Assay.

**Water** <2.48> Not more than 12.0% (0.2 g, volumetric titration, direct titration. Use a mixture of formamide for water determination and methanol for water determination (2:1) instead of methanol for water determination).

**Residue on ignition** <2.44> Not more than 1.0% (1 g).

**Assay** Weigh accurately an amount of Isepamicin Sulfate and Isepamicin Sulfate Reference Standard, equivalent to about 20 mg (potency), dissolve each in water to make exactly 100 mL, and use these solutions as the sample solution and the standard solution, respectively. Perform the test with exactly 5  $\mu$ L each of the sample solution and standard solution as directed under Liquid Chromatography <2.01> according to the following conditions, and calculate the peak areas,  $A_T$  and  $A_S$ , of isepamicin of the solutions.

$$\text{Amount } [\mu\text{g (potency)}] \text{ of isepamicin (C}_{22}\text{H}_{43}\text{N}_5\text{O}_{12}) \\ = W_S \times (A_T/A_S) \times 1000$$

$W_S$ : Amount [mg (potency)] of Isepamicin Sulfate Reference Standard

**Operating conditions—**

Apparatus: Consist of two pumps for the mobile phase and the reagent transport, inject port, column, reaction coil, detector and recorder. Use a reaction coil with thermostat.

Detector: Fluorometry (excitation wavelength: 360 nm, de-

tection wavelength: 440 nm).

Column: A stainless steel column 4 mm in inside diameter and 15 cm in length, packed with octadecylsilanized silica gel for liquid chromatography (5  $\mu$ m in particle diameter).

Column temperature: A constant temperature of about 25°C.

Reaction coil: A column 0.25  $\mu$ m in inside diameter and 5 m in length.

Mobile phase: Dissolve 28.41 g of anhydrous sodium sulfate and 5.23 g of sodium 1-pentane sulfonate in 900 mL of water, add 1 mL of acetic acid (100), and add water to make exactly 1000 mL.

Reagent: To 500 mL of boric acid-potassium chloride-sodium hydroxide buffer solution, pH 10.0, add 5 mL of a solution of *o*-phthalaldehyde in ethanol (95) (2 in 25), 1 mL of 2-mercaptoethanol and 2 mL of a solution of lauromacrogol (1 in 4).

Reaction temperature: A constant temperature of about 45°C.

Flow rate of the mobile phase: About 0.6 mL per minute.

Flow rate of the reagent: About 0.5 mL per minute.

**System suitability—**


System performance: Dissolve 2 mg of Gentamicin B in 10 mL of the standard solution. When the procedure is run with 5  $\mu$ L of this solution under the above operating conditions, isepamicin and gentamicin B are eluted in this order with the resolution between these peaks being not less than 1.0.

System repeatability: When the test is repeated 5 times with 5  $\mu$ L of the standard solution under the above operating conditions, the relative standard deviation of the peak areas of isepamicin is not more than 3.0%.

**Containers and storage** Containers—Tight containers.

## Isoflurane

イソフルラン



$\text{C}_3\text{H}_2\text{ClF}_5\text{O}$ : 184.49

(2*RS*)-2-Chloro-2-(difluoromethoxy)-1,1,1-trifluoroethane  
[26675-46-7]

Isoflurane contains not less than 99.0% and not more than 101.0% of  $\text{C}_3\text{H}_2\text{ClF}_5\text{O}$ , calculated on the anhydrous basis.

**Description** Isoflurane occurs as a clear, colorless fluid liquid.

It is miscible with ethanol (99.5), with methanol and with *o*-xylene.

It is slightly soluble in water.

It is volatile, and has no inflammability.

It shows no optical rotation.

Refractive index  $n_D^{20}$ : about 1.30

Boiling point: about 47 – 50°C

**Identification** (1) The test solution obtained by the Oxygen Flask Combustion Method <1.06> with 50  $\mu$ L of Isoflurane, using 40 mL of water as the absorbing liquid, responds

to the Qualitative Tests <1.09> for chloride and fluoride.

(2) Determine the infrared absorption spectrum of Isoflurane as directed in the liquid film method under Infrared Spectrophotometry <2.25>, and compare the spectrum with the Reference Spectrum or the spectrum of Isoflurane Reference Standard: both spectra exhibit similar intensities of absorption at the same wave numbers.

**Specific gravity** <2.56>  $d_{20}^{20}$ : 1.500 – 1.520

**Purity** (1) Acidity or alkalinity—To 10 mL of Isoflurane add 5 mL of freshly boiled and cooled water, and shake for 1 minute: the water layer is neutral.

(2) Soluble chloride—To 60 g of Isoflurane add 40 mL of water, shake thoroughly, and separate the water layer. To 20 mL of the layer add 6 mL of dilute nitric acid and water to make 50 mL. Perform the test with this solution as directed under Chloride Limit Test <1.03>. Prepare the control solution with 0.25 mL of 0.01 mol/L hydrochloric acid VS (not more than 3 ppm).

(3) Soluble fluoride—To 6 g of Isoflurane add 12 mL of diluted 0.01 mol/L sodium hydroxide TS (1 in 20), and shake for 10 minutes. Transfer 4.0 mL of the water layer into a Nessler tube, add 30 mL of a mixture of alizarin complexone TS, acetic acid-potassium acetate buffer solution, pH 4.3 and cerium (III) nitrate TS (1:1:1), add water to make 50 mL, allow to stand for 60 minutes, and use this solution as the sample solution. Separately, to 0.4 mL of the fluorine standard solution and 4.0 mL of diluted 0.01 mol/L sodium hydroxide TS (1 in 20) in a Nessler tube add 30 mL of the mixture of alizarin complexone TS, acetic acid-potassium acetate buffer solution, pH 4.3 and cerium (III) nitrate TS (1:1:1), then proceed in the same manner as for the preparation of the sample solution, and use the solution so obtained as the standard solution. Determine the absorbances of the sample solution and standard solution at 600 nm as directed under Ultraviolet-visible Spectrophotometry <2.24>, using a solution, obtained by proceeding in the same manner as above with 4.0 mL of diluted 0.01 mol/L sodium hydroxide TS (1 in 20), as the blank: the absorbance of the sample solution is not more than that of the standard solution (not more than 2 ppm).

Fluorine standard solution: Dissolve exactly 2.21 g of sodium fluoride in water to make exactly 1000 mL. Pipet 10 mL of this solution, and add water to make exactly 1000 mL. Each mL of this solution contains 0.01 mg of fluorine (F).

(4) Peroxide—To 10 mL of Isoflurane add 1 mL of a freshly prepared solution of potassium iodide (1 in 10), shake vigorously, and allow to stand in a dark place for 1 hour: the water layer is not yellow.

(5) Related substances—Use Isoflurane as the sample solution. To exactly 1 mL of the sample solution add *o*-xylene to make exactly 100 mL. Pipet 1 mL of this solution, add *o*-xylene to make exactly 100 mL, and use this solution as the standard solution. Perform the test with exactly 5  $\mu$ L each of the sample solution and standard solution as directed under Gas Chromatography <2.02> according to the following conditions, and determine each peak area by the automatic integration method: the area of the peak other than isoflurane is not more than the peak area of isoflurane from the standard solution, and the total area of the peaks other than isoflurane is not more than 3 times the peak area of isoflurane from the standard solution.

**Operating conditions—**

Detector, column, column temperature, carrier gas, and

flow rate: Proceed as directed in the operating conditions in the Assay.

Time span of measurement: About 5 times as long as the retention time of isoflurane after injection of the sample solution.

**System suitability—**

Test for required detectability: To exactly 1 mL of the standard solution add *o*-xylene to make exactly 2 mL. Confirm that the peak area of isoflurane obtained with 5  $\mu$ L of this solution is equivalent to 35 to 65% of that with 5  $\mu$ L of the standard solution.

System performance and system repeatability: Proceed as directed in the system suitability in the Assay.

(6) Residue on evaporation Pipet 65 mL of Isoflurane, evaporate on a water bath, and dry the residue at 105°C for 1 hour: not more than 1.0 mg.

**Water** <2.48> Not more than 0.1% (2 g, Courometric titration).

**Assay** To exactly 5 mL each of Isoflurane and Isoflurane Reference Standard (separately determined water content <2.48> in the same manner as Isoflurane), add exactly 3 mL of ethyl acetate as the internal standard, then add *o*-xylene to make exactly 50 mL each. To 5 mL each of these solutions add *o*-xylene to make 50 mL, and use these solutions as the sample solution and the standard solution, respectively. Perform the test with 2  $\mu$ L each of the sample solution and standard solution as directed under Gas Chromatography <2.02> according to the following conditions, and determine the ratios,  $Q_T$  and  $Q_S$ , of the peak area of isoflurane to that of the internal standard.

$$\begin{aligned} &\text{Amount (mg) of isoflurane (C}_3\text{H}_2\text{ClF}_5\text{O) in 5 mL of} \\ &\text{Isoflurane} \\ &= V_S \times (Q_T/Q_S) \times 1000 \times 1.506 \end{aligned}$$

$V_S$ : Amount (mL) of Isoflurane Reference Standard, calculated on the anhydrous basis

1.506: Specific gravity ( $d_{20}^{20}$ ) of isoflurane

**Operating conditions—**

Detector: A hydrogen flame-ionization detector.

Column: A stainless steel column 3 mm in inside diameter and 3.5 m in length, packed with siliceous earth for gas chromatography (125 – 149  $\mu$ m in particle diameter), coated in 10% with nonylphenoxypoly(ethyleneoxy)ethanol for gas chromatography and in 15% with polyalkylene glycol for gas chromatography.

Column temperature: A constant temperature of about 80°C.

Carrier gas: Nitrogen

Flow rate: Adjust the flow rate so that the retention time of isoflurane is about 7 minutes.

**System suitability—**

System performance: When the procedure is run with 2  $\mu$ L of the standard solution under the above operating conditions, isoflurane and the internal standard are eluted in this order with the resolution between these peaks being not less than 3


System repeatability: When the test is repeated 6 times with 2  $\mu$ L of the standard solution under the above operating conditions, the relative standard deviation of the peak area of isoflurane is not more than 1.0%.

**Containers and storage** Containers—Tight containers.

Storage—At a temperature not exceeding 30°C.

## L-Isoleucine

L-イソロイシン



$C_6H_{13}NO_2$ : 131.17

(2S,3S)-2-Amino-3-methylpentanoic acid [73-32-5]

L-Isoleucine, when dried, contains not less than 98.5% of  $C_6H_{13}NO_2$ .

**Description** L-Isoleucine occurs as white crystals or crystalline powder. It is odorless or has a faint characteristic odor, and has a slightly bitter taste.

It is freely soluble in formic acid, sparingly soluble in water, and practically insoluble in ethanol (95).

It dissolves in dilute hydrochloric acid.

**Identification** Determine the infrared absorption spectrum of L-Isoleucine, previously dried, as directed in the potassium bromide disk method under Infrared Spectrophotometry <2.25>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wave numbers.

**Optical rotation** <2.49>  $[\alpha]_D^{20}$ : +39.5 – +41.5° (after drying, 1 g, 6 mol/L hydrochloric acid TS, 25 mL, 100 mm).

**pH** <2.54> Dissolve 1.0 g of L-Isoleucine in 100 mL of water: the pH of this solution is between 5.5 and 6.5.

**Purity** (1) Clarity and color of solution—Dissolve 0.5 g of L-Isoleucine in 10 mL of 1 mol/L hydrochloric acid TS: the solution is clear and colorless.

(2) Chloride <1.03>—Perform the test with 0.5 g of L-Isoleucine. Prepare the control solution with 0.30 mL of 0.01 mol/L hydrochloric acid VS (not more than 0.021%).

(3) Sulfate <1.14>—Perform the test with 0.6 g of L-Isoleucine. Prepare the control solution with 0.35 mL of 0.005 mol/L sulfuric acid VS (not more than 0.028%).

(4) Ammonium—Perform the test with 0.25 g of L-Isoleucine. Prepare the control solution with 5.0 mL of Standard Ammonium Solution (not more than 0.02%).

(5) Heavy metals <1.07>—Dissolve 1.0 g of L-Isoleucine in 40 mL of water and 2 mL of dilute acetic acid by warming, cool, and add water to make 50 mL. Perform the test using this solution as the test solution. Prepare the control solution as follows: to 2.0 mL of Standard Lead Solution add 2 mL of dilute acetic acid and water to make 50 mL (not more than 20 ppm).

(6) Arsenic <1.11>—Prepare the test solution with 1.0 g of L-Isoleucine according to Method 2, and perform the test (not more than 2 ppm).

(7) Related substances—Dissolve 0.10 g of L-Isoleucine in 25 mL of water, and use this solution as the sample solution. Pipet 1 mL of the sample solution, and add water to make exactly 50 mL. Pipet 5 mL of this solution, add water to make exactly 20 mL, and use this solution as the standard solution. Perform the test with these solutions as directed un-

der Thin-layer Chromatography <2.03>. Spot 5  $\mu$ L each of the sample solution and standard solution on a plate of silica gel for thin-layer chromatography. Develop the plate with a mixture of 1-butanol, water and acetic acid (100) (3:1:1) to a distance of about 10 cm, and dry the plate at 80°C for 30 minutes. Spray evenly the plate with a solution of ninhydrin in acetone (1 in 50), and heat at 80°C for 5 minutes: the spots other than the principal spot from the sample solution are not more intense than the spot from the standard solution.

**Loss on drying** <2.41> Not more than 0.30% (1 g, 105°C, 3 hours).

**Residue on ignition** <2.44> Not more than 0.1% (1 g).


**Assay** Weigh accurately about 0.13 g of L-Isoleucine, previously dried, and dissolve in 3 mL of formic acid, add 50 mL of acetic acid (100), and titrate <2.50> with 0.1 mol/L perchloric acid VS (potentiometric titration). Perform a blank determination, and make any necessary correction.

Each mL of 0.1 mol/L perchloric acid VS  
= 13.12 mg of  $C_6H_{13}NO_2$

**Containers and storage** Containers—Tight containers.

## Isoniazid

イソニアジド



$C_6H_7N_3O$ : 137.14

Pyridine-4-carbohydrazide [54-85-3]

Isoniazid, when dried, contains not less than 98.5% of  $C_6H_7N_3O$ .

**Description** Isoniazid occurs as colorless crystals or a white, crystalline powder. It is odorless.

It is freely soluble in water, sparingly soluble in ethanol (95), slightly soluble in acetic anhydride, and very slightly soluble in diethyl ether.

**Identification** (1) Dissolve about 20 mg of Isoniazid in water to make 200 mL. To 5 mL of the solution add 1 mL of 0.1 mol/L hydrochloric acid TS and water to make 50 mL. Determine the absorption spectrum of the solution as directed under Ultraviolet-visible Spectrophotometry <2.24>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wavelengths.

(2) Determine the infrared absorption spectrum of Isoniazid, previously dried, as directed in the potassium bromide disk method under Infrared Spectrophotometry <2.25>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wave numbers.

**pH** <2.54> Dissolve 1.0 g of Isoniazid in 10 mL of freshly boiled and cooled water: the pH of this solution is between 6.5 and 7.5.

**Melting point** <2.60> 170 – 173°C

**Purity** (1) Clarity and color of solution—Dissolve 1.0 g of isoniazid in 20 mL of water: the solution is clear and colorless.

(2) Heavy metals <1.07>—Proceed with 1.0 g of Isoniazid according to Method 2, and perform the test. Prepare the control solution with 2.0 mL of Standard Lead Solution (not more than 20 ppm).

(3) Arsenic <1.11>—Prepare the test solution with 0.40 g of Isoniazid according to Method 3, and perform the test. In this case, add 10 mL of a solution of magnesium nitrate hexahydrate in ethanol (95) (1 in 50), then add 1.5 mL of hydrogen peroxide (30), and ignite the ethanol to burn (not more than 5 ppm).

(4) Hydrazine—Dissolve 0.10 g of isoniazid in 5 mL of water, add 0.1 mL of a solution of salicylaldehyde in ethanol (95) (1 in 20), shake immediately, and allow to stand for 5 minutes: no turbidity is produced.

**Loss on drying** <2.41> Not more than 0.5% (1 g, 105°C, 2 hours).

**Residue on ignition** <2.44> Not more than 0.1% (1 g).

**Assay** Weigh accurately about 0.3 g of isoniazid, previously dried, dissolve in 50 mL of acetic acid (100) and 10 mL of acetic anhydride, and titrate <2.50> with 0.1 mol/L perchloric acid VS until the color of the solution changes from yellow to green (indicator: 0.5 mL of *p*-naphtholbenzein TS). Perform a blank determination, and make any necessary correction.

Each mL of 0.1 mol/L perchloric acid VS  
= 13.71 mg of C<sub>6</sub>H<sub>7</sub>N<sub>3</sub>O

**Containers and storage** Containers—Tight containers.

Storage—Light-resistant.

## Isoniazid Injection

### イソニアジド注射液

Isoniazid Injection is an aqueous solution for injection.

It contains not less than 95% and not more than 105% of the labeled amount of isoniazid (C<sub>6</sub>H<sub>7</sub>N<sub>3</sub>O: 137.14).

**Method of preparation** Prepare as directed under Injections, with Isoniazid.

**Description** Isoniazid Injection occurs as a clear, colorless liquid.

pH: 6.5 – 7.5.

**Identification** To a volume of Isoniazid Injection, equivalent to 20 mg of Isoniazid according to the labeled amount, and add water to make 200 mL. To 5 mL of the solution add 1 mL of 0.1 mol/L hydrochloric acid TS and water to make 50 mL. Determine the absorption spectrum of this solution as directed under Ultraviolet-visible Spectrophotometry <2.24>: it exhibits a maximum between 264 nm and 268 nm.

**Extractable volume** <6.05> It meets the requirement.

**Assay** To an exactly measured volume of Isoniazid Injection,

equivalent to about 50 mg of isoniazid (C<sub>6</sub>H<sub>7</sub>N<sub>3</sub>O), add water to make exactly 100 mL. Pipet 5 mL of the solution, add exactly 5 mL of the internal standard and the mobile phase to make 50 mL, and use this solution as the sample solution. Separately, weigh accurately about 50 mg of isoniazid for assay, previously dried at 105°C for 2 hours, and dissolve in water to make exactly 100 mL. Pipet 5 mL of this solution, add exactly 5 mL of the internal standard and the mobile phase to make 50 mL, and use this solution as the standard solution. Perform the test with 5 µL each of the sample solution and standard solution as directed under Liquid Chromatography <2.01> according to the following conditions, and determine the ratios,  $Q_T$  and  $Q_S$ , of the peak area of isoniazid to that of the internal standard.

Amount (mg) of isoniazid (C<sub>6</sub>H<sub>7</sub>N<sub>3</sub>O) =  $W_S \times (Q_T/Q_S)$

$W_S$ : Amount (mg) of isoniazid for assay

**Internal standard solution**—A solution of propyl parahydroxybenzoate (1 in 4000).

**Operating conditions**—

Detector: An ultraviolet absorption photometer (wavelength: 265 nm).

Column: A stainless steel column 4.6 mm in inside diameter and 15 cm in length, packed with octadecylsilanized silica gel for liquid chromatography (5 µm in particle diameter).

Column temperature: A constant temperature of about 40°C.

Mobile phase: Dissolve 6.80 g of potassium dihydrogen phosphate in water to make 1000 mL. Separately, to 5.76 g of phosphoric acid add water to make 1000 mL. Mix these solutions to make a solution having pH 2.5. To 500 mL of this solution add 500 mL of methanol, and add 2.86 g of sodium tridecanesulfonate to dissolve.

Flow rate: Adjust the flow rate so that the retention time of isoniazid is about 5 minutes.

**System suitability**—

System performance: When the procedure is run with 5 µL of the standard solution under the above operating conditions, isoniazid and the internal standard are eluted in this order with the resolution between these peaks being not less than 10.

System repeatability: When the test is repeated 6 times with 5 µL of the standard solution under the above operating conditions, the relative standard deviation of the ratios of the peak area of isoniazid to that of the internal standard is not more than 1.3%.

**Containers and storage** Containers—Hermetic containers, and colored containers may be used.

Storage—Light-resistant.

## Isoniazid Tablets

### イソニアジド錠

Isoniazid Tablets contain not less than 95% and not more than 105% of the labeled amount of isoniazid (C<sub>6</sub>H<sub>7</sub>N<sub>3</sub>O: 137.14).

**Method of preparation** Prepare as directed under Tablets,

with Isoniazid.

**Identification** Take a quantity of powdered Isoniazid Tablets, equivalent to 0.02 g of Isoniazid according to the labeled amount, add 200 mL of water, shake well, and filter. To 5 mL of the filtrate add 1 mL of 0.1 mol/L hydrochloric acid TS and water to make 50 mL, and determine the absorption spectrum of this solution as directed under Ultraviolet-visible Spectrophotometry <2.24>: it exhibits a maximum between 264 nm and 268 nm.

**Dissolution** <6.10> Perform the test according to the following method: it meets the requirement.

Perform the test with 1 tablet of Isoniazid Tablets at 50 revolutions per minute according to the Paddle method, using 900 mL of water as the test solution. Take 20 mL or more of the dissolved solution 20 minutes after starting the test, and filter through a membrane filter with pore size of not more than 0.45  $\mu$ m. Discard the first 10 mL of the filtrate, pipet 5 mL of the subsequent, add water to make exactly 50 mL, and use this solution as the sample solution. Separately, weigh accurately about 0.1 g of isoniazid for assay, previously dried at 105°C for 2 hours, dissolve in water to make exactly 100 mL, then pipet 5 mL of this solution, add water to make exactly 50 mL, and then pipet 5 mL of this solution, add water to make exactly 50 mL, and use this solution as the standard solution. Determine the absorbances,  $A_T$  and  $A_S$ , of the sample solution and standard solution at 267 nm as directed under Ultraviolet-visible Spectrophotometry <2.24>. The dissolution rate of Isoniazid Tablets in 20 minutes is not less than 75%.

$$\begin{aligned} &\text{Dissolution rate (\%)} \text{ with respect to the labeled} \\ &\text{amount of isoniazid (C}_6\text{H}_7\text{N}_3\text{O)} \\ &= W_S \times (A_T/A_S) \times (90/C) \end{aligned}$$

$W_S$ : Amount (mg) of isoniazid for assay.

$C$ : Labeled amount (mg) of isoniazid (C<sub>6</sub>H<sub>7</sub>N<sub>3</sub>O) in 1 tablet.

**Assay** Weigh accurately and powder not less than 20 Isoniazid Tablets. Weigh accurately a quantity of the powder, equivalent to about 0.1 g of isoniazid (C<sub>6</sub>H<sub>7</sub>N<sub>3</sub>O), add 150 mL of water, shake for 30 minutes, then add water to make exactly 200 mL, and filter. Discard the first 10 mL of the filtrate, pipet 5 mL of the subsequent filtrate, add the mobile phase to make exactly 50 mL, and use this solution as the sample solution. Separately, weigh accurately about 50 mg of isoniazid for assay, previously dried at 105°C for 2 hours, dissolve in water to make exactly 100 mL. Pipet 5 mL of this solution, add the mobile phase to make exactly 50 mL, and use this solution as the standard solution. Perform the test with exactly 10  $\mu$ L each of the sample solution and standard solution as directed under Liquid Chromatography <2.01> according to the following conditions. Determine the peak areas,  $A_T$  and  $A_S$ , of isoniazid of the sample solution and standard solution.

$$\begin{aligned} &\text{Amount (mg) of isoniazid (C}_6\text{H}_7\text{N}_3\text{O)} \\ &= W_S \times (A_T/A_S) \times 2 \end{aligned}$$

$W_S$ : Amount (mg) of isoniazid for assay

**Operating conditions**—

Detector: An ultraviolet absorption photometer (wavelength: 265 nm).

Column: A stainless steel column 4.6 mm in inside di-

ameter and 25 cm in length, packed with octadecylsilanized silica gel for liquid chromatography (5  $\mu$ m in particle diameter).

Column temperature: A constant temperature of about 40°C.

Mobile phase: Dissolve 6.80 g of potassium dihydrogen phosphate in water to make 1000 mL. Separately, to 5.76 g of phosphoric acid add water to make 1000 mL. Mix these solutions to adjust the pH to 2.5. To 400 mL of this solution add 600 mL of methanol, and dissolve 2.86 g of sodium tridecanesulfonate in this.

Flow rate: Adjust the flow rate so that the retention time of isoniazid is about 5 minutes.

**System suitability**—

System performance: Dissolve 5 mg of Isoniazid and 5 mg of isonicotinic acid in 100 mL of the mobile phase. When the procedure is run with 10  $\mu$ L of this solution under the above operating conditions, isonicotinic acid and isoniazid are eluted in this order with the resolution between these peaks being not less than 1.5.


System repeatability: When the test is repeated 6 times with 10  $\mu$ L of the standard solution under the above operating conditions, the relative standard deviation of the peak area of isoniazid is not more than 1.0%.

**Containers and storage** Containers—Tight containers.

Storage—Light-resistant.

## *l*-Isoprenaline Hydrochloride

*l*-イソプレナリン塩酸塩



C<sub>11</sub>H<sub>17</sub>NO<sub>3</sub>·HCl: 247.72

4-[(1*R*)-1-Hydroxy-2-[(1-methylethyl)amino]ethyl] benzene-1,2-diol monohydrochloride [51-30-9]

*l*-Isoprenaline Hydrochloride, when dried, contains not less than 98.0% of C<sub>11</sub>H<sub>17</sub>NO<sub>3</sub>·HCl.

**Description** *l*-Isoprenaline Hydrochloride occurs as a white, crystalline powder. It is odorless.

It is freely soluble in water, sparingly soluble in ethanol (95), and practically insoluble in acetic acid (100), in acetic anhydride, in diethyl ether and in chloroform.

It gradually changes in color by air and by light.

**Identification** (1) Dissolve 0.01 g of *l*-Isoprenaline Hydrochloride in 5 mL of water, and add 1 drop of iron (III) chloride TS: a deep green color develops, and changes through yellow-green to brown on standing.

(2) Dissolve 1 mg each of *l*-Isoprenaline Hydrochloride in 1 mL of water in the test tubes A and B. Add 10 mL of potassium hydrogen phthalate buffer solution, pH 3.5 to A, and add 10 mL of phosphate buffer solution, pH 6.5 to B. To each of the test tubes add 1 mL of iodine TS, allow to stand for 5 minutes, and add 2 mL each of sodium thiosulfate TS: a red color develops in the test tube A, and a deep red color de-

velops in the test tube B.

(3) Dissolve 0.01 g of *l*-Isoprenaline Hydrochloride in 1 mL of water, and add 1 mL of phosphotungstic acid TS: a light brown precipitate is produced.

(4) Determine the absorption spectrum of a solution of *l*-Isoprenaline Hydrochloride in 0.1 mol/L hydrochloric acid TS (1 in 20,000) as directed under Ultraviolet-visible Spectrophotometry <2.24>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wavelengths.

(5) A solution of *l*-Isoprenaline Hydrochloride (1 in 10) responds to the Qualitative Tests <1.09> (2) for chloride.

**Optical rotation** <2.49>  $[\alpha]_D^{20}$ :  $-36 - -41^\circ$  (after drying, 0.25 g, water, 25 mL, 100 mm).

**pH** <2.54> Dissolve 0.01 g of *l*-Isoprenaline Hydrochloride in 10 mL of water: the pH of the solution is between 4.5 and 5.5.

**Purity** (1) Clarity and color of solution—Dissolve 1.0 g of *l*-Isoprenaline Hydrochloride in 20 mL of 0.1 mol/L hydrochloric acid TS: the solution is clear and colorless.

(2) Sulfate <1.14>—Perform the test with 0.10 g of *l*-Isoprenaline Hydrochloride. Prepare the control solution with 0.40 mL of 0.005 mol/L sulfuric acid VS (not more than 0.192%).

(3) Heavy metals <1.07>—Proceed with 1.0 g of *l*-Isoprenaline Hydrochloride according to Method 1, and perform the test. Prepare the control solution with 2.0 mL of Standard Lead Solution (not more than 20 ppm).

(4) Isoproterenone—Dissolve 50 mg of *l*-Isoprenaline Hydrochloride in 0.01 mol/L hydrochloric acid TS to make exactly 25 mL, and determine the absorbance of the solution at 310 nm as directed under Ultraviolet-visible Spectrophotometry <2.24>: not more than 0.040.

**Loss on drying** <2.41> Not more than 0.5% (1 g, in vacuum, silica gel, 4 hours).

**Residue on ignition** <2.44> Not more than 0.2% (1 g).

**Assay** Weigh accurately about 0.5 g of *l*-Isoprenaline Hydrochloride, previously dried, dissolve in 100 mL of a mixture of acetic acid (100) and acetic anhydride (3:2) by warming, cool, and titrate <2.50> with 0.1 mol/L perchloric acid VS (potentiometric titration). Perform a blank determination.


Each mL of 0.1 mol/L perchloric acid VS  
= 24.77 mg of  $C_{11}H_{17}NO_3 \cdot HCl$

**Containers and storage** Containers—Tight containers.  
Storage—Light-resistant.

## Isopropanol

### Isopropyl Alcohol

イソプロパノール



$C_3H_8O$ : 60.10

Propan-2-ol [67-63-0]

**Description** Isopropanol is a clear, colorless liquid. It has a characteristic odor.

It is miscible with water, with methanol, with ethanol (95), and with diethyl ether.

It is flammable and volatile.

**Identification** (1) To 1 mL of Isopropanol add 2 mL of iodine TS and 2 mL of sodium hydroxide TS, and shake: a light yellow precipitate is formed.

(2) To 5 mL of Isopropanol add 20 mL of potassium dichromate and 5 mL of sulfuric acid with caution, and warm gently on a water bath: the produced gas has the odor of acetone, and the gas turns the filter paper, previously wetted with a solution of salicylaldehyde in ethanol (95) (1 in 10) and with a solution of sodium hydroxide (3 in 10), to red-brown.

**Specific gravity** <2.56>  $d_{20}^{20}$ : 0.785 – 0.788

**Purity** (1) Clarity of solution—To 2.0 mL of Isopropanol add 8 mL of water, and shake: the solution is clear.

(2) Acidity—To 15.0 mL of Isopropanol add 50 mL of freshly boiled and cooled water and 2 drops of phenolphthalein TS, and add 0.40 mL of 0.01 mol/L sodium hydroxide VS: a red color develops.

(3) Residue on evaporation—Evaporate 20.0 mL of Isopropanol on a water bath to dryness, and dry at 105°C for 1 hour: the mass of the residue is not more than 1.0 mg.

**Water** <2.48> Not more than 0.75 w/v% (2 mL, direct titration).


**Distilling range** <2.57> 81 – 83°C, not less than 94 vol%.

**Containers and storage** Containers—Tight containers.  
Storage—Remote from fire.

## Isopropylantipyrene

### Propyphenazone

イソプロピルアンチピリン



$C_{14}H_{18}N_2O$ : 230.31

1,5-Dimethyl-4-(1-methylethyl)-2-phenyl-  
1,2-dihydro-3*H*-pyrazol-3-one [479-92-5]



Isopropylantipyrine, when dried, contains not less than 98.0% of  $C_{14}H_{18}N_2O$ .

**Description** Isopropylantipyrine occurs as white crystals or crystalline powder. It is odorless, and has a slightly bitter taste.

It is very soluble in acetic acid (100), freely soluble in ethanol (95) and in acetone, soluble in diethyl ether, and slightly soluble in water.

**Identification (1)** To 2 mL of a solution of Isopropylantipyrine (1 in 500) add 1 drop of iron (III) chloride TS: a light red color develops. Further add 3 drops of sulfuric acid to this solution: the color changes to pale yellow.

**(2)** Add 5 mL of a solution of Isopropylantipyrine (1 in 500) to a mixture of 5 mL of potassium hexacyanoferrate (III) TS and 1 to 2 drops of iron (III) chloride TS: a dark green color gradually develops.

**(3)** To 2 mL of a solution of Isopropylantipyrine (1 in 500) add 2 to 3 drops of tannic acid TS: a white precipitate is produced.

**Melting point** <2.60> 103 – 105°C

**Purity (1) Chloride** <1.03> —Dissolve 1.0 g of Isopropylantipyrine in 30 mL of dilute ethanol, and add 6 mL of dilute nitric acid and water to make 50 mL. Perform the test using this solution as the test solution. Prepare the control solution as follows: to 0.40 mL of 0.01 mol/L hydrochloric acid VS add 6 mL of dilute nitric acid, 30 mL of dilute ethanol and water to make 50 mL (not more than 0.014%).

**(2) Sulfate** <1.14>—Dissolve 1.0 g of Isopropylantipyrine in 30 mL of dilute ethanol, and add 1 mL of dilute hydrochloric acid and water to make 50 mL. Perform the test using this solution as the test solution. Prepare the control solution as follows: to 0.40 mL of 0.005 mol/L sulfuric acid VS add 1 mL of dilute hydrochloric acid and 30 mL of dilute ethanol, and dilute with water to make 50 mL (not more than 0.019%).

**(3) Heavy metals** <1.07>—Dissolve 1.0 g of Isopropylantipyrine in 25 mL of acetone, add 2 mL of dilute acetic acid and water to make 50 mL, and perform the test using this solution as the test solution. Prepare the control solution as follows: to 2.0 mL of Standard Lead Solution add 2 mL of dilute acetic acid, 25 mL of acetone, and dilute with water to make 50 mL (not more than 20 ppm).

**(4) Arsenic** <1.11>—Prepare the test solution with 1.0 g of Isopropylantipyrine according to Method 3, and perform the test (not more than 2 ppm).

**(5) Antipyrine**—Dissolve 1.0 g of Isopropylantipyrine in 10 mL of dilute ethanol, and add 1 mL of sodium nitrite TS and 1 mL of dilute sulfuric acid: no green color develops.

**Loss on drying** <2.41> Not more than 0.5% (1 g, in vacuum, silica gel, 5 hours).

**Residue on ignition** <2.44> Not more than 0.1% (1 g).

**Assay** Weigh accurately about 0.4 g of Isopropylantipyrine, previously dried, dissolve in 60 mL of a mixture of acetic acid (100) and acetic anhydride (2:1), and titrate <2.50> with 0.1 mol/L perchloric acid VS (potentiometric titration). Perform a blank determination.


Each mL of 0.1 mol/L perchloric acid VS

= 23.03 mg of  $C_{14}H_{18}N_2O$

**Containers and storage** Containers—Tight containers.

## Isosorbide

イソソルビド



$C_6H_{10}O_4$ : 146.14

1,4:3,6-Dianhydro-D-glucitol [652-67-5]

Isosorbide contains not less than 98.5% of  $C_6H_{10}O_4$ , calculated on the anhydrous basis.

**Description** Isosorbide occurs as white crystals or masses. It is odorless, or has a faint, characteristic odor, and has a bitter taste.

It is very soluble in water and in methanol, freely soluble in ethanol (95), and slightly soluble in diethyl ether.

It is hygroscopic.

**Identification (1)** To 0.1 g of Isosorbide add 6 mL of diluted sulfuric acid (1 in 2), and dissolve by heating in a water bath. After cooling, shake well with 1 mL of a solution of potassium permanganate (1 in 30), and heat in a water bath until the color of potassium permanganate disappears. To this solution add 10 mL of 2,4-dinitrophenylhydrazine TS, and heat in a water bath: an orange precipitate is formed.

**(2)** To 2 g of Isosorbide add 30 mL of pyridine and 4 mL of benzoyl chloride, boil under a reflux condenser for 50 minutes, cool, and pour gradually the solution into 100 mL of cold water. Filter the formed precipitate by suction through a glass filter (G3), wash with water, recrystallize twice from ethanol (95), and dry in a desiccator (in vacuum, silica gel) for 4 hours: it melts <2.60> between 102°C and 103°C.

**(3)** Determine the infrared absorption spectrum of Isosorbide as directed in the potassium bromide disk method under Infrared Spectrophotometry <2.25>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wave numbers.

**Optical rotation** <2.49>  $[\alpha]_D^{20}$ : +45.0 – +46.0° (5 g, calculated on the anhydrous basis, water, 50 mL, 100 mm).

**Purity (1) Clarity and color of solution**—Take 25 g of Isosorbide in a Nessler tube, and dissolve in 50 mL of water: the solution is clear, and has no more color than the following control solution.

Control solution: To a mixture of 1.0 mL of Cobaltous Chloride Stock CS, 3.0 mL of Ferric Chloride Stock CS and 2.0 mL of Cupric Sulfate Stock CS add water to make 10.0 mL. To 3.0 mL of this solution add water to make 50 mL.

**(2) Sulfate** <1.14>—Perform the test with 2.0 g of Isosorbide. Prepare the control solution with 1.0 mL of 0.005 mol/L sulfuric acid VS (not more than 0.024%).

**(3) Heavy metals** <1.07>—Proceed with 5.0 g of Isosorbide according to Method 1, and perform the test. Prepare the control solution with 2.5 mL of Standard Lead Solution

(not more than 5 ppm).

(4) **Arsenic** <1.11>—Prepare the test solution with 1.0 g of Isosorbide according to Method 1, and perform the test (not more than 2 ppm).

(5) **Related substances**—Dissolve 0.10 g of Isosorbide in 10 mL of methanol, and use this solution as the sample solution. Pipet 2 mL of the sample solution, add methanol to make exactly 100 mL, and use this solution as the standard solution. Perform the test with these solutions as directed under Thin-layer Chromatography <2.03>. Spot 10  $\mu$ L each of the sample solution and standard solution on a plate of silica gel for thin-layer chromatography. Develop the plate with a mixture of ethanol (95) and cyclohexane (1:1) to a distance of about 10 cm, and air-dry the plate. Spray evenly a mixture of ethanol (95) and sulfuric acid (9:1) on the plate, and heat at 150°C for 30 minutes; the spots other than the principal spot from the sample solution are not more intense than the spot from the standard solution.

**Water** <2.48> Not more than 1.5% (2 g, direct titration).

**Residue on ignition** <2.44> Not more than 0.1% (1 g).


**Assay** Weigh accurately about 10 g of Isosorbide, calculated on the anhydrous basis, and dissolve in water to make exactly 100 mL. Determine the optical rotation <2.49>,  $\alpha_D$ , of this solution at 20  $\pm$  1°C in a 100-mm cell.

$$\text{Amount (g) of } C_6H_{10}O_4 = \alpha_D \times 2.1978$$

**Containers and storage** Containers—Tight containers.

## Isosorbide Dinitrate

硝酸イソソルビド



$C_6H_8N_2O_8$ : 236.14

1,4:3,6-Dianhydro-D-glucitol dinitrate [87-33-2]

Isosorbide Dinitrate contains not less than 95.0% of  $C_6H_8N_2O_8$ , calculated on the anhydrous basis.

**Description** Isosorbide Dinitrate occurs as white crystals or crystalline powder. It is odorless or has a faint odor like that of nitric acid.

It is very soluble in *N,N*-dimethylformamide and in acetone, freely soluble in chloroform and in toluene, soluble in methanol, in ethanol (95) and in diethyl ether, and practically insoluble in water.

It explodes if heated quickly or subjected to percussion.

**Identification** (1) Dissolve 0.01 g of Isosorbide Dinitrate in 1 mL of water, and dissolve by adding 2 mL of sulfuric acid cautiously. After cooling, superimpose 3 mL of iron (II) sulfate TS, and allow to stand for 5 to 10 minutes: a brown ring is produced at the zone of contact.

(2) Dissolve 0.1 g of Isosorbide Dinitrate in 6 mL of

diluted sulfuric acid (1 in 2) by heating in a water bath. After cooling, add 1 mL of a solution of potassium permanganate (1 in 30), stir well, and heat in a water bath until the color of potassium permanganate disappears. Add 10 mL of 2,4-dinitro-phenylhydrazine TS, and heat in a water bath: an orange precipitate is produced.

**Optical rotation** <2.49>  $[\alpha]_D^{20}$ : +134 – +139° (1 g, calculated on the anhydrous basis, ethanol (95), 100 mL, 100 mm).

**Purity** (1) **Clarity and color of solution**—Dissolve 1.0 g of Isosorbide Dinitrate in 10 mL of acetone: the solution is clear and colorless.

(2) **Sulfate** <1.14>—Dissolve 1.5 g of Isosorbide Dinitrate in 15 mL of *N,N*-dimethylformamide, add 60 mL of water, cool, and filter. Wash the filter paper with three 20-mL portions of water, combine the washings with the filtrate, and add water to make 150 mL. To 40 mL of this solution add 1 mL of dilute hydrochloric acid and water to make 50 mL. Perform the test using this solution as the test solution. Prepare the control solution with 0.40 mL of 0.005 mol/L sulfuric acid VS (not more than 0.048%).

(3) **Nitrate**—Dissolve 0.05 g of Isosorbide Dinitrate in 30 mL of toluene, and extract with three 20-mL portions of water. Combine the aqueous layers, and wash with two 20-mL portions of toluene. To the aqueous layer add water to make 100 mL, and use this solution as the sample solution. Pipet 5.0 mL of Standard Nitric Acid Solution and 25 mL of the sample solution in each Nessler tube, and add water to make 50 mL, respectively. To each of them add 0.06 g of Griss-Romijn's nitric acid reagent, stir well, allow to stand for 30 minutes, and observe from the side of the Nessler tube: the sample solution has no more color than the standard solution.

(4) **Heavy metals** <1.07>—Dissolve 1.0 g of Isosorbide Dinitrate in 30 mL of acetone, and add 2 mL of dilute acetic acid and water to make 50 mL. Perform the test using this solution as the test solution. Prepare the control solution as follows: to 2.0 mL of Standard Lead Solution add 30 mL of acetone, 2 mL of dilute acetic acid and water to make 50 mL (not more than 20 ppm).

**Water** <2.48> Not more than 1.5% (0.3 g, direct titration).

**Assay** Weigh accurately about 0.1 g of Isosorbide Dinitrate in a Kjeldahl flask as described under the Nitrogen Determination <1.08>, dissolve in 10 mL of methanol, add 3 g of Devarda's alloy and 50 mL of water, and connect the flask with the distillation apparatus as described under the Nitrogen Determination <1.08>. Measure exactly 25 mL of 0.05 mol/L sulfuric acid VS in an absorption flask, add 5 drops of bromocresol green-methyl red TS, and immerse the lower end of the condenser tube in it. Add 15 mL of a solution of sodium hydroxide (1 in 2) through the funnel, cautiously rinse the funnel with 20 mL of water, immediately close the clamp attached to the rubber tubing, then begin the distillation with steam gradually, and continue the distillation until the distillate measures 100 mL. Remove the absorption flask, rinse the end of the condenser tube with a small quantity of water, and titrate <2.50> the distillate and the rinsings with 0.1 mol/L sodium hydroxide VS until the color of the solution changes from red through light red-purple to light blue-green. Perform a blank determination.

Each mL of 0.05 mol/L sulfuric acid VS

$$= 11.81 \text{ mg of } \text{C}_6\text{H}_8\text{N}_2\text{O}_8$$

**Containers and storage** Containers—Tight containers.

Storage—Light-resistant, and in a cold place.

## Isosorbide Dinitrate Tablets

硝酸イソソルビド錠

Isosorbide Dinitrate Tablets contain not less than 93% and not more than 107% of the labeled amount of isosorbide dinitrate ( $\text{C}_6\text{H}_8\text{N}_2\text{O}_8$ ; 236.14).

**Method of preparation** Prepare as directed under Tablets, with Isosorbide Dinitrate.

**Identification** Weigh a quantity of powdered Isosorbide Dinitrate Tablets, equivalent to 0.1 g of Isosorbide Dinitrate according to the labeled amount, add 50 mL of diethyl ether, shake well, and filter. Measure 5 mL of the filtrate, evaporate to dryness cautiously, add 1 mL of water to the residue, and dissolve by adding 2 mL of sulfuric acid cautiously. After cooling, superimpose 3 mL of iron (II) sulfate TS, and allow to stand for 5 to 10 minutes: a brown ring is produced at the zone of contact.

**Purity** Free nitrate ion—Weigh accurately a quantity of powdered Isosorbide Dinitrate Tablets, equivalent to 0.05 g of Isosorbide Dinitrate according to the labeled amount, transfer to a separator, add 30 mL of toluene, shake thoroughly, extract with three 20-mL portions of water, and proceed as directed in Purity (3) under Isosorbide Dinitrate.

**Disintegration** <6.09> It meets the requirement.

For sublingual tablets, the time limit of the test is 2 minutes, and omit the use of the disk.

**Assay** Weigh accurately and powder not less than 20 Isosorbide Dinitrate Tablets. Weigh accurately a portion of the powder, equivalent to about 5 mg of isosorbide dinitrate ( $\text{C}_6\text{H}_8\text{N}_2\text{O}_8$ ), add exactly 50 mL of acetic acid (100), shake for 15 minutes, filter, and use this filtrate as the sample solution. Separately, weigh accurately about 90 mg of potassium nitrate, previously dried at 105°C for 4 hours, dissolve in 5 mL of water, and add acetic acid (100) to make exactly 100 mL. Measure exactly 10 mL of this solution, add acetic acid (100) to make exactly 100 mL, and use this solution as the standard solution. Measure exactly 2 mL each of the sample solution and standard solution, add exactly 2.5 mL of salicylic acid TS to each, shake well, allow to stand for 15 minutes, and add 10 mL of water. Make them alkaline with about 12 mL of a solution of sodium hydroxide (2 in 5) while cooling in an ice bath, and add water to make exactly 50 mL. Perform the test with these solutions as directed under Ultraviolet-visible Spectrophotometry <2.24>, using a solution, prepared with 2 mL of glacial acetic in the same manner, as the blank. Determine the absorbances,  $A_T$  and  $A_S$ , of the subsequent solutions of the sample solution and standard solution at 412 nm, respectively.

$$\begin{aligned} &\text{Amount (mg) of isosorbide dinitrate } (\text{C}_6\text{H}_8\text{N}_2\text{O}_8) \\ &= W_S \times (A_T/A_S) \times (1/20) \times 1.1678 \end{aligned}$$

$W_S$ : Amount (mg) of potassium nitrate

**Containers and storage** Containers—Tight containers.

## Japanese Encephalitis Vaccine

日本脳炎ワクチン

Japanese Encephalitis Vaccine is a liquid for injection containing inactivated Japanese encephalitis virus.

It conforms to the requirements of Japanese Encephalitis Vaccine in the Minimum Requirements for Biological Products.

**Description** Japanese Encephalitis Vaccine is a clear or a slightly whitish turbid and colorless liquid.

## Freeze-dried Japanese Encephalitis Vaccine

乾燥日本脳炎ワクチン


Freeze-dried Japanese Encephalitis Vaccine is a preparation for injection which is dissolved before use. It contains inactivated Japanese encephalitis virus.

It conforms to the requirements of Freeze-dried Japanese Encephalitis Vaccine in the Minimum Requirements for Biological Products.

**Description** Freeze-dried Japanese Encephalitis Vaccine is a clear or a slightly whitish turbid and colorless liquid on addition of solvent.

## Josamycin

ジョサマイシン



$\text{C}_{42}\text{H}_{69}\text{NO}_{15}$ : 827.99

(3*R*,4*R*,5*S*,6*R*,8*R*,9*R*,10*E*,12*E*,15*R*)-3-Acetoxy-5-[2,6-dideoxy-4-*O*-(3-methylbutanoyl)-3-*C*-methyl- $\alpha$ -L-ribo-hexopyranosyl-(1 $\rightarrow$ 4)-3,6-dideoxy-3-dimethylamino- $\beta$ -D-glucopyranosyloxy]-6-formylmethyl-9-hydroxy-4-methoxy-8-methylhexadeca-10,12-dien-15-olide [16846-24-5]

Josamycin is a macrolide substance having antibacterial activity produced by the growth of *Streptomyces narboensis* var. *josamyceticus*.

It contains not less than 900  $\mu\text{g}$  (potency) and not more than 1100  $\mu\text{g}$  (potency) per mg, calculated on the

dried basis. The potency of Josamycin is expressed as mass (potency) of josamycin ( $C_{42}H_{69}NO_{15}$ ).

**Description** Josamycin occurs as a white to yellowish white powder.

It is very soluble in methanol and in ethanol (99.5), and very slightly soluble in water.

**Identification (1)** Determine the absorption spectrum of a solution of Josamycin in methanol (1 in 100,000) as directed under Ultraviolet-visible Spectrophotometry <2.24>, and compare the spectrum with the Reference Spectrum or the spectrum of a solution of Josamycin Reference Standard prepared in the same manner as the sample solution: both spectra exhibit similar intensities of absorption at the same wavelengths.

(2) Dissolve 5 mg each of Josamycin and Josamycin Reference Standard in 1 mL of methanol, add diluted methanol (1 in 2) to make 100 mL, and use these solutions as the sample solution and the standard solution, respectively. Perform the test with 10  $\mu$ L each of the sample solution and standard solution as directed under Liquid Chromatography <2.01> according to the following conditions: the retention time of the main peak obtained from the sample solution is the same as that of the peak of josamycin from the standard solution.

**Operating conditions—**

Detector, column, column temperature, mobile phase, and flow rate: Proceed as directed in the operating conditions in the Purity (2).

**Purity (1)** Heavy metals <1.07>—Proceed with 1.0 g of Josamycin according to Method 2, and perform the test. Prepare the control solution with 3.0 mL of Standard Lead Solution (not more than 30 ppm).

(2) Related substances—Dissolve 50 mg of Josamycin in 5 mL of methanol, add diluted methanol (1 in 2) to make 50 mL, and use this solution as the sample solution. Perform the test with 10  $\mu$ L of the sample solution as directed under Liquid Chromatography <2.01> according to the following conditions. Determine each peak area by the automatic integration method, and calculate the amounts of josamycin and the related substances by the area percentage method: the amounts of the peaks other than josamycin are not more than 6%, and the total of these peaks is not more than 20%.

**Operating conditions—**

Detector: An ultraviolet absorption photometer (wavelength: 231 nm).

Column: A stainless steel column 4.6 mm in inside diameter and 5 cm in length, packed with octadecylsilanized silica gel for liquid chromatography (3  $\mu$ m in particle diameter).

Column temperature: A constant temperature of about 40°C

Mobile phase: Dissolve 119 g of sodium perchlorate monohydrate in water to make 1000 mL, and adjust the pH to 2.5 with 1 mol/L hydrochloric acid TS. To 600 mL of this solution add 400 mL of acetonitrile.

Flow rate: Adjust the flow rate so that the retention time of josamycin is about 10 minutes.

Time span of measurement: About 4 times as long as the retention time of josamycin beginning after the solvent peak.

**System suitability—**

Test for required detectability: Measure exactly 3 mL of the sample solution, add diluted methanol (1 in 2) to make

exactly 50 mL, and use this solution as the solution for system suitability test. Measure exactly 2 mL of the solution for system suitability test, and add diluted methanol (1 in 2) to make exactly 20 mL. Confirm that the peak area of josamycin obtained from 10  $\mu$ L of this solution is equivalent to 8 to 12% of that from 10  $\mu$ L of the solution for system suitability test.

System performance: Dissolve about 0.05 g of Josamycin in 50 mL of 0.1 mol/L potassium dihydrogen phosphate TS, pH 2.0, and allow to stand at 40°C for 3 hours. Adjust the pH of this solution to 6.8 to 7.2 with 2 mol/L sodium hydroxide TS, and add 50 mL of methanol. This solution contains both josamycin and josamycin S1. When the procedure is run with 10  $\mu$ L of this solution under the above operating conditions, the resolution between the peaks of josamycin S1, which relative retention time to josamycin is about 0.9, and josamycin is not less than 1.5.

System repeatability: When the test is repeated 6 times with 10  $\mu$ L of the solution for system suitability test under the above operating conditions, the relative standard deviation of the peak areas of josamycin is not more than 1.5%.

**Loss on drying <2.41>** Not more than 1.0% (0.5 g, in vacuum, phosphorus (V) oxide, 60°C, 3 hours).

**Residue on ignition <2.44>** Not more than 0.1% (1 g).

**Assay** Perform the test according to the Cylinder-plate method as directed under Microbial Assay for Antibiotics <4.02> according to the following conditions.

(i) Test organism—*Bacillus subtilis* ATCC 6633

(ii) Culture medium—Use the medium ii in 3) Medium for other organisms under (1) Agar media for seed and base layer. Adjust the pH of the medium so that it will be 7.9 to 8.1 after sterilization.

(iii) Standard solutions—Weigh accurately an amount of Josamycin Reference Standard, equivalent to about 30 mg (potency), dissolve in 5 mL of methanol, add water to make exactly 100 mL, and use this solution as the standard stock solution. Keep the standard stock solution at 5°C or below, and use within 7 days. Take exactly a suitable amount of the standard stock solution before use, add water to make solutions so that each mL contains 30  $\mu$ g (potency) and 7.5  $\mu$ g (potency), and use these solutions as the high concentration standard solution and low concentration standard solution, respectively.


(iv) Sample solutions—Weigh accurately an amount of Josamycin, equivalent to about 30 mg (potency), dissolve in 5 mL of methanol, and add water to make exactly 100 mL. Take exactly a suitable amount of this solution, add water to make solutions so that each mL contains 30  $\mu$ g (potency) and 7.5  $\mu$ g (potency), and use these solutions as the high concentration sample solution and low concentration sample solution, respectively.

**Containers and storage** Containers—Tight containers.

Storage—Light-resistant.

## Josamycin Propionate

ジョサマイシンプロピオン酸エステル



$C_{45}H_{73}NO_{16}$ : 884.06

(3*R*,4*R*,5*S*,6*R*,8*R*,9*R*,10*E*,12*E*,15*R*)-3-Acetoxy-5-[2,6-dideoxy-4-*O*-(3-methylbutanoyl)-3-*C*-methyl- $\alpha$ -*L*-ribo-hexopyranosyl-(1 $\rightarrow$ 4)-3,6-dideoxy-3-dimethylamino- $\beta$ -*D*-glucopyranosyloxy]-6-formylmethyl-4-methoxy-8-methyl-9-propanoyloxyhexadeca-10,12-dien-15-olide [16846-24-5, Josamycin]

Josamycin Propionate is a derivative of josamycin.

It contains not less than 843  $\mu$ g (potency) and not more than 1000  $\mu$ g (potency) per mg, calculated on the dried basis. The potency of Josamycin Propionate is expressed as mass (potency) of josamycin ( $C_{42}H_{69}NO_{15}$ : 827.99).

**Description** Josamycin Propionate occurs as a white to light yellowish white crystalline powder.

It is very soluble in acetonitrile, freely soluble in methanol and in ethanol (99.5), and practically insoluble in water.

**Identification** (1) Determine the absorption spectrum of a solution of Josamycin Propionate in methanol (1 in 100,000) as directed under Ultraviolet-visible Spectrophotometry <2.24>, and compare the spectrum with the Reference Spectrum or the spectrum of a solution of Josamycin Propionate Reference Standard prepared in the same manner as the sample solution: both spectra exhibit similar intensities of absorption at the same wavelengths.

(2) Dissolve 5 mg each of Josamycin Propionate and Josamycin Propionate Reference Standard in 50 mL of diluted acetonitrile (1 in 2), and use these solutions as the sample solution and the standard solution, respectively. Perform the test with 10  $\mu$ L each of the sample solution and standard solution as directed under Liquid Chromatography <2.01> according to the following conditions: the retention time of the peak of josamycin propionate obtained from the sample solution is the same with that of the peak of josamycin propionate from the standard solution.

**Operating conditions**—

Detector, column, column temperature, mobile phase, and flow rate: Proceed as directed in the operating conditions in the Purity (2).

**Purity** (1) Heavy metals <1.07>—Proceed with 1.0 g of Josamycin Propionate according to Method 2, and perform the test. Prepare the control solution with 3.0 mL of Stan-

dard Lead Solution (not more than 30 ppm).

(2) Related substances—Dissolve 50 mg of Josamycin Propionate in the mobile phase to make 50 mL, and use this solution as the sample solution. Perform the test with 10  $\mu$ L of the sample solution as directed under Liquid Chromatography <2.01> according to the following conditions. Determine each peak area by the automatic integration method, and calculate the amounts of each peak other than josamycin propionate by the area percentage method: the amount of any peak other than josamycin is not more than 6 %, and the total of these peaks is not more than 22%.

**Operating conditions**—

Detector: An ultraviolet absorption photometer (wavelength: 234 nm).

Column: A stainless steel column 6 mm in inside diameter and 15 cm in length, packed with octadecylsilanized silica gel for liquid chromatography (5  $\mu$ m in particle diameter).

Column temperature: A constant temperature of about 40°C.

Mobile phase: To 10 mL of triethylamine add water to make 1000 mL, and adjust the pH to 4.3 with acetic acid (100). To 500 mL of this solution add 500 mL of acetonitrile.

Flow rate: Adjust the flow rate so that the retention time of josamycin propionate is about 24 minutes.

Time span of measurement: About 3.5 times as long as the retention time of josamycin propionate beginning after the solvent peak.

**System suitability**—

Test for required detectability: Measure exactly 3 mL of the sample solution, add the mobile phase to make exactly 50 mL, and use this solution as the solution for system suitability test. Measure exactly 2 mL of the solution for system suitability test, and add the mobile phase to make exactly 20 mL. Confirm that the peak area of josamycin propionate obtained from 10  $\mu$ L of this solution is equivalent to 8 to 12% of that from 10  $\mu$ L of the solution for system suitability test.

System performance: Dissolve 5 mg of josamycin propionate and 2 mg of josamycin in 50 mL of the mobile phase. When the procedure is run with 10  $\mu$ L of this solution under the above operating conditions, josamycin and josamycin propionate are eluted in this order with the resolution between these peaks being not less than 25.

System repeatability: When the test is repeated 6 times with 10  $\mu$ L of the solution for system suitability test under the above operating conditions, the relative standard deviation of the peak area of josamycin propionate is not more than 1.5%.

**Loss on drying** <2.41> Not more than 1.0% (1 g, in vacuum, phosphorus (V) oxide, 60°C, 3 hours).

**Residue on ignition** <2.44> Not more than 0.1% (1 g).

**Assay** Perform the test according to the Cylinder-plate method as directed under Microbial Assay for Antibiotics <4.02> according to the following conditions.

(i) Test organism—*Bacillus subtilis* ATCC 6633

(ii) Culture medium—Use the medium ii in 3) Medium for other organisms under (1) Agar media for seed and base layer. Adjust the pH of the medium so that it will be 7.9 to 8.1 after sterilization.

(iii) Standard solutions—Weigh accurately an amount of Josamycin Propionate Reference Standard, equivalent to about 20 mg (potency), dissolve in 10 mL of methanol, add

1/15 mol/L phosphate buffer solution, pH 5.6 to make exactly 50 mL, and use this solution as the standard stock solution. Keep the standard stock solution at 5°C or below, and use within 3 days. Take exactly a suitable amount of the standard stock solution before use, add 1/15 mol/L phosphate buffer solution, pH 5.6 to make solutions so that each mL contains 80 µg (potency) and 20 µg (potency), and use these solutions as the high concentration standard solution and low concentration standard solution, respectively.


(iv) Sample solutions—Weigh accurately an amount of Josamycin Propionate, equivalent to about 20 mg (potency), dissolve in 10 mL of methanol, add 1/15 mol/L phosphate buffer solution, pH 5.6 to make exactly 50 mL. Take exactly a suitable amount of this solution, add 1/15 mol/L phosphate buffer solution, pH 5.6 to make solutions so that each mL contains 80 µg (potency) and 20 µg (potency), and use these solutions as the high concentration sample solution and low concentration sample solution, respectively.

**Containers and storage** Containers—Tight containers.

Storage—Light-resistant.

## Kainic Acid Hydrate

カイニン酸水和物



$C_{10}H_{15}NO_4 \cdot H_2O$ : 231.25

(2*S*,3*S*,4*S*)-3-(Carboxymethyl)-4-(1-methylethenyl)pyrrolidine-2-carboxylic acid monohydrate [487-79-6, anhydride]

Kainic Acid Hydrate, when dried, contains not less than 99.0% of kainic acid ( $C_{10}H_{15}NO_4$ : 213.23).

**Description** Kainic Acid Hydrate occurs as white crystals or crystalline powder. It is odorless, and has an acid taste.

It is sparingly soluble in water and in warm water, very slightly soluble in acetic acid (100) and in ethanol (95), and practically insoluble in diethyl ether.

It dissolves in dilute hydrochloric acid and in sodium hydroxide TS.

The pH of its solution (1 in 100) is between 2.8 and 3.5.

Melting point: about 252°C (with decomposition).

**Identification** (1) To 5 mL of a solution of Kainic Acid Hydrate (1 in 5000) add 1 mL of ninhydrin TS, and warm in a water bath at a temperature between 60°C and 70°C for 5 minutes: a yellow color is produced.

(2) Dissolve 0.05 g of Kainic Acid Hydrate in 5 mL of acetic acid (100), and add 0.5 mL of bromine TS: the color of bromine disappears immediately.

**Optical rotation** <2.49>  $[\alpha]_D^{20}$ :  $-13 - -17^\circ$  (0.5 g, water, 50 mL, 200 mm).

**Purity** (1) Clarity and color of solution—Dissolve 0.10 g of Kainic Acid Hydrate in 10 mL of water: the solution is clear and colorless.

(2) Chloride <1.03>—Take 0.5 g of Kainic Acid Hydrate

in a platinum crucible, dissolve in 5 mL of sodium carbonate TS, and evaporate on a water bath to dryness. Heat the crucible slowly at first, and then ignite until the sample is almost incinerated. After cooling, add 12 mL of dilute nitric acid to the residue, dissolve by warming, and filter. Wash the residue with 15 mL of water, combine the washings and the filtrate, and add water to make 50 mL. Perform the test using this solution as the test solution.

Control solution: Add 5 mL of sodium carbonate TS to 0.30 mL of 0.01 mol/L hydrochloric acid VS, and proceed as directed above (not more than 0.021%).

(3) Sulfate <1.14>—Dissolve 0.5 g of Kainic Acid Hydrate in 40 mL of water by warming. Cool, add 1 mL of dilute hydrochloric acid and water to make 50 mL, and perform the test using this solution as the test solution. Prepare the control solution with 0.30 mL of 0.005 mol/L sulfuric acid VS (not more than 0.028%).

(4) Ammonium—Take 0.25 g of Kainic Acid Hydrate, and perform the test. Prepare the control solution with 5.0 mL of Standard Ammonium Solution (not more than 0.02 %).

(5) Heavy metals <1.07>—Proceed with 1.0 g of Kainic Acid Hydrate according to Method 2, and perform the test. Prepare the control solution with 2.0 mL of Standard Lead Solution (not more than 20 ppm).

(6) Arsenic <1.11>—Dissolve 1.0 g of Kainic Acid Hydrate in 5 mL of dilute hydrochloric acid, and perform the test with this solution as the test solution (not more than 2 ppm).

(7) Amino acid and other imino acid—Dissolve 0.10 g of Kainic Acid Hydrate in 10 mL of water, and use this solution as the sample solution. Pipet 2 mL of this solution, and add water to make exactly 100 mL. Pipet 1 mL of this solution, add water to make exactly 20 mL, and use this solution as the standard solution. Perform the test as directed under Thin-layer Chromatography <2.03> with these solutions. Spot 10 µL each of the sample solution and standard solution on a plate of silica gel for thin-layer chromatography. Develop the plate with the supernatant liquid of a mixture of water, 1-butanol and acetic acid (100) (5:4:1) to a distance of about 10 cm, and air-dry the plate. Spray evenly a solution of ninhydrin in acetone (1 in 50) on the plate, and dry the plate at 80°C for 5 minutes: the spots other than the principal spot from the sample solution are not more intense than the spot from the standard solution.

**Loss on drying** <2.41> 6.5 – 8.5% (1 g, 105°C, 4 hours).

**Residue on ignition** <2.44> Not more than 0.1% (0.5 g).

**Assay** Weigh accurately about 0.4 g of Kainic Acid Hydrate, previously dried, and dissolve in 50 mL of warm water, cool and titrate <2.50> with 0.1 mol/L sodium hydroxide VS (indicator: 10 drops of bromothymol blue TS).

Each mL of 0.1 mol/L sodium hydroxide VS  
= 21.32 mg of  $C_{10}H_{15}NO_4$

**Containers and storage** Containers—Tight containers.

## Kainic Acid and Santonin Powder

カイニン酸・サントニン散

Kainic Acid and Santonin Powder contains not less than 9.0% and not more than 11.0% of santonin ( $C_{15}H_{18}O_3$ : 246.30), and not less than 1.80% and not more than 2.20% of kainic acid hydrate ( $C_{10}H_{15}NO_4 \cdot H_2O$ : 231.25).

### Method of preparation

|  |                       |
|--|-----------------------|
| Santonin                                 | 100 g                 |
| Kainic Acid Hydrate                      | 20 g                  |
| Starch, Lactose Hydrate or their mixture | a sufficient quantity |
| To make 1000 g                           |                       |

Prepare as directed under Powders, with the above ingredients.

**Description** Kainic Acid and Santonin Powder occurs as a white powder.

**Identification (1)** Shake 1 g of Kainic Acid and Santonin Powder with 10 mL of chloroform, and filter [use the residue for the test (2)]. Distil off the chloroform of the filtrate, and dissolve the residue in 2 mL of potassium hydroxide-ethanol TS: a red color is produced (santonin).

(2) Shake the residue obtained in (1) with 20 mL of warm water, filter, and to 1 mL of the filtrate add 10 mL of water and 1 mL of ninhydrin-L-ascorbic acid TS. Warm in a water bath between 60°C and 70°C for 5 minutes: a yellow color is produced (kainic acid).

**Assay (1)** Santonin—Weigh accurately about 0.25 g of Kainic Acid and Santonin Powder, add 20 mL of ethanol (95), shake thoroughly for 5 minutes, and filter. Wash the residue with three 10-mL portions of ethanol (95), and filter. Combine the filtrate and the washings, and add ethanol (95) to make exactly 50 mL. Pipet 2 mL of this solution, add ethanol (95) to make exactly 100 mL, and use this solution as the sample solution. Weigh accurately about 25 mg of santonin for assay, proceed in the same manner as the sample solution, and use the obtained solution as the standard solution. Determine the absorbances,  $A_T$  and  $A_S$ , of these solutions at 240 nm as directed under Ultraviolet-visible Spectrophotometry <2.24>.

$$\begin{aligned} &\text{Amount (mg) of santonin (C}_{15}\text{H}_{18}\text{O}_3\text{)} \\ &= W_S \times (A_T/A_S) \end{aligned}$$

$W_S$ : Amount (mg) of santonin for assay

(2) Kainic acid—Weigh accurately about 1.25 g of Kainic Acid and Santonin Powder, add 20 mL of diluted pyridine (1 in 10), shake thoroughly for 5 minutes, and filter. Wash the residue with three 10-mL portions of diluted pyridine (1 in 10), and filter. Combine the filtrate and the washings, and add diluted pyridine (1 in 10) to make exactly 50 mL. Pipet 2 mL of this solution, add diluted pyridine (1 in 10) to make exactly 25 mL, and use this solution as the sample solution. Dissolve about 25 mg of kainic acid for assay, previously dried at 105°C for 4 hours and accurately weighed, in diluted

pyridine (1 in 10) to make exactly 50 mL. Pipet 2 mL of this solution, add diluted pyridine (1 in 10) to make exactly 25 mL, and use this solution as the standard solution. Pipet 2 mL each of the sample solution and standard solution, add 2 mL of ninhydrin-L-ascorbic acid TS, and heat on a water bath for 30 minutes. After cooling immediately, shake vigorously for 2 minutes, add water to make exactly 20 mL, and allow to stand for 15 minutes. Determine the absorbances,  $A_T$  and  $A_S$ , of these solutions at 425 nm as directed under Ultraviolet-visible Spectrophotometry <2.24>, using the solution prepared in the same manner with 2 mL of diluted pyridine (1 in 10) instead of the sample solution as the blank.

$$\begin{aligned} &\text{Amount (mg) of kainic acid hydrate (C}_{10}\text{H}_{15}\text{NO}_4 \cdot \text{H}_2\text{O)} \\ &= W_S \times (A_T/A_S) \times 1.0845 \end{aligned}$$

$W_S$ : Amount (mg) of kainic acid for assay

**Containers and storage** Containers—Well-closed containers.

Storage—Light-resistant.

## Kallidinogenase

カリジノゲナーゼ

[9001-01-8]

Kallidinogenase is an enzyme obtained from healthy porcine pancreas, and has kinin-releasing activity based on cleavage of kininogen.

It contains not less than 25 Kallidinogenase Units per mg. Usually, it is diluted with Lactose Hydrate or the like.

Kallidinogenase contains not less than 90% and not more than 110% of the labeled Units.

**Description** Kallidinogenase occurs as a white to light brown powder. It is odorless or has a faint, characteristic odor.

It is freely soluble in water, and practically insoluble in ethanol (95) and in diethyl ether.

The pH of a solution of Kallidinogenase (1 in 300) is between 5.5 and 7.5.

**Identification (1)** Weigh accurately an appropriate amount of Kallidinogenase according to the labeled Units, and dissolve in 0.05 mol/L phosphate buffer solution, pH 7.0 to prepare a solution containing 10 Kallidinogenase Units per mL. Pipet 5 mL of this solution, and add exactly 1 mL of trypsin inhibitor TS and 0.05 mol/L phosphate buffer solution, pH 7.0 to make exactly 10 mL. Pipet 4 mL each of this solution into two separate test tubes, add exactly 1 mL each of aprotinin TS and 0.05 mol/L phosphate buffer solution, pH 7.0 separately to each test tube, allow them to stand at room temperature for 20 minutes, and use these solutions as the sample solutions 1 and 2. Separately, pipet 1 mL of trypsin inhibitor TS, and add 0.05 mol/L phosphate buffer solution, pH 7.0 to make exactly 10 mL. Pipet 4 mL each of this solution into two separate test tubes, add exactly 1 mL each of aprotinin TS and 0.05 mol/L phosphate buffer solution, pH 7.0 separately to each tube, allow them to stand at room temperature for 20 minutes, and use these solutions as the

sample solutions 3 and 4. Then, pipet 2.5 mL of substrate TS for kallidinogenase assay (1), previously warmed at  $30.0 \pm 0.5^\circ\text{C}$  for 5 minutes, place in a 10-mm cell, add exactly 0.5 mL of the sample solution 1 warmed at  $30.0 \pm 0.5^\circ\text{C}$  for 5 minutes, and start simultaneously a chronograph. Perform the test at  $30.0 \pm 0.5^\circ\text{C}$  as directed under Ultraviolet-visible Spectrophotometry <2.24> using water as the blank, and determine the absorbances at 405 nm,  $A_{1-2}$  and  $A_{1-6}$ , of this solution, after having allowed it to stand for exactly 2 and 6 minutes. Perform the same test with the sample solutions 2, 3 and 4, and determine the absorbances,  $A_{2-2}$ ,  $A_{2-6}$ ,  $A_{3-2}$ ,  $A_{3-6}$ ,  $A_{4-2}$  and  $A_{4-6}$ , of these solutions. Calculate  $I$  by using the following equation: the value of  $I$  does not exceed 0.2.

$$I = \frac{(A_{1-6} - A_{1-2}) - (A_{3-6} - A_{3-2})}{(A_{2-6} - A_{2-2}) - (A_{4-6} - A_{4-2})}$$

(2) Pipet 2.9 mL of substrate TS for kallidinogenase assay (2), previously warmed at  $30.0 \pm 0.5^\circ\text{C}$  for 5 minutes, place in a 10-mm cell, add exactly 0.1 mL of the sample solution obtained in the Assay, and start simultaneously a chronograph. Perform the test at  $30.0 \pm 0.5^\circ\text{C}$  as directed under Ultraviolet-visible Spectrophotometry <2.24>, and determine the change of the absorbance at 253 nm for 4 to 6 minutes. Separately, pipet 1 mL of trypsin inhibitor TS, and add 0.05 mol/L phosphate buffer solution, pH 7.0 to make exactly 10 mL. Add exactly 0.1 mL of this solution to exactly 2.9 mL of substrate TS for kallidinogenase assay (2), previously warmed at  $30.0 \pm 0.5^\circ\text{C}$  for 5 minutes, and use this solution as the blank. If the rate of change in the absorbance remains constant, determine the change of absorbance per 1 minute,  $A$ , and calculate  $R$  by using the following equation: the value of  $R$  is between 0.12 and 0.16.

$$R = (A/0.0383) \times \{1/(a - b)\}$$

$a$ : Amount (mg) of Kallidinogenase in 1 mL of the sample solution.

$b$ : Amount (Unit) of kallidinogenase in 1 mg of Kallidinogenase obtained in the Assay.

**Specific activity** Perform the test with Kallidinogenase as directed under Nitrogen Determination <1.08> to determine the nitrogen content, convert 1mg of nitrogen (N:14.01) into 6.25 mg of protein, and calculate the specific activity using the amount (Units) of Kallidinogenase obtained in the Assay: it is not less than 100 Kallidinogenase Units per 1mg of protein.

**Purity** (1) Fat—To 1.0 g of Kallidinogenase add 20 mL of diethyl ether, extract with occasional shaking for 30 minutes, and filter. Wash the residue with 10 mL of diethyl ether, combine the washing with the filtrate, evaporate the diethyl ether, and dry the residue at  $105^\circ\text{C}$  for 2 hours: the mass of the residue is not more than 1 mg.

(2) Kininase—

(i) Bradykinin solution: Weigh an appropriate amount of bradykinin, and dissolve in gelatin-phosphate buffer solution, pH 7.4 to prepare a solution containing  $0.200 \mu\text{g}$  of bradykinin per mL.

(ii) Kallidinogenase solution: Weigh accurately a suitable amount of Kallidinogenase according to the labeled unit, dissolve in gelatin-phosphate buffer solution, pH 7.4 to make a solution containing 1 unit of kallidinogenase per mL.

(iii) Sample solution: Pipet 0.5 mL of bradykinin solu-

tion, warm at  $30 \pm 0.5^\circ\text{C}$  for 5 minutes, then add exactly 0.5 mL of kallidinogenase solution previously warmed at  $30 \pm 0.5^\circ\text{C}$  for 5 minutes, and mix immediately. After allow this solution to stand at  $30 \pm 0.5^\circ\text{C}$  for exactly 150 seconds, add exactly 0.2 mL of a solution of trichloroacetic acid (1 in 5), and shake. Boil for 3 minutes, then cool in ice immediately, centrifuge, and allow to stand at a room temperature for 15 minutes. Pipet 0.5 mL of the supernatant liquid, add exactly 0.5 mL of gelatin-tris buffer solution, pH 8.0, and mix. Pipet 0.1 mL of this solution, add exactly 0.9 mL of trichloroacetic acid-gelatin-tris buffer solution, and mix. Pipet 0.2 mL of this solution, add exactly 0.6 mL of trichloroacetic acid-gelatin-tris buffer solution, shake, and use this solution as the sample solution.

(iv) Control solution: Proceed with 0.5 mL of gelatin-phosphate buffer solution, pH 7.4 as described in (iii), and use the solution so obtained as the control solution.

(v) Procedure: Add 0.1 mL of anti-bradykinin antibody TS to anti-rabbit antibody-coated wells of a 96-well microplate, shake, and allow to stand at a constant temperature of about  $25^\circ\text{C}$  for 1 hour. Remove the anti-bradykinin antibody TS, add 0.3 mL of phosphate buffer solution for microplate washing to the wells, then remove. Repeat this procedure 3 times, take off the washings thoroughly, then add  $100 \mu\text{L}$  each of the sample solution and control solution, and  $50 \mu\text{L}$  of gelatin-phosphate buffer solution, pH 7.0, shake, and allow to stand at a constant temperature of about  $25^\circ\text{C}$  for 1 hour. Then add  $50 \mu\text{L}$  of peroxidase-labeled bradykinin TS, shake, and allow to stand in a cold place for a night.

Take off the solution, add 0.3 mL of phosphate buffer solution for microplate washing, and remove. Repeat this procedure more 4 times, take off the washings thoroughly, add  $100 \mu\text{L}$  of substrate solution for peroxidase determination, and allow to stand at a constant temperature of about  $25^\circ\text{C}$  for exactly 30 minutes while protecting from light. Then add  $100 \mu\text{L}$  of diluted sulfuric acid (23 in 500), shake, and determine the absorbance at 490 – 492 nm.

Separately, dissolve a suitable amount of bradykinin in gelatin-phosphate buffer solution, pH 7.0 to make solutions containing exactly 100 ng, 25 ng, 6.25 ng, 1.56 ng, 0.39 ng and 0.098 ng of bradykinin per mL, and use these solutions as the standard solution (1), the standard solution (2), the standard solution (3), the standard solution (4), the standard solution (5) and the standard solution (6), respectively. Use 1 mL of gelatin-phosphate buffer solution, pH 7.0 as the standard solution (7). To each of the well add  $50 \mu\text{L}$  each of the standard solutions and  $100 \mu\text{L}$  of trichloroacetic acid-gelatin-tris buffer solution, and proceed in the same manner as for the sample solution and for the control solution.

Prepare the standard curve from the amounts of bradykinin in the standard solutions and their absorbances, and determine the amount of bradykinin,  $B_T$  (pg) and  $B_S$  (pg), of the sample solution and the control solution.

The absorbance is usually determined by using a spectrophotometer for microplate. Since the wells are used as the cell for absorbance determination, take care for dirt and scratch of the well. Light pass length of the well is changeable by the amount of the liquid, exact addition of the liquid is necessary.

(vi) Judgment: The value  $R$  calculated by the following equation is not less than 0.8.

$$R = (B_T/B_S)$$



(3) Trypsin-like substances—Pipet 4 mL of the sample stock solution prepared for the Assay, add exactly 1 mL of trypsin inhibitor TS and 0.05 mol/L phosphate buffer solution, pH 7.0 to make exactly 10 mL, and use this solution as the sample solution. Pipet 2.5 mL of substrate TS for kallidinogenase assay (1), previously warmed at  $30 \pm 0.5^\circ\text{C}$  for 5 minutes, place in a 10-mm cell, add exactly 0.5 mL of the sample solution, warmed at  $30 \pm 0.5^\circ\text{C}$  for 5 minutes, and start simultaneously a chronograph. Perform the test at  $30 \pm 0.5^\circ\text{C}$  as directed under Ultraviolet-visible Spectrophotometry <2.24> using water as the blank, and determine the absorbances at 405 nm,  $A_2$  and  $A_6$ , of this solution after having allowed it to stand for exactly 2 and 6 minutes. Separately, pipet 4 mL of the sample stock solution prepared for the Assay, add 0.05 mol/L phosphate buffer solution, pH 7.0 to make exactly 10 mL, and use this solution as the control solution. Perform the same test with the control solution, and determine the absorbances,  $A'_2$  and  $A'_6$ . Calculate  $T$  by using the following equation: the value of  $T$  does not exceed 0.05.

$$T = \{(A'_6 - A_2) - (A_6 - A'_2)/(A'_6 - A'_2)\}$$

(4) Protease—Weigh accurately an appropriate amount of Kallidinogenase according to the labeled Units, dissolve in 0.05 mol/L phosphate buffer solution, pH 7.0 to prepare a solution containing 1 Kallidinogenase Unit per mL, and use this solution as the sample solution. Pipet 1 mL of the sample solution, place in a test tube, and allow to stand at  $35 \pm 0.5^\circ\text{C}$  for 5 minutes. Then, pipet 5 mL of substrate TS for kallidinogenase assay (3), previously warmed to  $35 \pm 0.5^\circ\text{C}$ , add quickly to the sample solution in the test tube, and allow to stand at  $35 \pm 0.5^\circ\text{C}$  for exactly 20 minutes. Then add exactly 5 mL of trichloroacetic acid TS, shake well, allow to stand at room temperature for 1 hour, and filter through a membrane filter (5  $\mu\text{m}$  in pore size). Discard the first 3 mL of the filtrate, and determine the absorbance,  $A$ , of the subsequent filtrate at 280 nm within 2 hours as directed under Ultraviolet-visible Spectrophotometry <2.24>, using water as the blank. Separately, pipet 1 mL of the sample solution, add exactly 5 mL of trichloroacetic acid TS, shake well, and add exactly 5 mL of the substrate TS for kallidinogenase assay (3). Proceed in the same manner as described for the sample solution, and determine the absorbance,  $A_0$ , of this solution. Calculate the value of  $(A - A_0)$ : it is not more than 0.2.

**Loss on drying** <2.41> Not more than 2.0% (0.5 g, in vacuum, phosphorus (V) oxide, 4 hours).

**Residue on ignition** <2.44> Not more than 3% (0.5g, 650 – 750°C).

#### Kinin-releasing activity

(i) Kallidinogenase solution: Weigh accurately a suitable amount of Kallidinogenase, according to the labeled unit, dissolve in 0.02 mol/L phosphate buffer solution, pH 8.0 to make a solution containing 0.1 unit of kallidinogenase per mL. Perform this procedure by using glassware.

(ii) Sample solution: Pipet 0.5 mL of kininogen TS, warm at  $30 \pm 0.5^\circ\text{C}$  for 5 minutes, then add exactly 0.5 mL of kallidinogenase solution previously warmed at  $30 \pm 0.5^\circ\text{C}$  for 5 minutes, and mix immediately. After allow this solution to stand at  $30 \pm 0.5^\circ\text{C}$  for exactly 2 minutes, add exactly 0.2 mL of a solution of trichloroacetic acid (1 in 5), and shake. Boil for 3 minutes, then cool in ice immediately, centrifuge, and allow to stand at a room temperature for 15 minutes.

Pipet 0.5 mL of the supernatant liquid, add exactly 0.5 mL of gelatin-tris buffer solution, pH 8.0, and shake. Pipet 0.1 mL of this solution, add exactly 1.9 mL of trichloroacetic acid-gelatin-tris buffer solution, shake, and use this solution as the sample solution.

(iii) Procedure: Perform the test with the sample solution as directed in the Purity (2), and determine the amount,  $B$  (pg), of kinin per well. The kinin-releasing activity per 1 unit of Kallidinogenase calculated by the following equation is not less than 500 ng bradykinin equivalent/min/unit.

$$\text{Kinin-releasing activity (ng bradykinin equivalent/min/unit) per 1 unit of Kallidinogenase} = B \times 4.8$$

**Assay** Weigh accurately an appropriate amount of Kallidinogenase according to the labeled Units, dissolve in 0.05 mol/L phosphate buffer solution, pH 7.0 to prepare a solution containing about 10 Kallidinogenase Units per mL, and use this solution as the sample stock solution. Pipet 4 mL of the sample stock solution, add exactly 1 mL of trypsin inhibitor TS and 0.05 mol/L phosphate buffer solution, pH 7.0 to make exactly 10 mL, and use this solution as the sample solution. Pipet 2.5 mL of substrate TS for kallidinogenase assay (1), previously warmed at  $30 \pm 0.5^\circ\text{C}$  for 5 minutes, place in a 1-cm cell, add exactly 0.5 mL of the sample solution, warmed at  $30 \pm 0.5^\circ\text{C}$  for 5 minutes, and start simultaneously a chronograph. Perform the test at  $30 \pm 0.5^\circ\text{C}$  as directed under the Ultraviolet-visible Spectrophotometry <2.24> using water as the blank, and determine the absorbances at 405 nm,  $A_{T2}$  and  $A_{T6}$ , of this solution after allowing to stand for exactly 2 and 6 minutes. Separately, dissolve Kallidinogenase Reference Standard in 0.05 mol/L phosphate buffer solution, pH 7.0 to make a solution so that each mL contains exactly 10 Units, and use this solution as the standard stock solution. Pipet 4 mL of the stock solution, add exactly 1 mL of trypsin inhibitor TS and 0.05 mol/L phosphate buffer solution, pH 7.0 to make exactly 10 mL, and use this solution as the standard solution. Take exactly 0.5 mL of the standard solution, perform the test in the same manner as described for the sample solution, and determine the absorbances,  $A_{S2}$  and  $A_{S6}$ , of the solution after allowing to stand for exactly 2 and 6 minutes. Separately, take exactly 1 mL of the trypsin inhibitor TS, and add 0.05 mol/L phosphate buffer solution, pH 7.0 to make exactly 10 mL. Pipet 0.5 mL of this solution, perform the test in the same manner as described for the sample solution, and determine the absorbances,  $A_{O2}$  and  $A_{O6}$ , of the solution after allowing to stand for exactly 2 and 6 minutes.

Units per 1 mg of Kallidinogenase

$$= \frac{(A_{T6} - A_{T2}) - (A_{O6} - A_{O2})}{(A_{S6} - A_{S2}) - (A_{O6} - A_{O2})} \times \frac{W_S}{a} \times \frac{1}{b}$$

$W_S$ : Amount (Units) of Kallidinogenase Reference Standard


$a$ : Volume (mL) of the standard stock solution

$b$ : Amount (mg) of Kallidinogenase in 1 mL of the sample stock solution

**Containers and storage** Containers—Tight containers.

## Kanamycin Monosulfate

カナマイシン—硫酸塩



$C_{18}H_{36}N_4O_{11} \cdot H_2SO_4$ : 582.58

3-Amino-3-deoxy- $\alpha$ -D-glucopyranosyl-(1 $\rightarrow$ 6)-  
[6-amino-6-deoxy- $\alpha$ -D-glucopyranosyl-(1 $\rightarrow$ 4)]-2-deoxy-  
D-streptamine monosulfate [25389-94-0]

Kanamycin Monosulfate is the sulfate of an aminoglycoside substance having antibacterial activity produced by the growth of *Streptomyces kanamyceticus*.

It contains not less than 750  $\mu$ g (potency) and not more than 832  $\mu$ g (potency) per mg, calculated on the dried basis. The potency of Kanamycin Monosulfate is expressed as mass (potency) of kanamycin ( $C_{18}H_{36}N_4O_{11}$ : 484.50).

**Description** Kanamycin Monosulfate occurs as a white crystalline powder.

It is freely soluble in water, and practically insoluble in ethanol (99.5).

**Identification** (1) Dissolve 50 mg of Kanamycin Monosulfate in 3 mL of water, and add 6 mL of anthrone TS: a blue-purple color develops.

(2) Dissolve 20 mg each of Kanamycin Monosulfate and Kanamycin Monosulfate Reference Standard in 1 mL of water, and use these solutions as the sample solution and the standard solution. Perform the test with these solutions as directed under Thin-layer Chromatography <2.03>. Spot 5  $\mu$ L each of the sample solution and standard solution on a plate of silica gel for thin-layer chromatography. Develop the plate with the supernatant layer of a mixture of chloroform, ammonia solution (28) and methanol (2:1:1) to a distance of about 10 cm, and air-dry the plate. Spray evenly a solution of 0.2% ninhydrin-water saturated 1-butanol TS on the plate, and heat at 100°C for 10 minutes: the principal spots obtained from the sample solution and the standard solution show a purple-brown color and the same *R<sub>f</sub>* value.

(3) To a solution of Kanamycin Monosulfate (1 in 5) add 1 drop of barium chloride TS: a white precipitate is formed.

**Optical rotation** <2.49>  $[\alpha]_D^{20}$ : +112 – +123° (0.2 g calculated on the dried basis, water, 20 mL, 100 mm).

**Sulfuric acid** Weigh accurately about 0.25 g of Kanamycin Monosulfate, dissolve in 100 mL of water, adjust the pH to

11.0 with ammonia solution (28), add exactly 10 mL of 0.1 mol/L barium chloride VS, and titrate <2.50> with 0.1 mol/L disodium dihydrogen ethylenediamine tetraacetate VS until the color of the solution, blue-purple, disappears (indicator: 0.5 mg of phthalein purple). At a near of the end-point add 50 mL of ethanol (99.5). Perform a blank determination in the same manner. The amount of sulfuric acid ( $SO_4$ ) is not less than 15.0% and not more than 17.0%, calculated on the dried basis.

Each mL of 0.1 mol/L barium chloride VS  
= 9.606 mg of  $SO_4$

**Purity** (1) Heavy metals <1.07>—Proceed with 2.0 g of Kanamycin Monosulfate according to Method 4, and perform the test. Prepare the control solution with 2.0 mL of Standard Lead Solution (not more than 10 ppm).

(2) Arsenic <1.11>—Prepare the test solution with 2.0 g of Kanamycin Monosulfate according to Method 4, and perform the test (not more than 1 ppm).

(3) Related substances—Dissolve 0.30 g of Kanamycin Monosulfate in water to make exactly 10 mL, and use this solution as the sample solution. Separately, dissolve 45 mg of Kanamycin Monosulfate Reference Standard in water to make exactly 50 mL, and use this solution as the standard solution. Perform the test with these solutions as directed under Thin-layer Chromatography <2.03>. Spot 1  $\mu$ L each of the sample solution and standard solution on a plate of silica gel for thin-layer chromatography. Develop the plate with a solution of potassium dihydrogen phosphate (3 in 40) to a distance of about 10 cm, and air-dry the plate. Spray evenly a solution of ninhydrin in 1-butanol (1 in 100) on the plate, and heat at 110°C for 10 minutes: the spot other than the principal spot obtained from the sample solution is not more intense than the spot from the standard solution.

**Loss on drying** <2.41> Not more than 4.0% (5 g, reduced pressure not exceeding 0.67 kPa, 60°C, 3 hours).

**Residue on ignition** <2.44> Not more than 0.5% (1 g).

**Assay** Perform the test according to the Cylinder-plate method as directed under Microbial Assay for Antibiotics <4.02> according to the following conditions.

(i) Test organism—*Bacillus subtilis* ATCC 6633

(ii) Culture medium—Use the medium i in 1) Medium for test organism [5] under (1) Agar media for seed and base layer.

(iii) Standard solutions—Weigh accurately an amount of Kanamycin Monosulfate Reference Standard, previously dried, equivalent to about 20 mg (potency), dissolve in diluted phosphate buffer solution, pH 6.0 (1 in 2) to make exactly 50 mL, and use this solution as the standard stock solution. Keep the standard stock solution between 5 and 15°C and use within 30 days. Take exactly a suitable amount of the standard stock solution before use, add 0.1 mol/L phosphate buffer solution, pH 8.0 to make solutions so that each mL contains 20  $\mu$ g (potency) and 5  $\mu$ g (potency), and use these solutions as the high concentration standard solution and low concentration standard solution, respectively.


(iv) Sample solutions—Weigh accurately an amount of Kanamycin Monosulfate, equivalent to about 20 mg (potency), and dissolve in water to make exactly 50 mL. Take exactly a suitable amount of this solution, add 0.1 mol/L phosphate buffer solution, pH 8.0 to make solutions so that each

mL contains 20  $\mu\text{g}$  (potency) and 5  $\mu\text{g}$  (potency), and use these solutions as the high concentration sample solution and low concentration sample solution, respectively.

**Containers and storage** Containers—Well-closed containers.

## Kanamycin Sulfate

カナマイシン硫酸塩



$\text{C}_{18}\text{H}_{36}\text{N}_4\text{O}_{11} \cdot x\text{H}_2\text{SO}_4$

3-Amino-3-deoxy- $\alpha$ -D-glucopyranosyl-(1 $\rightarrow$ 6)-  
[6-amino-6-deoxy- $\alpha$ -D-glucopyranosyl-(1 $\rightarrow$ 4)]-2-deoxy-  
D-streptamine sulfate [133-92-6]

Kanamycin Sulfate is the sulfate of an aminoglycoside substance having antibacterial activity produced by the growth of *Streptomyces kanamyceticus*.

It contains not less than 690  $\mu\text{g}$  (potency) and not more than 740  $\mu\text{g}$  (potency) per mg, calculated on the dried basis. The potency of Kanamycin Sulfate is expressed as mass (potency) of kanamycin ( $\text{C}_{18}\text{H}_{36}\text{N}_4\text{O}_{11}$ : 484.50).

**Description** Kanamycin Sulfate occurs as a white to yellowish white powder.

It is very soluble in water, and practically insoluble in ethanol (99.5).

**Identification** (1) Dissolve 20 mg each of Kanamycin Sulfate and Kanamycin Monosulfate Reference Standard in 1 mL of water, and use these solutions as the sample solution and the standard solution. Perform the test with these solutions as directed under Thin-layer Chromatography <2.03>. Spot 5  $\mu\text{L}$  each of the sample solution and standard solution on a plate of silica gel for thin-layer chromatography. Develop the plate with a mixture of chloroform, ammonia solution (28) and methanol (2:1:1) to a distance of about 10 cm, and air-dry the plate. Spray evenly 0.2% ninhydrin-water saturated 1-butanol TS on the plate, and heat at 100°C for 10 minutes: the principal spots obtained from the sample solution and the standard solution show a purple-brown color and the same  $R_f$  value.

(2) A solution of Kanamycin Sulfate (1 in 10) responds to the Qualitative Test <1.09> (1) for sulfate.

**Optical rotation** <2.49>  $[\alpha]_D^{20}$ : +103 – +115° (0.5 g calculated on the dried basis, water, 50 mL, 100 mm).

**pH** <2.54> The pH of a solution obtained by dissolving 1.0 g

of Kanamycin Sulfate in 20 mL of water is between 6.0 and 7.5.

**Purity** (1) Clarity and color of solution—Dissolve 1.5 g of Kanamycin Sulfate in 5 mL of water: the solution is clear and colorless to pale yellow.

(2) Heavy metals <1.07>—Proceed with 1.0 g of Kanamycin Sulfate according to Method 4, and perform the test. Prepare the control solution with 3.0 mL of Standard Lead Solution (not more than 30 ppm).

(3) Arsenic <1.11>—Prepare the test solution with 2.0 g of Kanamycin Sulfate according to Method 3, and perform the test (not more than 1 ppm).

(4) Related substances—Dissolve 0.30 g of Kanamycin Sulfate in water to make exactly 10 mL, and use this solution as the sample solution. Separately, dissolve 9.0 mg of Kanamycin Monosulfate Reference Standard in water to make exactly 10 mL, and use this solution as the standard solution. Perform the test with these solutions as directed under Thin-layer Chromatography <2.03>. Spot 1  $\mu\text{L}$  each of the sample solution and standard solution on a plate of silica gel for thin-layer chromatography. Develop the plate with a solution of potassium dihydrogen phosphate (3 in 40) to a distance of about 10 cm, and air-dry the plate. Spray evenly a solution of ninhydrin in 1-butanol (1 in 100) on the plate, and heat at 110°C for 10 minutes: the spot other than the principal spot obtained from the sample solution is not more intense than the spot from the standard solution.

**Loss on drying** <2.41> Not more than 5.0% (0.5 g, reduced pressure not exceeding 0.67 kPa, 60°C, 3 hours).

**Assay** Perform the test according to the Cylinder-plate method as directed under Microbial Assay for Antibiotics <4.02> according to the following conditions.

(i) Test organism—*Bacillus subtilis* ATCC 6633

(ii) Culture medium—Use the medium i in 1) Medium for test organism [5] under (1) Agar media for seed and base layer having pH 7.8 to 8.0 after sterilization.

(iii) Standard solutions—Weigh accurately an amount of Kanamycin Monosulfate Reference Standard, previously dried, equivalent to about 20 mg (potency), dissolve in diluted phosphate buffer solution, pH 6.0 (1 in 2) to make exactly 50 mL, and use this solution as the standard stock solution. Keep the standard stock solution at 5 to 15°C and use within 30 days. Take exactly a suitable amount of the standard stock solution before use, add 0.1 mol/L phosphate buffer solution, pH 8.0 to make solutions so that each mL contains 20  $\mu\text{g}$  (potency) and 5  $\mu\text{g}$  (potency), and use these solutions as the high concentration standard solution and low concentration standard solution, respectively.

(iv) Sample solutions—Weigh accurately an amount of Kanamycin Sulfate, equivalent to about 20 mg (potency), and dissolve in water to make exactly 50 mL. Take exactly a suitable amount of this solution, add 0.1 mol/L phosphate buffer solution, pH 8.0 to make solutions so that each mL contains 20  $\mu\text{g}$  (potency) and 5  $\mu\text{g}$  (potency), and use these solutions as the high concentration sample solution and low concentration sample solution, respectively.

**Containers and storage** Containers—Tight containers.

## Kaolin

カオリン

Kaolin is a native, hydrous aluminum silicate.

**Description** Kaolin occurs as white or nearly white, fragmentary masses or powder. It has a slightly clay-like odor.

It is practically insoluble in water, in ethanol (99.5) and in diethyl ether.

It is insoluble in dilute hydrochloric acid and in sodium hydroxide TS.

When moistened with water, it darkens and becomes plastic.

**Identification** (1) Heat 1 g of Kaolin with 10 mL of water and 5 mL of sulfuric acid in a porcelain dish, and evaporate the mixture nearly to dryness. Cool, add 20 mL of water, boil for 2 to 3 minutes, and filter: the color of the residue is gray.

(2) The filtrate obtained in (1) responds to the Qualitative Tests <1.09> (1), (2) and (4) for aluminum salt.

**Purity** (1) Acid or alkali—Add 25 mL of water to 1.0 g of Kaolin, agitate thoroughly, and filter: the pH <2.54> of the filtrate is between 4.0 and 7.5.

(2) Acid-soluble substances—Add 20 mL of dilute hydrochloric acid to 1.0 g of Kaolin, agitate for 15 minutes, and filter. Evaporate 10 mL of the filtrate to dryness, and heat strongly between 450°C and 550°C to constant mass: the mass of the ignited residue is not more than 0.010 g.

(3) Carbonate—Stir 1.0 g of Kaolin with 5 mL of water, then add 10 mL of diluted sulfuric acid (1 in 2): no effervescence occurs.

(4) Heavy metals <1.07>—Boil 1.5 g of Kaolin gently with 50 mL of water and 5 mL of hydrochloric acid for 20 minutes with frequent agitation, cool, centrifuge, and separate the supernatant liquid. Wash the precipitate twice with 10 mL of water, centrifuge each time, and combine the supernatant liquid and the washings. Add dropwise ammonia solution (28) to this solution until a slight precipitate occurs, then add dilute hydrochloric acid dropwise while agitating strongly to complete solution. Add 0.45 g of hydroxylammonium chloride, and heat. Cool, add 0.45 g of sodium acetate trihydrate and 6 mL of dilute acetic acid, filter if necessary, and wash with 10 mL of water. Combine the filtrate and the washings, and add water to make 150 mL. Perform the test using 50 mL of this solution as the test solution. To 2.5 mL of Standard Lead Solution add 0.15 g of hydroxylammonium chloride, 0.15 g of sodium acetate trihydrate, 2 mL of acetic acid (31) and water to make 50 mL, and use this solution as the control solution (not more than 50 ppm).

(5) Iron <1.10>—Add 10 mL of dilute hydrochloric acid to 40 mg of Kaolin, and heat for 10 minutes with shaking in a water bath. After cooling, add 0.5 g of L-tartaric acid, dissolve with shaking, prepare the test solution with this solution according to Method 2, and perform the test according to Method B. Prepare the control solution with 2.0 mL of Standard Iron Solution (not more than 500 ppm).

(6) Arsenic <1.11>—Add 5 mL of water and 1 mL of sulfuric acid to 1.0 g of Kaolin, and heat on a sand bath until white fumes begin to evolve. Cool, and add water to make 5

mL. Perform the test with this solution as the test solution (not more than 2 ppm).

(7) Foreign matter—Place 5 g of Kaolin in a beaker, add 100 mL of water, stir, and decant to leave sand. Repeat this procedure several times with 100-mL portions of water: no sandy residue remains.


**Loss on ignition** <2.43> Not more than 15.0% (1 g, 600°C, 5 hours).

**Plasticity** Add 7.5 mL of water to 5.0 g of Kaolin, and agitate thoroughly: the resultant mass has no remarkable fluidity.

**Containers and storage** Containers—Well-closed containers.

## Ketamine Hydrochloride

ケタミン塩酸塩



$C_{13}H_{16}ClNO \cdot HCl$ : 274.19

(2*RS*)-2-(2-Chlorophenyl)-2-(methylamino)cyclohexanone monohydrochloride [1867-66-9]

Ketamine Hydrochloride, when dried, contains not less than 99.0% of  $C_{13}H_{16}ClNO \cdot HCl$ .

**Description** Ketamine Hydrochloride occurs as white crystals or crystalline powder.

It is very soluble in formic acid, freely soluble in water and in methanol, sparingly soluble in ethanol (95) and in acetic acid (100), and practically insoluble in acetic anhydride and in diethyl ether.

A solution of Ketamine Hydrochloride (1 in 10) shows no optical rotation.

Melting point: about 258°C (with decomposition).

**Identification** (1) Determine the absorption spectrum of a solution of Ketamine Hydrochloride in 0.1 mol/L hydrochloric acid TS (1 in 3000) as directed under Ultraviolet-visible Spectrophotometry <2.24>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wavelengths.

(2) Determine the infrared absorption spectrum of Ketamine Hydrochloride, previously dried, as directed in the potassium bromide disk method under Infrared Spectrophotometry <2.25>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wave numbers.

(3) A solution of Ketamine Hydrochloride (1 in 10) responds to the Qualitative Tests <1.09> (2) for chloride.

**Absorbance** <2.24>  $E_{1\text{cm}}^{1\%}$  (269 nm): 22.0 – 24.5 (after drying, 0.03 g, 0.1 mol/L hydrochloric acid TS, 100 mL).

**pH** <2.54> Dissolve 1.0 g of Ketamine Hydrochloride in 10 mL of freshly boiled and cooled water: the pH of the solution is between 3.5 and 4.5.

**Purity** (1) Clarity and color of solution—Dissolve 1.0 g of

Ketamine Hydrochloride in 5 mL of water: the solution is clear and colorless.

(2) Heavy metals <1.07>—Proceed with 1.0 g of Ketamine Hydrochloride according to Method 1, and perform the test. Prepare the control solution with 2.0 mL of Standard Lead Solution (not more than 20 ppm).

(3) Arsenic <1.11>—Prepare the test solution with 1.0 g of Ketamine Hydrochloride, according to Method 1, and perform the test (not more than 2 ppm).

(4) Related substances—Dissolve 0.5 g of Ketamine Hydrochloride in 10 mL of methanol and use this solution as the sample solution. Pipet 1 mL of the sample solution, add methanol to make exactly 200 mL, and use this solution as the standard solution. Perform the test with these solutions as directed under Thin-layer Chromatography <2.03>. Spot 2  $\mu$ L each of the sample solution and standard solution on a plate of silica gel for thin-layer chromatography. Develop the plate with a mixture of cyclohexane and isopropylamine (49:1) to a distance of about 10 cm, and air-dry the plate. Spray evenly Dragendorff's TS for spraying on the plate, dry the plate, and then spray evenly hydrogen peroxide TS: the spots other than the principal spot from the sample solution is not more intense than the spot from the standard solution.

**Loss on drying** <2.41> Not more than 0.5% (1 g, 105°C, 3 hours).

**Residue on ignition** <2.44> Not more than 0.1% (1 g).


**Assay** Weigh accurately about 0.5 g of Ketamine Hydrochloride, previously dried, dissolve in 1 mL of formic acid, add 70 mL of a mixture of acetic anhydride and acetic acid (100) (6:1), and titrate <2.50> with 0.1 mol/L perchloric acid VS (potentiometric titration). Perform a blank determination, and make any necessary correction.

Each mL of 0.1 mol/L perchloric acid VS  
= 27.42 mg of  $C_{13}H_{16}ClNO.HCl$

**Containers and storage** Containers—Tight containers.

## Ketoprofen

ケトプロフェン



$C_{16}H_{14}O_3$ : 254.28

(2*RS*)-2-(3-Benzoylphenyl)propanoic acid [22071-15-4]

Ketoprofen, when dried, contains not less than 99.0% and not more than 100.5% of  $C_{16}H_{14}O_3$ .

**Description** Ketoprofen occurs as a white, crystalline powder.

It is very soluble in methanol, freely soluble in ethanol (95) and in acetone, and practically insoluble in water.

A solution of Ketoprofen in ethanol (99.5) (1 in 100) shows no optical rotation.

It is colored to pale yellow by light.

**Identification** (1) Determine the absorption spectrum of a solution of Ketoprofen in methanol (1 in 200,000) as directed under Ultraviolet-visible Spectrophotometry <2.24>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wavelengths.

(2) Determine the infrared absorption spectrum of Ketoprofen, previously dried, as directed in the potassium bromide disk method under Infrared Spectrophotometry <2.25>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wave numbers.

**Melting point** <2.60> 94 – 97°C

**Purity** (1) Clarity and color of solution—Dissolve 1.0 g of Ketoprofen in 10 mL of acetone: the solution is clear, and has no more color than the following control solution.

Control solution: To a mixture of 0.6 mL of Cobalt (II) Chloride Colorimetric Stock Solution and 2.4 mL of Iron (III) Chloride Colorimetric Stock Solution add diluted hydrochloric acid (1 in 10) to make 10 mL. To 5.0 mL of this solution add diluted hydrochloric acid (1 in 10) to make 100 mL.

(2) Heavy metals <1.07>—Proceed with 2.0 g of Ketoprofen according to Method 2, and perform the test. Prepare the control solution with 2.0 mL of Standard Lead Solution (not more than 10 ppm).

(3) Related substances—Conduct this procedure with a minimum of exposure to light, using light-resistant vessels. Dissolve 20 mg of Ketoprofen in 20 mL of the mobile phase, and use this solution as the sample solution. Pipet 1 mL of the sample solution, and add the mobile phase to make exactly 50 mL. Pipet 1 mL of this solution, add the mobile phase to make exactly 10 mL, and use this solution as the standard solution. Perform the test with exactly 20  $\mu$ L each of the sample solution and standard solution as directed under Liquid Chromatography <2.01> according to the following conditions, and determine each peak area by the automatic integration method: the areas of the peaks, having the relative retention time of about 1.5 and about 0.3 with respect to ketoprofen, are not larger than 4.5 times and not larger than 2 times the peak area of ketoprofen from the standard solution, respectively, the area of the peak other than ketoprofen and the peaks mentioned above is not larger than the peak area of ketoprofen from the standard solution, and the total area of these peaks is not larger than 2 times the peak area of ketoprofen from the standard solution.

**Operating conditions**—

Detector: An ultraviolet absorption photometer (wavelength: 233 nm).

Column: A stainless steel column 4.6 mm in inside diameter and 15 cm in length, packed with octadecylsilanized silica gel for liquid chromatography (5  $\mu$ m in particle diameter).

Column temperature: A constant temperature of about 25°C.

Mobile phase: Dissolve 68.0 g of potassium dihydrogen phosphate in water to make 1000 mL, and adjust the pH to 3.5 with phosphoric acid. To 20 mL of this solution add 430 mL of acetonitrile and 550 mL of water.

Flow rate: Adjust the flow rate so that the retention time of

ketoprofen is about 7 minutes.

Time span of measurement: About 7 times as long as the retention time of ketoprofen.

**System suitability**—

Test for required detectability: To exactly 1 mL of the standard solution add the mobile phase to make exactly 10 mL. Confirm that the peak area of ketoprofen obtained with 20  $\mu$ L of this solution is equivalent to 9 to 11% of that with 20  $\mu$ L of the standard solution.

System performance: When the procedure is run with 20  $\mu$ L of the standard solution under the above operating conditions, the number of theoretical plates and the symmetry factor of the peak of ketoprofen are not less than 8000 and not more than 1.5, respectively.

System repeatability: When the test is repeated 6 times with 20  $\mu$ L of the standard solution under the above operating conditions, the relative standard deviation of the peak area of ketoprofen is not more than 2.0%.

**Loss on drying** <2.41> Not more than 0.5% (0.5 g, in vacuum, 60°C, 24 hours).

**Residue on ignition** <2.44> Not more than 0.1% (1 g).


**Assay** Weigh accurately about 0.3 g of Ketoprofen, previously dried, dissolve in 25 mL of ethanol (95), add 25 mL of water, and titrate <2.50> with 0.1 mol/L sodium hydroxide VS (potentiometric titration). Perform a blank determination, and make any necessary correction.

Each mL of 0.1 mol/L sodium hydroxide VS  
= 25.43 mg of  $C_{16}H_{14}O_3$

**Containers and storage** Containers—Tight containers.  
Storage—Light-resistant.

## Ketotifen Fumarate

ケトチフェン fumarate



$C_{19}H_{19}NOS \cdot C_4H_4O_4$ : 425.50

4-(1-Methylpiperidin-4-ylidene)-4*H*-benzo[4,5]cyclohepta[1,2-*b*]thiophen-10(9*H*)-one monofumarate [34580-14-8]

Ketotifen Fumarate, when dried, contains not less than 99.0% and not more than 101.0% of ketotifen fumarate ( $C_{19}H_{19}NOS \cdot C_4H_4O_4$ ).

**Description** Ketotifen Fumarate occurs as a white to light yellowish white crystalline powder.

It is sparingly soluble in methanol and in acetic acid (100), and slightly soluble in water, in ethanol (99.5) and in acetic anhydride.

Melting point: about 190°C (with decomposition).

**Identification** (1) Prepare the test solution with 0.03 g of

Ketotifen Fumarate as directed under Oxygen Flask Combustion Method using 20 mL of water as the absorbing liquid: the test solution responds to the Qualitative Tests <1.09> for sulfate.

(2) Determine the absorption spectrum of a solution of Ketotifen Fumarate in methanol (1 in 50,000) as directed under Ultraviolet-visible Spectrophotometry <2.24>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wavelengths.

(3) Determine the infrared absorption spectrum of Ketotifen Fumarate, previously dried, as directed in the potassium bromide disk method under Infrared Spectrophotometry <2.25>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wave numbers.

**Purity** (1) Chloride <1.03>—Dissolve 0.6 g of Ketotifen Fumarate in 2.5 mL of sodium carbonate TS in a crucible, heat on a water bath to dryness, and ignite at about 500°C. Dissolve the residue in 15 mL of water, filter if necessary, neutralize with diluted nitric acid (3 in 10), and add 6 mL of dilute nitric acid and water to make 50 mL. Perform the test using this solution as the test solution. Prepare the control solution as follows: To 0.25 mL of 0.01 mol/L hydrochloric acid VS add 2.5 mL of sodium carbonate TS, the used amount of diluted nitric acid (3 in 10) for the neutralization, 6 mL of dilute nitric acid and water to make 50 mL (not more than 0.015%).

(2) Heavy metals <1.07>—Proceed with 1.0 g of Ketotifen Fumarate according to Method 2, and perform the test. Prepare the control solution with 2.0 mL of Standard Lead Solution (not more than 20 ppm).

(3) Related substances—Dissolve 0.10 g of Ketotifen Fumarate in 10 mL of a mixture of methanol and ammonia TS (99:1), and use this solution as the sample solution. Pipet 1 mL of the sample solution, and add a mixture of methanol and ammonia TS (99:1) to make exactly 25 mL. Pipet 1 mL of this solution, add a mixture of methanol and ammonia TS (99:1) to make exactly 20 mL, and use this solution as the standard solution. Perform the test with these solutions as directed under Thin-layer Chromatography <2.03>. Spot 10  $\mu$ L each of the sample solution and standard solution on a plate of silica gel for thin-layer chromatography. Develop the plate with a mixture of acetonitrile, water and ammonia solution (28) (90:10:1) to a distance of about 15 cm, and air-dry the plate. Spray evenly Dragendorff's TS for spraying and then hydrogen peroxide TS on the plate: the number of the spot other than the principal spot obtained from the sample solution is not more than four, and they are not more intense than the spot from the standard solution.

**Loss on drying** <2.41> Not more than 0.5% (1 g, 105°C, 4 hours).

**Residue on ignition** <2.44> Not more than 0.1% (1 g).

**Assay** Weigh accurately about 0.35 g of Ketotifen Fumarate, previously dried, dissolve in 80 mL of a mixture of acetic anhydride and acetic acid (100) (7:3), and titrate <2.50> with 0.1 mol/L perchloric acid VS (potentiometric titration). Perform a blank determination, and make any necessary correction.

Each mL of 0.1 mol/L perchloric acid VS


= 42.55 mg of  $C_{19}H_{19}NOS \cdot C_4H_4O_4$

**Containers and storage** Containers—Tight containers.

## Kitasamycin

### Leucomycin

キタマイシン



(Leucomycins A<sub>1</sub>, A<sub>5</sub>, A<sub>7</sub>, A<sub>9</sub> and A<sub>13</sub>)  
(3*R*,4*R*,5*S*,6*R*,8*R*,9*R*,10*E*,12*E*,15*R*)-5-[4-*O*-Acyl-2,6-dideoxy-3-*C*-methyl- $\alpha$ -*L*-ribo-hexopyranosyl-(1 $\rightarrow$ 4)-3,6-dideoxy-3-dimethylamino- $\beta$ -*D*-glucopyranosyloxy]-6-formylmethyl-3,9-dihydroxy-4-methoxy-8-methylhexadeca-10,12-dien-15-olide

Leucomycin A<sub>1</sub>: acyl = 3-methylbutanoyl  
Leucomycin A<sub>5</sub>: acyl = butanoyl  
Leucomycin A<sub>7</sub>: acyl = propanoyl  
Leucomycin A<sub>9</sub>: acyl = acetyl  
Leucomycin A<sub>13</sub>: acyl = hexanoyl

(Leucomycins A<sub>3</sub>, A<sub>4</sub>, A<sub>6</sub> and A<sub>8</sub>)  
(3*R*,4*R*,5*S*,6*R*,8*R*,9*R*,10*E*,12*E*,15*R*)-3-Acetoxy-5-[4-*O*-acyl-2,6-dideoxy-3-*C*-methyl- $\alpha$ -*L*-ribo-hexopyranosyl-(1 $\rightarrow$ 4)-3,6-dideoxy-3-dimethylamino- $\beta$ -*D*-glucopyranosyloxy]-6-formylmethyl-9-hydroxy-4-methoxy-8-methylhexadeca-10,12-dien-15-olide

Leucomycin A<sub>3</sub>: acyl = 3-methylbutanoyl  
Leucomycin A<sub>4</sub>: acyl = butanoyl  
Leucomycin A<sub>6</sub>: acyl = propanoyl  
Leucomycin A<sub>8</sub>: acyl = acetyl

[1392-21-8, Kitasamycin]

Kitasamycin is a mixture of macrolide substances having antibacterial activity produced by the growth of *Streptomyces kitasatoensis*.

It contains not less than 1450  $\mu$ g (potency) and not more than 1700  $\mu$ g (potency) per mg, calculated on the anhydrous basis. The potency of Kitasamycin is expressed as mass (potency) of kitasamycin corresponding to the mass of leucomycin A<sub>5</sub> ( $C_{39}H_{65}NO_{14}$ : 771.93). One mg (potency) of kitasamycin is equivalent to 0.530 mg of leucomycin A<sub>5</sub> ( $C_{39}H_{65}NO_{14}$ ).

**Description** Kitasamycin occurs as a white to light yellow-white powder.

It is very soluble in acetonitrile, in methanol and in ethanol (95), and practically insoluble in water.

**Identification** Determine the absorption spectrum of a solution of Kitasamycin in methanol (1 in 40,000) as directed under Ultraviolet-visible Spectrophotometry <2.24>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wavelengths.

**Content ratio of the active principle** Dissolve 0.02 g of Kitasamycin in diluted acetonitrile (1 in 2) to make 20 mL, and use this solution as the sample solution. Perform the test with 5  $\mu$ L of the sample solution as directed under Liquid Chromatography <2.01> according to the following conditions, and measure each peak area by the automatic integration method. Calculate the amounts of leucomycin A<sub>5</sub>, leucomycin A<sub>4</sub> and leucomycin A<sub>1</sub> by the area percentage method: the amounts of leucomycin A<sub>5</sub>, leucomycin A<sub>4</sub> and leucomycin A<sub>1</sub> are 40 to 70%, 5 to 25% and 3 to 12%, respectively. Relative retention times of leucomycin A<sub>4</sub> and leucomycin A<sub>1</sub> to that of leucomycin A<sub>5</sub> are 1.2 and 1.5, respectively.

**Operating conditions—**

**Detector:** An ultraviolet absorption photometer (wavelength: 232 nm).

**Column:** A stainless steel column 4.0 mm in inside diameter and 15 cm in length, packed with octylsilanized silica gel for liquid chromatography (5  $\mu$ m in particle diameter).

**Column temperature:** A constant temperature of about 40°C.

**Mobile phase:** To a volume of a solution of ammonium acetate (77 in 500) add diluted phosphoric acid (1 in 150) to adjust to pH 5.5. To 370 mL of this solution add 580 mL of methanol and 50 mL of acetonitrile.

**Flow rate:** Adjust the flow rate so that the retention time of leucomycin A<sub>5</sub> is about 8 minutes.

**Time span of measurement:** About 3 times as long as the retention time of leucomycin A<sub>5</sub>.

**System suitability—**

**System performance:** Dissolve about 20 mg each of Leucomycin A<sub>5</sub> Reference Standard and Josamycin Reference Standard in 20 mL of diluted acetonitrile (1 in 2). When the procedure is run with 5  $\mu$ L of this solution under the above operating conditions, leucomycin A<sub>5</sub> and josamycin are eluted in this order with the resolution between these peaks being not less than 5.

**System repeatability:** When the test is repeated 6 times with 5  $\mu$ L of the sample solution under the above operating conditions, the relative standard deviation of the peak area of leu-

comycin A<sub>5</sub> is not more than 1.0%.

**Water** <2.48> Not more than 3.0% (0.1 g, volumetric titration, direct titration).

**Assay** Perform the test according to the Cylinder-plate method as directed under Microbial Assay for Antibiotics <4.02> according to the following conditions.

(i) Test organism—*Bacillus subtilis* ATCC 6633

(ii) Culture medium—Use the medium i in 1) Medium for test organism [5] under (1) Agar media for seed and base layer.

(iii) Standard solutions—Weigh accurately an amount of Leucomycin A<sub>5</sub> Reference Standard equivalent to about 30 mg (potency), dissolve in 10 mL of methanol, add water to make exactly 100 mL, and use this solution as the standard stock solution. Keep the standard stock solution at 5°C or below and use within 3 days. Take exactly a suitable amount of the standard stock solution before use, add phosphate buffer solution, pH 8.0 to make solutions so that each mL contains 30 μg (potency) and 7.5 μg (potency), and use these solutions as the high concentration standard solution and low concentration standard solution, respectively.



(iv) Sample solutions—Weigh accurately an amount of Kitasamycin equivalent to about 0.03 g (potency), dissolve in 10 mL of methanol, and add water to make exactly 100 mL. Take exactly a suitable amount of the solution, add phosphate buffer solution, pH 8.0 to make solutions so that each mL contains 30 μg (potency) and 7.5 μg (potency), and use these solutions as the high concentration sample solution and low concentration sample solution, respectively.

**Containers and storage** Containers—Tight containers.

## Kitasamycin Acetate

### Leucomycin Acetate

キタマイシン酢酸エステル



(3*R*,4*R*,5*S*,6*R*,8*R*,9*R*,10*E*,12*E*,15*R*)-3,9-Diacetoxy-5-[4-*O*-acyl-2,6-dideoxy-3-*C*-methyl- $\alpha$ -*L*-ribo-hexopyranosyl-(1→4)-2-*O*-acetyl-3,6-dideoxy-3-dimethylamino- $\beta$ -*D*-glucopyranosyloxy]-6-formylmethyl-4-methoxy-8-methylhexadeca-10,12-dien-15-olide

Leucomycin A<sub>1</sub> and A<sub>3</sub> Acetates: acyl = 3-methylbutanoyl

Leucomycin A<sub>4</sub> and A<sub>5</sub> Acetates: acyl = butanoyl

Leucomycin A<sub>6</sub> and A<sub>7</sub> Acetates: acyl = propanoyl

[178234-32-7, Kitasamycin Acetate]

Kitasamycin Acetate is a derivative of kitasamycin.

It contains not less than 680 μg (potency) and not more than 790 μg (potency) per mg, calculated on the anhydrous basis. The potency of Kitasamycin Acetate is expressed as mass (potency) of kitasamycin corresponding to the mass of leucomycin A<sub>5</sub> (C<sub>39</sub>H<sub>65</sub>NO<sub>14</sub>: 771.93). One mg (potency) of kitasamycin is equivalent to 0.530 mg of leucomycin A<sub>5</sub> (C<sub>39</sub>H<sub>65</sub>NO<sub>14</sub>).

**Description** Kitasamycin Acetate occurs as a white to light yellow-white powder.

It is very soluble in methanol and in ethanol (95), and practically insoluble in water.

**Identification** (1) Determine the absorption spectrum of a solution of Kitasamycin Acetate in methanol (1 in 40,000) as directed under Ultraviolet-visible Spectrophotometry <2.24>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wavelengths.



(2) Determine the infrared absorption spectrum of Kitasamycin Acetate as directed in the potassium bromide disk method under Infrared Spectrophotometry <2.25>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wave numbers.

**Water** <2.48> Not more than 5.0% (0.1 g, volumetric titration, direct titration).

**Assay** Perform the test according to the Cylinder-plate method as directed under Microbial Assay for Antibiotics <4.02> according to the following conditions.


- (i) Test organism—*Bacillus subtilis* ATCC 6633
- (ii) Culture medium—Use the medium i in 1) under (1) Agar media for seed and base layer.
- (iii) Standard solution—Weigh accurately an amount of Leucomycin A<sub>5</sub> Reference Standard equivalent to about 30 mg (potency), dissolve in 10 mL of methanol, add water to make exactly 100 mL, and use this solution as the standard stock solution. Keep the standard stock solution at 5°C or below and use within 3 days. Take exactly a suitable amount of the standard stock solution before use, add 0.1 mol/L phosphate buffer solution, pH 8.0 to make solutions so that each mL contains 30 µg (potency) and 7.5 µg (potency), and use these solutions as the high concentration standard solution and low concentration standard solution, respectively.
- (iv) Sample solution—Weigh accurately an amount of Kitasamycin Acetate equivalent to about 30 mg (potency), dissolve in 25 mL of methanol, add water to make exactly 50 mL, shake well, and allow to stand at 37 ± 2°C for 24 hours. Take exactly a suitable amount of the solution, add 0.1 mol/L phosphate buffer solution, pH 8.0 to make solutions so that each mL contains 30 µg (potency) and 7.5 µg (potency), and use these solutions as the high concentration sample solution and low concentration sample solution, respectively.

**Containers and storage** Containers—Tight containers.

## Kitasamycin Tartrate

### Leucomycin Tartrate

キタマイシン酒石酸塩



(Leucomycin A<sub>1</sub>, A<sub>5</sub>, A<sub>7</sub>, A<sub>9</sub> and A<sub>13</sub> Tartrates)  
 (3*R*,4*R*,5*S*,6*R*,8*R*,9*R*,10*E*,12*E*,15*R*)-5-[4-*O*-Acyl-2,6-dideoxy-3-*C*-methyl- $\alpha$ -*L*-ribo-hexopyranosyl-(1→4)-3,6-dideoxy-3-dimethylamino- $\beta$ -*D*-glucopyranosyloxy]-6-formylmethyl-3,9-dihydroxy-4-methoxy-8-methylhexadeca-10,12-dien-15-olide mono-(2*R*,3*R*)-tartrate  
 Leucomycin A<sub>1</sub> Tartrate: acyl = 3-methylbutanoyl  
 Leucomycin A<sub>5</sub> Tartrate: acyl = butanoyl  
 Leucomycin A<sub>7</sub> Tartrate: acyl = propanoyl  
 Leucomycin A<sub>9</sub> Tartrate: acyl = acetyl  
 Leucomycin A<sub>13</sub> Tartrate: acyl = hexanoyl

(Leucomycin A<sub>3</sub>, A<sub>4</sub>, A<sub>6</sub> and A<sub>8</sub> Tartrates)  
 (3*R*,4*R*,5*S*,6*R*,8*R*,9*R*,10*E*,12*E*,15*R*)-3-Acetoxy-5-[4-*O*-acyl-2,6-dideoxy-3-*C*-methyl- $\alpha$ -*L*-ribo-hexopyranosyl-(1→4)-3,6-dideoxy-3-dimethylamino- $\beta$ -*D*-glucopyranosyloxy]-6-formylmethyl-9-hydroxy-4-methoxy-8-methylhexadeca-10,12-dien-15-olide mono-(2*R*,3*R*)-tartrate  
 Leucomycin A<sub>3</sub> Tartrate: acyl = 3-methylbutanoyl  
 Leucomycin A<sub>4</sub> Tartrate: acyl = butanoyl  
 Leucomycin A<sub>6</sub> Tartrate: acyl = propanoyl  
 Leucomycin A<sub>8</sub> Tartrate: acyl = acetyl  
 [37280-56-1, Kitasamycin Tartrate]

Kitasamycin Tartrate is the tartrate of kitasamycin. It contains not less than 1300 µg (potency) per mg,

calculated on the anhydrous basis. The potency of Kitasamycin Tartrate is expressed as mass (potency) of kitasamycin based on the amount of leucomycin A<sub>5</sub> (C<sub>39</sub>H<sub>65</sub>NO<sub>14</sub>: 771.93). One mg (potency) of Kitasamycin Tartrate is equivalent to 0.530 mg of leucomycin A<sub>5</sub> (C<sub>39</sub>H<sub>65</sub>NO<sub>14</sub>).

**Description** Kitasamycin Tartrate occurs as a white to light yellowish white powder.

It is very soluble in water, in methanol and in ethanol (99.5).

**Identification (1)** Determine the absorption spectrum of a solution of Kitasamycin Tartrate in methanol (1 in 40,000) as directed under Ultraviolet-visible Spectrophotometry <2.24>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wavelengths.

(2) Determine the infrared absorption spectrum of Kitasamycin Tartrate as directed in the potassium bromide disk method under Infrared Spectrophotometry <2.25>, and compare the spectrum with the Reference Spectrum: both spectra exhibit similar intensities of absorption at the same wave numbers.

(3) Dissolve 1 g of Kitasamycin Tartrate in 20 mL of water, add 3 mL of sodium hydroxide TS, add 20 mL of *n*-butyl acetate, shake well, and discard the *n*-butyl acetate layer. To the aqueous layer add 20 mL of *n*-butyl acetate, and shake well. The aqueous layer so obtained responds to the Qualitative Tests <1.09> (1) for tartrate.

**pH <2.54>** Dissolve 3.0 g of Kitasamycin Tartrate in 100 mL of water: the pH of the solution is between 3.0 and 5.0.

**Content ratio of the active principle** Dissolve 20 mg of Kitasamycin Tartrate in diluted acetonitrile (1 in 2) to make 20 mL, and use this solution as the sample solution. Perform the test with 5  $\mu$ L of the sample solution as directed under Liquid Chromatography <2.01> according to the following conditions, determine the peak areas by the automatic integration method, and calculate the amounts of leucomycin A<sub>5</sub>, leucomycin A<sub>4</sub> and leucomycin A<sub>1</sub> by the area percentage method: the amount of leucomycin A<sub>5</sub> is 40 – 70%, leucomycin A<sub>4</sub> is 5 – 25%, and leucomycin A<sub>1</sub> is 3 – 12%. The relative retention times of leucomycin A<sub>4</sub> and leucomycin A<sub>1</sub> with respect to leucomycin A<sub>5</sub> are 1.2 and 1.5, respectively.

**Operating conditions—**

**Detector:** An ultraviolet absorption photometer (wavelength: 232 nm).

**Column:** A stainless steel column 4.0 mm in inside diameter and 15 cm in length, packed with octylsilanized silica gel for liquid chromatography (5  $\mu$ m in particle diameter).

**Column temperature:** A constant temperature of about 40°C.

**Mobile phase:** To a suitable amount of a solution of ammonium acetate (77 in 500) add diluted phosphoric acid (1 in 150) to adjust the pH to 5.5. To 370 mL of this solution add 580 mL of methanol and 50 mL of acetonitrile.

**Flow rate:** Adjust the flow rate so that the retention time of leucomycin A<sub>5</sub> is about 8 minutes.

**Time span of measurement:** About 3 times as long as the retention time of leucomycin A<sub>5</sub>.

**System suitability—**

**System performance:** Dissolve about 20 mg of Leucomycin A<sub>5</sub> Reference Standard and about 20 mg of Josamycin Refer-

ence Standard in 20 mL of diluted acetonitrile (1 in 2). When the procedure is run with 5  $\mu$ L of this solution under the above operating conditions, leucomycin A<sub>5</sub> and josamycin are eluted in this order with the resolution between these peaks being not less than 5.

**System repeatability:** When the test is repeated 6 times with 5  $\mu$ L of the sample solution under the above operating conditions, the relative standard deviation of the peak area of leucomycin A<sub>5</sub> is not more than 1.0%.

**Purity (1)** Clarity and color of solution—Dissolve 1.0 g of Kitasamycin Tartrate in 10 mL of water: the solution is clear and colorless or light yellow.

(2) Heavy metals <1.07>—Proceed with 1.0 g of Kitasamycin Tartrate according to Method 2, and perform the test. Prepare the control solution with 3.0 mL of Standard Lead Solution (not more than 30 ppm).

**Water <2.48>** Not more than 3.0% (0.1 g, volumetric titration, direct titration).

**Assay** Perform the test according to the Cylinder-plate method as directed under Microbial Assay for Antibiotics <4.02> according to the following conditions.

(i) Test organism—*Bacillus subtilis* ATCC 6633

(ii) Culture medium—Use the medium i in 1) Medium for test organism [5] under (1) Agar media for seed and base layer.


(iii) Standard solutions—Weigh accurately an amount of Leucomycin A<sub>5</sub> Reference Standard, equivalent to about 30 mg (potency), dissolve in 10 mL of methanol, add water to make exactly 100 mL, and use this solution as the standard stock solution. Keep the standard stock solution at not exceeding 5°C, and use within 3 days. Take exactly a suitable amount of the standard stock solution before use, add phosphate buffer solution, pH 8.0 to make solutions so that each mL contains 30  $\mu$ g (potency) and 7.5  $\mu$ g (potency), and use these solutions as the high concentration standard solution and low concentration standard solution, respectively.

(iv) Sample solutions—Weigh accurately an amount of Kitasamycin Tartrate, equivalent to about 30 mg (potency), and dissolve in water to make exactly 100 mL. Take exactly a suitable amount of this solution, add phosphate buffer solution, pH 8.0 to make solutions so that each mL contains 30  $\mu$ g (potency) and 7.5  $\mu$ g (potency), and use these solutions as the high concentration sample solution and low concentration sample solution, respectively.

**Containers and storage** Containers—Tight containers.

## Lactic Acid

乳酸



C<sub>3</sub>H<sub>6</sub>O<sub>3</sub>: 90.08

(2*RS*)-2-Hydroxypropanoic acid [50-21-5]

Lactic Acid is a mixture of lactic acid and lactic anhydride.

It contains not less than 85.0% and not more than 92.0% of C<sub>3</sub>H<sub>6</sub>O<sub>3</sub>.