



This English version is intended to be a reference material to provide convenience for users. In the event of inconsistency between the Japanese original and this English translation, the former shall prevail.

## Revision of Precautions Clonazepam

March 21, 2017

## Non-proprietary name

Clonazepam

## Safety measure

Precautions should be revised in the package insert.

In the Clinically Significant Adverse Reactions subsection regarding dependence of the Adverse Reactions section, the following text should be revised (underlined parts are revised):

Dependence: Drug dependence may occur due to prolonged use. Patients should be carefully monitored and <u>caution should be exercised for dosage and treatment</u> <u>duration</u> when administering this drug. In addition, withdrawal symptoms such as seizure, delirium, tremor, insomnia, anxiety, hallucination, or delusion may occur due to rapid dose reduction or discontinuation of this drug during prolonged use. Discontinuation of administration should be carefully carried out such as by gradual dose reduction.

In the Clinically Significant Adverse Reactions subsection regarding irritable excitation and confusion of the Adverse Reactions section, the following text should be revised:

Irritable excitation, confusion, etc.: Irritable excitation or confusion, etc., may occur. Patients should be carefully monitored. If any abnormalities are observed, appropriate measures should be taken. Discontinuation of administration should be carefully carried out such as by gradual dose reduction.

\* This drug is designated as a drug requiring preparation of a Drug Guide for Patients.