



This English version is intended to be a reference material to provide convenience for users. In the event of inconsistency between the Japanese original and this English translation, the former shall prevail.

Revision of Precautions Antimicrobials indicated for any of pharyngitis/laryngitis, tonsillitis, acute bronchitis, infectious enteritis, or sinusitis

March 27, 2018

Non-proprietary name

Antimicrobials indicated for any of pharyngitis/laryngitis, tonsillitis, acute bronchitis, infectious enteritis, or sinusitis

Safety measure

Precautions should be revised in the package insert.

In the Precaution Concerning Indications section, the following language should be added (revised language is underlined):

For the treatment of pharyngitis/laryngitis, tonsillitis, acute bronchitis, infectious enteritis, or sinusitis*, the Guidance for Appropriate Use of Antimicrobials should be consulted to decide whether administration of antimicrobials is necessary, and then this drug should be administered in cases where treatment with this drug is determined to be appropriate.

<Reference> Guidance for Appropriate Use of Antimicrobials compiled by the Tuberculosis and Infectious Diseases Control Division, of the Health Bureau, at the Ministry of Health, Labour and Welfare

Published by Ministry of Health, Labour and Welfare

Translated by Pharmaceuticals and Medical Devices Agency





This English version is intended to be a reference material to provide convenience for users. In the event of inconsistency between the Japanese original and this English translation, the former shall prevail.

*In the part of "pharyngitis/laryngitis, tonsillitis, acute bronchitis, infectious enteritis, or sinusitis," approved indications should be listed exactly as listed in the Approval document.