



Module 3

Selection of doses for use in confirmatory MRCTs

ICH E17: General principles for planning and design of Multi-Regional Clinical Trials

International Council for Harmonisation of Technical Requirements
for Pharmaceuticals for Human Use

Legal Notice

- This presentation is protected by copyright and may, with the exception of the ICH logo, be used, reproduced, incorporated into other works, adapted, modified, translated or distributed under a public license provided that ICH's copyright in the presentation is acknowledged at all times. In case of any adaption, modification or translation of the presentation, reasonable steps must be taken to clearly label, demarcate or otherwise identify that changes were made to or based on the original presentation. Any impression that the adaption, modification or translation of the original presentation is endorsed or sponsored by the ICH must be avoided.
- The presentation is provided "as is" without warranty of any kind. In no event shall the ICH or the authors of the original presentation be liable for any claim, damages or other liability arising from the use of the presentation.
- The above-mentioned permissions do not apply to content supplied by third parties. Therefore, for documents where the copyright vests in a third party, permission for reproduction must be obtained from this copyright holder.

Outline

- **Introduction and purpose**
- **Points to consider for selection of doses**
- **Examples of using different dose regimens in a confirmatory MRCT**
- **Concluding remarks**

Introduction and objectives

- **In principle, the same dose regimens should generally be used for all participating ethnic populations in a confirmatory MRCT**
- **The purpose of this Module is:**
 - To provide points to consider to scientifically justify setting different dose regimens
 - To provide illustrative examples where it may be appropriate to use different dose regimens in a confirmatory MRCT


Points to consider

- **Available information on pharmacokinetics (PK), pharmacodynamics (PD), efficacy and safety of the drug should be collected as early as possible, to identify potential regional differences**
- **Dose-exposure and exposure-response relationships in different ethnic populations should be examined to**
 - Identify intrinsic and extrinsic factors which may affect the treatment effect
 - Investigate possible mechanisms for any observed ethnic differences in PK and/or treatment effect
- **Strategy for dose selection in an MRCT should be determined**
 - If early data shows a clear difference in dose-response and/or exposure-response relationships among ethnic populations, it may be appropriate to use different dosing regimens in a confirmatory MRCT

Example 1: Drug X


- **Chemical entity**
- **Orally administered, low bioavailability**
- **PK differences were observed between ethnic populations A and B, due to differences in diet and body weight**
- **The PK differences resulted in a difference in treatment effects.**
- **Safety was exposure-related**

PK difference between populations



- Concentration of Drug X in plasma was 2 times higher in population A than population B
- Concentrations of Drug X achieved in population A at the 10 mg dose level were comparable to concentrations in population B at the 20 mg dose level

Dose-response difference between populations



- 10 mg in population A has similar treatment effect as 20 mg in population B
- Taken together with the previous PK data of Drug X in populations A and B, the differences in PK resulted in a difference in treatment effect

Conclusion for example 1: Drug X

- Early trial data for Drug X showed a difference in PK that resulted in a difference in treatment effect; lower dose in population A achieved similar treatment effect to full dose in population B
- As safety of Drug X was exposure-dependent, from a safety perspective, a lower dose was preferable in population A
- In such a case, it may be appropriate to consider the use of different dose regimens in a confirmatory MRCT
- However, if PK difference between ethnic populations is caused by a specific factor (e.g., poor metabolizer status), it would be more appropriate to select dose regimens based on the factor instead of ethnic population


Example 2: Drug Y

- **Chemical entity**
- **Orally administered**
- **PK differences observed between ethnic populations**
- **Cause for PK differences examined but remained unknown**
- **Inter-individual variability in PK and PD found to be large in both ethnic populations**
 - Fixed dose would not be appropriate

PK difference between populations


Phase I study

Population	Dose	C _{max}	AUC
A	50 mg	5.7 [3.5 - 9.4]	81.0 [41.2 - 153]
B	50 mg	3.4 [2.5 - 3.8]	45.0 [38.7 - 48.8]



- Concentration of Drug Y was 1.8 times higher in ethnic population A than ethnic population B
- Population PK (PPK) analysis suggested that covariates significantly affecting total body clearance (CL/F) included body weight, race, and health condition (i.e., healthy subject or patient).

Exposure-response relationship between populations



Similarity in exposure-response relationship between populations A and B

- PD parameter tended to increase with increasing AUC in both ethnic population A and B
 - Exposure-response relationship is similar in ethnic populations A and B
- Response-guided titration and monitoring of efficacy and safety would be necessary because of large inter-individual variability in PK and PD

Conclusion for example 2: Drug Y

- PK differences observed, but exposure-response relationship was similar between ethnic populations A and B
- In such a case, it may be appropriate to consider a lower starting dose in ethnic population A than that in ethnic population B
- Due to large inter-individual variability in PK, response-guided titration and PD monitoring recommended
- For titration dose setting, the setting of the starting, increment and maximum dose should be based on the PK/PD and safety data from exploratory trials

Concluding remarks

- If substantial differences in dose-response and/or exposure-response relationships between ethnic populations are observed, it may be appropriate to consider a different dose regimen for the impacted ethnic population.
- Early trial data are useful to determine if the selected dose regimens are expected to produce similar therapeutic effects with an acceptable safety margin.
- Scientific justifications for selected dose regimens should be described in study protocol, including clinical relevance of regional differences
- Impact of the use of different dose regimens on the logistics of MRCTs should be considered
- Dose selection strategies in confirmatory MRCT should be discussed in advance with relevant regulatory authorities