

Published by
Ministry of Health, Labour and Welfare



Translated by
Pharmaceuticals and Medical Devices Agency



This English version is intended to be a reference material to provide convenience for users. In the event of inconsistency between the Japanese original and this English translation, the former shall prevail.

Revision of Precautions

Ampiroxicam

Piroxicam (oral dosage form)

February 25, 2021

Therapeutic category

Antipyretics, analgesics and anti-inflammatory agents

Non-proprietary name

Ampiroxicam, piroxicam

Safety measure

Precautions should be revised in the package insert.

Pharmaceuticals and Medical Devices Agency

3-3-2 Kasumigaseki, Chiyoda-ku, Tokyo 100-0013 Japan
E-mail: safety.info@pmda.go.jp

Revision in line with the Instructions for Package Inserts of Prescription Drugs, PAB Notification No. 606 by the Director General of Pharmaceutical Affairs Bureau, MHW, dated April 25, 1997 (Old instructions): Revised language is underlined.

Current	Revision
<p>Use during Pregnancy, Delivery or Lactation</p> <p>The safety of this drug administered during pregnancy has not been established. Pregnant women or women who may be pregnant should be administered this drug only when the therapeutic benefits are considered to outweigh the risks.</p>	<p>Use during Pregnancy, Delivery or Lactation</p> <p>The safety of this drug administered during pregnancy has not been established. Pregnant women <u>(excluding those in their term)</u> or women who may be pregnant should be administered this drug only when the therapeutic benefits are considered to outweigh the risks. <u>If such administration is deemed necessary, caution should be exercised such as limiting to the minimum effective use and monitoring amniotic fluid as necessary. Renal impairment and decreased urine output in fetuses as well as accompanying oligohydramnios have been reported following use of cyclooxygenase inhibitors (oral dosage form or suppository) in pregnant women.</u></p>

Revision in line with the Instructions for Package Inserts of Prescription Drugs, etc. PSEHB Notification No. 0608-1 by the Director of Pharmaceutical Safety and Environmental Health Bureau, MHLW, dated June 8, 2017 (New instructions): Revised language is underlined.

Current	Revision
<p>9. PRECAUTIONS CONCERNING PATIENTS WITH SPECIFIC BACKGROUNDS</p> <p>9.5 Pregnant Women</p> <p>Pregnant women or women who may be pregnant <u>(excluding those in their third trimester)</u></p>	<p>9. PRECAUTIONS CONCERNING PATIENTS WITH SPECIFIC BACKGROUNDS</p> <p>9.5 Pregnant Women</p> <p>Pregnant women <u>(excluding those in their third trimester)</u> or women who may be pregnant</p>

Pharmaceuticals and Medical Devices Agency

3-3-2 Kasumigaseki, Chiyoda-ku, Tokyo 100-0013 Japan
 E-mail: safety.info@pmda.go.jp

This drug should be administered only when the therapeutic benefits are considered to outweigh the risks.

This drug should be administered only when the therapeutic benefits are considered to outweigh the risks. If such administration is deemed necessary, caution should be exercised such as limiting to the minimum effective use and monitoring amniotic fluid as necessary. Renal impairment and decreased urine output in foetuses as well as accompanying oligohydramnios have been reported following use of cyclooxygenase inhibitors (oral dosage form or suppository) in pregnant women.

Pharmaceuticals and Medical Devices Agency

3-3-2 Kasumigaseki, Chiyoda-ku, Tokyo 100-0013 Japan
E-mail: safety.info@pmda.go.jp