Published by Ministry of Health, Labour and Welfare

Translated by Pharmaceuticals and Medical Devices Agency





This English version is intended to be a reference material to provide convenience for users. In the event of inconsistency between the Japanese original and this English translation, the former shall prevail.

Revision of Precautions Sulpyrine hydrate

February 25, 2021

Therapeutic category

Antipyretics, analgesics and anti-inflammatory agents

Non-proprietary name

Sulpyrine hydrate

Safety measure

Precautions should be revised in the package insert.

Pharmaceuticals and Medical Devices Agency

3-3-2 Kasumigaseki, Chiyoda-ku, Tokyo 100-0013 Japan E-mail: safety.info@pmda.go.jp

Revision in line with the Instructions for Package Inserts of Prescription Drugs, PAB Notification No. 606 by the Director General of Pharmaceutical Affairs Bureau, MHW, dated April 25, 1997 (Old instructions):

Revised language is underlined.

	·
Current	Revision
Use during Pregnancy, Delivery or Lactation This drug should be administered to pregnant women or women who may be pregnant only when the therapeutic benefits are considered to outweigh the risks.	Use during Pregnancy, Delivery or Lactation This drug should be administered to pregnant women or women who may be pregnant only when the therapeutic benefits are considered to outweigh the risks. If such administration is deemed necessary, caution should be exercised such as limiting to the minimum effective use and monitoring amniotic fluid as necessary. Renal impairment and decreased urine output in foetuses as well as accompanying oligohydramnios have been reported following use of cyclooxygenase inhibitors (oral dosage form or suppository) in pregnant women.
	in program women.