

Published by  
Ministry of Health, Labour and Welfare



Translated by  
Pharmaceuticals and Medical Devices Agency



---

*This English version is intended to be a reference material to provide convenience for users. In the event of inconsistency between the Japanese original and this English translation, the former shall prevail.*

# Revision of Precautions

## Salicylamide/acetaminophen/anhydrous caffeine/chlorpheniramine maleate, salicylamide/acetaminophen/anhydrous caffeine/promethazine methylenedisalicylate

February 25, 2021

### **Therapeutic category**

Agents used for common cold

### **Non-proprietary name**

Salicylamide/acetaminophen/anhydrous caffeine/chlorpheniramine maleate, salicylamide/acetaminophen/anhydrous caffeine/promethazine methylenedisalicylate

### **Safety measure**

Precautions should be revised in the package insert.

**Pharmaceuticals and Medical Devices Agency**

3-3-2 Kasumigaseki, Chiyoda-ku, Tokyo 100-0013 Japan  
E-mail: [safety.info@pmda.go.jp](mailto:safety.info@pmda.go.jp)

Revision in line with the Instructions for Package Inserts of Prescription Drugs, PAB Notification No. 606 by the Director General of Pharmaceutical Affairs Bureau, MHW, dated April 25, 1997 (Old instructions): Revised language is underlined.

Current	Revision
<p>Use during Pregnancy, Delivery or Lactation</p> <p>This drug should be administered to pregnant women <u>(those within 12 weeks before due date or in their third trimester)</u> or women who may be pregnant only when the therapeutic benefits are considered to outweigh the risks.</p>	<p>Use during Pregnancy, Delivery or Lactation</p> <p>This drug should be administered to pregnant women or women who may be pregnant only when the therapeutic benefits are considered to outweigh the risks. <u>If such administration is deemed necessary, caution should be exercised such as limiting to the minimum effective use and monitoring amniotic fluid as necessary. Renal impairment and decreased urine output in fetuses as well as accompanying oligohydramnios have been reported following use of cyclooxygenase inhibitors (oral dosage form or suppository) in pregnant women.</u></p>

**Pharmaceuticals and Medical Devices Agency**

3-3-2 Kasumigaseki, Chiyoda-ku, Tokyo 100-0013 Japan  
E-mail: [safety.info@pmda.go.jp](mailto:safety.info@pmda.go.jp)