

## To patients who receive and their caregivers [Joyclu 30mg intra-articular injection]

- **Anaphylaxis may occur during a few hours following administration of this drug.**
- Patients should be careful for any abnormalities immediately following administration until and after they reach home. When at home, family members should also be alert for any abnormalities patients may experience.
- Please seek medical attention immediately if you experience any of the symptoms below. Tell the medical institution you visit that you have received Joyclu intra-articular injection.

### ◆Skin symptoms

Itchy skin,  
Urticarial,  
Erythema/redness, etc.



### ◆Respiratory symptoms

Hoarseness,  
Sneezing,  
Itchy throat,  
Swollen lips,  
Difficulty in breathing,  
Dyspnoea, etc



### ◆Cardiovascular symptoms

Tachycardia,  
Palpitations,  
Pale look  
Loss of consciousness,  
Staggering,  
Dizziness, etc.



### ◆Gastrointestinal symptoms

Abdominal pain,  
Feeling queasy,  
Diarrhea, etc.



- Shock symptoms such as pale look and loss of consciousness, and difficulty in breathing are a medical emergency. Please act quickly by calling an ambulance or

### 【Contact point】

Ono Pharmaceutical Co., Ltd Medicine Consultation Service      Toll free: 0120-886-336  
Business hours: 9:00~17:00 (Excluding Sat., Sun., holidays and other company's holidays)