Published by Ministry of Health, Labour and Welfare Translated by Pharmaceuticals and Medical Devices Agency





This English version is intended to be a reference material to provide convenience for users. In the event of inconsistency between the Japanese original and this English translation, the former shall prevail.

## **Revision of Precautions**

# Coronavirus modified uridine RNA vaccine (SARS-CoV-2)

(COVID-19 Vaccine Moderna Intramuscular Injection)

October 15, 2021

## Therapeutic category

Vaccines

### Non-proprietary name

Coronavirus modified uridine RNA vaccine (SARS-CoV-2)

#### Safety measure

Precautions should be revised in the package insert.

**Pharmaceuticals and Medical Devices Agency** 

Revision in line with the Instructions for Electronic Package Inserts of Prescription Drugs, etc. PSEHB Notification No. 0611-1 by the Director of Pharmaceutical Safety and Environmental Health Bureau, MHLW, dated June 11, 2021 (New instructions): Revised language is underlined.

Current	Revision
15. OTHER PRECAUTIONS	15. OTHER PRECAUTIONS
15.1 Information Based on Clinical Use	15.1 Information Based on Clinical Use
(N/A)	It is suggested that the frequency of myocarditis and pericarditis
	was higher in the male adolescents and young adults inoculated
	with this vaccine by comparing the reporting rates of myocarditis
	and pericarditis in the domestic suspected adverse reaction reports
	after the start of vaccination and the estimated background
	incidence rates of myocarditis and pericarditis in the general
	population utilizing a domestic medical information database.

N/A: Not Applicable. No corresponding language is included in the current package insert.

(References) Material for the Side Effect Subcommittee of the Immunization and Vaccine Section Meeting in the Health Science Council (the 70th meeting), and the 2021 Subcommittee on Drug Safety of the Committee on Drug Safety in the Pharmaceutical Affairs and Food Sanitation Council (the 19th meeting) (joint meeting)