Published by Ministry of Health, Labour and Welfare Translated by Pharmaceuticals and Medical Devices Agency





This English version is intended to be a reference material to provide convenience for users. In the event of inconsistency between the Japanese original and this English translation, the former shall prevail.

Revision of Precautions

Coronavirus modified uridine RNA vaccine (SARS-CoV-2)

(Spikevax Intramuscular Injection)

April 25, 2022

Therapeutic category

Vaccines

Non-proprietary name

Coronavirus modified uridine RNA vaccine (SARS-CoV-2)

Safety measure

Precautions should be revised in the package insert.

Pharmaceuticals and Medical Devices Agency

Revision in line with the Instructions for Electronic Package Inserts of Prescription Drugs, etc. PSEHB Notification No. 0611-1 by the Director of Pharmaceutical Safety and Environmental Health Bureau, MHLW, dated June 11, 2021 (New instructions):

Revised language is underlined.

Current	Revision
7. PRECAUTIONS CONCERNING DOSAGE AND ADMINISTRATION	7. PRECAUTIONS CONCERNING DOSAGE AND ADMINISTRATION
Booster dose	Booster dose
Timing of vaccination	Timing of vaccination
The third dose may be administered as a booster dose at least <u>6</u>	The third dose may be administered as a booster dose at least <u>5</u>
months after the second dose.	months after the second dose.
No clinical studies have been conducted on the booster dose	For the fourth dose, the vaccination may be considered in elderly
(0.25 mL) of this vaccine in people who have received the primary	people, etc. based on the benefit/risk balance at least 5 months
series of other SARS-CoV-2 vaccines.	after the third dose.
	The effectiveness and safety on the booster dose (0.25 mL) of this
	vaccine in people who have received other SARS-CoV-2 vaccines
	have not been established.

N/A: Not Applicable. No corresponding language is included in the current package insert.