Published by Ministry of Health, Labour and Welfare

62

Translated by Pharmaceuticals and Medical Devices Agency



This English version is intended to be a reference material to provide convenience for users. In the event of inconsistency between the Japanese original and this

English translation, the former shall prevail.

## Revision of Precautions Eszopiclone

July 20, 2022

Therapeutic category

Hypnotics and sedatives, antianxietics

Non-proprietary name

Eszopiclone

**Safety measure** Precautions should be revised.

Pharmaceuticals and Medical Devices Agency

Revision in line with the Instructions for Package Inserts of Prescription Drugs, PAB Notification No. 606 by the Director General ofPharmaceutical Affairs Bureau, MHW, dated April 25, 1997 (Old instructions):Revised language is underlined.

Current	Revision
Careful Administration	Careful Administration
(N/A)	Patients who have experienced abnormal behavior as parasomnia
	(somnambulism, etc.) with administration of this drug [Parasomnia
	leading to serious self/other-injuries, accidents, etc. may occur.
	Discontinuation of this drug should be considered.]
Adverse Reactions	Adverse Reactions
Clinically Significant Adverse Reactions	Clinically Significant Adverse Reactions
Psychiatric symptom, disturbed consciousness:	Psychiatric symptom, disturbed consciousness:
Psychiatric symptoms or disturbed consciousness such as	Psychiatric symptoms or disturbed consciousness such as
nightmare (abnormal dreams), depressed level of consciousness,	nightmare (abnormal dreams), depressed level of consciousness,
excitement (agitation), confusion (confusional state), hallucination.	excitement (agitation), confusion (confusional state), hallucination,
somnambulism, aggression, delirium, and abnormal behavior may	aggression, delirium, and abnormal behavior may occur. Patients
occur. Patients should be carefully monitored. If any abnormalities	should be carefully monitored. If any abnormalities are observed,
are observed, administration of this drug should be discontinued.	administration of this drug should be discontinued.
Transient anterograde amnesia, twilight state:	Transient anterograde amnesia, twilight state <u>, parasomnia</u>
Transient anterograde amnesia (failure to remember events during	(somnambulism, etc.):
nocturnal awakening, etc.) and twilight state may occur. Careful	Transient anterograde amnesia (failure to remember events during
administration of this drug is required, such as starting with a low	nocturnal awakening, etc.), twilight state, and parasomnia
dose. With respect to zopiclone preparations, there have been	(somnambulism, etc.) may occur. Careful administration of this drug

Pharmaceuticals and Medical Devices Agency

reports of patients driving a car, eating, etc. without being fully awake	is required, such as starting with a low dose. With respect to
and not remembering the events. If any abnormalities are observed,	zopiclone preparations, there have been reports of patients driving
administration of this drug should be discontinued.	a car, eating, etc. without being fully awake and not remembering
	the events. If any abnormalities are observed, administration of this
	drug should be discontinued.

N/A: Not Applicable. No corresponding language is included in the current Precautions.

Revision in line with the Instructions for Electronic Package Inserts of Prescription Drugs, etc. PSEHB Notification No. 0611-1 by the Director of Pharmaceutical Safety and Environmental Health Bureau, MHLW, dated June 11, 2021 (New instructions): Revised language is underlined.

Current	Revision
9. PRECAUTIONS CONCERNING PATIENTS WITH SPECIFIC	9. PRECAUTIONS CONCERNING PATIENTS WITH SPECIFIC
BACKGROUNDS	BACKGROUNDS
9.1 Patients with Complication or History of Diseases, etc.	9.1 Patients with Complication or History of Diseases, etc.
(N/A)	Patients who have experienced abnormal behavior as parasomnia
	(somnambulism, etc.) with administration of this drug
	Discontinuation of this drug should be considered. Parasomnia
	leading to serious self/other-injuries, accidents, etc. may occur.
11. ADVERSE REACTIONS	11. ADVERSE REACTIONS
11.1 Clinically Significant Adverse Reactions	11.1 Clinically Significant Adverse Reactions
Psychiatric symptom, disturbed consciousness	Psychiatric symptom, disturbed consciousness
Nightmare (abnormal dreams), depressed level of consciousness,	Nightmare (abnormal dreams), depressed level of consciousness,
excitement (agitation), confusion (confusional state), hallucination,	excitement (agitation), confusion (confusional state), hallucination,
somnambulism, aggression, delirium, and abnormal behavior, etc.	aggression, delirium, and abnormal behavior, etc. may occur.

Pharmaceuticals and Medical Devices Agency

may occur.	
Transient anterograde amnesia, twilight state	Transient anterograde amnesia, twilight state <u>, parasomnia</u>
Careful administration of this drug is required, such as starting with	<u>(somnambulism, etc.)</u>
a low dose. There have been reports of patients driving, eating, etc.	Careful administration of this drug is required, such as starting with
without being fully awake and not remembering the events.	a low dose. There have been reports of patients driving, eating, etc.
	without being fully awake and not remembering the events.

N/A: Not Applicable. No corresponding language is included in the current Precautions.

Pharmaceuticals and Medical Devices Agency