

Published by
Ministry of Health, Labour and Welfare



Translated by
Pharmaceuticals and Medical Devices Agency



This English version is intended to be a reference material to provide convenience for users. In the event of inconsistency between the Japanese original and this English translation, the former shall prevail.

Revision of Precautions

Preparations containing loxoprofen sodium hydrate (oral dosage form) (OTC drugs)

October 12, 2022

Therapeutic category

Antipyretics and analgesics

Non-proprietary name

Loxoprofen sodium hydrate

Safety measure

Precautions should be revised.

Pharmaceuticals and Medical Devices Agency

3-3-2 Kasumigaseki, Chiyoda-ku, Tokyo 100-0013 Japan
E-mail: safety.info@pmda.go.jp

Revised language is underlined.

| Current | | Revision | | | | | | | | | |
|--|---|---|----------|---|--|--|--|------------------|----------|--|---|
| <p>Consultation</p> <p>If the following symptoms are observed after taking this drug, these may be adverse reactions. In such a case, the use of this drug should be immediately discontinued, and a physician, dentist or pharmacist should be consulted with this document.</p> <p>The following serious symptoms may occur rarely. In such a case, medical attention should be sought immediately.</p> | | <p>Consultation</p> <p>If the following symptoms are observed after taking this drug, these may be adverse reactions. In such a case, the use of this drug should be immediately discontinued, and a physician, dentist or pharmacist should be consulted with this document.</p> <p>The following serious symptoms may occur rarely. In such a case, medical attention should be sought immediately.</p> | | | | | | | | | |
| <table border="1"> <thead> <tr> <th>Name of symptoms</th> <th>Symptoms</th> </tr> </thead> <tbody> <tr> <td>Oculomucocutaneous syndrome (Stevens-Johnson syndrome), toxic epidermal necrolysis, erythema multiforme</td> <td>Some symptoms, such as hyperthermia, ocular hyperaemia, eye discharge, sore lips, pharynx pain, widespread skin rash/redness, and blisters on reddened skin, may persist or suddenly worsen.</td> </tr> </tbody> </table> | | Name of symptoms | Symptoms | Oculomucocutaneous syndrome (Stevens-Johnson syndrome), toxic epidermal necrolysis, erythema multiforme | Some symptoms, such as hyperthermia, ocular hyperaemia, eye discharge, sore lips, pharynx pain, widespread skin rash/redness, and blisters on reddened skin, may persist or suddenly worsen. | <table border="1"> <thead> <tr> <th>Name of symptoms</th> <th>Symptoms</th> </tr> </thead> <tbody> <tr> <td>Oculomucocutaneous syndrome (Stevens-Johnson syndrome), toxic epidermal necrolysis, erythema multiforme, <u>acute generalized exanthematous pustulosis</u></td> <td>Some symptoms, such as hyperthermia, ocular hyperaemia, eye discharge, sore lips, pharynx pain, widespread skin rash/redness, blisters on reddened skin, <u>small pimples (small pustules) on reddened skin, general</u> malaise, and anorexia, may persist or suddenly worsen.</td> </tr> </tbody> </table> | | Name of symptoms | Symptoms | Oculomucocutaneous syndrome (Stevens-Johnson syndrome), toxic epidermal necrolysis, erythema multiforme, <u>acute generalized exanthematous pustulosis</u> | Some symptoms, such as hyperthermia, ocular hyperaemia, eye discharge, sore lips, pharynx pain, widespread skin rash/redness, blisters on reddened skin, <u>small pimples (small pustules) on reddened skin, general</u> malaise, and anorexia, may persist or suddenly worsen. |
| Name of symptoms | Symptoms | | | | | | | | | | |
| Oculomucocutaneous syndrome (Stevens-Johnson syndrome), toxic epidermal necrolysis, erythema multiforme | Some symptoms, such as hyperthermia, ocular hyperaemia, eye discharge, sore lips, pharynx pain, widespread skin rash/redness, and blisters on reddened skin, may persist or suddenly worsen. | | | | | | | | | | |
| Name of symptoms | Symptoms | | | | | | | | | | |
| Oculomucocutaneous syndrome (Stevens-Johnson syndrome), toxic epidermal necrolysis, erythema multiforme, <u>acute generalized exanthematous pustulosis</u> | Some symptoms, such as hyperthermia, ocular hyperaemia, eye discharge, sore lips, pharynx pain, widespread skin rash/redness, blisters on reddened skin, <u>small pimples (small pustules) on reddened skin, general</u> malaise, and anorexia, may persist or suddenly worsen. | | | | | | | | | | |

Pharmaceuticals and Medical Devices Agency

3-3-2 Kasumigaseki, Chiyoda-ku, Tokyo 100-0013 Japan
 E-mail: safety.info@pmda.go.jp