Published by Ministry of Health, Labour and Welfare Translated by Pharmaceuticals and Medical Devices Agency





This English version is intended to be a reference material to provide convenience for users. In the event of inconsistency between the Japanese original and this English translation, the former shall prevail.

# **Revision of Precautions**

# **Acetaminophen (injections)**

## Isopropylantipyrine/acetaminophen/allylisopropylacetylurea/anhydrous caffeine

January 17, 2023

#### Therapeutic category

Antipyretics, analgesics and anti-inflammatory agents

### Non-proprietary name

Acetaminophen

Isopropylantipyrine/acetaminophen/allylisopropylacetylurea/anhydrous caffeine

### Safety measure

Precautions should be revised.

**Pharmaceuticals and Medical Devices Agency** 

Revision in line with the Instructions for Electronic Package Inserts of Prescription Drugs, etc. PSEHB Notification No. 0611-1 by the Director of Pharmaceutical Safety and Environmental Health Bureau, MHLW, dated June 11, 2021 (New instructions): Revised language is underlined.

Current	Revision
11. ADVERSE REACTIONS	11. ADVERSE REACTIONS
11.1 Clinically Significant Adverse Reactions	11.1 Clinically Significant Adverse Reactions
(N/A)	Drug-induced hypersensitivity syndrome
	Initial symptoms of rash and pyrexia, followed by serious delayed
	symptoms of hypersensitivity accompanied by hepatic impairment,
	swollen lymph nodes, increased white blood cell, eosinophilia, and
	appearance of atypical lymphocytes may occur. Symptoms are
	often accompanied by virus reactivation, such as human herpes
	virus type 6 (HHV-6). Caution is required for recurrence or
	prolongation of rash, pyrexia, and hepatic impairment, etc. that may
	occur even after discontinuation of administration.

N/A: Not Applicable. No corresponding language is included in the current Precautions.