

Published by  
Ministry of Health, Labour and Welfare



Translated by  
Pharmaceuticals and Medical Devices Agency



---

*This English version is intended to be a reference material to provide convenience for users. In the event of inconsistency between the Japanese original and this English translation, the former shall prevail.*

# Revision of Precautions

## Preparations containing acetaminophen (oral dosage form, suppositories) (OTC drugs)

January 17, 2023

### **Therapeutic category**

Cold medicines

Antipyretics and analgesics

### **Non-proprietary name**

Preparations containing acetaminophen

### **Safety measure**

Precautions should be revised.

**Pharmaceuticals and Medical Devices Agency**

3-3-2 Kasumigaseki, Chiyoda-ku, Tokyo 100-0013 Japan  
E-mail: [safety.info@pmda.go.jp](mailto:safety.info@pmda.go.jp)

Current	Revision				
<p>Consultation</p> <p>If the following symptoms are observed after taking this drug, these may be adverse reactions. In such a case, the use of this drug should be immediately discontinued, and a physician, dentist, pharmacist or registered sales clerk should be consulted with this document.</p> <p>The following serious symptoms may occur rarely. In such a case, medical attention should be sought immediately.</p> <p>(N/A)</p> <p>*The highlighted part should be listed only in the preparations containing ibuprofen among antipyretics and analgesics.</p>	<p>Consultation</p> <p>If the following symptoms are observed after taking this drug, these may be adverse reactions. In such a case, the use of this drug should be immediately discontinued, and a physician, dentist, pharmacist or registered sales clerk should be consulted with this document.</p> <p>The following serious symptoms may occur rarely. In such a case, medical attention should be sought immediately.</p> <table border="1" data-bbox="1133 587 1962 927"> <thead> <tr> <th data-bbox="1133 587 1469 636">Name of symptoms</th> <th data-bbox="1473 587 1962 636">Symptoms</th> </tr> </thead> <tbody> <tr> <td data-bbox="1133 636 1469 927"><u>Drug-induced hypersensitivity syndrome</u></td> <td data-bbox="1473 636 1962 927"><u>Some symptoms, such as redness over large part of the skin, generalised exanthema, pyrexia, malaise, swollen lymph nodes (neck, armpits, groin, etc.) may occur.</u></td> </tr> </tbody> </table> <p>*The highlighted part should be listed only in the preparations containing ibuprofen among antipyretics and analgesics.</p>	Name of symptoms	Symptoms	<u>Drug-induced hypersensitivity syndrome</u>	<u>Some symptoms, such as redness over large part of the skin, generalised exanthema, pyrexia, malaise, swollen lymph nodes (neck, armpits, groin, etc.) may occur.</u>
Name of symptoms	Symptoms				
<u>Drug-induced hypersensitivity syndrome</u>	<u>Some symptoms, such as redness over large part of the skin, generalised exanthema, pyrexia, malaise, swollen lymph nodes (neck, armpits, groin, etc.) may occur.</u>				

N/A: Not Applicable. No corresponding language is included in the current Precautions.

**Pharmaceuticals and Medical Devices Agency**

3-3-2 Kasumigaseki, Chiyoda-ku, Tokyo 100-0013 Japan  
E-mail: [safety.info@pmda.go.jp](mailto:safety.info@pmda.go.jp)