

Published by  
Ministry of Health, Labour and Welfare



Translated by  
Pharmaceuticals and Medical Devices Agency



---

*This English version is intended to be a reference material to provide convenience for users. In the event of inconsistency between the Japanese original and this English translation, the former shall prevail.*

# Revision of Precautions

## Kanamycin sulfate

## Streptomycin sulfate

March 23, 2023

### **Therapeutic category**

Antibiotic preparations acting mainly on acid-fast bacteria

### **Non-proprietary name**

Kanamycin sulfate

Streptomycin sulfate

### **Safety measure**

Precautions should be revised.

**Pharmaceuticals and Medical Devices Agency**

3-3-2 Kasumigaseki, Chiyoda-ku, Tokyo 100-0013 Japan  
E-mail: [safety.info@pmda.go.jp](mailto:safety.info@pmda.go.jp)

Revision in line with the Instructions for Package Inserts of Prescription Drugs, PAB Notification No. 606 by the Director General of Pharmaceutical Affairs Bureau, MHW, dated April 25, 1997 (Old instructions): Revised language is underlined.

Current	Revision
Important Precautions (N/A)	Important Precautions <u>A paradoxical drug reaction may be observed due to the treatment with antitubercular agents including this drug. If worsening of existing tuberculosis or new onset of tuberculosis symptoms is observed after initiating the treatment, whether to continue administration should be determined on the basis of drug susceptibility tests, etc.</u>

Note: Kanamycin sulfate is designated as a drug requiring preparation of a Drug Guide for Patients.

N/A: Not Applicable. No corresponding language is included in the current Precautions.

Revision in line with the Instructions for Electronic Package Inserts of Prescription Drugs, etc. PSEHB Notification No. 0611-1 by the Director of Pharmaceutical Safety and Environmental Health Bureau, MHLW, dated June 11, 2021 (New instructions): Revised language is underlined.

Current	Revision
8. IMPORTANT PRECAUTIONS (N/A)	8. IMPORTANT PRECAUTIONS <u>&lt;Pulmonary tuberculosis and other tuberculosis&gt;</u> <u>A paradoxical drug reaction may be observed due to the treatment with antitubercular agents including this drug. If worsening of existing tuberculosis or new onset of tuberculosis symptoms is observed after initiating the treatment, whether to continue administration should be determined on the basis of drug susceptibility tests, etc.</u>

**Pharmaceuticals and Medical Devices Agency**

3-3-2 Kasumigaseki, Chiyoda-ku, Tokyo 100-0013 Japan  
 E-mail: [safety.info@pmda.go.jp](mailto:safety.info@pmda.go.jp)

Note: Kanamycin sulfate is designated as a drug requiring preparation of a Drug Guide for Patients.

N/A: Not Applicable. No corresponding language is included in the current Precautions.

**Pharmaceuticals and Medical Devices Agency**

3-3-2 Kasumigaseki, Chiyoda-ku, Tokyo 100-0013 Japan  
E-mail: [safety.info@pmda.go.jp](mailto:safety.info@pmda.go.jp)