Published by Ministry of Health, Labour and Welfare Translated by Pharmaceuticals and Medical Devices Agency



This English version is intended to be a reference material to provide convenience for users. In the event of inconsistency between the Japanese original and this

English translation, the former shall prevail.

Revision of Precautions Rifampicin Ethambutol hydrochloride Levofloxacin hydrate (oral dosage form)

March 23, 2023

Therapeutic category Antibiotic preparations acting mainly on acid-fast bacteria Anti-tuberculous agents Synthetic antibacterials

62

Pharmaceuticals and Medical Devices Agency

3-3-2 Kasumigaseki, Chiyoda-ku, Tokyo 100-0013 Japan E-mail: <u>safety.info@pmda.go.jp</u>

Non-proprietary name

Rifampicin Ethambutol hydrochloride Levofloxacin hydrate

Safety measure

Precautions should be revised.

Pharmaceuticals and Medical Devices Agency

3-3-2 Kasumigaseki, Chiyoda-ku, Tokyo 100-0013 Japan E-mail: <u>safety.info@pmda.go.jp</u> Revision in line with the Instructions for Package Inserts of Prescription Drugs, PAB Notification No. 606 by the Director General of Pharmaceutical Affairs Bureau, MHW, dated April 25, 1997 (Old instructions): Revised language is underlined.

Current	Revision
Important Precautions	Important Precautions
(N/A)	A paradoxical drug reaction may be observed due to the treatment
	with antitubercular agents including this drug. If worsening of
	existing tuberculosis or new onset of tuberculosis symptoms is
	observed after initiating the treatment, whether to continue
	administration should be determined on the basis of drug
	susceptibility tests, etc.

N/A: Not Applicable. No corresponding language is included in the current Precautions.

Revision in line with the Instructions for Electronic Package Inserts of Prescription Drugs, etc. PSEHB Notification No. 0611-1 by the Director of Pharmaceutical Safety and Environmental Health Bureau, MHLW, dated June 11, 2021 (New instructions): Revised language is underlined.

Current	Revision
8. IMPORTANT PRECAUTIONS	8. IMPORTANT PRECAUTIONS
<pulmonary and="" other="" tuberculosis=""></pulmonary>	<pulmonary and="" other="" tuberculosis=""></pulmonary>
(N/A)	A paradoxical drug reaction may be observed due to the treatment
	with antitubercular agents including this drug. If worsening of
	existing tuberculosis or new onset of tuberculosis symptoms is
	observed after initiating the treatment, whether to continue
	administration should be determined on the basis of drug
	susceptibility tests, etc.

N/A: Not Applicable. No corresponding language is included in the current Precautions.

Pharmaceuticals and Medical Devices Agency