

1 **Blonanserin Powder**

2 ブロナンセリン散

4 Blonanserin Powder contains not less than 95.0%
5 and not more than 105.0% of the labeled amount of
6 blonanserin ($C_{23}H_{30}FN_3$; 367.50).

7 **Method of preparation** Prepare as directed under Gran-
8 ules or Powders, with Blonanserin.

9 **Identification** To an amount of powdered Blonanserin
10 Powder, equivalent to 1.4 mg of Blonanserin, add 1 mL of
11 water to moisten. Then, add 60 mL of methanol, shake for 20
12 minutes, add methanol to make 100 mL, and centrifuge. De-
13 termine the absorption spectrum of the supernatant liquid as
14 directed under Ultraviolet-visible Spectrophotometry <2.24>:
15 it exhibits maxima between 234 nm and 238 nm, between 251
16 nm and 255 nm, and between 312 nm and 316 nm.

17 **Dissolution** <6.10> When the test is performed at 50 revo-
18 lutions per minute according to the Paddle method, using 900
19 mL of a solution prepared by adding 0.05 mol/L disodium
20 hydrogen phosphate TS to 0.05 mol/L potassium dihydrogen
21 phosphate TS and adjusted to pH 6.0 as the dissolution me-
22 dium, the dissolution rate in 15 minutes of Blonanserin Pow-
23 der is not less than 75%.

24 Start the test with an accurately weighed amount of
25 Blonanserin Powder, equivalent to about 4 mg of blonanserin
26 ($C_{23}H_{30}FN_3$), withdraw not less than 30 mL of the medium at
27 the specified minute after starting the test, and filter through
28 a membrane filter with a pore size not exceeding 0.45 μ m.
29 Discard not less than 20 mL of the first filtrate, pipet 4 mL of
30 the subsequent filtrate, add exactly 1 mL of 0.1 mol/L hydro-
31 chloric acid TS, and use this solution as the sample solution.
32 Separately, weigh accurately about 20 mg of Blonanserin RS,
33 previously dried at 105°C for 2 hours, and dissolve in meth-
34 anol to make exactly 100 mL. Pipet 4 mL of this solution, add
35 a mixture of the dissolution medium and 0.1 mol/L hydro-
36 chloric acid TS (4:1) to make exactly 250 mL, and use this
37 solution as the standard solution. Perform the test with ex-
38 actly 40 μ L each of the sample solution and standard solution
39 as directed under Liquid Chromatography <2.01> according
40 to the following conditions, and determine the peak areas, A_T
41 and A_S , of blonanserin in each solution.

42 Dissolution rate (%) with respect to the labeled amount of
43 blonanserin ($C_{23}H_{30}FN_3$)

$$44 = M_S / M_T \times A_T / A_S \times 1 / C \times 9$$

45 M_S : Amount (mg) of Blonanserin RS taken

46 M_T : Amount (g) of Blonanserin Powder taken

47 C : Labeled amount (mg) of blonanserin ($C_{23}H_{30}FN_3$) in 1 g

48 **Operating conditions**—

49 Proceed as directed in the operating conditions in the As-
50 say under Blonanserin.

51 **System suitability**—

52 System performance: When the procedure is run with 40
53 μ L of the standard solution under the above operating condi-
54 tions, the number of theoretical plates and the symmetry fac-
55 tor of the peak of blonanserin are not less than 8000 and not
56 more than 2.0, respectively.

57 System repeatability: When the test is repeated 6 times
58 with 40 μ L of the standard solution under the above operating
59 conditions, the relative standard deviation of the peak area of
60 blonanserin is not more than 2.0%.

61 **Assay** Weigh accurately an amount of powdered Blonan-
62 serin Powder, equivalent to about 4 mg of blonanserin
63 ($C_{23}H_{30}FN_3$), add 4 mL of water to moisten, add 60 mL of
64 methanol, and sonicate for 10 minutes. Further, shake for 20
65 minutes, add methanol to make exactly 100 mL, and centri-
66 fuge. Pipet 8 mL of the supernatant liquid, add exactly 2 mL
67 of the internal standard solution, and use this solution as the
68 sample solution. Separately, weigh accurately about 40 mg
69 of Blonanserin RS, previously dried at 105°C for 2 hours, and
70 dissolve in methanol to make exactly 100 mL. Pipet 4 mL of
71 this solution, add exactly 10 mL of the internal standard so-
72 lution, add methanol to make 50 mL, and use this solution as
73 the standard solution. Proceed as directed in the Assay under
74 Blonanserin.

$$75 \text{ Amount (mg) of blonanserin (} C_{23}H_{30}FN_3 \text{)} \\ 76 = M_S \times Q_T / Q_S \times 1 / 10$$

77 M_S : Amount (mg) of Blonanserin RS taken

78 **Internal standard solution**—A solution of isoamyl benzoate
79 in methanol (1 in 8000).

80 **Containers and storage** Containers—Tight containers.

81 **Add the following to 9.01 Reference**
82 **Standards (1).**

83 Blonanserin RS

84