

This document is an English-translated version of an attachment of a notification for Revision of PRECAUTIONS issued by the Ministry of Health, Labour and Welfare.

This English version is intended to be a reference material to provide convenience for users.

In the event of inconsistency between the Japanese original and this English translation, the former shall prevail.

Revision of PRECAUTIONS

Triamcinolone acetonide (ophthalmic injection)

November 13, 2024

Therapeutic category

Agents for ophthalmic use

Non-proprietary name

Triamcinolone acetonide (ophthalmic injection)

Safety measure

PRECAUTIONS should be revised.

Revised language is underlined.

Current	Revision
<p>8. IMPORTANT PRECAUTIONS</p> <p><Subtenon injection></p> <p>(N/A)</p> <p>11. ADVERSE REACTIONS</p> <p>11.1 Clinically Significant Adverse Reactions</p> <p>(N/A)</p> <p><Subtenon injection></p> <p>Eye disorders</p> <p>Cataract, increased intraocular pressure, <u>or</u> glaucoma may occur.</p> <p>Surgical intervention may be needed.</p>	<p>8. IMPORTANT PRECAUTIONS</p> <p><Subtenon injection></p> <p><u>Infective scleritis may occur. Patients should be closely monitored after administration of this drug. The patients must be instructed to immediately consult physicians if any abnormalities are observed.</u></p> <p>11. ADVERSE REACTIONS</p> <p>11.1 Clinically Significant Adverse Reactions</p> <p><u><Intravitreal injection: Visualization of the vitreous body during vitreous surgery></u></p> <p><u>Eye disorders</u></p> <p><u>Endophthalmitis may occur. Surgical intervention may be needed.</u></p> <p><Subtenon injection></p> <p>Eye disorders</p> <p>Cataract, increased intraocular pressure, glaucoma, <u>or infective scleritis</u> may occur. Surgical intervention may be needed.</p>

N/A: Not Applicable. No corresponding language is included in the current PRECAUTIONS.