This document is an English-translated version of an attachment of a notification for Revision of PRECAUTIONS issued by the Ministry of Health, Labour and Welfare.

This English version is intended to be a reference material to provide convenience for users.

In the event of inconsistency between the Japanese original and this English translation, the former shall prevail.

Revision of PRECAUTIONS

Ensitrelvir fumaric acid

December 17, 2024

Therapeutic category

Anti-virus agents

Non-proprietary name

Ensitrelyir fumaric acid

Safety measure

PRECAUTIONS should be revised.

Revised language is underlined.

| Current | Revision |
|--------------------------|---|
| 8. IMPORTANT PRECAUTIONS | 8. IMPORTANT PRECAUTIONS |
| (N/A) | The necessity of administering this drug to women of childbearing |
| | potential should be carefully considered. If administration is deemed |
| | necessary, attention should be paid to the following points. |
| | Prior to administering this drug, the absence of pregnancy and |
| | the absence of the possibility of pregnancy should be confirmed |
| | through sufficient patient interviews. |
| | The following should be explained to patients before starting |
| | administration of this drug: |
| | •This drug can cause foetal harm when administered to a pregnant |
| | woman. |
| | •If pregnancy is detected or suspected during administration of this |
| | drug, this drug should be discontinued immediately. |
| | •If pregnancy is detected or suspected during administration of this |
| | drug or within 2 weeks after the last administration of this drug, a |
| | physician, pharmacist, etc. should be consulted promptly. |

N/A: Not Applicable. No corresponding language is included in the current PRECAUTIONS.