This document is an English-translated version of an attachment of a notification for Revision of PRECAUTIONS issued by the Ministry of Health, Labour and Welfare.

This English version is intended to be a reference material to provide convenience for users.

In the event of inconsistency between the Japanese original and this English translation, the former shall prevail.

## **Revision of PRECAUTIONS**

## **Eplerenone**

May 20, 2025

Therapeutic category

Antihypertensives

Non-proprietary name

Eplerenone

Safety measure

PRECAUTIONS should be revised.

Nevisea language le unacimilea.						
Current				Revision		
2. CONTRAINDICATIONS (This drug is contraindicated to the				2. CONTRAINDICATIONS (This drug is contraindicated to the		
following patients.)				following patients.)		
<common all="" indications="" to=""></common>				<common all="" indications="" to=""></common>		
Patients receiving the following drugs: Itraconazole, preparations				Patients receiving the following drugs: Itraconazole, voriconazole,		
containing ritonavir, and ensitrelvir fumaric acid				posaconazole, preparations containing ritonavir, and ensitrelvir fumaric acid		
10. INTERACTIONS 10.1 Contraindications for Co-administration (Do not co-administer with the following.)				10. INTERACTIONS 10.1 Contraindications for Co-administration (Do not co-administer with the following.)		
<common all="" indications="" to=""></common>				<common all="" indications="" to=""></common>		
Drugs	Signs, symptoms, and treatment	Mechanism/risk factors		Drugs	Signs, symptoms, and treatment	Mechanism/risk factors
Itraconazole Preparations containing ritonavir Ensitrelvir fumaric acid	The plasma concentration of eplerenone may increase, and an increase in serum potassium levels may be induced.	The metabolism of eplerenone is suppressed by a strong CYP3A4-inhibitor.		Itraconazole Voriconazole Posaconazole Preparations containing ritonavir Ensitrelvir fumaric acid	The plasma concentration of eplerenone may increase, and an increase in serum potassium levels may be induced.	The metabolism of eplerenone is suppressed by a strong CYP3A4-inhibitor.