

*This document is an English-translated version of an attachment of a notification for Revision of PRECAUTIONS issued by the Ministry of Health, Labour and Welfare.*

*This English version is intended to be a reference material to provide convenience for users.*

*In the event of inconsistency between the Japanese original and this English translation, the former shall prevail.*

# Revision of PRECAUTIONS

**Olmesartan medoxomil**

**Olmesartan medoxomil/azelnidipine**

**Valsartan**

**Valsartan/amlodipine besilate**

**Valsartan/cilnidipine**

**Valsartan/hydrochlorothiazide**

September 9, 2025

## **Therapeutic category**

Antihypertensives

## **Non-proprietary name**

Olmesartan medoxomil

Olmesartan medoxomil/azelnidipine

Valsartan

Valsartan/amlodipine besilate

Valsartan/cilnidipine

Valsartan/hydrochlorothiazide

*This document is an English-translated version of an attachment of a notification for Revision of PRECAUTIONS issued by the Ministry of Health, Labour and Welfare.*

*This English version is intended to be a reference material to provide convenience for users.*

*In the event of inconsistency between the Japanese original and this English translation, the former shall prevail.*

## **Safety measure**

PRECAUTIONS should be revised.

*This document is an English-translated version of an attachment of a notification for Revision of PRECAUTIONS issued by the Ministry of Health, Labour and Welfare.*

*This English version is intended to be a reference material to provide convenience for users.*

*In the event of inconsistency between the Japanese original and this English translation, the former shall prevail.*

Revised language is underlined.

Current	Revision
11. ADVERSE REACTIONS 11.1 Clinically Significant Adverse Reactions Angioedema Symptoms such as swelling of the face, lip, pharynx, and tongue may occur.	11. ADVERSE REACTIONS 11.1 Clinically Significant Adverse Reactions Angioedema Symptoms such as swelling of the face, lip, pharynx, and tongue may occur. <u>Of note, intestinal angioedema accompanied by abdominal pain, queasy, vomiting, diarrhoea, etc. may occur.</u>