

事務連絡
令和8年3月4日

各都道府県衛生主管部（局）薬務主管課 御中

厚生労働省医薬局医薬品審査管理課

第十八改正日本薬局方（英文版）正誤表の送付について（その6）

第十八改正日本薬局方（令和3年厚生労働省告示第220号）の英文版につきまして、一部に誤植等がありましたので別紙のとおり正誤表を送付いたします。

JP18 table of errata

March 4, 2026

Official Monographs

Dextran 40 デキストラン 40

Page	Line	Correction	Error
838	left ↑ 24	After cooling, add 1 mL of a solution of potassium <u>iodide</u> (1 in 40) and 1.5 mL of dilute sulfuric acid, and titrate <2.50> with 0.005 mol/L sodium thiosulfate VS (indicator: 2 mL of starch TS).	After cooling, add 1 mL of a solution of potassium <u>iodine</u> (1 in 40) and 1.5 mL of dilute sulfuric acid, and titrate <2.50> with 0.005 mol/L sodium thiosulfate VS (indicator: 2 mL of starch TS).
838	right ↓ 16	Determine the optical rotation α_D with the sample solution as directed under Optical Rotation Determination <2.49> in a <u>100-mm</u> cell at $20 \pm 1^\circ\text{C}$.	Determine the optical rotation α_D with the sample solution as directed under Optical Rotation Determination <2.49> in a <u>100-mL</u> cell at $20 \pm 1^\circ\text{C}$.

Stearic Acid ステアリン酸

Page	Line	Correction	Error
1743	right ↑ 23	Add 10 mL of a solution of potassium <u>iodide</u> (1 in 10) and 100 mL of water.	Add 10 mL of a solution of potassium <u>iodine</u> (1 in 10) and 100 mL of water.