

【Therapeutic category】 Gastrointestinal medicine

【Non-proprietary name】 Preparations containing rabeprazole sodium (guidance-mandatory drugs)

【Safety measure】 PRECAUTIONS should be revised.

A revised description is underlined.

Current		Revision	
CONSULTATION If the following symptoms are observed after taking this drug, they may be adverse reactions, so immediately discontinue the use of this drug, and show this document to your physician or pharmacist for a consultation.		CONSULTATION If the following symptoms are observed after taking this drug, they may be adverse reactions, so immediately discontinue the use of this drug, and show this document to your physician or pharmacist for a consultation.	
Relevant region	Symptoms	Relevant region	Symptoms
Neuropsychiatric system	Headache, dizziness, light-headedness, sleepiness, weakness of limbs, hypoesthesia (reduced sensation), grip strength decreased, slurred speech, disorientation (being confused about the place, time, name, etc.), delirium (mild clouding of consciousness with illusion or hallucinations), coma (complete loss of consciousness), <u>temporary mental dullness,</u> <u>consciousness decreased, syncope</u>	Neuropsychiatric system	Headache, dizziness, light-headedness, sleepiness, weakness of limbs, hypoesthesia (reduced sensation), grip strength decreased, slurred speech, disorientation (being confused about the place, time, name, etc.), delirium (mild clouding of consciousness with illusion or hallucinations), coma (complete loss of consciousness)
Others	Anemia, blurred/flickering vision, edema, malaise, fever, alopecia, numbness, <u>twitching muscles of the</u>	Others	Anemia, blurred/flickering vision, edema, malaise, fever, alopecia, numbness, arthralgia, myalgia, gynecomastia

	<u>face or limbs, muscle stiffness with coarse jerky tremor of limbs</u> arthralgia, myalgia, gynecomastia		
The following serious symptoms may occur in rare cases. In such cases, immediately seek medical aid: (N/A)		The following serious symptoms may occur in rare cases. In such cases, immediately seek medical aid:	
		Name of symptoms	Symptoms
		<u>Hypomagnesaemia</u>	<u>Symptoms such as nausea, vomiting, Loss of appetite, feeling of weariness, muscle twitching in the face and limbs, temporary drowsiness, decreased consciousness, stiffening of the limbs and trembling, numbness, dizziness, palpitations, loss of consciousness may occur.</u>

N/A: Not Applicable. No corresponding description is included in the current PRECAUTIONS.

This English version is intended to be a reference material to provide convenience for users. In the event of inconsistency between the Japanese original and this English translation, the former shall prevail.