

【Therapeutic category】 Other agents for epidermis

【Non-proprietary name】 Tacrolimus hydrate (pediatric ointment)

【Safety measure】 PRECAUTIONS should be revised.

A revised description is underlined.

Current	Revision																		
2. CONTRAINDICATIONS (This drug is contraindicated to the following patients.) Low birth weight infants, neonates, <u>infants, or children under 2 years of age</u>	2. CONTRAINDICATIONS (This drug is contraindicated to the following patients.) Low birth weight infants, neonates, <u>or infants under 6 months of age</u>																		
7. PRECAUTIONS CONCERNING DOSAGE AND ADMINISTRATION The following table should be used as a guide for the maximum amount per application.	7. PRECAUTIONS CONCERNING DOSAGE AND ADMINISTRATION The following table should be used as a guide for the maximum amount per application.																		
<table> <tr> <th>Age (body weight)</th><th>Maximum amount per application</th></tr> <tr> <td>2 to 5 years (less than 20 kg)</td><td>1 g</td></tr> <tr> <td>6 to 12 years (20 kg or more and less than 50 kg)</td><td>2 to 4 g</td></tr> <tr> <td>13 years or older (50 kg or more)</td><td>5 g</td></tr> </table>	Age (body weight)	Maximum amount per application	2 to 5 years (less than 20 kg)	1 g	6 to 12 years (20 kg or more and less than 50 kg)	2 to 4 g	13 years or older (50 kg or more)	5 g	<table> <tr> <th>Age (body weight)</th><th>Maximum amount per application</th></tr> <tr> <td><u>6 months to under 2 years (less than 10 kg)</u></td><td><u>0.5 g</u></td></tr> <tr> <td>2 to 5 years (<u>10 kg or more and less than 20 kg</u>)</td><td>1 g</td></tr> <tr> <td>6 to 12 years (20 kg or more and less than 50 kg)</td><td>2 to 4 g</td></tr> <tr> <td>13 years or older (50 kg or more)</td><td>5 g</td></tr> </table>	Age (body weight)	Maximum amount per application	<u>6 months to under 2 years (less than 10 kg)</u>	<u>0.5 g</u>	2 to 5 years (<u>10 kg or more and less than 20 kg</u>)	1 g	6 to 12 years (20 kg or more and less than 50 kg)	2 to 4 g	13 years or older (50 kg or more)	5 g
Age (body weight)	Maximum amount per application																		
2 to 5 years (less than 20 kg)	1 g																		
6 to 12 years (20 kg or more and less than 50 kg)	2 to 4 g																		
13 years or older (50 kg or more)	5 g																		
Age (body weight)	Maximum amount per application																		
<u>6 months to under 2 years (less than 10 kg)</u>	<u>0.5 g</u>																		
2 to 5 years (<u>10 kg or more and less than 20 kg</u>)	1 g																		
6 to 12 years (20 kg or more and less than 50 kg)	2 to 4 g																		
13 years or older (50 kg or more)	5 g																		
Reference: Doses used in clinical studies																			

9. PRECAUTIONS CONCERNING PATIENTS WITH SPECIFIC BACKGROUNDS 9.7 Pediatric Patients This drug should not be used in low birth weight infants, neonates, <u>infants, or children under 2 years of age</u>. No clinical studies have been conducted in <u>children under 2 years of age</u>.	Reference: Doses used in clinical studies 9. PRECAUTIONS CONCERNING PATIENTS WITH SPECIFIC BACKGROUNDS 9.7 Pediatric Patients This drug should not be used in low birth weight infants, neonates, <u>or infants under 6 months of age</u>. No clinical studies have been conducted in <u>infants under 6 months of age</u>.
---	---

This English version is intended to be a reference material to provide convenience for users. In the event of inconsistency between the Japanese original and this English translation, the former shall prevail.